



Hyperviscosity, Galerkin Truncation, and Bottlenecks in Turbulence

Uriel Frisch, Susan Kurien, Rahul Pandit, Walter Pauls, Samriddhi Sankar Ray, Achim Wirth, Jian-Zhou Zhu

► To cite this version:

Uriel Frisch, Susan Kurien, Rahul Pandit, Walter Pauls, Samriddhi Sankar Ray, et al.. Hyperviscosity, Galerkin Truncation, and Bottlenecks in Turbulence. Physical Review Letters, 2008, 101, pp.144501. 10.1103/PhysRevLett.101.144501 . hal-00390765

HAL Id: hal-00390765

<https://hal.science/hal-00390765>

Submitted on 6 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hyperviscosity, Galerkin Truncation, and Bottlenecks in Turbulence

Uriel Frisch,¹ Susan Kurien,² Rahul Pandit,³ Walter Pauls,⁴ Samriddhi Sankar Ray,³ Achim Wirth,⁵ and Jian-Zhou Zhu²

¹*Laboratoire Cassiopée, OCA, UNS, CNRS, BP 4229, 06304 Nice cedex 4, France*

²*CNLS & Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

³*Centre for Condensed Matter Theory, Department of Physics, Indian Institute of Science, Bangalore, India*

⁴*Max Planck Institute for Dynamics and Self-Organization, Goettingen, Germany*

⁵*LEGI, CNRS, Grenoble, France*

(Received 29 March 2008; published 29 September 2008)

It is shown that the use of a high power α of the Laplacian in the dissipative term of hydrodynamical equations leads asymptotically to truncated inviscid conservative dynamics with a finite range of spatial Fourier modes. Those at large wave numbers thermalize, whereas modes at small wave numbers obey ordinary viscous dynamics [C. Cichowlas *et al.*, Phys. Rev. Lett. **95**, 264502 (2005)]. The energy bottleneck observed for finite α may be interpreted as incomplete thermalization. Artifacts arising from models with $\alpha > 1$ are discussed.

DOI: 10.1103/PhysRevLett.101.144501

PACS numbers: 47.27.Gs, 05.20.Jj

A single Maxwell daemon embedded in a turbulent flow would hardly notice that the fluid is not exactly in thermal equilibrium because incompressible turbulence, even at very high Reynolds numbers, constitutes a tiny perturbation on thermal molecular motion. Dissipation in real fluids is just the transfer of macroscopically organized (hydrodynamic) energy to molecular thermal energy. Artificial microscopic systems can act just like the real ones as far as the emergence of hydrodynamics is concerned; for instance, in lattice gases the “molecules” are discrete Boolean entities [1] and thermalization is easily observed at high wave numbers [2]. Another example has been found recently by Cichowlas *et al.* [3] wherein the Euler equations of ideal nondissipative flow are (Galerkin) truncated by keeping only a finite—but large—number of spatial Fourier harmonics. The modes with the highest wave numbers k then rapidly thermalize through a mechanism discovered by Lee [4] and studied further by Kraichnan [5], leading in three dimensions (3D) to an equipartition energy spectrum $\propto k^2$. The thermalized modes act as a fictitious microworld on modes with smaller wave numbers in such a way that the usual dissipative Navier–Stokes dynamics is recovered at large scales [6].

All the known systems presenting thermalization are conservative. As we shall show thermalization may be present in dissipative hydrodynamic systems when the dissipation rate increases so fast with the wave number that it mimics ideal hydrodynamics with a Galerkin truncation. This is best understood by considering hydrodynamics with hyperviscosity: the usual momentum diffusion operator (a Laplacian) is replaced by the α th power of the Laplacian, where $\alpha > 1$ is the dissipativity. Hyperviscosity is frequently used in turbulence modeling to avoid wasting numerical resolution by reducing the range of scales over which dissipation is effective [7].

The unforced hyperviscous 1D Burgers and multidimensional incompressible Navier–Stokes (NS) equations are

$$\partial_t \mathbf{v} + \mathbf{v} \partial_x \mathbf{v} = -\mu k_G^{-2\alpha} (-\partial_x^2)^\alpha \mathbf{v}; \quad (1)$$

$$\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} = -\nabla p - \mu k_G^{-2\alpha} (-\nabla^2)^\alpha \mathbf{v}; \quad \nabla \cdot \mathbf{v} = 0. \quad (2)$$

The equations must be supplemented with suitable initial and boundary conditions. We employ 2π -periodic boundary conditions in space, so that we can use Fourier decompositions such as $\mathbf{v}(\mathbf{x}) = \sum_k \hat{\mathbf{v}}_k e^{ik \cdot \mathbf{x}}$. Note that minus the Laplacian is a positive operator, with Fourier transform k^2 , which can be raised to an arbitrary power α . The coefficient μ is taken positive to make the hyperviscous operator dissipative. The Galerkin wave number $k_G > 0$ is chosen off-lattice so that no wave number is exactly equal to k_G . In Fourier space the hyperdissipation rate is $\mu(k/k_G)^{2\alpha}$, where $k \equiv |\mathbf{k}|$.

If we now hold μ and k_G fixed and let $\alpha \rightarrow \infty$ we see that the hyperdissipation rate tends to zero, for $k < k_G$, and to infinity, for $k > k_G$. This implies that in the limit of infinite dissipativity, the solution of a hyperviscous hydrodynamical equation converges to that of the corresponding inviscid equations Galerkin-truncated at wave number k_G .

To define inviscid Galerkin truncation precisely, we rewrite Eqs. (1) and (2) in the abstract form $\partial_t \mathbf{v} = B(\mathbf{v}, \mathbf{v}) + L_\alpha \mathbf{v}$, where B is a quadratic form representing the nonlinear term (including the pressure p in the NS case). The truncation projector P_{k_G} is the linear, low-pass filtering operator that, when applied to $\mathbf{v}$, sets all Fourier harmonics with $k > k_G$ to zero. The inviscid, Galerkin-truncated equation, with initial condition $\mathbf{v}_0$, is

$$\partial_t \mathbf{u} = P_{k_G} B(\mathbf{u}, \mathbf{u}), \quad \mathbf{u}_0 = P_{k_G} \mathbf{v}_0. \quad (3)$$

Since $\mathbf{u}$ can be written in terms of a finite number of modes with $k < k_G$, Eq. (3) is a dynamical system of finite dimension. In addition to momentum, it conserves the energy and other quadratic invariants for the inviscid equations [5]. There is good numerical evidence—but no rigorous

proof—that the solutions of the Galerkin-truncated inviscid Burgers and 3D Euler equations tend, at large times, to statistical equilibria defined by their respective invariants.

A rigorous proof of the convergence, as $\alpha \rightarrow \infty$, of solutions of the hyperviscous Burgers equation (1) and of the hyperviscous NS equation (2) in any dimension to those of the associated Galerkin-truncated, inviscid equation will be given elsewhere. It uses standard tools of functional analysis; note that the formidable mathematical difficulties that beset the ordinary ($\alpha = 1$) 3D NS equation disappear for $\alpha \geq 5/4$ [8].

From a physicist's point of view the convergence result looks rather obvious, though it has hardly been noted before (see, however, Refs. [9–11]): as $\alpha \rightarrow \infty$ all the modes with $k > k_G$ are immediately suppressed by an infinite dissipation, whereas those with $k < k_G$ obey inviscid truncated dynamics. Not surprisingly, the fate of couplings between triads of modes whose wave numbers straddle k_G is a delicate point. In a Galerkin truncation any such triad should be left out. It may be shown that for $\alpha \rightarrow \infty$ such straddling couplings are suppressed, not only for the Burgers and NS equation but also for the hyperviscous magnetohydrodynamical equations and for some turbulence closures, specifically, the direct interaction approximation (DIA) [9] and the eddy-damped-quasi-normal-markovian (EDQNM) approximation [12]. Hence the convergence to the corresponding Galerkin-truncated equations holds for all the aforementioned equations in any dimension of space.

There are, however, interesting exceptions among hydrodynamical equations for which the result does not hold. They include the kinetic theory of resonant wave interactions [13] and the Markovian random coupling model [14].

Indeed, the resonant wave interaction theory arises in the limit when the period of the waves goes to zero and this limit does not commute with the limit of a vanishing damping time for modes having $k > k_G$; a similar remark can be made about the MRCM equation.

Let us stress that systems with a finite dissipativity—however large—are quite different from Galerkin-truncated systems. For example, consider the 3D NS equation with a random force, delta-correlated in time, for which we know the mean energy input ε per unit volume. It is still true that, for $\alpha \rightarrow \infty$, the solution of this equation converges to that of the Galerkin-truncated equation, but this time with a random force. If E_0 is the initial energy, this solution has a mean energy $E(t) = E_0 + \varepsilon t$, which grows indefinitely in time. But, as soon as α is given a finite value, however large, a statistical steady state, in which energy input and hyperviscous energy dissipation balance, is achieved at large times. Such a steady state presents an interesting interplay of thermalization and dissipation, when α is large, as we show below.

The direct numerical simulation (DNS) of the Galerkin-truncated 3D Euler equations in Ref. [3] used 1600^3 Fourier modes. Large- α simulations of Eq. (2) would require significantly higher resolution to identify the various spectral ranges that we can expect, namely, inertial, thermalized, and far-dissipation ranges and transition regimes between these. Fortunately, Bos and Bertoglio [15] have shown that key features of the Galerkin-truncated Euler equations, such as the presence of inertial and thermalized ranges, can be reproduced by the two-point EDQNM closure [12] for the energy spectrum. For Eq. (2), with stochastic, white-in-time, homogeneous, and isotropic forcing with spectrum $F(k)$, the hyperviscous EDQNM equations are

$$\left[\partial_t + 2\mu \left(\frac{k}{k_G} \right)^{2\alpha} \right] E(k, t) = \iint_{\Delta_k} dp dq \theta_{kpq} b(k, p, q) \frac{k}{pq} E(q, t) [k^2 E(p, t) - p^2 E(k, t)] + F(k), \quad (4)$$

$$\theta_{kpq} = \frac{1}{\mu_k + \mu_p + \mu_q}, \quad b(k, p, q) = \frac{p}{k} (xy + z^3), \quad \mu_k = \mu \left(\frac{k}{k_G} \right)^{2\alpha} + \lambda \left[\int_0^k p^2 E(p, t) dp \right]^{1/2}.$$

Here $E(k, t)$ is the energy spectrum, Δ_k defines the set of $p \geq 0$ and $q \geq 0$ such that k, p, q can form a triangle, x, y, z are the cosines of its angles and the eddy-damping parameter λ is expressed in terms of the Kolmogorov constant. The EDQNM equations have been studied numerically for more than three decades [16], but their hyperviscous versions Eq. (4) present new difficulties that we overcome as follows. Since we are interested in the steady state we use an iterative method: the emission term, $E(p, t)E(q, t)$ in Eq. (4), is considered as a renormalization of the force $F(k)$; the absorption term $E(q, t)E(k, t)$ is treated as a renormalization of the hyperviscous damping [17]. We then construct a sequence of energy spectra that, at stage $n + 1$, is just the renormalized force divided by the renormalized damping, both based on stage n . This gives rapid convergence to the steady state at low wave numbers,

but, beyond a certain (α -dependent) wave number, convergence slows down dramatically and it is better to use time marching to obtain the steady state. At large values of k and α the problem becomes very stiff, so we use a slaved fourth-order Runge–Kutta scheme [18]. We discretize k logarithmically, with N_c collocation points per octave. Triad interactions involving wave number ratios significantly larger than N_c are poorly represented [19]; so, since wave number ratios of up to 50 play an important role for large α , we have used $N_c \leq 90$; this is computationally demanding because the complexity of the code is $O(N_c^3)$. We force at the lowest wave number ($k = 1$) in our numerical study of Eq. (4) with $k_G = 10^5$, $\lambda = 0.36$, and $1 \leq \alpha \leq 729$. The resulting compensated, steady-state energy spectra $k^{+5/3} E(k)$ are shown in Fig. 1; flat regions, extending over two to five decades of k (depending on α), are

close to the Kolmogorov inertial range; for large α there is a distinct thermalized range with $E(k) \sim k^2$ (also found in the transition between classical and quantum superfluid turbulence [20]), as we expect from our discussion of the Galerkin-truncated Euler equations. In the far-dissipation range $k > k_G$ the spectra fall off very rapidly. For all values of α the far-dissipation range is preceded by a bump or bottleneck. It is also observed, in some experiments [21] and DNS of Navier-Stokes, with a shape that is quite independent of the Reynolds number [22]. The bottleneck for $\alpha = 1$ has previously been explained as the inhibition of the energy cascade from low to high wave numbers because of viscous suppression of the cascade in the dissipation range [23]. Our work provides an alternative explanation: the usual bottleneck may be viewed as incomplete thermalization.

At large values of α the thermalized range gives rise to an eddy viscosity ν_{eddy} . This acts on modes with wave numbers lower than those in the thermalized range; the corresponding damping rate is $\nu_{\text{eddy}} k^2$. The eddy viscosity can be expressed as an integral over the thermalized range [15,19]. As α grows, so does ν_{eddy} and, eventually, the renormalized viscous damping overwhelms the hyperviscous damping for modes at low wave numbers (below those in the thermalized range). The dynamics of these modes is then governed by the usual $\alpha = 1$ equation. Not surprisingly, then, we find a pseudodissipation range around $k \approx 10^4$ that is shown in an expanded scale in the inset of Fig. 1; a similar range for the Galerkin-truncated case is discussed in Ref. [15] and is already visible in the DNS of Ref. [3]. For large α the inset of Fig. 1 also shows a secondary bottleneck range for $10^3 < k < 10^4$; this may be viewed as the usual ($\alpha = 1$) EDQNM bottleneck stemming from ν_{eddy} .

Our results apply to compressible flows also. We have studied the simplest instance, that is the unforced hyperviscous 1D Burgers equation (1). Its solution converges to the entropy solution, i.e., the standard solution with shocks, obtained when $k_G \rightarrow \infty$ for any $\alpha \geq 1$ [24]. Here we are interested in the large- α behavior at fixed k_G . We do not




FIG. 1 (color online). Log-log plots of the compensated spectrum $k^{5/3}E(k)$ versus k from a numerical integration of the hyperviscous EDQNM Eq. (4) for different values of α ; inset: enlarged spectra showing a secondary bottleneck (see text).

have to resort to closure now since we can solve the primitive equation (1) directly by a pseudospectral method. If we choose a single initial condition the resulting spectrum is noisy because, unlike the ordinary Burgers equation, its Galerkin-truncated version and thus also the high- α versions are believed to be chaotic dynamical systems [25]. So we solve (1) with the two-mode random initial condition $v_0(x) = \sin x + \sin(2x + \phi)$, where ϕ is distributed uniformly in the interval $[-\pi, \pi]$. We use 2^{14} collocation points and set $\mu = 1$, $k_G = 342.1$, and $\alpha = 1000$. In Fig. 2 we show the Burgers energy spectrum $E(k) = |\tilde{v}(k)|^2$, averaged over 20 realizations of the phase ϕ at various times. At the latest output times the spectrum is almost completely flat, i.e., thermalized, with equipartition of the energy between all the Fourier modes. At earlier times $E(k)$ behaves approximately as k^{-2} in an inertial range that corresponds to shocks in physical space; there is a thermalized range at higher wave numbers up to k_G ; for $k > k_G$ the spectrum falls very rapidly. No pseudodissipation range is observed here between the inertial and thermalized ranges as seen in the 3D NS case (Fig. 1). Perhaps the data are too noisy, but a careful examination of $v(x)$ in physical space indicates that this phenomenon might arise from the compressible nature of the Burgers dynamics: thermalization begins over the whole physical range (as high-frequency noise with wave number $\approx k_G$); noise generated close to shocks is absorbed by them and not enough is left to produce any appreciable eddy viscosity that could broaden the shocks.

We now summarize our main findings from the study of hyperviscous hydrodynamical equations with powers α of the Laplacian ranging from unity to very large values.

The simplest results are obtained for very large α . The solutions of the 1D Burgers equation or the Navier-Stokes equations in any space dimension d are then very close to the solutions of the corresponding Galerkin-truncated equations, displaying thermalization at wave numbers below k_G . The detailed scenario will of course be affected by the dimension of space. In 3D, with enough resolution, we may be able to observe up to five ranges: an inertial range,




FIG. 2 (color online). Log-log plots of averaged energy spectra (see text) versus wave number k for the hyperviscous Burgers equation (1) at various times.

a secondary bottleneck, a pseudodissipation range, a thermalized range, and a far-dissipation range. Because of enstrophy conservation and of the predominance of Fourier-space nonlocal interactions, the 2D case is rather special and deserves a separate study.

The most relevant case is of course that of ordinary dissipation ($\alpha = 1$). The energy-spectrum bottleneck generally observed at high Reynolds numbers in 3D incompressible turbulence may be viewed as an incomplete thermalization: as we increase α larger and larger bottlenecks are present, eventually displaying thermalization on their rising side.

We finally deal with the case of moderately large α of the sort used in many simulations [7]. How safe is this procedure and what kind of artifacts can we expect?

Using large values of α in simulations to “avoid wasting resolution” is hardly advocated by anybody, but we now understand what goes wrong: a huge thermalized bottleneck will develop at high wave numbers, whose action on smaller wave numbers is an ordinary $\alpha = 1$ dissipation with an eddy viscosity much larger than what would be permissible in a normal $\alpha = 1$ simulation.

When α is chosen just a bit larger than unity (e.g., $\alpha = 2$ which is standard in oceanography [7]) the advantage of widening the inertial range may be offset by artifacts at bottleneck scales; indeed, even an incomplete thermalization will bring the statistical properties of such scales closer to Gaussian, thereby reducing the rather strong intermittency which would otherwise be expected [26]. For similar reasons spurious isotropization can be expected for problems with an anisotropic constraint, such as rapidly rotating or stratified flow or MHD with a strong uniform magnetic field.

We thank J. Bec, R. Benzi, M. E. Brachet, W. Bos, J. R. Herring, D. Mitra, K. Khanin, A. Majda, S. Nazarenko, D. Schertzer, E. S. Titi, and V. Yakhot for useful discussions. J. Z. Z. acknowledges many interactions at CNLS. U. F. and W. P. were partially supported by ANR “OTARIE” BLAN07-2_183172. R. P. and S. S. R. thank DST, JNCASR, and UGC (India) for support and SERC (IISc) for computational resources. U. F., R. P., and W. P. are part of the International Collaboration for Turbulence Research (ICTR). S. K. and J. Z. Z. were funded by the Center for Nonlinear Studies LDRD program and the DOE Office of Science Advanced Scientific Computing Research (ASCR) Program in Applied Mathematics Research.

-
- [1] U. Frisch, B. Hasslacher, and Y. Pomeau, Phys. Rev. Lett. **56**, 1905 (1986).
 - [2] S. Succi, P. Santangelo, and R. Benzi, Phys. Rev. Lett. **60**, 2738 (1988).
 - [3] C. Cichowlas, P. Bonaiti, F. Debbasch, and M. Brachet, Phys. Rev. Lett. **95**, 264502 (2005).
 - [4] T. D. Lee, Q. Appl. Math. **10**, 69 (1952).
 - [5] R. H. Kraichnan, Phys. Fluids **10**, 2080 (1967).

- [6] A similar mechanism, involving sound waves, explains why superfluid and ordinary turbulence can be similar; see C. Nore, M. Abid, and M. E. Brachet, Phys. Rev. Lett. **78**, 3896 (1997).
- [7] G. Holloway, J. Phys. Oceanogr. **22**, 1033 (1992); V. Borue and S. A. Orszag, Europhys. Lett. **29**, 687 (1995); N. E. L. Haugen and A. Brandenburg, Phys. Rev. E **70**, 026405 (2004).
- [8] J. L. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites non Linéaires* (Gauthier-Villars, Paris, 1969).
- [9] R. H. Kraichnan, Phys. Rev. **109**, 1407 (1958). While introducing the DIA here, Kraichnan wrote the following about Galerkin truncation: “We may regard this as the introduction of infinite damping (infinite resistance) for the degrees of freedom removed.”
- [10] J. Jiménez, J. Fluid Mech. **279**, 169 (1994), notes that “An interesting property of the hyperviscous solutions is the sharpness of their spectral cutoff, which goes from smooth Gaussian . . . , to a step function in the ultraviscosity limit.”
- [11] J. P. Boyd, J. Sci. Comput. **9**, 81 (1994), obtained a truncation result for the linear hyperdiffusion equation.
- [12] S. A. Orszag, in *Fluid Dynamics*, Proceedings of the Les Houches Summer School, Session XXIV, edited by R. Balian and J. L. Peube (Gordon and Breach, New York, 1977), p. 237.
- [13] V. E. Zakharov, V. S. L’vov, and G. Falkovich, *Kolmogorov Spectra of Turbulence I* (Springer, New York, 1992).
- [14] U. Frisch, M. Lesieur, and A. Brissaud, J. Fluid Mech. **65**, 145 (1974).
- [15] W. J. T. Bos and J.-P. Bertoglio, Phys. Fluids **18**, 071701 (2006).
- [16] J. R. Herring and R. H. Kraichnan, J. Fluid Mech. **91**, 581 (1979).
- [17] R. H. Kraichnan, Phys. Fluids **7**, 1163 (1964).
- [18] Scheme ETD4RK of C. M. Cox and P. C. Matthews, J. Comput. Phys. **176**, 430 (2002).
- [19] M. Lesieur and D. Schertzer, J. Méc. (Paris) **17**, 609 (1978).
- [20] V. L’vov, S. Nazarenko, and O. Rudenko, Phys. Rev. B **76**, 024520 (2007).
- [21] S. G. Saddoughi and S. V. Veeravalli, J. Fluid Mech. **268**, 333 (1994). As shown by W. Dobler, N. E. L. Haugen, T. A. Yousef, and A. Brandenburg, Phys. Rev. E **68**, 026304 (2003), the 1D spectrum, generally measured in experiments, may not display a bottleneck even if the 3D spectrum does.
- [22] Z. S. She, G. Doolen, R. H. Kraichnan, and S. A. Orszag, Phys. Rev. Lett. **70**, 3251 (1993); Y. Kaneda, T. Ishihara, M. Yokokawa, K. Itakura, and A. Uno, Phys. Fluids **15**, L21 (2003); S. Kurien, M. A. Taylor, and T. Matsumoto, Phys. Rev. E **69**, 066313 (2004); J. Schumacher, Europhys. Lett. **80**, 54001 (2007); P. D. Mininni, A. Alexakis, and A. Pouquet, Phys. Rev. E **77**, 036306 (2008).
- [23] G. Falkovich, Phys. Fluids **6**, 1411 (1994); D. Lohse and A. Müller-Groeling, Phys. Rev. Lett. **74**, 1747 (1995).
- [24] E. Tadmor, Commun. Math. Sci. **2**, 317 (2004).
- [25] R. V. Abramov, G. Kovacic, and A. J. Majda, Commun. Pure Appl. Math. **56**, 1 (2003).
- [26] For $\alpha = 1$ a lull in the growth of intermittency at bottleneck scales may already be observed; cf. Fig. 2, panel R4 of J.-Z. Zhu, Chin. Phys. Lett. **23**, 2139 (2006).