



## Draft Genome Sequence of *Methylobacterium buryatense* Strain 5G, a Haloalkaline-Tolerant Methanotrophic Bacterium.

Valentina N Khmelenina, David a C Beck, Christine Munk, Karen Davenport, Hajnalka Daligault, Tracy Erkkila, Lynne Goodwin, Wei Gu, Chien-Chi Lo, Matthew Scholz, et al.

### ► To cite this version:

Valentina N Khmelenina, David a C Beck, Christine Munk, Karen Davenport, Hajnalka Daligault, et al.. Draft Genome Sequence of *Methylobacterium buryatense* Strain 5G, a Haloalkaline-Tolerant Methanotrophic Bacterium.. *Genome Announc*, 2013, 1 (4), epub ahead of print. 10.1128/genomeA.00053-13 . hal-00880783

**HAL Id: hal-00880783**

**<https://hal.science/hal-00880783>**

Submitted on 6 Nov 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Draft Genome Sequence of *Methylobacterium buryatense* Strain 5G, a Haloalkaline-Tolerant Methanotrophic Bacterium

Valentina N. Khmelenina,<sup>a</sup> David A. C. Beck,<sup>b,c</sup> Christine Munk,<sup>e</sup> Karen Davenport,<sup>e</sup> Hajnalka Daligault,<sup>e</sup> Tracy Erkkila,<sup>e</sup> Lynne Goodwin,<sup>e</sup> Wei Gu,<sup>e</sup> Chien-Chi Lo,<sup>e</sup> Matthew Scholz,<sup>e</sup> Hazuki Teshima,<sup>e</sup> Yan Xu,<sup>e</sup> Patrick Chain,<sup>e</sup> Francoise Bringel,<sup>f</sup> Stéphane Vuilleumier,<sup>f</sup> Alan DiSpirito,<sup>g</sup> Peter Dunfield,<sup>h</sup> Mike S. M. Jetten,<sup>i</sup> Martin G. Klotz,<sup>j</sup> Claudia Knief,<sup>k</sup> J. Colin Murrell,<sup>l</sup> Huub J. M. Op den Camp,<sup>i</sup> Yasuyoshi Sakai,<sup>m</sup> Jeremy Semrau,<sup>n</sup> Mette Svenning,<sup>o</sup> Lisa Y. Stein,<sup>p</sup> Yuri A. Trotsenko,<sup>a</sup> Marina G. Kalyuzhnaya<sup>d</sup>

GK Skryabin Institute of Biochemistry and Physiology of Microorganisms, Russian Academy of Sciences, Pushchino, Russia<sup>a</sup>; Department of Chemical Engineering, University of Washington, Seattle, Washington, USA<sup>b</sup>; eScience Institute, University of Washington, Seattle, Washington, USA<sup>c</sup>; Department of Microbiology, University of Washington, Seattle, Washington, USA<sup>d</sup>; Los Alamos National Laboratory, Joint Genome Institute, Biosciences Division Genome Science B6, Los Alamos, New Mexico, USA<sup>e</sup>; Equipe Adaptations et Interactions Microbiennes dans l'Environnement, UMR 7156 UDS-CNRS Génétique Moléculaire, Génomique, Microbiologie, Université de Strasbourg, Strasbourg, France<sup>f</sup>; Roy J. Carver Department of Biochemistry, Biophysics and Molecular Biology, Iowa State University, Ames, Iowa, USA<sup>g</sup>; Department of Biological Sciences, University of Calgary, Calgary, Canada<sup>h</sup>; Department of Microbiology, Institute of Water and Wetland Research, Radboud University Nijmegen, Nijmegen, the Netherlands<sup>i</sup>; Department of Biology, University of North Carolina, Charlotte, North Carolina, USA<sup>j</sup>; Institute of Crop Science and Resource Conservation—Molecular Biology of the Rhizosphere, University of Bonn, Bonn, Germany<sup>k</sup>; School of Environmental Sciences, University of East Anglia, Norwich, United Kingdom<sup>l</sup>; Division of Applied Life Sciences, Graduate School of Agriculture, Kyoto University, Kyoto, Japan<sup>m</sup>; Department of Civil & Environmental Engineering, the University of Michigan, Ann Arbor, Michigan, USA<sup>n</sup>; Department of Arctic and Marine Biology, University of Tromsø, Tromsø, Norway<sup>o</sup>; OMeGA, the Organization for Methanotrophic Genome Analysis, Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada<sup>p</sup>

**Robust growth of the gammaproteobacterium *Methylobacterium buryatense* strain 5G on methane makes it an attractive system for CH<sub>4</sub>-based biocatalysis. Here we present a draft genome sequence of the strain that will provide a valuable framework for metabolic engineering of the core pathways for the production of valuable chemicals from methane.**

Received 23 January 2013 Accepted 8 May 2013 Published 27 June 2013

**Citation** Khmelenina VN, Beck DAC, Munk C, Davenport K, Daligault H, Erkkila T, Goodwin L, Gu W, Lo C-C, Scholz M, Teshima H, Xu Y, Chain P, Bringel F, Vuilleumier S, DiSpirito A, Dunfield P, Jetten MSM, Klotz MG, Knief C, Murrell JC, Op den Camp HJM, Sakai Y, Semrau J, Svenning M, Stein LY, Trotsenko YA, Kalyuzhnaya MG. 2013. Draft genome sequence of *Methylobacterium buryatense* strain 5G, a haloalkaline-tolerant methanotrophic bacterium. *Genome Announc.* 1(4):e00053-13. doi:10.1128/genomeA.00053-13.

**Copyright** © 2013 Khmelenina et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Marina G. Kalyuzhnaya, mkalyuzh@uw.edu.

Microbial utilization of methane is a key step in the carbon cycle (1–3). Methanotrophs provide an attractive platform for production of commodity chemicals and biofuels from natural gas/renewable biogas (4–6). Over the last 10 years, several novel methane-utilizing microbes have been isolated in pure culture but only a few, including *Methylobacterium buryatense* strain 5G, show robust growth on methane (5–7).

The draft genome was generated at the Department of Energy (DOE) Joint Genome Institute using Illumina sequencing (8). A short-insert paired-end library (insert size of 270 bp) generated 7.25 Mbp of data (<http://www.jgi.doe.gov/>). The initial draft data were assembled with Allpaths version 39750 and computationally shredded into 10-kbp overlapping fake reads (9). The initial data were also assembled with Velvet, version 1.1.05 (10), computationally shredded into 1.5-kbp overlapping fake reads, reassembled with Velvet, and shredded into 1.5-kbp overlapping fake reads. The fake reads from the Allpaths and two Velvet assemblies, as well as a subset of the Illumina CLIP paired-end reads, were assembled using parallel Phrap, version 4.24 (High Performance Software, LLC). Possible misassemblies were corrected by manual editing in Consed (11–13). Gap closure was accomplished using repeat resolution software and sequencing of bridging PCR fragments with Sanger and/or PacBio technologies (C. Han, W. Gu, unpublished). Fifty-three PCR PacBio consensus sequences were used to close gaps and to improve the quality of the final sequence.

The total estimated size of the genome is 5.4 Mb with an average coverage of 1,343×.

Comparative genome analysis of strain 5G and two other *Methylobacterium* species, *M. album* strain BG8 and *M. alcaliphilum* strain 20Z, revealed that 5G was most similar to *M. alcaliphilum*, sharing approximately 70% of its proteome at 90% protein sequence identity.

We identified genes encoding membrane-associated methane monooxygenase, soluble methane monooxygenase and an associated chaperon and a transcriptional activator (14), pyrroloquinoline quinone (PQQ)-dependent methanol dehydrogenase, an associated *c*-cytochrome, genes for enzyme assembly and PQQ biosynthesis, tetrahydromethanopterin- and tetrahydrofolate-linked C<sub>1</sub>-transfer pathways, two formate dehydrogenases, and the ribulose monophosphate pathway. The Embden-Meyerhof-Parnas pathway, the Entner–Doudoroff pathway, and the pentose phosphate pathway (transaldolase variant) are predicted. As with the genomes of other gammaproteobacterial methanotrophs, the genome of *M. buryatense* 5G encodes all genes essential for operation of the citric acid cycle and the serine cycle, except for phosphoenolpyruvate carboxylase, isocitrate lyase, and the ethylmalonyl pathway (15,16).

Genes for urea uptake and hydrolysis, assimilatory nitrate/nitrite reduction, dissimilatory nitric oxide reduction, and ammonium uptake were identified. A gene homologous to hy-

droxylamine oxidoreductase is present (17, 18). The ammonium assimilation inventory includes genes for glutamate and alanine dehydrogenases, glutamate synthase/glutamine synthetase, serine-pyruvate/serine-glyoxylate, and aspartate aminotransferases (19). Genes essential for ectoine biosynthesis were identified.

**Nucleotide sequence accession numbers.** The *Methylobacterium buryatense* 5G genome sequence was deposited in GenBank/EMBL under the accession numbers [AOTL01000000](#) and [KB455575](#) and [KB455576](#).

## ACKNOWLEDGMENTS

This work was supported by the National Science Foundation (MCB-0842686, to M.G.K.), NSERC (L.Y.S.), and the Russian Foundation for Basic Research (RFBR 11-04-00801, to V.N.K.). The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-05CH11231.

## REFERENCES

- Op den Camp HJM, Islam T, Stott MB, Harhangi HR, Hynes A, Schouten S, Jetten MSM, Birkeland N-K, Pol A, Dunfield PF. 2009. Environmental, genomic and taxonomic perspectives on methanotrophic *Verrucomicrobia*. *Environ. Microbiol. Rep* 1:293–306.
- Semrau JD, DiSpirito AA, Yoon S. 2010. Methanotrophs and copper. *FEMS Microbiol. Rev.* 34:496–531.
- Murrell JC, Jetten MSM. 2009. The microbial methane cycle. *Environ. Microbiol. Rep.* 1:279–284.
- Trotsenko YA, Doronina NV, Khmelenina VN. 2005. Biotechnological potential of aerobic methylotrophic bacteria: a review of current state and future prospects. *Appl. Biochem. Microbiol.* 41:433–441.
- Ojala DS, Beck DA, Kalyuzhnaya MG. 2011. Genetic systems for moderately halo(alkali)philic bacteria of the genus *Methylobacterium*. *Meth. Enzymol.* 495:99–118.
- Kaluzhnaya MG, Khmelenina VN, Eshinimaev B, Suzina N, Nikitin D, Solonin A, Lin JL, McDonald I, Murrell C, Trotsenko YA. 2001. Taxonomic characterization of new alkaliphilic and alkalitolerant methanotrophs from soda lakes of the southeastern Transbaikalian region and description of *Methylobacterium buryatense* sp. nov. *Syst. Appl. Microbiol.* 24:166–176.
- Kalyuzhnaya MG, Khmelenina V, Eshinimaev B, Sorokin D, Fuse H, Lidstrom M, Trotsenko Y. 2008. Reclassification and emended description of halo(alkali)philic and halo(alkali)tolerant methanotrophs provisionally assigned to the genera *Methylobacterium* and *Methylobacter* and emended description of the genus *Methylobacterium*. *Int. J. Syst. Evol. Microbiol.* 58:591–596.
- Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438.
- Butler J, MacCallum I, Kleber M, Shlyakhter IA, Belmonte MK, Lander ES, Nusbaum C, Jaffe DB. 2008. ALLPATHS: de novo assembly of whole-genome shotgun microreads. *Genome Res.* 18:810–820.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
- Ewing B, Green P. 1998. Base-calling of automated sequencer traces using Phred. II. Error probabilities. *Genome Res.* 8:186–194.
- Ewing B, Hillier L, Wendt MC, Green P. 1998. Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res.* 8:175–185.
- Gordon D, Abajian C, Green P. 1998. Consed: a graphical tool for sequence finishing. *Genome Res.* 8:195–202.
- Csáki R, Bodrossy L, Klem J, Murrell JC, Kovács KL. 2003. Genes involved in the copper-dependent regulation of soluble methane monooxygenase of *Methylococcus capsulatus* (Bath): cloning, sequencing and mutational analysis. *Microbiology* 149:1785–1795.
- Svenning MM, Hestnes AG, Warttinen I, Stein LY, Klotz MG, Kalyuzhnaya MG, Spang A, Bringel F, Vuilleumier S, Lajus A, Médigue C, Bruce DC, Cheng JF, Goodwin L, Ivanova N, Han J, Han CS, Hauser L, Held B, Land ML, Lapidus A, Lucas S, Nolan M, Pitluck S, Woyke T. 2011. Genome sequence of the Arctic methanotroph *Methylobacter tundripaludum* SV96. *J. Bacteriol.* 193:6418–6419.
- Ward N, Larsen Ø, Sakwa J, Bruseth L, Khouri H, Durkin AS, Dimitrov G, Jiang L, Scanlan D, Kang KH, Lewis M, Nelson KE, Methé B, Wu M, Heidelberg JF, Paulsen IT, Fouts D, Ravel J, Tettelin H, Ren Q, Read T, DeBoy RT, Seshadri R, Salzberg SL, Jensen HB, Birkeland NK, Nelson WC, Dodson RJ, Grindhaug SH, Holt I, Eidhammer I, Jonassen I, Vanaken S, Utterback T, Feldblyum TV, Fraser CM, Lillehaug JR, Eisen JA. 2004. Genomic insights into methanotrophy: the complete genome sequence of *Methylococcus capsulatus* (Bath). *PLoS Biol.* 2:1616–1628.
- Campbell MA, Nyerges G, Kozłowski JA, Poret-Peterson AT, Stein LY, Klotz MG. 2011. Model of the molecular basis for hydroxylamine oxidation and nitrous oxide production in methanotrophic bacteria. *FEMS Microbiol. Lett.* 322:82–89.
- Simon J, Klotz MG. 2013. Diversity and evolution of bioenergetic systems involved in microbial nitrogen compound transformations. *Biochim. Biophys. Acta* 1827:114–135.
- Boden R, Cunliffe M, Scanlan J, Moussard H, Kits KD, Klotz MG, Jetten MS, Vuilleumier S, Han J, Peters L, Mikhailova N, Teshima H, Tapia R, Kyrpides N, Ivanova N, Pagani I, Cheng JF, Goodwin L, Han C, Hauser L, Land ML, Lapidus A, Lucas S, Pitluck S, Woyke T, Stein L, Murrell JC. 2011. Complete genome sequence of the aerobic marine methanotroph *Methylobacterium methanica* MC09. *J. Bacteriol.* 193:7001–7002.