

From apparent competition to facilitation, impacts of consumer niche construction on the coexistence and stability of consumer-resource communities

Aurore Picot, Thibaud Monnin, Nicolas Loeuille

▶ To cite this version:

Aurore Picot, Thibaud Monnin, Nicolas Loeuille. From apparent competition to facilitation, impacts of consumer niche construction on the coexistence and stability of consumer-resource communities. Functional Ecology, 2019, 33 (9), pp.1746-1757. 10.1111/1365-2435.13378 hal-02174329

HAL Id: hal-02174329 https://cnrs.hal.science/hal-02174329

Submitted on 8 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 From apparent competition to facilitation, impacts of consumer

2 niche construction on the coexistence and stability of consumer-

3 resource communities

4	Aurore Picot ^{*1} , Thibaud Monnin ¹ , Nicolas Loeuille ¹			
5	¹ Sorbonne Université, Université Paris Est Créteil, Université Paris Diderot,			
6	CNRS, INRA, IRD, Institute of Ecology and Environmental sciences - Paris, iEES-Paris,			
7	75005 Paris, France			
8	*Corresponding author: aurore.picot@normalesup.org			

9 Summary

In addition to their direct trophic effects, some consumers have a positive indirect
 effect on their resource, due to niche construction. A predator can facilitate its prey
 resource acquisition, through prey transport, or through modifications of nutrient cycling.
 Other predators defend their prey against other predators, or actively manage it, as in
 agriculture, which is found in numerous taxa such as humans, but also ants, beetles, fishes
 and microbes.

16 2. Here we investigate the ecological consequences of considering such positive 17 effects in a simple two resource–one predator module, in which the consumer has a 18 positive effect on one of the resources.

19 3. We consider several scenarios, in which the positive effect of the resource is 20 either non costly, ie resulting from a by-product of the consumer phenotype such as 21 nutrient cycling, or costly. The cost either decreases the exploitation of the helped 22 resource or the opportunity to forage the alternative resource.

4. We show that by modifying the trophic interactions in the module, niche
construction alters the apparent competition between the resources, thereby impacting
their coexistence.

5. We investigate how the intensity of niche construction impacts species coexistence, the distribution of biomass among the three species, and the stability of the community. When niche construction has little or no cost, it leads to higher consumer and helped resource densities, while decreasing the alternative resource density. Alternatively, when niche construction has a strong cost, the alternative resource can increase in density, niche construction thereby leading to the emergence of facilitative interactions among resource species.

33 Keywords

- 34 niche construction, consumer-resource theory, apparent competition, facilitation,
- 35 agriculture, trophic module

36 Introduction

37 While most studies in network ecology consider only one interaction type (ie, 38 either food webs or mutualistic networks), the co-occurrence of different interaction types 39 within networks has been of increasing interest in the last decade, in both theoretical and 40 empirical studies (Fontaine et al., 2011; Kéfi et al., 2012; Mougi & Kondoh, 2012). In particular, positive interactions that co-occur with antagonistic interactions may originate 41 42 from mutualism, facilitation (Bruno, Stachowicz, & Bertness, 2003), ecosystem engineering (Jones, Lawton, & Shachak, 1994) and niche construction (Odling-Smee, 43 44 Laland, & Feldman, 1996). They can either directly affect the partner of interaction (as in 45 mutualism) or alter the environment (as in ecological engineering) (Kéfi et al., 2012).

46 Importantly, the co-occurrence of multiple interaction types may affect predictions 47 on general ecological questions such as ecosystem functioning, community stability and 48 persistence (Fontaine et al., 2011; Kéfi et al., 2012). For instance, positive interactions 49 can increase the persistence of a consumer under resource-limited conditions (Kylafis & 50 Loreau, 2008) thus facilitating the coexistence and increasing community diversity 51 (Gross, 2008). Positive interactions can notably occur when a given species has both 52 trophic and non-trophic effects on the same partner of interaction. For instance, ant 53 species that simultaneously tend aphids and prey on them can provide benefits to aphids 54 by eating honeydew and protecting them from predators (Stadler & Dixon, 2005). Note 55 that the frontier between a consumer helping its resource and a mutualist exploiting its 56 partner is not clear, and the examples we cite later on could often belong to either 57 categories (Bronstein, 2001; Offenberg, 2001). The net demographic effect of the 58 interaction theoretically allows a complete classification but is not always measurable.

Indirect positive effects can emerge from a variety of consumer behaviours that improve some components of the resource growth rate (Brown, Ferris, Fu, & Plant, 2004 and references within). A first intuitive case would be a consumer facilitating its resource

62 nutrient acquisition. For instance, while grazing, herbivores recycle nutrients to the soil, 63 and may under some conditions increase primary productivity. An intermediate level of 64 herbivory then leads to an optimal primary productivity (grazing optimization hypothesis, 65 (de Mazancourt, Loreau, & Abbadie, 1998)). Predators may also reduce prev mortality, 66 when they protect it against other predators (through interference with or predation of 67 alternative predators) or inhibits the prey's competitors. For instance, in devil's gardens, ants kill competitors of their host plant species (Frederickson, Greene, & Gordon, 2005). 68 69 Finally, a consumer may help its resource dispersal and reproduction (eg, seed dispersal 70 linked to granivory, (Davidson, 1977)), or reduce prey intraspecific competition. The 71 nematode *Caenorhabditis elegans* for instance transports its prev bacteria and inoculate 72 them to unexploited resource pools (Ingham, Trofymow, Ingham, & Coleman, 1985; 73 Thutupalli et al., 2017).

74 Such helping behaviours can occur at a cost for the consumer or not. This 75 distinction ultimately changes the way the positive effect affects the demography of the 76 consumer species in the system, altering the prey-consumer feedback. We call the positive 77 effect passive if it results from a by-product of the consumer phenotype, with no direct 78 metabolic cost. Nutrient cycling, as in the grazing optimization hypothesis (de 79 Mazancourt et al., 1998), seed dispersal or passive transport of the resource to 80 unexploited areas as in the nematode-bacteria interaction would all fall in this category 81 (Thutupalli et al., 2017). Dissuading other predators from attacking the resource through 82 the mere presence of the consumer could also be a passive positive effect.

When the positive effect on the resource is active, its negatively affects another consumer fitness component. For example, active resource management, that we refer loosely to as "agriculture", incurs a cost in terms of time and energy. Agriculture cultivation of plants, algae, fungi and animal herding - is found in humans, but also ants, beetles, fishes and even microbes (Boomsma, 2011; Hata & Kato, 2006; Mueller, Gerardo, Aanen, Six, & Schultz, 2005; Rowley-Conwy & Layton, 2011; Smith, 2016).

The cost of agriculture can be envisioned through foraging theory (Charnov, 1976), where spending time on one resource reduces the available time to forage another resource, leading to an "opportunity cost". Another type of cost can emerge if actively defending a resource against other predators or competitors implies moving away from this resource site or decreasing its consumption (eg, by allocating time to defense rather than consumption). Then, an "exploitation cost" scenario emerges with a trade-off between resource consumption and resource protection.

96 We here investigate the ecological consequences of considering such positive 97 effects in a simple trophic module. We consider a consumer that feeds on a helped 98 resource while also foraging on a second (non-helped) resource. Such simple trophic 99 modules have been extensively used in ecology to understand mechanisms promoting 100 coexistence and stability in ecological networks (Bascompte & Melián, 2005; Holt, 1997; 101 Stouffer & Bascompte, 2010). We assume the consumer has a positive effect on one of the 102 resources (the "helped" one). We consider three scenarios that cover passive positive 103 effects ("no cost" scenario) and two types of active positive effects ("exploitation cost" 104 and "opportunity cost" scenarios). "No cost" scenarios assume that niche construction 105 only has an effect on the resource growth rate with no allocative cost (nutrient recycling, 106 cross-feeding). In "exploitation cost" scenarios, we assume that investment in niche 107 construction decreases the direct consumption of the helped resource. This may occur 108 when time devoted to defense against predators constrains consumption (for instance, ants 109 protecting aphids against ladybirds (Stadler & Dixon, 2005). "Opportunity cost" scenarios 110 assume that niche construction decreases consumption of the alternative resource. This 111 scenario is tightly linked to optimal foraging theory (Charnov, 1976; Pyke, Pulliam, & 112 Charnov, 1977)(Charnov, 1976; Pyke et al., 1977) and exploitation-exploration trade-offs 113 (Monk et al., 2018). It relates to transitions between predation and breeder behaviours, 114 found in numerous species that specialize partly (facultative aphid rearing ants) or fully 115 on cultivated resources (humans, obligate aphid rearing ants (Ivens, von Beeren,

Blüthgen, & Kronauer, 2016), fungus growing ants (Chapela, Rehner, Schultz, & Mueller,117 1994)).

118 We investigate how niche construction impacts species coexistence, the distribution 119 of biomass among the three species, and the stability of the community. Predictions can be made considering the indirect effects occurring in the system. In our module, the two 120 121 resource species do not directly compete but engage in apparent competition through their 122 interactions with the shared predator (Holt, 1977). Any increase in biomass of a given 123 prey has an indirect negative effect on the other prey as it increases predator density. Previous works show that the winner of the competition is the species that sustain the 124 highest density of predator, leading to a P^* rule similar to the R^* of exploitative 125 126 competition (Holt, Grover, & Tilman, 1994). Because we consider that one species 127 receives an additional positive effect from the consumer, indirect effects are altered. If the 128 net effect of the consumer on the helped resource is positive, then the apparent 129 competition may become an apparent antagonism as the alternative resource has a positive indirect effect on the helped resource (see figure 1). In "no cost" scenarios, we 130 predict that niche construction increases the growth rate of the helped resource, hence 131 132 negatively impacting the alternative resource through increased apparent competition. 133 Eventually, such an effect may lead to the loss of coexistence. Considering a trade-off 134 with the consumption of either resource may modify these predictions by affecting the balance of indirect effects. In the "exploitation cost" scenario, niche construction 135 136 decreases the consumption of the helped resource, hence makes it less vulnerable to 137 predation: we predict that it would win the competition because it suffers less from the indirect negative effects received from the alternative resource. We therefore predict that 138 niche construction should eventually negatively impact coexistence. In the "opportunity 139 140 cost" scenario, niche construction decreases the consumption of the alternative resource, 141 hence the effects of apparent competition should be more balanced among prey species, 142 promoting coexistence.

143 Predictions can also be made regarding the effects of niche construction on the stability of the system. In the "exploitation cost" scenario, niche construction should 144 145 reduce the energy flux from the resource to the consumer, relative to the consumer loss. 146 Such limited energy fluxes should be stabilizing, as long as the net interaction remains 147 trophic (Rip & McCann, 2011). Considering the whole three-species module, niche 148 construction is expected to be stabilizing when it increases interaction heterogeneity 149 among prev species (McCann, Hastings, & Huxel, 1998), for instance due to costs on the 150 consumption of either species. However, if the net interaction between the consumer and 151 the helped resource becomes mutualistic due to large positive effects, we expect it to be 152 destabilizing, because a negative trophic feedback loop then becomes a positive feedback 153 loop (May, 1973).



Figure 1: Presentation of the three-species model and indirect effects occurring between the resources. **a** Direct interactions: the consumer C consumes both resources R_1 and R_2 depending on its specialization on each resource (s_1 and s_2). It increases the growth of resource R_1 by a factor x, which is the niche construction investment trait. **b** When the outcome of the C- R_1 interaction is negative ($s_1 > x$), resources are limited by apparent competition. **c** When the outcome of the C- R_1 interaction is positive ($s_1 < x$), the resources engage in apparent antagonism.

161 Model presentation

1

162

Ecological dynamics are modeled using ordinary differential equations (eq 1):

$$\begin{cases} \frac{dC}{dt} = C(e_1s_1R_1 + e_2s_2R_2 - m) \\ \frac{dR_1}{dt} = R_1(b_1 - g_1R_1 - s_1C + wxC) \\ \frac{dR_2}{dt} = R_2(b_2 - g_2R_2 - s_2C) \end{cases}$$
1

163 The consumer C per capita growth rate depends on its consumption of resources 164 (modelled by specialization on resource R_i , s_i , and conversion efficiency of resource R_i , e_i) 165 and on its *per capita* death rate *m*. Resources have a logistic growth in the absence of the 166 consumer (allowing for their coexistence in such situations): b_i is the *per capita* birth rate 167 and q_i is the intraspecific competition rate for resource R_i . We do not consider any direct 168 competition between resources so that their coexistence can be entirely understood based 169 on variations of indirect effects inherent to the consumer species. To this logistic growth 170 is added a consumption rate scaled by the specialization of consumer on the resource, and 171 a niche construction effect for resource R_1 . All interactions are linear. In particular, in this 172 first model, niche construction is proportional to the investment trait x, and the consumer 173 density, modulated by a niche construction efficiency w. Such simple linear functions 174 allow for an analytical study of the system. In the Supplementary Information, we 175 consider a saturating response for the niche construction effects and discuss the 176 robustness of the results we present in the main text. We assume that e_i , s_i , b_i , q_i , x are positive, so that niche construction is facultative for the maintenance of resource 1. 177

The ecological analysis of the system focuses on the feasibility and linear stability criteria applied on the different possible equilibria. Feasibility conditions require the positivity of all equilibrium densities. The stability analysis relies on the analysis of the

- 181 Jacobian matrix (thereby assessing the return time to equilibrium following a small
- 182 disturbance) and of the invasibility of considered equilibrium by species that are not
- 183 present at equilibrium. All figures and computations were made using Mathematica 11.

184 Results

The model displays community states (equilibria) at which resources can subsist without the consumer, or the consumer coexists with one or both resources. We sum up the general stability and feasibility conditions for the "no cost" scenario, and study how niche construction impacts these coexistence and stability conditions. We then investigate how the addition of a cost modifies those results in the "exploitation cost" and "opportunity cost" scenarios. The detailed mathematical analysis can be found in the Supplementary Information.

192 1) Coexistence and stability under the "no cost" scenario $(s_1'(x)=s_2'(x)=0)$

In this scenario, the positive effect of the consumer on its resource happens through a passive effect, as a by-product of metabolism or activity of the consumer. There is no direct cost for the consumer. For instance, the large effects on nutrient recycling by the wildebeest in the Serengeti ecosystem (McNaughton, 1976) could be considered as a motivation for such a scenario. The positive effect only impacts the growth rate of resource 1 through the consumer density-dependent factor, and predation rates s_1 and s_2 are constant.

200 We show that the consumer-helped resource $(C-R_1)$ equilibrium is feasible and 201 stable when the interaction between the two species remains trophic, (ie, positive effects 202 are not too high). The ratio between the resource birth rates and their vulnerabilities 203 determines the invasion potential for resource R_2 : if it has a high birth to vulnerability 204 ratio, it can invade the consumer-helped resource system. In this $(C-R_1)$ subsystem, niche 205 construction has no effect on the resource density but increases the consumer density. The 206 increase in growth rate is compensated by an increase in predation rate, so that the 207 resource remains top-down controlled.

Regarding the $(C-R_2)$ equilibrium, our analysis reveals that niche construction does not have an impact on the equilibrium densities, as the helped resource is absent. Niche construction is destabilizing, as large *x* allows an invasion of the equilibrium by resource R_1 . Conditions of coexistence of the three species can therefore be expressed as upper and lower limits of the positive effect intensity. Hence, niche construction favors coexistence at intermediate values.

Niche construction affects the distribution of species densities. Given the dynamical
system 1, species densities at the coexistence equilibrium in the "no cost" scenario can be
written:

$$C^{*} = \frac{b_{1}e_{1}g_{2}s_{1}+b_{2}e_{2}g_{1}s_{2}-g_{1}g_{2}m}{e_{1}g_{2}s_{1}(s_{1}-wx)+e_{2}g_{1}s_{2}^{2}}$$

$$R_{1}^{*} = \frac{b_{1}-(s_{1}-wx)C^{*}}{g_{1}}$$

$$R_{2}^{*} = \frac{b_{2}-s_{2}C^{*}}{g_{2}}$$

$$2$$

217 It is then possible to show how species densities vary depending on the intensity of niche218 construction:

$$\frac{\partial C^{*}}{\partial x} = \frac{g_{2}C^{*}(x)e_{1}s_{1}}{g_{2}e_{1}s_{1}(s_{1}-wx)+e_{2}g_{1}s_{2}^{2}} > 0$$

$$\frac{\partial R_{1}^{*}}{\partial x} = \frac{e_{2}C^{*}(x)s_{2}^{2}}{g_{2}e_{1}s_{1}(s_{1}-wx)+e_{2}g_{1}s_{2}^{2}} > 0$$

$$\frac{\partial R_{2}^{*}}{\partial x} = \frac{-C^{*}(x)e_{1}s_{1}s_{2}}{g_{2}e_{1}s_{1}(s_{1}-wx)+e_{2}g_{1}s_{2}^{2}} < 0$$

$$3$$

As predicted, niche construction has a positive effect on the helped resource density, leading to a positive bottom-up effect on its consumer, negatively affecting the alternative resource density through apparent competition (figure 2). Equilibrium densities at the C- R_2 equilibrium do not vary with the intensity of niche construction, but

223 the invasion potential of R_1 increases up until R_1 can invade, eventually leading to 224 coexistence.

225 Niche construction also affect the resilience of the system (measured as the 226 negative real part of the dominant eigenvalue (Pimm & Lawton, 1977)). The stability 227 measure varies abruptly around the ecological states frontiers (see Rip & McCann, 228 (2011)), and we focus on variation for intermediate densities. For the coexistent 229 equilibrium (as predicted from the mathematical conditions, see (S12)), niche 230 construction is initially destabilizing then stabilizing, as it eventually increases interaction heterogeneity (McCann et al., 1998). Concerning the $C-R_1$ equilibrium, niche 231 232 construction is stabilizing, as it decreases the *per capita* energy flux from the resource 233 to the consumer (Rip & McCann, 2011).



Figure 2: Effects of niche construction on the stable equilibrium densities and stability in the "no cost" scenario. The consumer density is in brown, resource 1 density in orange, resource 2 density in green. Above the plot, the different states of the module are represented. An empty box means that density decreases, light box means that density does not vary, while a dark box means that density increases. Within the plot, the background shows stability variation: no variation (yellow), destabilization (red),

240 stabilization (blue). Stability is non-monotonous in white areas (transition between

241 ecological states). $e_1=e_2=0.5$, $g_1=g_2=0.8$, m=1, $b_1=2$, $b_2=4$, $s_1=2$, $s_2=2$.

242 2) Effects of costly niche construction on coexistence and stability

243 We now assume that niche construction is costly for the consumer. This cost may 244 decrease the consumption of either the helped resource ("exploitation cost"), or the alternative resource ("opportunity cost"). "Exploitation cost" scenarios include situations 245 246 where higher positive effects on the helped resource (eg agriculture) decreases the 247 exploitation of the same resource. This can happen because the time or energy devoted to 248 protection of the resource against predators cannot be used for consumption (for instance, 249 ants protecting aphids against ladybirds (Stadler and Dixon 2005)). "Opportunity cost" scenarios correspond to situations where the foraging on an alternative resource 250 251 decreases, implying a trade-off between predation and agriculture activities.

From eq 1, it is easy to show that niche construction affects the distribution of species densities as (see Supplementary Information (S14)):

$$R_{1}^{*'}(x) = \frac{\left(\left(-s_{1}+x\right)C^{*}\right)'(x)}{g_{1}} = -\frac{\left(s_{1}(x)-x\right)C^{*'}(x) + \left(s_{1}'(x)-1\right)C^{*}(x)}{g_{1}}$$

$$R_{2}^{*'}(x) = \frac{\left(-s_{2}C^{*}\right)'(x)}{g_{2}} = -\frac{s_{2}(x)C^{*'}(x) + s_{2}'(x)C^{*}(x)}{g_{2}}$$

$$4$$

These derivatives are made of two terms. The first one shows the ecological consequences of niche construction (its impact on consumer population $C^*(x)$), while the second term embodies the effect of each cost, $s_1'(x)$ and $s_2'(x)$.

257 a) "Exploitation cost" scenario $(S_1'(x) < 0, S_2'(x) = 0)$

In the "exploitation cost" scenario, niche construction directly lowers the consumption of R_1 . Intuitively, such a decrease in consumption reinforces the positive

260 effect of niche construction on R_1 . In turn, this should increase apparent competition and

harm R_2 . In such a scenario, we obtain that:

$$\frac{\partial C^{*}}{\partial x} = \left[\frac{\partial C^{*}}{\partial x}\right]^{(1)} + s_{1}'(x)\frac{g_{2}e_{1}(-C^{*}(x)s_{1}(x) + g_{1}R_{1}^{*}(x))}{g_{2}e_{1}s_{1}(x)(s_{1}(x) - x) + e_{2}g_{1}s_{2}^{2}}$$

$$\frac{\partial R_{1}^{*}}{\partial x} = \left[\frac{\partial R_{1}^{*}}{\partial x}\right]^{(1)} - s_{1}'(x)\frac{e_{2}C^{*}(x)s_{2}^{2} + g_{2}e_{1}R_{1}^{*}(x)(s_{1}(x) - x)}{g_{2}e_{1}s_{1}(x)(s_{1}(x) - x) + e_{2}g_{1}s_{2}^{2}}$$

$$\frac{\partial R_{2}^{*}}{\partial x} = \left[\frac{\partial R_{2}^{*}}{\partial x}\right]^{(1)} + s_{1}'(x)\frac{e_{1}s_{2}(C^{*}(x)s_{1}(x) - g_{1}R_{1}^{*}(x))}{g_{2}e_{1}s_{1}(x)(s_{1}(x) - x) + e_{2}g_{1}s_{2}^{2}}$$
5

262 $\left[\frac{\partial R_1^*}{\partial x}\right]^{(1)}$, $\left[\frac{\partial R_2^*}{\partial x}\right]^{(1)}$ and $\left[\frac{\partial C^*}{\partial x}\right]^{(1)}$ are density variations in the "no cost" scenario 263 (see eq. 3). These derivatives can either be all positive or all negative. Because niche 264 construction comes at a cost for the consumer, the overall effect on *C** may be negative. 265 Consumption is then relaxed on the alternative resource so that it may also benefit from 266 niche construction.

On figure 3, we illustrate these outcomes assuming a linear trade-off: $s_1(x) = s_1 - \beta_1$ 267 x, with different trade-off strengths (β_1) and interaction intensity s_2 . At low niche 268 269 construction, when the alternative interaction is low (panels a and b), variations are 270 similar to the no cost scenario (equation 3 and 5). On panel a, note that intermediate niche 271 construction does not allow the stable maintenance of a coexistent system (grey area). 272 Starting from a coexistent state, dynamics lead to the extinction of resource R_2 and the 273 exponential growth of the C- R_1 system. An initial C- R_1 subsystem (R_2 =0) would lead to 274 the extinction of the consumer. This may be explained by the fact that in the first case, the 275 transient dynamics during which R_2 is present allows large consumer populations that 276 ever increase due to the positive feedback with R_1 (the interaction C- R_1 is mutualistic, as 277 $s_1 < x$ in this area). In the second case, the initial state does not allow such an infinite 278 growth as populations are too small. Because x is intermediate, the cost on the consumer 279 growth rate is not high enough to stabilize the dynamics: the energy intake from the

280 consumption of resource 1 is still high. Hence, we obtain typical unstable mutualistic 281 dynamics with exponential growth of the consumer and the helped resource (those 282 dynamics are illustrated in supplementary information, figure S1). When niche 283 construction x is high (fig 3a, right), it decreases consumer density (reduced energy 284 intake). Since the interaction is mutualistic, this also decreases R_1 density. From the 285 helped resource perspective, the positive effect of niche construction (increasing x and 286 reducing attack rate s_1 is compensated by the negative density-dependent effect on the 287 consumer. Considering equation 4 helps visualizing this balance: the first part of the equation is negative because $C^*(x) < 0$ and $s_1(x) - x < 0$ (the interaction is mutualistic). 288 289 The second part corresponds to a positive effect stemming from the reduction in attack 290 rate $s_1(x)$. On the contrary, variations in R_2 density is only driven by the consumer density 291 effect (equation 4). We would like to draw the attention on an important consideration 292 here: for high values of niche construction, the $C-R_1$ interaction is mutualistic but it is not 293 symmetrical. In particular, when x is high enough, $s_1(x)$ tends towards 0 so that the 294 interaction tends to commensalism (null effect of R_1 on C while R_1 benefits from niche 295 construction). In panel c, because the alternative interaction is high, R_1 dominates the 296 apparent competition when there is low niche construction. Increasing x first increases 297 both R_1 and C but eventually heightens the cost of niche construction. R_2 can eventually 298 invade the system when consumer density goes below its P*. Niche construction then 299 favors both resources densities while harming the consumer density.

Effects of niche construction on stability are largely consistent with the "no cost" scenario (compare Fig3a,b vs Fig2). However, note that for higher x values, niche construction is stabilizing. Niche construction costs there reduce the attack rate on R_1 and makes the interaction tend to commensalism. This reduces the positive feedback between R_1 and *C* thereby explaining the stabilization of the system. As a corollary, intermediate niche construction levels lead to unstable dynamics such as in the grey area of Fig3a.



Figure 3: Effect of niche construction in the "exploitation cost" scenario for cost values and alternative resource interactions. The color code is the same as for figure 1. The grey area corresponds to no stable equilibria. The purple line indicates the value of niche construction where interaction between C and R₁ switches from trophic to mutualistic. $e_1=e_2=1, g_1=g_2=0.8, m=1, b_1=2, b_2=4$. In panel a, $s_1(x)=0.5-0.5 x, s_2=0.5$. In b, $s_1(x)=0.5-x, s_2=0.5$. In c, $s_1(x)=0.5-x, s_2=2$.

313 b) "Opportunity cost" scenario $(s_2'(x) < 0, s_1'(x) = 0)$

Given "opportunity costs", niche construction decreases predation on the alternative resource. We can expect this to dampen the effects of apparent competition: as x increases, benefits on resource R_1 increase, and predation on R_2 is relaxed. We thus expect coexistence to be facilitated in this scenario compared to the previous ones.

318 We obtain that:

$$\frac{\partial C^{*}}{\partial x} = \left[\frac{\partial C^{*}}{\partial x}\right]^{(1)} + s_{2}'(x) \frac{e_{2}g_{1}(g_{2}R_{2}^{*}(x) - s_{2}(x)C^{*}(x))}{g_{2}e_{1}s_{1}(s_{1}-x) + e_{2}g_{1}s_{2}(x)^{2}}$$

$$\frac{\partial R_{1}^{*}}{\partial x} = \left[\frac{\partial R_{1}^{*}}{\partial x}\right]^{(1)} - s_{2}'(x) \frac{e_{2}(s_{1}-x)(g_{2}R_{2}^{*}(x) - s_{2}(x)C^{*}(x))}{g_{2}e_{1}s_{1}(s_{1}-x) + e_{2}g_{1}s_{2}(x)^{2}}$$

$$\frac{\partial R_{2}^{*}}{\partial x} = \left[\frac{\partial R_{2}^{*}}{\partial x}\right]^{(1)} - s_{2}'(x) \frac{e_{2}g_{1}R_{2}^{*}(x)s_{2}(x) + e_{1}s_{1}C^{*}(x)(s_{1}-x)}{g_{2}e_{1}s_{1}(s_{1}-x) + e_{2}g_{1}s_{2}(x)^{2}}$$

$$6$$

As in the "exploitation scenario", we illustrate the effects of the cost, assuming a 319 320 linear trade-off $s_2(x) = s_2 - \beta_2 x$ (Fig4), varying the intensity of the cost and interaction s_1 . 321 From system 6, $s_2'(x)$ modifies the density variations expected from the "no cost 322 scenario". When this cost is low ($\beta_2 \sim 0$), qualitative patterns are consistent with the no cost 323 scenario, both in terms of density and stability variations (Fig4a). Higher costs benefit all 324 densities, as predicted (Fig4b). Niche construction also stabilizes the system, as it 325 increases the asymmetry between the two interactions (Fig4b,c). Results in Fig4c, where 326 both the cost and the alternative interaction are high, are consistent with those of figure 327 3c, that also assumed a high cost and a high alternative interaction. R_1 can only invade 328 when high niche construction reduces the consumer density under its P^* . The two 329 resource densities then increase with x, while consumer density slightly decreases. At low 330 x values, for the consumer- R_2 equilibrium, niche construction is first stabilizing then 331 destabilizing (eventually allowing the invasion of R_1). Results are also similar at higher 332 levels of niche construction (fig 4c vs 3c).



333 Figure 4: Effects of niche construction in the "opportunity cost" scenario. The color code

- 334 is the same as in figure 3. $e_1=e_2=1$, $g_1=g_2=0.8$, m=1, $b_1=2$, $b_2=4$. Panel a: $s_2(x) = 0.5 10^{-1}$
- 335 0.5 x, $s_1 = 0.5$, panel b: $s_2(x) = 0.5 x$, $s_1 = 0.5$, panel c: $s_2(x) = 0.5 x$, $s_1 = 1$

The general results for the stock variations of the densities with *x* in the three scenarios are summed up in the Table 1. We note that, when variations can be determined, the consumer is usually positively affected by niche construction, except in high cost scenarios. Similarly, niche construction is often positive for the helped resource, and may create positive facilitative effects on the second resource, when costs of niche construction are included.

	"No cost" scenario	"Exploitation cost" scenario	"Opportunity cost" scenario
C-R ₁	$\frac{\partial R_1^*}{\partial x} = 0, \frac{\partial C^*}{\partial x} > 0$	$\frac{\partial R_1^*}{\partial x} > 0, \frac{\partial C^*}{\partial x} > 0 \text{ or } < 0$	$\frac{\partial R_1^*}{\partial x} = 0, \frac{\partial C^*}{\partial x} > 0$
	C R ₁	C R ₁	C R ₁
$C-R_2$	$\frac{\partial R_2^*}{\partial x} = 0, \frac{\partial C^*}{\partial x} = 0$	$\frac{\partial R_2^*}{\partial x} = 0, \frac{\partial C^*}{\partial x} = 0$	$\frac{\partial R_2^*}{\partial x} > 0, \frac{\partial C^*}{\partial x} > 0 \text{ or } < 0$
	C R ₂	C R ₂	C R ₂
$C-R_1-R_2$	$\frac{\partial R_1^*}{\partial R_2^*} > 0, \frac{\partial R_2^*}{\partial R_2^*} < 0,$	low <i>x</i> and low cost: same as in	low x and low cost: same as
	$\frac{\partial x}{\partial C^*} > 0$	the no cost scenario	no cost scenario
	∂x	high <i>x</i> , low cost:	high x , high cost, low C - R_1 :
		R ₁ R ₂	R ₁ R ₂
		high <i>x</i> , high cost :	high x, high cost, high $C-R_1$:
		R	C R ₁ R ₂

- 342 Table 1: General effects of niche construction on density distributions. Empty, dark, light
- 343 and no boxes mean decrease, increase, no variation and indetermination, respectively.

344 3) Robustness analysis

While we assumed effects of niche construction to be proportional to average investment *x* and consumer density, little information exists to link niche construction with species demography. In the supplementary material, we study an alternative model where niche construction eventually saturates. Effects of niche construction on the distribution of biomasses are largely consistent with the linear model, though species density variations are more moderate and stabilize because of the saturating response.

351 Discussion

The motivation for our study was to investigate how the addition of a non trophic positive effect to a trophic interaction impacts the structure and functioning of a oneconsumer-two-resources module. We investigate three scenarios that differ in whether and how the positive effect entails an allocative cost for the consumer. We discuss the three questions that we addressed in the introduction: the effect of niche construction on coexistence, on the distribution of densities within the module, and on stability.

358 When niche construction has no or little cost, we find that niche construction 359 generally favors coexistence when the alternative resource initially dominates the 360 apparent competition (for instance because it has a higher intrinsic growth rate or suffers 361 less predation) (Holt, 1977). Niche construction then benefits the helped resource and 362 allows coexistence, though increasing further eventually leads to the competitive 363 exclusion of the alternative resource. Such a pattern may be linked to various empirical 364 examples. For instance, devil's gardens are almost pure plantations of one tree species 365 maintained by ants killing other, more competitive plants (Frederickson et al., 2005). On 366 the contrary, if the two resources are initially equally competitive or the alternative 367 resource is less competitive, niche construction increases the asymmetry between the two 368 resources thereby limiting coexistence. These results are consistent with classic trophic 369 module studies (Holt & Lawton, 1994), with niche construction simply modulating 370 apparent competition.

Niche construction also alters the abundance distribution of the different species. Intuitively, we expect that niche construction behaviour may benefit the consumer population as well as the abundance of the helped species. When niche construction has no or little cost, we indeed find such a pattern, and also find that the alternative resource density is decreased. Such results are largely consistent with the direct effects of niche construction: it has a positive effect on the helped resource density, while increasing

377 consumer density through bottom-up effects (increased resource availability). This 378 increase in consumer biomass leads to a negative top-down effect on the alternative 379 resource density. Consistent with such a mechanism, some studies where ant tend aphids 380 have noted an increased predation from ants on alternative non-tended aphid species 381 (Warrington & Whittaker, 1985). It is however not clear whether this negative effect 382 occurs from a consumer density increase or through changes in the foraging pattern. 383 Similarly, Wimp & Whitham (2001) show that the experimental removal of an aggressive 384 aphids-tending ant strongly increases the biodiversity of other arthropod species, 385 suggesting that the ant-aphid association has a negative impact on the abundance of other 386 species and may indeed limit species abundances and coexistence.

387 Interestingly, larger costs can strongly modify this intuitive pattern. The main effect 388 of a strong cost is, counter-intuitively, not dependent on where this cost occurs in our 389 trophic module. Whether this cost involves the exploitation of the helped resource (strong 390 "exploitation costs"), or of the alternative resource (strong "opportunity costs"), it leads to 391 a negative effect of niche construction on consumer density, and a positive effect on the 392 alternative resource. The effect on the helped resource varies depending on the balance 393 between the direct niche construction effect (through trait x) and the indirect effect on 394 consumer density. This general pattern can be interpreted in the following way: if the 395 interaction between the consumer and the helped resource becomes globally mutualistic 396 (ie, the positive effect is larger than the trophic effect of the consumer), then the consumer 397 and helped resource densities covary with the intensity of niche construction. Given a 398 high cost, niche construction then decreases both densities, helping the alternative resource through relaxed predation (either because of the decrease in consumer density or, 399 400 in the case of opportunity costs, because of lower predation rates). If the interaction 401 remains mainly trophic, then niche construction generally decreases the consumer density 402 but increases both of the resource densities due to relaxed predation. Such overall positive 403 effects of niche construction on all resources can be linked to the notion of prudent

404 predation (Goodnight et al., 2008; Slobodkin, 1974) We also note, in the "exploitation
405 cost" scenarios, that consumer density reaches an optimum at intermediate niche
406 construction.

407 We note that such positives effect on the alternative resource, not directly helped, 408 may be seen as a facilitation of this species by the $C-R_1$ interaction. Such a facilitation 409 emerges when the interaction between the consumer and the helped resources becomes 410 mostly mutualistic. Assuming strong costs of niche construction, apparent competition is 411 then replaced by a dominant facilitation effect. Assuming strong costs, niche construction 412 leads to an increase in both resources because it relaxes predation impacts not only on the 413 helped resource, but also on the alternative resource species (either through a consumer 414 density-dependent effect, or through a decrease in the *per capita* attack rate). Our model 415 thus allows a continuum between competitive and facilitative interactions, whose 416 importance in ecology is increasingly recognized (Bruno et al., 2003; He, Bertness, & 417 Altieri, 2013; Kéfi et al., 2012). Because the balance between competition and facilitation 418 in our system depends on the levels of cost, this highlights the importance of investigating the possible trade-offs associated to niche construction. We stress that the non-cultivated 419 420 resource is facilitated though it receives no direct benefit. Niche construction can modify 421 a foraging pattern in a way that is not necessarily costly: for instance, fire ants forage on 422 the ground but when they tend aphids they also forage on the arthropods present on the 423 host plant (Kaplan & Eubanks, 2005). Foraging on plants for prevs only is not profitable 424 but foraging for honeydew makes foraging for nearby preys profitable. We here assume 425 that costs of niche construction act on predation rates. This assumption, based on 426 allocation of time and energy between different functions (niche construction on one side, 427 foraging on the other), could be generalized, for instance assuming a continuum between 428 our "exploitation cost" and "opportunity cost" scenarios.

429 Effects of niche construction on stability are less intuitive. When costs are low, 430 niche construction is first destabilizing then stabilizing. This can be explained thanks to

431 previous works on stability in trophic modules or networks. Notably, when it increases the 432 heterogeneity of interaction strengths (eg, due to costs), niche construction is stabilizing, 433 as found in food web modules (McCann et al., 1998). When only the helped resource is 434 present, increasing niche construction is also stabilizing, because it decreases the *per* 435 *capita* energy flux from the resource to the consumer, consistent with classical "paradox 436 of enrichment" results (Rip & McCann, 2011; Rosenzweig, 1971).

437 We chose to keep our model linear to allow for a better analytical tractability. We 438 assume no direct competition among resources focusing on apparent competition instead. 439 We believe that adding direct (or exploitative) competition between the resources would 440 give predictions similar to the P^*-R^* coexistence rule (Holt et al., 1994) with niche 441 construction modulating the *P** value of the resource species. We analyze the robustness 442 of the model regarding the assumption of a linear niche construction effect, investigating 443 a scenario where effects of niche construction progressively saturate. In terms of 444 equilibrium densities response, we find similar patterns in the linear and the saturating model. The saturating response is generally stabilizing, as expected from previous studies 445 446 showing the stabilization of mutualism when considering saturating functional responses 447 (Holland & DeAngelis, 2010; Holland, DeAngelis, & Bronstein, 2002).

448 The explicit consideration of trophic interaction modifications has recently received 449 increased attention (Terry, Morris, & Bonsall, 2017 and references within). In this line of work, our model investigates how positive niche construction effects interfere with 450 451 apparent competition, to constrain species coexistence and community stability. While 452 these small modules are by essence simplified compared to larger, natural networks, 453 ecology has a long tradition of using them to propose and test predictions on coexistence 454 or stability. We also assume that the consumer positively affected one of the two resource 455 species only. Note however that a consumer often has positive effects on several of its 456 resources, a case we can reasonably expect in the cases of nutrient cycling (Cargill & 457 Jefferies, 1984; de Mazancourt et al., 1998) and seed dispersal (Serrano-Cinca, Fuertes-

458 Callén, & Mar-Molinero, 2005; Thutupalli et al., 2017). Explicit simulations of these 459 more complex (ie, more species or more diffuse effects) scenarios are beyond the scope of 460 this article, but they would help to get a better understanding of the role of non trophic 461 interactions in larger networks, by allowing the accumulation of more indirect effects of 462 different types (Kéfi et al., 2012).

463 While the present study would benefit from being extended to multispecies 464 networks, we believe that even in such situations, some of our results would hold. For 465 instance, if a herbivore has a positive effect on many plants through recycling, such 466 positive effects will be larger for the most nutrient limited species. Our model should then be seen as an approximation of such heterogeneities in the positive effects. Next to the 467 468 extension to complex networks, another important issue would be to explicitly consider 469 spatial dynamics. Indeed, local niche construction processes may create spatial 470 heterogeneities in nature. Obvious examples include ant gardens (Frederickson et al., 471 2005) in which whole plant species communities are modified by the combination of 472 trophic and non trophic effects of ant species, or large scale patterns of vegetation as in 473 the termite-driven hypothesis for Namibian fairy circles (Juergens, 2013; Pringle & 474 Tarnita, 2017).

475 References

- 476 Bascompte, J., & Melián, C. J. (2005). Simple trophic modules for complex food webs.
- 477 *Ecology*, *86*(11), 2868–2873. doi:10.1890/05-0101
- 478 Boomsma, J. J. (2011). Evolutionary biology: Farming writ small. *Nature*, 469(7330),
- 479 308–309. doi:10.1038/469308a
- 480 Bronstein, J. (2001). The exploitation of mutualisms. *Ecology Letters*, 4, 277–287.
- 481 Brown, D. H., Ferris, H., Fu, S., & Plant, R. (2004). Modeling direct positive feedback
- 482 between predators and prey. *Theoretical Population Biology*, 65(2), 143–152.
- 483 doi:10.1016/j.tpb.2003.09.004
- 484 Bruno, J. F., Stachowicz, J. J., & Bertness, M. D. (2003). Inclusion of facilitation into
- 485 ecological theory. Trends in Ecology & Evolution, 18(3), 119–125. doi:10.1016/S0169-
- 486 5347(02)00045-9
- 487 Cargill, S. M., & Jefferies, R. L. (1984). The Effects of Grazing by Lesser Snow Geese on
- 488 the Vegetation of a Sub-Arctic Salt Marsh. *The Journal of Applied Ecology*, *21*(2), 669.
- 489 doi:10.2307/2403437
- 490 Chapela, I. H., Rehner, S. A., Schultz, T. R., & Mueller, U. G. (1994). Evolutionary
- 491 History of the Symbiosis Between Fungus-Growing Ants and Their Fungi. Science,
- 492 266(5191), 1691–1694. doi:10.1126/science.266.5191.1691
- 493 Charnov, E. (1976). Optimal foraging, the marginal value theorem. *Theoretical*
- 494 *Population Biology*, 9(2), 129–136.
- 495 Davidson, D. (1977). Foraging ecology and community organization in desert seed-eating
- 496 ants. *Ecology*, 58(4), 725–737.
- 497 de Mazancourt, C., Loreau, M., & Abbadie, L. (1998). Grazing Optimization and Nutrient
- 498 Cycling : When Do Herbivores Enhance Plant Production ? *Ecology*, *7*9(7), 2242–2252.
- 499 Fontaine, C., Guimarães, P. R., Kéfi, S., Loeuille, N., Memmott, J., van der Putten, W. H.,
- 500 ... Thébault, E. (2011). The ecological and evolutionary implications of merging different

- 501 types of networks. *Ecology Letters*, 14(11), 1170–1181. doi:10.1111/j.1461-
- 502 0248.2011.01688.x
- 503 Frederickson, M. E., & Gordon, D. M. (2007). The devil to pay: a cost of mutualism with
- 504 Myrmelachista schumanni ants in "devil's gardens" is increased herbivory on Duroia
- 505 hirsuta trees. Proceedings of the Royal Society B: Biological Sciences, 274(1613), 1117–
- 506 1123. doi:10.1098/rspb.2006.0415
- 507 Frederickson, M. E., Greene, M. J., & Gordon, D. M. (2005). 'Devil's gardens' bedevilled
- 508 by ants. *Nature*, 437(7058), 495–496. doi:10.1038/437495a
- 509 Goodnight, C., Rauch, E., Sayama, H., De Aguiar, M. A. M., Baranger, M., & Bar-Yam,
- 510 Y. (2008). Evolution in spatial predator prey models and the "prudent predator": The
- 511 inadequacy of steady-state organism fitness and the concept of individual and group
- 512 selection. *Complexity*, *13*(5), 23–44. doi:10.1002/cplx.20209
- 513 Gross, K. (2008). Positive interactions among competitors can produce species-rich
- 514 communities. *Ecology Letters*, 11(9), 929–936. doi:10.1111/j.1461-0248.2008.01204.x
- 515 Hata, H., & Kato, M. (2006). A novel obligate cultivation mutualism between damselfish
- 516 and Polysiphonia algae. *Biology Letters*, *2*(4), 593 LP-596.
- 517 He, Q., Bertness, M. D., & Altieri, A. H. (2013). Global shifts towards positive species
- 518 interactions with increasing environmental stress. *Ecology Letters*, *16*(5), 695–706.
- 519 doi:10.1111/ele.12080
- 520 Holland, J. N., & DeAngelis, D. L. (2010). A consumer-resource approach to the density-
- 521 dependent population dynamics of mutualism. *Ecology*, 91(5), 1286–1295.
- 522 doi:10.1890/09-1163.1
- 523 Holland, J. N., DeAngelis, D. L., & Bronstein, J. L. (2002). Population Dynamics and
- 524 Mutualism: Functional Responses of Benefits and Costs. The American Naturalist,
- 525 159(3), 231–244. doi:10.1086/338510
- 526 Holt, R. D. (1977). Predation, apparent competition, and the structure of prey
- 527 communities. Theoretical Population Biology, 12(2), 197–229. doi:10.1016/0040-
- 528 5809(77)90042-9

- 529 Holt, R. D. (1997). Community Modules. In Multitrophic interactions in terrestrial
- 530 systems (pp. 333–350).
- 531 Holt, R. D., Grover, J., & Tilman, D. (1994). Simple rules for interspecific dominance in
- 532 systems with exploitative and apparent competition. American Naturalist, 144(5), 741–
- 533 771.
- Holt, R. D., & Lawton, J. . (1994). The ecological consequences of shared natural
- enemies. Annual Review of Ecology and Systematics, 25(1), 495–520.
- 536 doi:10.1146/annurev.es.25.110194.002431
- 537 Ingham, R. E., Trofymow, J. A., Ingham, E. R., & Coleman, D. C. (1985). Interactions of
- 538 Bacteria, Fungi, and their Nematode Grazers: Effects on Nutrient Cycling and Plant
- 539 Growth. Ecological Monographs, 55(1), 119–140. doi:10.2307/1942528
- 540 Ivens, A. B. F., von Beeren, C., Blüthgen, N., & Kronauer, D. J. C. (2016). Studying the
- 541 Complex Communities of Ants and Their Symbionts Using Ecological Network Analysis.
- 542 Annual Review of Entomology, 61(1), 353–371. doi:10.1146/annurev-ento-010715-
- 543 023719
- Jones, C., Lawton, J., & Shachak, M. (1994). Organisms as ecosystem engineers. *Oikos*,
 69, 373–86.
- 546 Juergens, N. (2013). The Biological Underpinnings of Namib Desert Fairy Circles.
- 547 Science, 339(6127), 1618–1621. doi:10.1126/science.1222999
- 548 Kaplan, I., & Eubanks, M. D. (2005). Aphids Alter the Community-Wide Impact of Fire
- 549 Ants. Ecology, 86(6), 1640–1649. doi:10.2307/3450789
- 550 Kéfi, S., Berlow, E. L., Wieters, E. A., Navarrete, S. A., Petchey, O. L., Wood, S. A., ...
- Brose, U. (2012). More than a meal... integrating non-feeding interactions into food
- 552 webs. *Ecology Letters*, 15(4), 291–300. doi:10.1111/j.1461-0248.2011.01732.x
- 553 Kylafis, G., & Loreau, M. (2008). Ecological and evolutionary consequences of niche
- 554 construction for its agent. *Ecology Letters*, *11*(10), 1072–81. doi:10.1111/j.1461-
- 555 0248.2008.01220.x

- 556 May, R. M. (1973). Qualitative Stability in Model Ecosystems. *Ecology*, 54(3), 638–641.
- 557 doi:10.2307/1935352
- 558 McCann, K. S., Hastings, A., & Huxel, G. R. (1998). Weak trophic interactions and the
- balance of nature. *Nature*, 395(6704), 794–798. doi:10.1038/27427
- 560 McNaughton, S. J. (1976). Serengeti Migratory Wildebeest: Facilitation of Energy Flow
- 561 by Grazing. Science, 191(4222), 92–94. doi:10.1126/science.191.4222.92
- 562 Monk, C. T., Barbier, M., Romanczuk, P., Watson, J. R., Alós, J., Nakayama, S., ...
- 563 Arlinghaus, R. (2018). How ecology shapes exploitation: A framework to predict the
- 564 behavioural response of human and animal foragers along exploration-exploitation trade-
- 565 offs. Ecology Letters. doi:10.1111/ele.12949
- 566 Mougi, A., & Kondoh, M. (2012). Diversity of interaction types and ecological
- 567 community stability. *Science*, 337(6092), 349–351. doi:10.1126/science.1220529
- 568 Mueller, U. G., Gerardo, N. M., Aanen, D. K., Six, D. L., & Schultz, T. R. (2005). The
- 569 Evolution of Agriculture in Insects. Annual Review of Ecology, Evolution, and
- 570 *Systematics*, 36(1), 563–595. doi:10.1146/annurev.ecolsys.36.102003.152626
- 571 Odling-Smee, F., Laland, K., & Feldman, M. (1996). Niche construction. American
- 572 Naturalist, 147(4), 641–648.
- 573 Offenberg, J. (2001). Balancing between mutualism and exploitation: The symbiotic
- 574 interaction between Lasius and aphids. Behavioral Ecology and Sociobiology, 49(4),
- 575 304–310. doi:10.1007/s002650000303
- 576 Pimm, S. L., & Lawton, J. H. (1977). Number of tropic levels in ecological communities.
- 577 Nature a Weekly Journal of Science, 268(5618), 329–331.
- 578 Pringle, R. M., & Tarnita, C. E. (2017). Spatial Self-Organization of Ecosystems:
- 579 Integrating Multiple Mechanisms of Regular-Pattern Formation. Annual Review of
- 580 *Entomology*, 62(1), 359–377. doi:10.1146/annurev-ento-031616-035413
- 581 Pyke, G., Pulliam, H., & Charnov, E. (1977). Optimal foraging: a selective review of
- theory and tests. *Quarterly Review of Biology*, 52(2), 138–154.

- 583 Rip, J. M. K., & McCann, K. S. (2011). Cross-ecosystem differences in stability and the
- 584 principle of energy flux. *Ecology Letters*, 14(8), 733–740. doi:10.1111/j.1461-
- 585 0248.2011.01636.x
- 586 Rosenzweig, M. (1971). Paradox of enrichment: destabilization of exploitation
- 587 ecosystems in ecological time. *Science*, *171*(Issue: 969), 385–387.
- 588 Rowley-Conwy, P., & Layton, R. (2011). Foraging and farming as niche construction:
- 589 stable and unstable adaptations. *Philosophical Transactions of the Royal Society B*:
- 590 Biological Sciences, 366(1566), 849–862. doi:10.1098/rstb.2010.0307
- 591 Serrano-Cinca, C., Fuertes-Callén, Y., & Mar-Molinero, C. (2005). Measuring DEA
- 6592 efficiency in Internet companies. Decision Support Systems, 38(4), 557–573. doi:10.1016/
- 593 j.dss.2003.08.004
- 594 Slobodkin, L. B. (1974). Prudent predation does not require group selection. American
- 595 Naturalist, 108(963), 665–678. doi:10.1086/282942
- 596 Smith, B. D. (2016). Neo-Darwinism, niche construction theory, and the initial
- 597 domestication of plants and animals. *Evolutionary Ecology*, 30(2), 307–324. doi:10.1007/
- 598 s10682-015-9797-0
- 599 Stadler, B., & Dixon, A. F. G. (2005). Ecology and Evolution of Aphid-Ant Interactions.
- 600 Annual Review of Ecology, Evolution, and Systematics, 36(1), 345–372.
- 601 doi:10.1146/annurev.ecolsys.36.091704.175531
- 602 Stouffer, D. B., & Bascompte, J. (2010). Understanding food-web persistence from local
- 603 to global scales. *Ecology Letters*, 13(2), 154–61. doi:10.1111/j.1461-0248.2009.01407.x
- 604 Terry, J. C. D., Morris, R. J., & Bonsall, M. B. (2017). Trophic interaction modifications:
- an empirical and theoretical framework. *Ecology Letters*, *20*(10), 1219–1230.
- 606 doi:10.1111/ele.12824
- 607 Thutupalli, S., Uppaluri, S., Constable, G. W. A., Levin, S. A., Stone, H. A., Tarnita, C. E.,
- 608 & Brangwynne, C. P. (2017). Farming and public goods production in *Caenorhabditis*
- 609 elegans populations. Proceedings of the National Academy of Sciences, 114(9), 2289–
- 610 2294. doi:10.1073/pnas.1608961114

- 611 Warrington, S., & Whittaker, J. B. (1985). An Experimental Field Study of Different
- 612 Levels of Insect Herbivory Induced By Formica rufa Predation on Sycamore (Acer
- 613 pseudoplatanus) I. Lepidoptera Larvae. *The Journal of Applied Ecology*, 22(3), 775.
- 614 doi:10.2307/2403228
- 615 Wimp, G. M. ., & Whitham, T. G. . (2001). Biodiversity Consequences of Predation and
- 616 Host Plant Hybridization on an Aphid-Ant Mutualism. *Ecology*, *82*(2), 440–452.