

Reinterpretation of the enigmatic Ordovician genus Bolboporites (Echinodermata).

Emeric Gillet, Bertrand Lefebvre, Véronique Gardien, Emilie Steimetz, Christophe Durlet, Frédéric Marin

▶ **To cite this version:**

Emeric Gillet, Bertrand Lefebvre, Véronique Gardien, Emilie Steimetz, Christophe Durlet, et al.. Reinterpretation of the enigmatic Ordovician genus Bolboporites (Echinodermata).. Zoosymposia, 2019, 15 (1), pp.44-70. 10.11646/zoosymposia.15.1.7 . hal-02333918

HAL Id: hal-02333918
https://hal.science/hal-02333918

Submitted on 13 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reinterpretation of the Enigmatic Ordovician Genus *Bolboporites* (Echinodermata)

EMERIC GILLET[1], BERTRAND LEFEBVRE[1,3], VERONIQUE GARDIEN[1], EMILIE STEIMETZ[2], CHRISTOPHE DURLET[2] & FREDERIC MARIN[2]

[1] *Université de Lyon, UCBL, ENSL, CNRS, UMR 5276 LGL-TPE, 2 rue Raphaël Dubois, F-69622 Villeurbanne, France*

[2] *Université de Bourgogne - Franche Comté, CNRS, UMR 6282 Biogéosciences, 6 boulevard Gabriel, F-2100 Dijon, France*

[3] *Corresponding author, E-mail: bertrand.lefebvre@univ-lyon1.fr*

Abstract

Bolboporites is an enigmatic Ordovician cone-shaped fossil, the precise nature and systematic affinities of which have been controversial over almost two centuries. For the first time, a wide range of techniques (CT-scan, SEM, cathodoluminescence, XPL, UV epifluorescence, EBSD, FT-IR and XRF spectrometry) were applied to well-preserved specimens of *Bolboporites* from Norway and Russia. Our main finding confirms its echinoderm affinities, as shown by its stereomic microstructure and by the first definitive evidence of its monocrystalline nature. Each cone consists in a single, microporous calcitic crystal with a narrow longitudinal internal canal. These results are combined with all previous data on *Bolboporites* to critically discuss five alternative interpretations of this fossil, namely theca, basal cone, spine, columnal, and holdfast, respectively. The most parsimonious scenario considers *Bolboporites* as an isolated spine, which was articulated in life by a short biserial appendage to the body wall of an unknown echinoderm, possibly of echinozoan affinities.

Introduction

The endoskeleton of echinoderms is a complex, multi-element structure typically consisting of several thousands of individual plates bound together in life by soft tissues (including collagen fibres) and each consisting of monocrystalline calcite. Taphonomic experiments suggest that collagen decays relatively soon after the death of the organism (within days or weeks), thus leading to the collapse and rapid disarticulation of the skeleton into isolated plates and/or sometimes, more resistant modules (Donovan 1991; Brett *et al*. 1997). The assignment of an isolated skeletal element to the phylum Echinodermata is generally straightforward and relies on the presence of a typical three-dimensional meshlike

microstructure, the stereom (Smith 1980a; Kouchinsky *et al.* 2012). However, both the potential diagenetic alteration of the stereom and the high morphological disparity of skeletal elements within a same individual (e.g., columnals, holdfast, spines) make it often difficult to identify, and sometimes interpret, such isolated plates, especially in the case of Palaeozoic taxa that have no current representatives (Berg-Madsen 1986; Pisera 1994; Zamora *et al*. 2013).

The situation is further complicated by the existence of numerous morphological convergences in echinoderms. For example, Palaeozoic deposits have yielded several relatively similar-looking, small, bowl- to cone-shaped structures (e.g., *Cymbionites* Whitehouse, 1941; *Oryctoconus* Colchen & Ubaghs, 1969; *Peridionites* Whitehouse, 1941; *Timorocidaris* Wanner, 1920). Their nature and precise taxonomic assignment (at class level) has often been strongly debated (Bather 1920; Gislén 1947; Schmidt 1951; Ubaghs 1968b, 1978a; Smith 1982; Alvaro & Colchen 2002; Seilacher & MacClintock 2005; Zamora *et al*. 2009). However, most of them are now convincingly interpreted either as pelmatozoan holdfasts (e.g., *Oryctoconus*; Alvaro & Colchen 2002; Seilacher & MacClintock 2005; Zamora *et al*. 2009), highly specialized columnals (e.g., Sumrall *et al*. 1997), basal thecal plates of eocrinoids (e.g., *Cymbionites*, *Peridionites*; Smith 1982) or highly derived crinoid calices made of few, tightly sutured (‘fused’) plates (e.g., *Timorocidaris*; Bather 1920; Ubaghs 1978a).

On the other hand, the interpretation of some other isolated echinoderm elements remains problematic and controversial. This is the case for the enigmatic Ordovician genus *Bolboporites* Pander, 1830, which corresponds to centimetric cone-shaped calcitic fossils, with a typical honeycomb-like ornamentation on their external lateral surface (Fig. 1B). The base of the cone is smooth, flat to strongly convex, and bears two adjoining, shallow depressions (Fig. 1A). Within this depressed area, a tiny orifice (Fig. 1A) opens into a narrow, longitudinal canal extending internally towards the apex of the cone (Yakovlev 1921; Yeltysheva 1955; Clark & Hofmann 1961; Rozhnov & Kushlina 1994a).

Bolboporites is particularly widespread and abundant in Baltica, where it is recorded from the Dapingian to the Darriwilian (e.g., Estonia, Russia; Pander 1830; Eichwald 1857; Bassler 1911; Yakovlev 1921; Yeltysheva 1955; Smith 1988; Kushlina, 1995, 2007; Federov 2003; Rozhnov & Kushlina 1994a; Rozhnov 2005) and locally to the Sandbian (e.g., Norway, Sweden; Kjerulf 1865; Lindström 1883; Kushlina 1995). This genus also occurs in the late Darriwilian of Laurentia (e.g., New York, Quebec, Virginia; Hall 1847; Billings 1859; Logan *et al*. 1863; Brainerd & Seely 1888, 1896; Miller 1889; Brainerd 1891; Ami 1896; White

1896; Ruedemann 1901; Raymond 1905, 1906, 1913; Bassler 1915; Twenhofel 1938; Butts 1940; Clark 1944, 1952; Oxley & Kay 1959; Clark & Hofmann 1961; Shaw & Bolton 2011). *Bolboporites* was also reported in the Tramore Limestone Formation of Ireland (Avalonia; Reed 1899), in deposits recently assigned to the late Darriwilian (Wyse Jackson *et al*. 2002).

This genus was originally described based on material from the Saint-Petersburg area (Russia) by Pander (1830), who considered that it was closely related to *Dactylopora* Lamarck, 1816 (see also Milne-Edwards & Haime 1851), then interpreted either as a bryozoan or as a foraminiferan, and now assigned to the algae (Dasycladales; see, e.g., Génot & Granier 2011). In North America, the first specimens of *Bolboporites* were reported in Quebec by Hall (1847), who described them as *Chaetetes* Fischer von Waldheim, 1829 (i.e., a genus of hypercalcified sponges; see Stanton *et al*. 2016). This Canadian material was later reidentified as *Bolboporites* by Billings (1859), who interpreted it as a zoophyte. Affinities with anthozoans, and in particular with tabulate corals close to *Favosites* Lamarck, 1816, were frequently suggested for *Bolboporites* (Bronn 1849; Bronn 1851–1856; Eichwald 1857, 1860; Fromentel 1861; Kjerulf 1865; Zittel 1879; Ruedemann 1901; Butts 1940). Yakovlev (1921) made the first sections through specimens of *Bolboporites,* thus demonstrating the presence of the longitudinal axial canal and internal growth lines. Based on these new observations, he concluded that *Bolboporites* was a highly derived stromatoporoid.

Possible echinoderm affinities for *Bolboporites* were first questioned by Logan *et al.* (1863), Quenstedt (1881) and Lindström (1883), based on the observation of stereom microstructure. Miller (1889) interpreted Russian specimens of *Bolboporites* as probable echinoderms, but North American ones as corals. With few exceptions (see above), the assignment of *Bolboporites* to echinoderms was finally accepted by most authors in the late 19th century. However, its nature and precise taxonomic affinities remained largely debated and enigmatic (Jaekel 1899; Bassler 1911; Régnell 1956, 1982; Smith 1988). Quenstedt (1881) was the first to point out that the morphology of *Bolboporites* was similar to that of isolated asteroid or echinoid spines. This interpretation was followed by several authors, who interpreted *Bolboporites* as probable spines belonging to various groups of echinoderms: asteroids (Lindström 1883; Yeltysheva 1955; Régnell 1956); echinoids (Wanner 1920); or ‘cystoids’ close to *Palaeocystites tenuiradiatus* (Hall, 1847) (Clarke & Hoffman 1961).

An alternative hypothesis was proposed by Ami (1896), who considered that *Bolboporites* was not an isolated spine, but the internal mould of the theca of an unknown ‘cystoid’. Following this interpretation, von Wöhrmann (*in* Jaekel 1899) suggested that *Bolboporites* was possibly the internal mould of a cheirocrinid rhombiferan (see also Régnell 1956). This

interpretation of *Bolboporites* as corresponding to the body capsule (theca) of a 'cystoid' was further elaborated by Rozhnov & Kushlina (1994a, b), based on the observation of two sets of small skeletal elements articulated into the two depressions on the convex surface of some well-preserved specimens. These two series of plates were interpreted as elements of a single feeding appendage (brachiole), which was inserting onto the convex (oral) surface of a highly derived eocrinoid (Rozhnov & Kushlina 1994a, b; Kushlina 1995, 2006, 2007; Rozhnov 2005, 2009). In this interpretation, the central longitudinal canal corresponds to the stem, which was entirely surrounded by massive, fused thecal plates.

Recently, the examination by one of us (BL) of numerous individuals of *Coelosphaeridium* Roemer, 1885 (a genus of Ordovician calcareous green algae; see Kato *et al*. 1987; Spjeldnaes & Nitecki 1990; Baarli 2008) from the Sandbian of Norway showed remarkable similarities in size, morphology and external ornamentation with co-occurring specimens of *Bolboporites* from the same levels and localities. This observation thus questioned the echinoderm affinities of *Bolboporites* (Lefebvre 2014, 2017). As a consequence, the aims of this paper were to apply for the first time a wide range of techniques (e.g., cathodoluminescence, CT-scan, FT-IR analyses, SEM, EBSD) on well-preserved specimens of *Bolboporites*, so as to test their putative echinoderm affinities and, if confirmed, to discuss the nature of *Bolboporites* (isolated skeletal element *vs*. body capsule), as well as its systematic position within the phylum Echinodermata.

Material and Methods

Material. This study is based on 28 specimens of *Bolboporites* from Norway and Russia. The ten Norwegian specimens were selected within the abundant material of *Bolboporites* sp. (about 100 individuals) belonging to the collections of the Paleontologisk Museum, Oslo (acronym: PMO). All Norwegian specimens of *Bolboporites* were originally collected in 1975 by J.F. Bockelie in bioclastic deposits of the Fossum Formation (Sandbian), at Gravastranda, Herøya (Skien-Langesund area, Norway). These levels are generally interpreted as relatively shallow-water deposits yielding abundant and diverse benthic assemblages regularly smothered by storm deposits (Bockelie 1981; Owen *et al*. 1990). The fauna is dominated by brachiopods, bryozoans, echinoderms (caryocystitid and cheirocrinid rhombiferans, eocrinoids, crinoids, edrioasteroids) and trilobites, associated to rare cephalopods and graptolites (Bockelie 1981; Owen *et al*. 1990).

Eighteen individuals of *Bolboporites mitralis* Pander, 1830, from Russia were made available for this study by S.V. Rozhnov, who donated them to the palaeontological collections of Lyon 1 University (acronym: UCBL). This material was collected on the banks of the Lynna river (Saint-Petersburg area) in the upper member (BIIγ, Frizy Limestone) of the Volkhov Formation. This ~3 m thick stratigraphic unit is dated as early Darriwilian, based on the occurrence of both conodonts typical of the *Baltoniodus norrlandicus* Zone and trilobites characteristic of the Scandinavian *Megistaspis simon* Zone (Federov 2003; Dronov 2005). The upper member of the Volkhov Formation consists predominantly of nodular, glauconitic limestones, with several intercalated levels of shales. The bioclastic limestones are generally interpreted as storm-generated deposits, in a shallow-water, temperate setting (Dronov 2005). In these levels, faunal assemblages are dominated by brachiopods, ostracods and isolated pelmatozoan remains, associated to bryozoans, conulariids, graptolites and trilobites (Federov 2003).

Methods. As Norwegian specimens are preserved as mouldic impressions in the rock, their original external aspect was revealed by making latex casts, which were coated with ammonium chloride (NH_4Cl) for observation and photographic purposes. External morphological features of *Bolboporites* sp. from Norway were observed at Lyon 1 University, with a Zeiss SteREO Discovery.V8 stereomicroscope binocular and captured with a Zeiss AxioCam MRc5 digital camera.

In contrast to the Norwegian material, the Russian specimens are preserved as three-dimensional fossils, thus allowing the application of a wider range of techniques of observation and analyses. Several Russian specimens were embedded in hydrophilic acrylic resin of low viscosity (LR white resin), allowing longitudinal and transverse sections with a Leica SP 1600 saw microtome. Other Russian specimens were set in epoxy resin and cut to make polished thin sections with a thickness of 30 μm and 130 μm. Both sections were observed with a Hitachi TM-100 scanning electron microscope (SEM) at the Université de Bourgogne, Dijon, so as to document putative internal structures.

Thin sections were observed under plane polarized light (PPL), cross-polarized light (XPL) and epifluorescence UV using a Nikon AZ100 microscope, equipped with a 360 nm exciting source and a Zeiss Axiocam MRc5 (Université de Bourgogne, Dijon). Polished thin sections were also observed under cathodoluminescence using a Leica MZ12 binocular microscope equipped with a Luminoscope ELM-3R device and a Zeiss Axiocam MRc5 camera (Université de Bourgogne, Dijon). Cathodoluminescence (CL) was successfully applied by

Gorzelak & Zamora (2013), so as to reveal the internal stereomic microstructure preserved in skeletal elements of various Cambrian echinoderms. This technique was usually shown to be efficient, even in the case of relatively strongly recrystallized specimens.

For FT-IR investigations, thin sections were used to extract *in situ* - with the tip of a scalpel blade - small chips of material from different sampling points located both inside the fossil and in the surrounding resin. The extracted materials were then reduced into powders with an agate mini mortar and pestle (< 10 μm) and the powder was subsequently analyzed by Fourier transform infrared (FT-IR) spectroscopy, to identify the different mineralogical phases, on a ALPHA FT-IR BRUKER device equipped with an ALPHA-P module. Data acquisition was performed in the 4000-500 cm^{-1} wavenumber range (12 scans at a spectral resolution of 4 cm^{-1}), in ATR mode (Attenuated Total Reflectance) with a single reflection diamond crystal adapted to solids. Blank spectra were acquired on resin alone. The qualitative assignment of absorption bands was performed by comparison with known IR spectra found in the literature (Jones & Jackson, 1993).

The magnesium content of some Russian specimens of *Bolboporites* was checked with a Bruker S1 Titan spectrometer equipped with a collimated beam and incorporated in a laboratory console. Every measurement consisted of two successive beam phases of 60 seconds, with energies of 45 kV and 15 kV. This protocol allows measuring equivalent MgO mass concentrations above 1% in relatively small windows (2 mm in diameter) and directly in the thicker thin sections.

The internal structure of one well-preserved *Bolboporites* was also analyzed with electron back-scattered diffraction (EBSD), a technique that permits the characterization of complex polycrystalline materials at nanoscale. In short, it allows measuring and representing - via 2D-coloured maps and pole figures - the crystallographic orientation of individual nanograins with respect to each other. This technique, currently used in materials science, is particularly adapted for calcium carbonate fossil and non-fossil biominerals (Checa *et al*. 2009; Cusack 2016). To this end, an embedded sample was manually mirror-polished on 0.05μm aluminium oxide powder and further processed on a vibratory polisher. The sample was fixed on a sample holder and analyzed on a JEOL JSM 760 F field emission scanning electron microscope, from which coloured maps were produced, with a step of 250 nm. Measurements were performed at the periphery of the sample and also in different areas of the central zone separated from each other by a few millimeters; this allowed detection of potential crystallographic disorientations at milli-metric scale.

Microtomographic observations were performed by using a Bruker CT-scan (Skyscan 1174 model) at the Université de Bourgogne, Dijon (Morphoptics Service), to obtain virtual cross-sections through some specimens, and also to reconstruct a three-dimensional model of *Bolboporites*. In recent years, tomography has become a routine technique of imagery, so as to reveal internal structures in various fossils and, in particular, Palaeozoic echinoderms (Sutton *et al*. 2005; Rahman & Clausen 2009; Rahman & Zamora 2009; Rahman *et al*. 2010, 2015; Briggs *et al*. 2017). Data acquisition was obtained at 50 kV and 800 µA. Two images per position (number of frames: 2) were obtained, each of them after an exposure time of 2500 ms. The rotation step of the sample was 0.7° and the total acquisition time was 75 minutes.

Other individuals of *B. mitralis* were kept intact, so as to explore minute details of their external morphology. However, in most specimens, the base of the cone and/or the honeycomb cells on the lateral walls were partly concealed by a thin layer of sedimentary rock. Consequently, fossils were placed in an ultrasonic cleaner containing a solution of dilute ethylenediaminetetraacetic acid (EDTA, 1% wt/vol, pH 8) to remove all pieces of surrounding rock and better expose the external aspect of the specimens. Once cleaned, fossils were rinsed with water, then with ethanol (C_2H_5OH), and finally dried with a hair-dryer. Observation of the external aspect of the specimens was made using both a Zeiss SteREO Discovery.V8 binocular stereomicroscope, equipped with a Zeiss AxioCam MRc5 digital camera, at Université Lyon 1, and a Hitachi TM-100 Scanning Electron Microscope (SEM) at the Université de Bourgogne, Dijon.

Finally, three dried specimens of the Recent asteroid *Pentaceraster mammilatus* (Audouin, 1826) were examined and prepared for morphological comparison purposes. This material belongs to the R. Koehler collections, which are part of the zoological collections of Lyon 1 University (acronym: UCBL). The specimens were collected between 1895 and 1930 (precise date of sampling not reported on labels) from an unknown locality, possibly in the Red Sea or the western part of the Indian Ocean (Clark & Rowe 1971). Dissection and extraction of some aboral spines was made with a scalpel. Photographs were made with a Nikon D5000 camera in the palaeontological collections of Lyon 1 university (CERESE).

Results

Chemical and mineralogical analyses.

The FT-IR spectroscopy performed on transverse sections of *Bolboporites mitralis* included in LR white resin generated two contrasting sets of infrared spectra, depending on the position of the sampling points, outside or inside the fossil (Fig. 2). The surrounding LR white resin produced a characteristic reference IR spectrum (Fig. 2B), while infrared spectra obtained in sampling points located within the fossils all showed the three absorption bands that are characteristic of calcite (Fig. 2C-E) at 711-712 cm^{-1}, 871 cm^{-1}, and 1395 cm^{-1}, respectively. This spectrum is clearly distinct from that of aragonite (not shown), characterized by a doublet at 700-713 cm^{-1} and two bands at 858 and 1477 cm^{-1}, in addition to a sharp one at 1083 cm^{-1}. No dolomite (identified by absorption bands at 729, 882 and 1441 cm^{-1}) was detected in the central zone, Our data unambiguously showed that the Russian specimens of *Bolboporites* are entirely made of calcite.

A slightly more complex infrared spectrum was observed in sampling points located on lateral edges of the fossils, with the three absorption bands typical of calcite, but also some additional minor bands (Fig. 2C, G). Comparison with the 'control' spectrum shows that this signal corresponds to the combination of both calcite and LR white resin infrared spectra, resulting from either the irregular external morphology of the body wall (ornamentation consisting of honeycomb cells), and/or a limited penetration of the resin into micropores and microfractures of the specimens.

Finally, a large absorption band of low amplitude in the 3700-3100 cm^{-1} range was observed in one spectrum corresponding to a sampling spot located within the fossil (Fig. 2F). This signal did not result from any contamination from the resin, but more likely corresponded to the vibrations of OH bond, suggesting the occurrence of water, putatively in the form of small fluid inclusions within the calcite.

The XRF measurements were made in central and exterior parts of *Bolboporites* sections. Magnesium was only detected, with MgO values between 1 and 5% (mass concentration), in most peripheral parts of the skeleton, where analyzed windows encroach the surrounding sedimentary rock. Close examination of these parts, under SEM and cathodoluminescence (see below) revealed the presence of small dolomite rhombs, thus driving the Mg content. In all other parts of the skeleton, Mg was never found, being below the 1% detection limit. Thus, a low magnesium calcite (LMC) is deduced for their current composition.

Microstructures.

Observation of thin sections of *B. mitralis* with a polarizing microscope shows that whole cones are affected by conjugate cleavage planes (Fig. 3A,C), and have a single and right

crystal extinction (Fig 3B), typical of monocrystals. This important result is confirmed by our investigations using electron backscattered diffraction (EBSD), the outcome of which is synthesized in Figure 4. When performed on the periphery of the section (Fig. 4A-D), EBSD mapping shows that the crown interface is constituted of a mixture of micritic grains (2 µm or less) surrounding sparitic crystals of about tens to more than 100 microns in diameter (Fig. 4B, D) while the upper left corner of the map corresponds to the sample itself, which is symbolized by one unique colour, and is, consequently, monocrystalline. The EBSD mapping performed on the central zone of the sample (Fig. 4E-G) shows that the analyzed area is uniform, with no detectable grain limit. This demonstrates clearly that the structure is monocrystalline. Interestingly, when maps are produced in different central zones distant from each other by few millimetres, one notices a slight change of crystallographic orientation, symbolized by a minor colour change: for example, two zones 1 mm apart (either in X or Y) show a grain orientation spread of about 0.3 - 0.5°. When the distance is larger between two analyzed zones (such as 4 mm between Fig. 4F and G), the disorientation is more important, around 3.73° in the present example. In summary, EBSD mapping confirms that the calcite that constitutes a specimen of *Bolboporites* is monocrystalline, with a very minor and gradual crystalline disorientation at millimetric scale, without any detectable grain limit.

Under cathodoluminescence, both transverse and longitudinal sections of *B. mitralis* revealed the same mottled, orange-to-brown luminescent microstructure (Fig. 5C). Orange irregular dots and axes are aligned within a browner sealing calcite crystal, forming a CL pattern usually detected for ancient echinoderm stereoms (e.g., Gorzelak & Zamora 2013). These orange parts probably represent the former porous stereom that has been secondary cemented by a syntaxial brown-luminescent cement. Under XPL and PPL, the same stereom-like microstructure is also detected in some parts of the cones, with the tenuous occurrence of small dark spots aligned along cleavage planes (Fig. 3C). At a larger scale, the stereom of *Bolboporites* exhibits a slightly differentiated, darker, narrow peripheral rim in CL (Fig. 5A,B), which probably results either from a diagenetic effect or from a distinct lighter density of the former microstructure. A similar observation was made on virtual cross-sections of *B. mitralis* obtained by CT-scan, which show a diffuse peripheral rim, consistently lighter than the darker central part of the fossils (Fig. 6A). In CT-scan imagery, intensity restitution largely depends on the density of the materials. Consequently, the central part of *Bolboporites* is currently made of slightly more porous (less dense) skeleton than the periphery. Views by SEM confirm this deduction, with numerous micropores occuring everywhere in the skeleton,

except in its peripheral parts (Fig. 5D), where initial micropores are probably cemented by the diagenetic brown-luminescent calcite. As shown by cathodoluminescence (Fig. 5A-C), the internal canal is also cemented by the brown syntaxial calcite.

Finally, contrary to these important observations, no structures have been detected within the cones using UV epifluorescence technique. This is probably due to the absence of fluorescent organic matter within the stereomic calcite microstructure, in probable relation with an intense thermal alteration of the organic matter during the burial history of these Palaeozoic fossils.

Internal morphological features.

Sections made through specimens of *B. mitralis* confirm previous reports of a narrow, straight to gently curved, longitudinal canal, extending from the base of the cone towards the apex (Fig. 5A,C; Yakovlev 1921; Yeltysheva 1955; Clark & Hofmann 1961; Rozhnov & Kushlina 1994a, b). The observation of this longitudinal canal in all specimens, consistently in the same position, indicates that it corresponds to an original internal structure.

In contrast, both transverse and longitudinal sections also demonstrated the presence of additional, randomly distributed tubulars and incurved holes, opening either on the base or on the lateral surface of the cone, and penetrating more or less deeply into it (Fig. 5A,B,D). Optical observations made on thin sections of *B. mitralis* show that these structures are filled by a wackestone to packstone sediment, containing diverse organic fragments (e.g., shell debris, echinoderm skeletal elements) and glauconite grains (Fig. 3). In unprepared Russian specimens (i.e., without ultrasonic cleaning), the nature of this infilling appears to be similar to the surrounding matrix. Consequently, the random distribution of these holes, their variable size and depth, as well as their infilling by sedimentary rock, all indicate that they are not original internal structures, but rather correspond to borings (*Trypanites* isp.) made by unknown drilling organisms. They are morphologically different from the less penetrative, though superficially extensive traces already reported on the basal surface of *B. mitralis* (Kushlina 2006).

Apart from the central longitudinal canal and randomly located *Trypanites*-like borings, no other internal morphological macrostructure was apparent in sections of *B. mitralis*. Observations made with a polarizing microscope and EBSD show that each specimen of *Bolboporites* currently consists in a single crystal of calcite (see above). Although the calcite is recrystallized, the diverse observations of an extensive stereom-like microstructure in all parts of a sectioned specimen of *Bolboporites* indicates that each cone was originally a

342 microporous, but unique, monocrystal of calcite, with a central canal forming the unique
343 internal macro-cavity.
344
345 *External morphological features.*
346 Similar external morphological features were observed in all available Norwegian and
347 Russian specimens of *Bolboporites* (Figs 6B,7,8). These features are in good agreement with
348 previous descriptions of the external aspect of *Bolboporites* made on specimens from Baltica
349 (e.g., Yeltysheva 1955; Rozhnov & Kushlina 1994a; Kushlina 1995) and Laurentia (e.g.,
350 Clark & Hofmann 1961).
351 Cells on the lateral walls of the cones typically display rounded to hexagonal outlines (Fig.
352 8A) and their diameter decreases in the apical direction (Fig. 7A-C). In contrast, cells located
353 towards the base tend to become larger and more elliptical in shape. Each cell corresponds to
354 a shallow (less than 1 mm deep), gently concave, smooth depression delimited by raised rims,
355 thus forming a characteristic honeycomb pattern on the lateral walls of the cone (Figs 7A-C,
356 8A). After ultrasonic cleaning, SEM observation of Russian specimens revealed the presence
357 of the typical porous, stereomic microstructure on the external surface of the cells (Fig. 8A).
358 No specimens express an orifice at the apex of the cones, thus suggesting that their internal
359 central canal (see above) terminated close to, though not opening at, the apex.
360 The base of the cone is entirely smooth, with the exception of two adjoining circular
361 depressions (or lunules), each surrounded by a low C-shaped ridge (Figs 7D, 8B). These two
362 shallow cavities are not in central positions, but more or less displaced laterally (Fig. 7D).
363 Ultrasonic cleaning of Russian specimens shows that the floor of each depression is gently
364 concave, entirely smooth and displays a well-preserved, porous stereom microstructure (Fig.
365 8B). A small orifice occurs at the junction between the two depressions (Figs 7D, 8B). This
366 opening represents the basal extremity of the internal longitudinal canal (see above). No other
367 conspicuous external morphological structure (e.g., orifice) could be observed on the basal,
368 flat to convex surface of the cones.
369
370
371 **Discussion**
372
373 Cathodoluminescence, applied here for the first time to *Bolboporites*, confirms previous
374 reports of the presence of a former stereomic microporous structure in this fossil (e.g., Logan
375 *et al*. 1863; Lindström 1883; Yeltysheva 1955; Clark & Hofmann 1961). Currently blocked

by a diagenetic syntaxial calcite, this microstructure is also detected under SEM and PPL observations. A second important result is that each specimen of *Bolboporites* corresponds to a single, formerly porous, monocrystal of calcite. Taken together, these two points provide definitive evidence for echinoderm affinities. Consequently, overall similarities in shape or ornamentation with algae (e.g., *Coelosphaeridium*), bryozoans, corals or sponges (e.g., stromatoporoids) are merely superficial.

The initial calcite stereom is now entirely cemented and recrystallized into an orange-to-brown luminescent LMC. This low Mg content does not preclude an initial low Mg content that may have been much higher during the formation of this skeletal element, but modified (and lowered) during diagenesis. Such a diagenetic pathway is well documented for most ancient echinoderm stereoms (Gorzelak *et al.* 2016) and for biogenic HMCs in general (Bischoff *et al*. 1993). Despite this recrystallization, the extensive distribution of stereomic microstructure everywhere within the cones suggests that, originally, *Bolboporites* did not contain any internal macro-structure, with the exception of the narrow longitudinal central canal, which opens within the two lunules, on the base of the cone.

These results, combined with all available evidence obtained during this study and/or from the literature, make it possible to critically evaluate all plausible interpretations concerning the nature and systematic position of *Bolboporites*. Comparison with other Palaeozoic echinoderm cone-shaped elements suggests that *Bolboporites* can be interpreted in five different ways: (1) internal mould or body wall of a blastozoan theca (by analogy with, e.g., *Timorocidaris*; Ami 1896; Rozhnov & Kushlina 1994a, b; Kushlina 1995); (2) large infrabasal cone of the body capsule (calyx or theca) of a stemless pelmatozoan (by comparison with, e.g., *Cymbionites*); (3) isolated spine or tuberculated plate of an unknown echinoderm (Quenstedt 1881; Lindström 1883; Wanner 1920; Yeltysheva 1955; Clark & Hofmann 1961); (4) highly modified columnal of a pelmatozoan; and (5) distal holdfast of a stemmed echinoderm ('*Oryctoconus* scenario').

Bolboporites *as a theca.*

In this interpretation, *Bolboporites* represents either the internal mould (Ami 1896; von Wöhrmann *in* Jaekel 1899; Régnell 1956) or the external wall of a blastozoan theca (Rozhnov & Kushlina 1994a, b; Kushlina 1995, 2006, 2007). In blastozoans, the theca was polyplated and entirely made of extraxial skeletal elements (i.e., deriving from the pre-metamorphic larva; David *et al*. 2000; Sprinkle & Guensburg 2001; Nardin *et al*. 2009, 2017). The theca was bearing all main body orifices (anus, hydropore, mouth) and, when present, various kinds

of respiratory structures (e.g., epispires, diplopores, rhombs; Kesling 1968; Sprinkle 1973; David *et al*. 2000). In life, the theca housed the main body cavity, which contained all internal organs (e.g., gut; Kesling 1968; Sprinkle 1973; Rahman *et al*. 2015).

The interpretation of *Bolboporites* as the internal mould of a blastozoan theca relies on several arguments: (1) its overall morphology is compatible with the cone-shaped aspect of the theca of various blastozoans, such as *Rhopalocystis* Ubaghs, 1963; (2) the honeycomb ornamentation could represent the imprint of thick polygonal thecal plates on the internal mould; (3) the smooth aspect of the basal surface could result from a more finely-plated oral surface; and (4) the lunules could correspond to the imprint left by one (or two) body opening(s). The main problem with this interpretation is that, by definition, the internal mould of a blastozoan theca corresponds to the infilling of this theca by sedimentary rock. This implies that the mineralogical composition of the internal mould should be similar to that of the surrounding rock. However, our results clearly demonstrate that (1) the mineralogical composition of *Bolboporites* is very distinct from that of the surrounding sedimentary rock; and (2) this fossil corresponds to a single echinoderm plate (stereom, monocrystal of calcite). Consequently, *Bolboporites* does not represent the internal mould of a blastozoan theca.

The second interpretation considers *Bolboporites* as the theca of an eocrinoid blastozoan, preserved either as an external mould (e.g., Norwegian specimens) or as a recrystalised three-dimensional fossil (e.g., Russian material). The main argument supporting this interpretation is the observation of a partially-preserved appendage-like structure inserted onto the lunules, on the smooth basal surface of some specimens from Russia (Rozhnov & Kushlina 1994a, pl. 6; Kushlina 2007, pl. 1 fig. 3). When present, this structure occurs consistently in the same location (i.e., perfectly fitting into the lunules), and it always shows a biserial pattern, with two opposite sets of small, thick, semi-circular skeletal elements. A longitudinal groove is running on one side of the appendage, along the suture between the two series of opposite ossicles (see Rozhnov & Kushlina 1994a, pl. 6 fig. 1d). This groove is apparently leading (proximally) into the small orifice located at the junction between the two lunules.

This biserial structure was interpreted by Rozhnov & Kuslina (1994a, b) as a feeding appendage (brachiole; see also Kushlina 1995, 2006; Rozhnov 2005, 2009). This identification implies that the groove borne by brachiolar plates probably housed a single ray of the ambulacral system and that, consequently, the mouth was located at the proximal extremity of this ray. The mouth would thus correspond to the small orifice located within the lunule. In their interpretation of *Bolboporites*, Rozhnov & Kushlina (1994a, b) further suggested that: (1) the internal longitudinal canal of *Bolboporites* was probably homologous

to the axial canal (lumen) of a pelmatozoan stem, thus implying that the stem was present, but entirely encased within the theca; and (2) all organs (e.g., gut) were not located within the theca, but outside of it, on the smooth surface of the cone.

No articulated appendage was preserved in our study material. However, its interpretation as a brachiole is plausible: in echinoderms, axial (ambulacral) flooring plates typically display a comparable biserial, zigzag pattern, resulting from their appearance through ontogeny, alternatively on the left and on the right of a growing ray of the ambulacral system ('ocular plate rule'; David & Mooi 1996, 1999; Mooi & David 1997, 1998, 2008). This identification has several implications: (1) the longitudinal groove borne by the flooring plates is an ambulacral food groove; (2) the orifice located at the proximal extremity of this groove is the mouth; (3) no ambulacral cover plates are apparently present (or preserved) above this groove; and (4) *Bolboporites* possessed one single brachiole. The reduction of the number of feeding appendages is relatively common in Palaeozoic echinoderms. It was documented both in some crinoids (e.g., *Monobrachiocrinus granulatus* Wanner, 1920; Ausich *et al*. 1999) and also in various groups of vagile, epibenthic taxa, such as pleurocystitid rhombiferans (Paul 1967; Kesling 1968; Parsley 1970), solutans (Ubaghs 1981; David *et al.* 2000; Lefebvre & Lerosey-Aubril 2018) and stylophorans (Ubaghs 1968a; David *et al*. 2000; Lefebvre 2003).

Our results demonstrate that *Bolboporites* corresponds to a single, massive cone-shaped echinoderm skeletal element. This observation has two major implications: if *Bolboporites* is interpreted as the theca of an eocrinoid, then this theca (1) is entirely made of one single plate; and (2) with the exception of the narrow central canal, it does not contain any internal structure. Reduction of the number of plates forming the body capsule is a trend described in various groups of Palaeozoic echinoderms, such as in solutans (e.g., Late Ordovician belemnocystitids from North America; Parsley & Caster 1965; Caster 1968) or in stylophorans (e.g., *Jaekelocarpus* Kolata, Frest & Mapes, 1991 from the Pennsylvanian of Oklahoma; Dominguez *et al*. 2002). Particularly drastic examples of such a reduction in the number of plates can be documented in the calices of several derived, Late Palaeozoic, stemless crinoids, all characterized by convergent, similar-looking cone-shaped morphologies, as, for example, *Agassizocrinus lobatus* Springer, 1926, from the Mississipian of Kentucky (Ettensohn 1975) or *Edriocrinus sacculus* Hall, 1859, from the Lower Devonian of New York (Moore 1978; Seilacher & MacClintock 2005; Herbert & Ettensohn 2018). The stemless Permian crinoid *Timorocidaris* probably represents the most extreme case of reduction in the number of plates, with its bowl-shaped calyx possibly made of a single

skeletal element (Ubaghs 1978a; Hess 1999). Consequently, *Bolboporites* could represent a case of convergent acquisition of a single plated body capsule in blastozoans.

In all echinoderms, including the most extreme crinoid morphologies, the body capsule always contains an internal cavity housing the viscera. However, our observations show that *Bolboporites* is a massive skeletal element, without any body cavity. If *Bolboporites* was a single-plated theca, then its mouth (i.e., the small orifice located within the lunules, at the proximal extremity of the ambulacral groove) would open into the narrow, distally closed and tapering central canal. This implies that (1) this internal canal cannot be homologous to the axial canal of a stem (in pelmatozoans, the mouth never opens into the stem canal); and (2) the absence of an internal body cavity and of any anal opening both suggest that, from a functional point of view, this interpretation is not valid. The suggestion that soft parts were lying in life over the smooth basal surface of the cone, that is, outside of the theca (see e.g., Rozhnov & Kushlina 1994a, b), is incompatible with the body plan of the phylum Echinodermata. In all echinoderms, the viscera are always housed within the body capsule, independently of whether it is loosely (e.g., holothurians) or more strongly calcified (most taxa).

In summary, our results do not confirm the identification of *Bolboporites* as the theca of an eocrinoid. Although most requirements of this interpretation are plausible (e.g., the biserial pattern of the appendage is similar to that of ambulacral flooring plates; echinoderms with a single feeding appendage did exist, as well as body capsules consisting of a reduced number of elements), this hypothesis has to be rejected, because the implied anatomy would be neither functional (no internal cavity, no viscera, no anus, no hydropore) nor compatible with the echinoderm body plan (extra-thecal viscera, mouth opening into the stem axial canal).

Bolboporites *as an infrabasal cone*.

The overall morphology of *Bolboporites* is strongly reminiscent of similar-looking, cone-shaped, massive skeletal elements forming the aboral part of the body capsule in some blastozoans and crinoids. Such aboral (or infrabasal) cones can be made of several tightly sutured plates, as, for example, in *Cymbionites* and *Peridionites*, both from the Cambrian of Australia (Whitehouse 1941; Smith 1982). However, single-plated, massive infrabasal cones have been described in several Cambro-Ordovician eocrinoids (Ubaghs 1963; Clausen 2004; Allaire *et al*. 2017), as well as in some Late Palaeozoic crinoids (Ettensohn 1975, 1980; Seilacher & MacClintock 2005; Webster & Kues 2006). As in *Bolboporites*, massive aboral pelmatozoan cones (plates) also display: (1) a very wide morphological disparity within a

same assemblage (see, e.g., Ettensohn 1980; Clausen 2004); and (2) a central canal, which is tapering proximally (i.e., towards the apex of the cone) in stemless taxa (e.g., in the Pennsylvanian crinoid genus *Paragassizocrinus* Moore & Plummer, 1940; Ettensohn 1980). In contrast, in stemmed taxa, the central canal extends throughout the infrabasal cone and leads proximally into the axial canal of the stem (see, e.g., Ubaghs 1963; Clausen 2004).

However, it seems unlikely that *Bolboporites* corresponds to the infrabasal cone of a stemless pelmatozoan. The main difficulty is the smooth aspect of its basal surface: in all pelmatozoans possessing an infrabasal cone, its upper (distal) surface is divided into several concave areas (facets), separated by ridges and corresponding to the insertion of the plates (i.e., basals) belonging to the overlying circlet (Ubaghs 1963; Ettensohn 1975, 1980; Clausen 2004; Webster & Kues 2006). The absence of such facets in *Bolboporites* implies that its basal surface was not sutured to any overlying plates and, thus, that this fossil does not represent an infrabasal cone. This interpretation is further supported by the strongly convex and particularly high morphology of the basal surface in some specimens of *Bolboporites* (in particular, in *B. americanus* Billings, 1859; see Clark & Hofmann 1961), which is incompatible with the presence of a putative overlying basal circlet. Finally, the biserial appendage articulated to the basal surface of some Russian specimens of *Bolboporites* (Rozhnov & Kushlina 1994a; Kushlina 2007) clearly demonstrates that this surface was not in contact with overlying thecal (or calyx) plates.

Bolboporites *as a spine*

. Spines are highly differentiated, mobile skeletal elements articulated to the body wall of echinoderms (see, e.g., Durham *et al*. 1966; Smith 1980b). Although this character gave its name to the phylum Echinodermata (‘spiny skin’), moveable spines indeed occur only in asterozoans, echinozoans (echinoids), edrioasteroids and stylophorans. In asterozoans, echinoids and edrioasteroids, the articulation of spines to the body wall is complex, typically consisting of (1) a concave socket (acetabulum) at the base of the spine; and (2) a corresponding convex ball (mamelon) located at the summit of a tubercle (Durham *et al*. 1966; Spencer & Wright 1966; Smith 1980b; Holloway & Jell 1983; Guensburg 1988; Lebrun 1998). Attachment and mobility of spines to the body wall are achieved by muscles and/or a ligamentary catch apparatus (Smith 1980b; Lebrun 1998). Spine movements are controlled by nerves, forming either a ring around the tubercle (e.g., in echinoids; Durham *et al*. 1966; Smith 1980b; Lebrun 1998) and/or extending inside the spine itself (e.g., in some ophiuroids; Lebrun 1998). In cornute stylophorans (e.g., chauvelicystids, *Thoralicystis griffei* (Ubaghs,

1970)), spines are articulated to various parts of the body wall (proximal aulacophore, supracentral area, thecal margin) by rudimentary balls and sockets (Ubaghs 1970, 1983; Lee *et al*. 2005). In most mitrate stylophorans, the articulation of posterior spines (digital, glossal) consists of complex balls and sockets (e.g., in anomalocystitids; Ubaghs 1968a; Parsley 1991; Ruta & Bartels 1998). In some other taxa (e.g., *Balanocystites primus* (Barrande, 1872)), the posterior spine is connected to the theca by a small column of articulated plates (Lefebvre 1999).

Spines are thus distinct from spine-shaped tubercles, which are non-articulated (fixed), protruding external structures that can be produced by various elements of the body wall (e.g., anal plates, calyx or thecal plates, columnals). Tubercles are particularly widespread at phylum-scale. In blastozoans, spine-shaped expansions of thecal plates have been described in aristocystitids (e.g., *Calix sedgwicki* Rouault, 1851; *Lepidocalix pulcher* Termier & Termier, 1950; see Chauvel 1941; Makhlouf *et al*. 2017), blastoids (e.g., *Thaumatoblastus* Wanner, 1924; *Pteratoblastus* Wanner, 1924; see Beaver *et al.* 1968), eocrinoids (e.g., *Rhopalocystis havliceki* Chauvel, 1978; see Chauvel & Régnault 1986; Allaire *et al*. 2017) and, to a lesser extent, in glyptocystitids (e.g., *Schizocystis* Jaekel, 1895; Kesling 1968). Strong tubercles also occur in both cinctans (e.g., *Undatacinctus quadricornuta* (Friedrich, 1993); see Smith & Zamora 2009) and solutans (e.g., *Girvanicystis batheri* Caster, 1968; see Daley 1992). Elongate, spine-shaped elements of the body wall also occur in some edrioasteroids (Guensburg 1988), but they are particularly widespread in Palaeozoic crinoids. In this class, such a spiny ornamentation has been described on anal plates (e.g., *Stenopecrinus* Strimple, 1961; *Uperocrinus* Meek & Worthen, 1865; see Ubaghs 1978b; Ausich *et al*. 1999), on brachials (e.g., *Eirmocrinus* Strimple & Watkins, 1969; *Separocrinus* Knapp, 1969; see Moore *et al*. 1978; Ausich *et al.* 1999), on tegmental plates (e.g., *Batocrinus* Casseday, 1854; see Ubaghs 1978b), on calyx plates (e.g., *Calceolispongia* Etheridge, 1915; *Dorycrinus* Roemer, 1854; see Moore *et al*. 1978; Ubaghs 1978b; Brett 1999), and to a lesser extent, on stem elements (e.g., *Aethocrinus moorei* Ubaghs, 1969). Elongate spine-shaped tubercles occur on the proximal brachials of some cornute (e.g., *Reticulocarpos hanusi* Jefferies & Prokop, 1972; *Nanocarpus milnerorum* Ruta, 1999; Jefferies, 1986; Lefebvre 2003) and most mitrate stylophorans (e.g., *Chinianocarpos thorali* Ubaghs, 1961; *Rhenocystis latipedunculata* Dehm, 1932; Ubaghs 1970; Jefferies 1986; Ruta & Bartels 1998; Lefebvre 2003). Strong spine-shaped skeletal elements are also present on the aboral surface of some Recent oreasterid asteroids (e.g., *Pentaceraster mammilatus*, *Protoreaster nodosus* (Linnaeus, 1758); see Yeltysheva 1955).

579 The cone-shaped morphology of *Bolboporites* is compatible with its interpretation either as
580 an isolated spine or a tuberculated plate of an echinoderm (Quenstedt 1881; Lindström 1883;
581 Wanner 1920; Yeltysheva 1955; Régnell 1956; Clark & Hoffman 1961). Both interpretations
582 imply that (1) the smooth basal surface was facing towards the organism and was either
583 articulated to it (spine) or part of its body wall (tubercle); and (2) the lateral, strongly
584 ornamented walls of the cone were external and directed away from the organism. These two
585 interpretations could explain the wide disparity in size and shape observed in individuals of
586 *Bolboporites* from a same level (Yeltysheva 1955; Clark & Hoffman 1961). Both
587 interpretations would be also in good agreement with the non-random distribution of biofilms
588 produced by encrusting organisms on Russian specimens (Kushlina 2007). When present,
589 thick putative algal-bacterial biofilms are consistently encrusting the lateral sides of
590 *Bolboporites*. They never occur on the smooth, convex surface (Kushlina 2007). This pattern
591 suggests that the biofilms formed when the organism was alive, otherwise biofilms would be
592 present on all surfaces. The distribution of these encrusting biofilms also suggests that the
593 lateral sides of the cones were directly in contact with sea water (i.e., lateral sides were
594 neither buried in the substrate, nor in contact with the body wall). Conversely, the absence of
595 biofilms on the smooth surface supports the view that the basal part of the cones was not
596 exposed to the external medium and, thus, probably in contact with or part of the body wall.
597 As pointed out by Yeltysheva (1955), the overall morphology of *Bolboporites* shows
598 several similarities with the cone-shaped aboral elements of some oreasterid asteroids. For
599 example, the observation and dissection of Recent specimens of *Pentaceraster mammilatus*
600 (Fig. 9D,E) showed that a honeycomb pattern is present on the lateral walls of their cone-
601 shaped abaxial tubercles (Fig. 9A). This sculpture, which is similar to that observed in
602 *Bolboporites*, was produced by the thick granulose membrane, which forms the aboral part of
603 the body wall and extends over the cones. However, the dissection of Recent specimens of *P.*
604 *mammilatus* also showed some major morphological differences between *Bolboporites* and
605 oreasterid cone-shaped abaxial elements. The most important one is that the basal surface of
606 oreasterid cones is not smooth and gently convex, but subdivided into several diverging
607 branches (Fig. 9B), connecting the cone with surrounding plates (Fig. 9C). This situation is
608 not unique to oreasterid abaxial cones. All echinoderm spine-shaped tubercles are borne by
609 elements, which are part of the body wall. This implies that tuberculated plates are necessarily
610 in contact with neighbouring skeletal elements and thus always display facets along their
611 sutures. As the basal surface of *Bolboporites* is entirely smooth and does not show any

612 evidence of facets, its interpretation as a putative oreasterid-like cone-shaped element has to
613 be rejected.
614 Finally, although the interpretation of *Bolboporites* as a spine is plausible (cone-shaped
615 morphology, smooth surface in contact with the body wall, wide morphological disparity,
616 encrusting organisms restricted to lateral walls), this hypothesis has implications that can be
617 tested: (1) skeletal evidence supporting an articulation should be present; and (2) soft parts
618 (muscles, ligaments and/or nerves) were very likely involved, too, and should have left some
619 traces. Clearly, the basal surface of *Bolboporites* is entirely smooth and does not show any
620 skeletal evidence suggesting the presence of a socket for articulation onto a putative tubercle
621 on the body wall. Consequently, if *Bolboporites* was a spine, its articulation was different
622 from the most widespread mechanism (balls and sockets) occurring in echinoderms (see
623 above). The only structures occurring on the smooth surface are the two lunules. However,
624 these two concave areas were apparently the place of insertion for a biserial appendage (see
625 discussion above; Rozhnov & Kushlina 1994a; Kushlina 2007). Consequently, if *Bolboporites*
626 was a spine, the small biserial appendage would then have been directed towards the body
627 wall. In the mitrate stylophoran *Balanocystites primus*, a similar-looking short appendage,
628 made of a single column of tiny plates connects the single posterior spine (glossal) to the
629 theca (Lefebvre 1999). By comparison with the situation in *B. primus*, the possibility that
630 *Bolboporites* was a spine, which was connected to the body wall by a short, biserial,
631 articulated appendage cannot be ruled out. If this interpretation is correct, it is thus likely that
632 the smooth aspect of the basal surface of *Bolboporites* is related to the insertion of soft parts
633 (probably muscles or ligaments) on it. Moreover, by analogy with the situation in some
634 ophiuroids (see above; Lebrun 1998), it is then possible to interpret the external longitudinal
635 groove running on the small appendage and extending internally into the central canal of the
636 cone as the probable course of a nerve.
637 The interpretation of *Bolboporites* as a spine is thus plausible and cannot be refuted on
638 available evidence. However, its systematic position remains an open question. The frequent
639 association of *Bolboporites* with skeletal remains of various blastozoans (e.g., cheirocrinids,
640 *Palaeocystites*) questioned the possibility that it could represent isolated spines of one of them
641 (Clark & Hoffman 1961). However, none of the blastozoans found in the same localities as
642 *Bolboporites* in both Baltica and Laurentia shows any evidence suggesting that spines were
643 articulated to its body wall. Indeed, spines are only known in asterozoans, echinozoans,
644 edrioasteroids and stylophorans (see above), thus ruling out any putative blastozoan affinities
645 for *Bolboporites*. The size of *Bolboporites* (from about 2 to 12 mm in height in both North

American and Russian specimens; Clark & Hoffman 1961; Rozhnov & Kushlina 1994a) suggests that putative stylophoran affinities are highly unlikely. In this class, the size of the theca is generally comprised between 5 and 30 mm, and it rarely exceeds 30 to 40 mm (e.g., the largest known mitrate, *Diamphidiocystis drepanon* Kolata & Guensburg, 1979, is about 40 mm wide; Lefebvre 1999). The large size and cone-shaped morphology of *Bolboporites* seem to be also incompatible with putative edrioasteroid affinities. When preserved, edrioasteroid spines are consistently consisting of narrow, elongate elements, typically less than 3 mm in length, which were articulated to ambulacral cover plates and/or skeletal elements of the pedunculate zone (Holloway & Jell 1983; Guensburg 1988; Guensburg & Sprinkle 1994).

Because of their stratigraphic range and palaeobiogeographic distribution in the Ordovician, asterozoans are more likely candidates: their oldest known representatives have been documented in Lower Ordovician deposits (Thoral 1935; Blake 2013; Jell 2014; Blake & Guensburg 2015), and their presence is recorded in Baltica at least from the Dapingian, i.e., as early as the oldest known occurrence of *Bolboporites* (Pisera 1994; Hansen *et al*. 2005; Rozhnov 2005; Blake & Rozhnov 2007; Tinn & Ainsaar 2014). However, *Bolboporites* clearly does not show any character supporting its interpretation as an isolated asterozoan spine. The skeletal morphology of Ordovician asterozoans is strongly constrained and, even when preserved as isolated remains, their plates are highly diagnostic (Pisera 1994; Tinn & Ainsaar 2014).

Echinozoans possibly appeared and diversified in Baltica during the Middle Ordovician, before spreading to Laurentia in Late Ordovician times (Reich 1999; Smith & Savill 2001; Lefebvre *et al*. 2013). Baltica has yielded the oldest known occurrences of echinoids (Darriwilian; Bockelie & Briskeby 1980; Pisera 1994), holothurians (Darriwilian; Reich 2010) and ophiocistioids (Dapingian; Reich 2001; Rozhnov 2005; Reich & Smith 2009). The stratigraphic range and palaeobiogeographic distribution of echinozoans are thus compatible with those of *Bolboporites*. However, the presence of typical tubercles in the oldest known echinoids (including the isolated plates record; Pisera 1994) suggests the existence of mechanisms for spine articulation comparable to those occurring in younger taxa (i.e., balls and sockets). Putative echinoid affinities are thus unlikely for *Bolboporites*. Although spines have not been documented so far in holothurians and ophiocistioids, it cannot be entirely excluded that *Bolboporites* corresponds to isolated spines of a yet unknown primitive echinozoan.

Bolboporites as a columnal.

The general aspect of *Bolboporites* reminds in some respects the cone-shaped morphology of some late Cambrian–Early Ordovician pelmatozoan columnals from Utah (Sumrall *et al.* 1997) and Spain (Zamora *et al*. 2009), and thus questions its possible identification as a highly differentiated stem plate. This interpretation is in good agreement with (1) the existence of a longitudinal internal canal; (2) the articulation of a biserial appendage on its basal surface (Rozhnov & Kuslina 1994a; Kushlina 2007); and (3) the morphology of the lunules. The biconcave depressed area formed by the lunules, as well as the presence of a tiny orifice opening in between them are morphological features which are reminiscent of synarthrial articulations in crinoids (Ubaghs 1978a; Donovan 1988; Ausich *et al*. 1999). Interestingly, synarthrial articulations were present on the distal columnals of some Ordovician crinoids (e.g., *Ristnacrinus* Öpik, 1934) co-occurring with *Bolboporites* in Baltica (Donovan 1984).

It seems, however, difficult to interpret *Bolboporites* as a highly differentiated, massive columnal for several reasons. First, in all pelmatozoan echinoderms, columnals display articulatory facets on their two opposite (proximal and distal) sides, whereas *Bolboporites* would display only one facet (i.e., on its basal surface). A second difficulty is that columnals always display a central canal (lumen), which opens on their two opposite sides: such a canal is present in *Bolboporites*, but it is not in central position and, more importantly, it opens only on one side. Finally, the strongly convex morphology of the basal surface in many North American specimens of *Bolboporites* (see Clark & Hofmann 1961) makes their interpretation as columnals highly improbable.

Bolboporites *as a holdfast.*

Holdfasts are anchoring structures occurring at the distalmost extremity of the stem in various blastozoans and crinoids (Ubaghs 1972, 1978a; Brett 1981; Ausich *et al*. 1999; Seilacher & MacClintock 2005). The massive, cone-shaped morphology of *Bolboporites* shows many similarities with similarly-shaped, isolated pelmatozoan elements (e.g., *Oryctoconus*), generally interpreted as holdfasts (Colchen & Ubaghs 1969; Alvaro & Colchen 2002; Seilacher & MacClintock 2005; Zamora *et al*. 2009). The identification of *Bolboporites* as a putative discoidal holdfast was discussed, but rejected by Rozhnov & Kushlina (1994a). Their main argument was that, if this fossil was a holdfast, its basal surface would then be attached (fixed) to the substrate: this orientation is incompatible with the presence of an appendage articulated to the basal surface of *Bolboporites*. However, it should be stressed that this base-down orientation occurs only in the case of pelmatozoan discoidal holdfasts tightly

714 and permanently encrusted on firmgrounds and hardgrounds (Ubaghs 1978a; Brett 1981;
715 Brett *et al.* 1983; Sumrall *et al*. 1997; Rozhnov 2002). The opposite (base-up) orientation of
716 the cone is observed in most pelmatozoans living on soft substrates and using their distal
717 holdfasts as an anchor or a grapnel, as, for example, the Ordovician eocrinoid *Balantiocystis*
718 Chauvel, 1966, and the Devonian crinoid *Ancyrocrinus* Hall, 1862 ('kite strategy'; Ubaghs
719 1972; Brett 1981; Le Menn 1985; Ausich *et al*. 1999; Alvaro & Colchen 2002; Seilacher &
720 MacClintock 2005; Zamora *et al*. 2009).
721 All above-listed arguments agreeing with the interpretation of *Bolboporites* as a columnal
722 remain valid if this fossil is interpreted as a discoidal terminal holdfast (i.e., internal canal;
723 biserial appendage inserted into the basal surface; lunules forming a facet with a synarthrial-
724 like articulation). However, if *Bolboporites* is a holdfast, the various issues raised for its
725 interpretation as a columnal are no longer problematic: it then makes sense that (1) a single
726 facet is present (on the basal surface); (2) the internal canal does not open distally into the
727 apex of the holdfast; and (3) the inflated morphology of the basal surface in some specimens
728 of *B. americanus* is not incompatible with their interpretation as distal holdfasts. Moreover,
729 the wide morphological disparity observed between specimens of *Bolboporites* from a same
730 level (Clark & Hofmann 1961; Kushlina 1995) is also in good agreement with its
731 interpretation as a holdfast: similar large variabilities in shape have been reported in
732 assemblages of, for example, *Oryctoconus* and grapnel-like holdfasts of *Ancyrocrinus* (Le
733 Menn 1985; Alvaro & Colchen 2002; Zamora *et al*. 2009). Finally, this interpretation is also
734 compatible with the occurrence of *Bolboporites* in deposits corresponding to shallow, storm-
735 generated deposits (see above; Clark & Hofmann 1961; Bockelie 1981; Dronov 2005). In
736 such environmental conditions, stemmed echinoderms and their anchoring structures are
737 generally preserved separately (Brett 1981). Organisms were detached from their anchoring
738 structures probably by autotomy rather than breakages and transported away by storm
739 currents, whereas their holdfasts were preserved in situ (Donovan 2012). Apart from some
740 rare exceptions, such as the eocrinoid *Balantiocystis* or the crinoid *Ancyrocrinus* (Ubaghs
741 1972; Le Menn 1985; Ausich *et al.* 1999), the distalmost part of the stem is unknown in most
742 pelmatozoans and, conversely, most holdfasts cannot been assigned to any specific taxa (e.g.,
743 *Aspidocrinus scutelliformis* Hall, 1859, *Oryctoconus*; Ubaghs 1978a; Brett *et al*. 1983;
744 Sumrall *et al*. 1997; Alvaro & Colchen 2002; Seilacher & MacClintock 2005; Zamora *et al*.
745 2009).
746 The interpretation of *Bolboporites* as a discoidal holdfast has also several implications, that
747 can be tested: (1) the biserial appendage inserting on the lunules would thus probably

correspond to the distal-most columnals of a pelmatozoan stem; and (2) if *Bolboporites* was used as an anchor, it was thus at least partly buried into the sediment. If *Bolboporites* was a distal holdfast, the presence of two lunules in all specimens suggests that a dimeric stem was articulated to it. This interpretation is further supported by the observation of a biserial appendage in at least some better preserved individuals from Russia (Rozhnov & Kushlina 1994a; Kushlina 2007). Although most Ordovician pelmatozoans possessed holomeric stems (i.e., formed by a single column of plates), tetra-, penta- and hexameric appendages have been also described in several crinoids (e.g., *Aethocrinus moorei*, *Ramseyocrinus* Bates, 1968; Ubaghs 1969, 1983; Donovan 1984, 1985), as well as in some echinosphaeritid and hemicosmitid blastozoans (Jaekel 1899; Bockelie 1981, 1982; Parsley 1998). Dimeric distal stems are the rule in Ordovician solutans (Caster 1968; Ubaghs 1970; Lefebvre *et al*. 2012; Noailles *et al*. 2014). Although they possibly retained an attached post-metamorphic stage, Ordovician solutans were vagile and their stem did not possess any distal discoidal holdfast. With the exception of solutan elements, only few occurrences of tri- and dimeric columnals were documented in Ordovician deposits, and all of them have been assigned to crinoids possibly related to *Ectenocrinus* Miller, 1889 (e.g., Donovan 1985). Consequently, the existence of Ordovician pelmatozoans with a stem comprising dimeric columnals supports the identification of *Bolboporites* as a possible distal holdfast articulated to a biserial appendage.

However, contrary to the situation in all echinoderm stem-like appendages, the biserial structure articulated to the basal surface of *Bolboporites* does not contain any lumen (internal central canal), but an external groove (Rozhnov & Kushlina 1994a). This external groove, which probably housed soft parts, communicates with the longitudinal internal canal of *Bolboporites*. In all stemmed echinoderms, the lumen contains coeloms associated with the extraxial part of the body wall (i.e., somatocoels) and, generally, extensions of the nervous system (Ubaghs 1978a; Heinzeller & Welsh 1994; David *et al*. 2000; Mooi & David 2008). The topology observed in *Bolboporites* and its associated appendage thus strongly departs from the situation in pelmatozoan stems (external *vs*. internal soft parts). This implies that the biserial structure articulated to *Bolboporites* cannot be interpreted as (part of) a stem-like appendage and, consequently, that *Bolboporites* was not a distal holdfast. This conclusion is confirmed by the distribution of biofilms produced by encrusting organisms over the body wall of *Bolboporites* (see above; Kushlina 2007). If this fossil was a discoidal terminal holdfast, comparison with similar structures in pelmatozoans (Ausich *et al*. 1999; Seilacher & MacClintock 2005) suggests that in life, a large part of the cone would have been at least partly buried into and/or in permanent contact with the substrate. This life orientation is not

compatible with the observed distribution of epibionts, which produced extensive films on the lateral walls of *Bolboporites*, but are absent from its basal surface (Kushlina 2007).

Consequently, although the interpretation of *Bolboporites* as a discoidal distal holdfast is plausible (e.g., massive cone-shaped morphology, wide morphological disparity, articulation to a dimeric appendage; see above), this identification has to be rejected because the structure articulated to its basal surface is not a stem-like appendage (no lumen, external groove housing soft parts). Further, the implied life orientation is not confirmed by the distribution of epibionts on the cones.

Conclusions

Our results not only confirmed the presence of stereomic microstructure in *Bolboporites* (and thus its echinoderm affinities), but they also showed that this fossil is a single, previously microporous, calcitic skeletal element, without any internal macrostructure, except a narrow longitudinal canal opening through a tiny orifice on the basal surface. These results combined with all previous descriptions of *Bolboporites* have made it possible to critically discuss several hypotheses about its nature (e.g., theca, basal cone, spine, columnal, holdfast) and its putative affinities within echinoderms (e.g., asterozoans, blastozoans, crinoids, echinozoans, stylophorans). Most interpretations could be rejected, because they comply with only part of available evidences. Although the identification of *Bolboporites* as a spine remains questionable, it represents the most parsimonious - and likely - interpretation. The precise affinities of *Bolboporites* remain difficult to assess and it is tentatively assigned here to an unknown, possibly basal echinozoan. As this was the case for other problematic fossils (e.g., conodonts, machaeridians), future discoveries of fully articulated specimens showing *Bolboporites* elements in connection with their host organism will probably help in revealing their actual nature and affinities within echinoderms.

Acknowledgments

This paper is a contribution to the International Geoscience Programme (IGCP) Project 653 – The onset of the Great Ordovician Biodiversification Event, and of the team 'Biosignatures, Vie Primitive' of UMR CNRS 5276 LGLTPE. The paper is also a contribution to the revision

of volumes S and U of the *Treatise on Invertebrate Paleontology*. The authors are particularly grateful to Sergei V. Rozhnov (Palaeontological Institute of the Russian Academy of Sciences, Moscow) for providing numerous, well-preserved specimens of *Bolboporites* sp. for this study, and for insightful and constructive discussions on their systematic affinities. Hans-Arne Nakrem and Franz-Josef Lindemann (Natural History Museum, Oslo) are thanked for access to the Norwegian material of *Bolboporites* and *Coelosphaeridium*, and Michel Creuzé des Châtelliers and Blandine Bärtschi (Lyon 1 University, Villeurbanne) for access and dissection of specimens of the Recent spiny starfish *Pentaceraster mammilatus*. Jih-Pai Lin (National Taïwan University) is also acknowledged for having made available specimens of *Timorocidaris* for comparison. Frédéric Herbst (UMR6303 ICB, Burgundy University, Dijon) acquired the EBSD data. This study also benefited from useful comments from and/or discussions with Daniel B. Blake (University of Illinois, Champaign), Gilles Cuny and Vincent Perrier (Lyon 1 University, Villeurbanne), Georgy Mirantsev (Palaeontological Institute of the Russian Academy of Sciences, Moscow), Elise Nardin (Observatoire Midi-Pyrénées, Toulouse), Colin D. Sumrall (University of Tennessee, Knoxville), and Gary D. Webster (Washington State University, Pullman). Finally, Thomas E. Guensburg (Field Museum, Chicago) and Stephen K. Donovan (Naturalis, Leiden) are greatly thanked for their constructive and helpful reviews.

References

Allaire, N., Lefebvre, B., Nardin, E., Martin, E.L.O., Vaucher, R. & Escarguel, G. (2017) Morphological disparity analysis and systematic revision of the eocrinoid genus *Rhopalocystis* (Echinodermata, Blastozoa) from the Lower Ordovician of the central Anti-Atlas (Morocco). *Journal of Paleontology*, 91, 685–714.

Alvaro, J.J. & Colchen, M. (2002) Earliest Ordovician pelmatozoan holdfasts from western Europe: the *Oryctoconus* problem revisited. *Eclogae Geologicae Helvetiae*, 95, 451–459.

Ami, H.M. (1896) Preliminary lists of the organic remains occurring in the various geological formations comprised in the southwest quartersheet map of the eastern townships of the Province of Quebec. *Geological Survey of Canada, Annual Report*, 7, 113–157.

Audouin, V. (1826) Explication des planches d'échinodermes de l'Egypte et de la Syrie publiées par Jules-César Savigny, membre de l'Institut. In: *Description de l'Egypte ou Recueil des Observations et des Recherches qui ont été faites pendant l'Expédition de l'Armée Française*. Imprimerie impériale (1809), Paris, pp. 203-212

Ausich, W.I., Brett, C.E., Hess, H. & Simms, M.J. (1999) Crinoid form and function. *In*: Hess, H., Ausich, W.I., Brett, C.E. & Simms, M.J. (Eds.), *Fossil Crinoids*. Cambridge University Press, Cambridge, pp. 3–30.

Baarli, B.G. (2008) *Fossilboka*. Forlaget Vett & Viten AS, Nesbru, 367 pp.
Barrande, J. (1872) *Système Silurien du Centre de la Bohême. Supplément au Volume I. Trilobites, Crustacés divers et Poissons*. Bellman, Prague, 647 pp.
Bassler, R.S. (1911) The Early Paleozoic Bryozoa of the Baltic provinces. *United States National Museum Bulletin*, 77, 1–49.
Bassler, R.S. (1915) Bibliographic index of American Ordovician and Silurian fossils. Volume 1. *United States National Museum Bulletin*, 92, 1–718.
Bates, D.E.B. (1968) On '*Dendrocrinus' cambriensis* Hicks, the earliest known crinoid. *Palaeontology*, 11, 406–409.
Bather, F.A. (1920) Echinoid or crinoid? *Geological Magazine*, 57, 371–372.
Beaver, H.H., Fay, R.O., Macurda, D.B., Moore, R.C. & Wanner, J. (1968) Blastoids. *In*: Moore, R.C (Ed.), *Treatise on Invertebrate Paleontology, Echinodermata 1(2)*. Geological Society of America, Boulder & University of Kansas Press, Lawrence, pp. S297–S455.
Berg-Madsen, V. (1986) Middle Cambrian cystoid (*sensu lato*) columnals from Bornholm, Denmark. *Lethaia*, 19, 67–80.
Billings, E. (1859) The fossils of the Chazy Limestone with descriptions of new species. *The Canadian Naturalist and Geologist*, 4, 426–470.
Bischoff, W.D., Mackenzie, F.T., Bishop, F.C. (1993) Diagenetic stabilization pathways of magnesian calcites. *Carbonates & Evaporites*, 8, 82–89.
Blake, D.B. (2013) Early asterozoan (Echinodermata) diversification: a paleontologic quandary. *Journal of Paleontology*, 87, 353–372.
Blake, D.B. & Guensburg, T.E. (2015) The class Somasteroidea (Echinodermata, Asterozoa): morphology and occurrence. *Journal of Paleontology*, 89, 465–486.
Blake, D.B. & Rozhnov, S.V. (2007) Aspects of life mode among Ordovician asteroids: implications of new specimens from Baltica. *Acta Palaeontologica Polonica*, 52, 519–533.
Bockelie, J.F. (1981) The Middle Ordovician of the Oslo region, Norway, 30. The eocrinoid genera *Cryptocrinites*, *Rhipidocystis* and *Bockia*. *Norsk Geologisk Tidsskrift*, 61, 123–147.
Bockelie, J.F. (1982) Morphology, growth and taxonomy of the Ordovician rhombiferan *Caryocystites*. *Geologiska Förenings I Stockholm Förandlingar*, 103, 491–498.
Bockelie, J.F. & Briskeby, P.I. (1980) The presence of a bothriocidarid (Echinoid) in the Ordovician of Norway. *Norsk Geologisk Tidsskrift*, 60, 89–91.
Brainerd, E. (1891) The Chazy Formation in the Champlain Valley. *Bulletin of the Geological Society of America*, 2, 293–300.
Brainerd, E. & Seely, H.M. (1888) The original Chazy rocks. *American Geologist*, 2, 323–330.
Brainerd, E. & Seely, H.M. (1896) The Chazy of Lake Champlain. *American Museum of Natural History Bulletin*, 8, 305–315.
Brett, C.E. (1981) Terminology and functional morphology of attachment structures in pelmatozoan echinoderms. *Lethaia*, 14, 343–370.
Brett, C.E. (1999) Middle Devonian Windom Shale of Vincent, New York, USA. *In*: Hess, H., Ausich, W.I., Brett, C.E. & Simms, M.J. (Eds.), *Fossil Crinoids*. Cambridge University Press, Cambridge, pp. 122–128.
Brett, C.E., Liddell, W.D. & Derstler, K.L. (1983) Late Cambrian hard substrate communities from Montana/Wyoming: the oldest known hardground encrusters. *Lethaia*, 16, 281–289.
Brett, C.E., Moffat, H.A. & Taylor, W.L. (1997) Echinoderm taphonomy, taphofacies, and Lagerstätten. *Paleontological Society Papers*, 3, 147–190.
Briggs, D.E.G., Siveter, D.J., Siveter, D.J., Sutton, M.F. & Rahman, I.A. (2017) An edrioasteroid from the Silurian Herefordshire Lagerstätte of England reveals the nature of the water vascular system in an extinct echinoderm. *Proceedings of the Royal Society B*, 284, 20171189.
Bronn, H.G. (1849) *Index Palaeontologicus oder Übersicht der bis jetzt bekannten fossilen Organismen*. E. Schweizerbart'sche Verlagshandlung und Druckerei, Stuttgart, 1381 pp.

Bronn, H.G. (1851–1856) Systematische Übersicht der fossilen Pflanzen und Thiere nach ihrer geologischen Verbreitung. Schlüssel-Tabellen oder Claves einzelner Klassen. Alphabetisches Register. *In*: Bronn, H.G. & Roemer, F. (Eds.), *Lethaea Geognostica oder Abbildung und Beschreibung der für die Gebirgs-Formationen Bezeichnendsten Versteinerungen*, Volume 1. Schweizerbart'sche Verlagsbuchhandlung und Druckerei, Stuttgart, pp. 1–204.

Butts, C. (1940) Geology of the Appalachian Valley in Virginia, part 1. Geologic text and illustrations. *Virginia Geological Survey Bulletin*, 52, 1–568.

Casseday, S.A. (1854) Beschreibung eines neuen Crinoideengeschlechts Nordamerika. *Zeitschrift der Deutschen Geologischen Gesellschaft*, 6, 237–242.

Caster, K.E. (1968) Homoiostelea. *In*: Moore, R.C (Ed.), *Treatise on Invertebrate Paleontology, Echinodermata 1(2)*. Geological Society of America, Boulder & University of Kansas Press, Lawrence, pp. S581–S627.

Chauvel, J. (1941) Recherches sur les cystoïdes et les carpoïdes armoricains. *Mémoires de la Société géologique et minéralogique de Bretagne*, 5, 1–286.

Chauvel, J. (1966) *Echinodermes de l'Ordovicien du Maroc*. Editions du CNRS, Paris, 120 pp.

Chauvel, J. (1978) Compléments sur les échinodermes du Paléozoïque marocain (diploporites, éocrinoïdes, édrioastéroïdés). *Notes du Service géologique du Maroc*, 39, 27–78.

Chauvel, J. & Régnault, (1986) Variabilité du genre *Rhopalocystis* Ubaghs, éocrinoïde du Trémadocien de l'Anti-Atlas marocain. *Geobios*, 19, 863–870.

Checa, A.G., Esteban-Delgado, F.J., Ramirez-Rico, J., Rodriguez-Navarro, A.B. (2009) Crystallographic reorganization of the calcitic prismatic layer of oysters. *Journal of Structural Biology*, 167, 261–270.

Clark, A.M. & Rowe, F.W.E. (1971) *Monograph of Shallow-Water Indo-West Pacific Echinoderms*. The Natural History Museum, London, 238 pp.

Clark, T.H. (1944) Unfolded Palaeozoic rocks of the St. Lawrence Lowlands. *In*: Dresser, J.A. & Denis, T.C. (Eds.), *The Geology of Quebec, Volume 2. Descriptive Geology*. Quebec Department of Mines, Geological Report, 20, pp. 250–291.

Clark, T.H. (1952) Montreal area, Laval and Lachine map areas. *Quebec Department of Mines, Geological Report*, 46, 1–159.

Clark, T.H. & Hofmann, H.J. (1961) *Bolboporites americanus* in the Chazy of Southern Quebec. *Transactions of the Royal Society of Canada*, 55, 13–28.

Clausen, S. (2004) New Early Cambrian eocrinoids from the Iberian Chains (NE Spain) and their role in nonreefal benthic communities. *Eclogae Geologicae Helvetiae*, 97, 371–379.

Colchen, M. & Ubaghs, G. (1969) Sur des restes d'échinodermes(?) du Cambro-Ordovicien de la Sierra de la Demanda (Burgos-Logrono, Espagne). *Bulletin de la Société géologique de France*, 11, 649–654.

Cusack, M (2016) Biomineral electron backscatter diffraction for palaeontology. *Palaeontology*, 59, 171–179.

Daley, P.E.J. (1992) The anatomy of the solute *Girvanicystis batheri* (?Chordata) from the Upper Ordovician of Scotland and a new species of *Girvanicystis* from the Upper Ordovician of south Wales. *Zoological Journal of the Linnean Society*, 105, 353–375.

David, B. & Mooi, R. (1996) Embryology supports a new theory of skeletal homologies for the phylum Echinodermata. *Comptes-Rendus de l'Académie des Sciences, Paris*, *Sciences de la vie/Life sciences*, 319, 577–584.

David, B. & Mooi, R. (1999) Comprendre les échinodermes: la contribution du modèle extraxial-axial. *Bulletin de la Société géologique de France*, 170, 91–101.

David, B., Lefebvre, B., Mooi, R. & Parsley, R. (2000) Are homalozoans echinoderms? An answer from the extraxial-axial theory. *Paleobiology*, 26, 529–555.

Dehm, R. (1932) Cystoideen aus dem rheinischen Unterdevon. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Beilige-Band, Abteilung A,* 69, 63–93.

Dominguez, P., Jacobson, A.G. & Jefferies, R.P.S. (2002) Paired gill slits in a fossil with a calcite skeleton. *Nature*, 417, 841–844.

954 Donovan, S.K. (1984) *Ramseyocrinus* and *Ristnacrinus* from the Ordovician of Britain. *Palaeontology*, 27,
955 623–634.
956 Donovan, S.K. (1985) Biostratigraphy and evolution of crinoid columnals from the Ordovician of Britain.
957 *In*: Keegan, B.F. & O'Connor, B.D.S. (Eds.), *Echinodermata: Proceedings of the Fifth International*
958 *Echinoderm Conference, Galway, 24-29 September, 1984*. Balkema, Rotterdam, pp. 19–24.
959 Donovan, S.K. (1988) Functional morphology of synarthrial articulations in the crinoid stem. *Lethaia*, 21,
960 169–175.
961 Donovan, S.K. (1991) The taphonomy of echinoderms: calcareous multi-element skeletons in the marine
962 environment. *In*: Donovan, S.K. (Ed.), *The Processes of Fossilization*. Belhaven Press, London, pp.
963 241–269.
964 Donovan, S.K. (2012) Was autotomy a pervasive adaptation of the crinoid stalk during the Paleozoic?
965 *Geology*, 40, 867–870.
966 Donovan, S.K. (2018) The internal morphology of primary spines of extant regular echinoids in the tropical
967 western Atlantic: a SEM atlas. *Swiss Journal of Palaeontology*, 137, 363–377.
968 Dronov, A. (2005) Introduction to the geology of the St. Petersburg region. *In*: Dronov, A., Tolmacheva,
969 T., Raevskaya, E. & Nestell, M. (Eds.), *Cambrian and Ordovician of St. Petersburg area. Guidebook of*
970 *the pre-conference field trip*. St. Petersburg State University, Saint-Petersburg, pp. 2–15.
971 Durham, J.W, Fell, H.B., Fischer, A.G., Kier, P.M., Melville, R.V., Pawson, D.L. & Wagner, C.D. (1966)
972 Echinoids. *In*: Moore, R.C (Ed.), *Treatise on Invertebrate Paleontology, Echinodermata 3(1–2).*
973 Geological Society of America, Boulder & University of Kansas Press, Lawrence, pp. U211–U640.
974 Eichwald, E. von (1857) *Beitrag zur geographischen Verbreitung der fossilen Thiere Russlands*. *Alte*
975 *Period*. Buchdrückerei der Kaiserlichen Universität, Moscow, 242 pp.
976 Eichwald, E. von (1860) *Lethaea Rossica ou Paléontologie de la Russie. Premier Volume. Première*
977 *Section de l'Ancienne Période*. Schweitzerbart, Stuttgart, 1657 pp.
978 Etheridge, R. (1915) Western Australian Carboniferous fossils, chiefly from Mount Marmion, Lennard
979 River, West Kimberley. *Bulletin - Geological Survey of Western Australia*, 58, 7-49.
980 Ettensohn, F.R. (1975) The autecology of *Agassizocrinus lobatus*. *Journal of Paleontology*, 49, 1044–1061.
981 Ettensohn, F.R. (1980) *Paragassizocrinus*: systematics, phylogeny and ecology. *Journal of Paleontology*,
982 54, 978–1007.
983 Federov, P. (2003) Lower Ordovician mud mounds from the St. Petersburg region, northwestern Russia.
984 *Bulletin of the Geological Society of Denmark*, 50, 125–137.
985 Fischer von Waldheim, G. (1829) Prodromus Petromatognosiae Animalium Systematicae Continens,
986 Bibliographiam Animalium Fossilium. *Nouveaux Mémoires de la Société Impériale des Naturalistes de*
987 *Moscou*, 1, 301–374.
988 Friedrich, W.P. (1993) Systematik und Funktionsmorphologie mittelkambrischer Cincta (Carpoidea,
989 Echinodermata). *Beringeria*, 7, 1–190.
990 Fromentel, E. de (1861) Introduction à l'étude des Polypiers fossiles. *Mémoires de la Société d'Emulation*
991 *du Département du Doubs*, 5, 1–357.
992 Génot, P. & Granier, B. (2011) Cenozoic Dasycladales. A photo-atlas of Thanetian, Ypresian and
993 Bartonian species from the Paris Basin. *Carnets de Géologie*, *Special Paper*, 2011/1, 1–44.
994 Gislén, T. (1947) On the Haplozoa and the interpretation of *Peridionites*. *Zoologiska Bidrag från Uppsala*,
995 25, 402–408.
996 Gorzelak, P. & Zamora, S. (2013) Stereom microstructures of Cambrian echinoderms revealed by
997 cathodoluminescence (CL). *Palaeontologia Electronica*, 16.3.32A, 1–17.
998 Gorzelak, P., Krzykawski, T., Stolarski, J. (2016) Diagenesis of echinoderm skeletons: constraints on
999 paleoseawater Mg/Ca reconstructions. *Global and Planetary Change*, 144, 142–157.
1000 Guensburg, T.E. (1988) Systematics, functional morphology, and life modes of Late Ordovician
1001 edrioasteroids, Orchard Creek Shale, southern Illinois. *Journal of Paleontology*, 62, 110–126.

1002 Guensburg, T.E. & Sprinkle, J. (1994) Revised phylogeny and functional interpretation of the
1003 Edrioasteroidea based on new taxa from the Early and Middle Ordovician of western Utah. *Fieldiana*,
1004 *Geology*, 29, 1–43.
1005 Hall, J. (1847) *Paleontology of New York. Volume 1. Containing Descriptions of the Organic Remains of*
1006 *the Lower Division of the New York System (equivalent to the Lower Silurian Rocks of Europe).* Van
1007 Benthuysen, New York, 338 pp.
1008 Hall, J. (1859) Descriptions and figures of the organic remains of the lower Helderberg group and the
1009 Oriskany sandstone. *New York Geological Survey*, 3, 1–532.
1010 Hall, J. (1862) Preliminary notice of some species of Crinoidea from the Waverly sandstone series of
1011 Summit Co., Ohio, supposed to be of the age of the Chemung group of New York. *Annual report on the*
1012 *New York State Museum of Natural History*, 17, 50–60.
1013 Hansen, T., Bruton, D.L. & Jakobsen, S.L. (2005) Starfish from the Ordovician of the Oslo region,
1014 Norway. *Norwegian Journal of Geology*, 85, 209–216.
1015 Heinzeller, T. & Welsch, U. (1994) Crinoidea. *In*: Harrison, F.W. & Ruppert, E.W. (Eds.), *Microscopic*
1016 *Anatomy of Invertebrates*. Wiley, New York, pp. 9–148.
1017 Herbert, B. & Ettensohn, F.R. (2018) What is *Edriocrinus*? *Geological Society of America Abstracts with*
1018 *Programs*, 50, 36–9.
1019 Hess, H. (1999) Permian. *In*: Hess, H., Ausich, W.I., Brett, C.E. & Simms, M.J. (Eds.), *Fossil Crinoids*.
1020 Cambridge University Press, Cambridge, pp. 160–163.
1021 Holloway, D.J. & Jell, P.A. (1983) Silurian and Devonian edrioasteroids from Australia. *Journal of*
1022 *Paleontology*, 57, 1001–1016.
1023 Jaekel, O. (1895) Über die Organisation der Cystoideen. *Deutsche Zoologische Gesellschaft*
1024 *Verhandlungen*, 5, 109-121.
1025 Jaekel, O. (1899) *Stammgeschichte der Pelmatozoen. Erster Band: Thecoidea und Cystoidea*. Springer,
1026 Berlin, 442 pp.
1027 Jefferies, R.P.S. (1986) *The Ancestry of the Vertebrates*. British Museum (Natural History), London, 376
1028 pp.
1029 Jefferies, R.P.S. & Prokop, R.J. (1972) A new calcichordate from the Ordovician of Bohemia and its
1030 anatomy, adaptations and relationships. *Biological Journal of the Linnean Society*, 4, 69–115.
1031 Jell, P.A. (2014) A Tremadocian asterozoan from Tasmania and a late Llandovery edrioasteroid from
1032 Victoria. *Alcheringa*, 38, 528–540.
1033 Jones, G. C. & Jackson, B. (1993) *Infrared Transmission Spectra of Carbonate Minerals*. Chapman & Hall,
1034 London, 254 pp.
1035 Kato, M., Goel, R.K. & Srivastava, S.S. (1987) Ordovician Algae from Spiti, India. *Journal of the Faculty*
1036 *of Sciences, Hokkaido University*, 22, 313–323.
1037 Kesling, R.V. (1968) Cystoids. *In*: Moore, R.C (Ed.), *Treatise on Invertebrate Paleontology,*
1038 *Echinodermata 1(1).* Geological Society of America, Boulder & University of Kansas Press, Lawrence,
1039 pp. S85–S267.
1040 Kjerulf, T. (1865) *Veiviser Ved Geologiske Excursioner i Christiania Omegn*. Brøgger & Christie's
1041 Bogtrykkeri, Christiania, 43 pp.
1042 Knapp, W.D. (1969) Declinida, a new order of late Paleozoic inadunate crinoids. *Journal of Paleontology*,
1043 43, 340–391.
1044 Kolata, D.R. & Guensburg, T.E. (1979) *Diamphidiocystis*, a new mitrate "carpoid" from the Cincinnatian
1045 (Upper Ordovician) Maquoketa Group in southern Illinois. *Journal of Paleontology*, 53,1121–1135.
1046 Kolata, D.R., Frest, T.J. & Mapes, R.H. (1991) The youngest carpoid: occurrence, affinities and life mode
1047 of a Pennsylvanian (Morrowan) mitrate from Oklahoma. *Journal of Paleontology,* 65, 844–855.
1048 Kouchinsky, A., Bengtson, S., Runnegar, B., Skovsted, C., Steiner, M. & Vedrasco, M. (2012) Chronology
1049 of early Cambrian biomineralization. *Geological Magazine*, 149, 221–251.
1050 Kushlina, V.B. (1995) The systematic position and composition of the genus *Bolboporites* (Echinodermata,
1051 Eocrinoidea). *Paleontological Journal*, 29, 46–61.

Kushlina, V.B. (2006) Biting traces on echinoderms from the Ordovician of the St. Petersburg region (Russia). *In*: Mikuláš, R. & Rindsberg, A.K. (Eds.), *Abstract Book: Workshop on Ichnotaxonomy – III, Prague and Moravia, Czech Republic*. Institute of Geology & Academy of Sciences of the Czech Republic, Prague, pp. 12–14.

Kushlina, V.B. (2007) Possible algal-bacterial biofilms on eocrinoids from the Ordovician of the St. Petersburg region (Russia). *Acta Palaeontologica Sinica*, 46 (Suppl.), 237–240.

Lamarck, J.B. M. de (1816) *Histoire Naturelle des Animaux sans Vertèbres. Tome Second*. Verdière, Paris, 568 pp.

Lebrun, P. (1998) Oursins. *Minéraux & Fossiles, hors-série* 8, 1–112.

Lee, S.B., Lefebvre, B. & Choi, D.K. (2005) Latest Cambrian cornutes (Echinodermata, Stylophora) from the Taebaeksan Basin, Korea. *Journal of Paleontology*, 79, 139–151.

Lefebvre, B. (1999) *Stylophores (Cornuta, Mitrata) : Situation au sein du Phylum des Echinodermes et Phylogenèse*, Unpublished PhD thesis, Lyon 1 University, 630 pp.

Lefebvre, B. (2003) Functional morphology of stylophoran echinoderms. *Palaeontology*, 46, 511–555.

Lefebvre, B. (2014) Reinterpretation of the problematic Ordovician genus *Bolboporites* (?Echinodermata) as a calcareous alga. *Programme and Abstracts, 58th Annual Meeting of the Palaeontological Association, Leeds*, 79.

Lefebvre, B. (2017) Réinterprétation de *Bolboporites*, un fossile ordovicien énigmatique. *Journal de l'Association Paléontologique Française*, 72, 30.

Lefebvre, B. & Lerosey-Aubril, R. (2018) Laurentian origin of solutan echinoderms: new evidence from the Guzhangian (Cambrian Series 3) Weeks Formation of Utah, USA. *Geological Magazine*, 155, 1190–1204.

Lefebvre, B., Derstler, K. & Sumrall, C.D. (2012) A reinterpretation of the solutan *Plasiacystis mobilis* (Echinodermata) from the Middle Ordovician of Bohemia. *In*: Kroh, A. & Reich, M. (Eds.), *Echinoderm Research 2010. Proceedings of the Seventh European Conference on Echinoderms, Göttingen, Germany, 2-9 October 2010*. *Zoosymposia*, 7, 287–306.

Lefebvre, B., Sumrall, C.D., Shroat-Lewis, R.A., Reich, M., Webster, G.D., Hunter, A.W., Nardin, E., Rozhnov, S.V., Guensburg, T.E., Touzeau, A., Noailles, F. & Sprinkle, J. (2013) Palaeobiogeography of Ordovician echinoderms. *In*: Harper, D.A.T & Servais, T. (Eds.), *Early Palaeozoic Biogeography and Palaeogeography*. *Geological Society, London, Memoirs*, 38, 173–198.

Le Menn, J. (1985) Les crinoïdes du Dévonien inférieur et moyen du Massif armoricain. Systématique, paléobiologie, évolution, biostratigraphie. *Mémoires de la Société Géologique et Minéralogique de Bretagne*, 30, 1–268.

Lindström, G. (1883) Index to generic names applied to the corals of the Palaeozoic formations. *Bihang till Kongliga Svenska vetenskaps-akademiens Handlingar*, 8, 3–14.

Linnaeus, C. (1758) *Systema Naturae per Regna Tria Naturae. Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. Tomus 1. Pars II*. Editio Duodecima Reformata Laurentii Salvii, Holmiae, pp. 533–1327.

Logan, W.E., Murray, A., Hunt, T.S. & Billings, E. (1863) *Report of Progress from its commencement to 1863*. Geological Survey of Canada, Montreal, 983 pp.

Makhlouf, Y., Lefebvre, B., Nardin, E., Nedjari, A. & Paul, C.R.C. (2017) *Lepidocalix pulcher* Termier and Termier, 1950 (Echinodermata, Diploporita) from the Middle Ordovician of northern Algeria: taxonomic revision and palaeoecological implications. *Acta Palaeontologica Polonica,* 62, 299–310.

Meek, F.B. & Worthen, A.H. (1865) Description of new species of Crinoidea, etc., from the Palaeozoic rocks of Illinois and some of the adjoining states. *Proceedings of the National Academy of Sciences*, 17, 143–155.

Miller, S.A. (1889) *North American Geology and Palaeontology for the Use of Amateurs, Students, and Scientists*. Western Methodist Book Concern, Cincinnati, 793 pp.

1100 Milne-Edwards, H. & Haime, J. (1851) Monographie des Polypiers fossiles des terrains paléozoïques,
1101 précédée d'un tableau général de la classification des Polypes. *Archives du Muséum d'Histoire*
1102 *Naturelle*, 5, 1–502.
1103 Mooi, R. & David, B. (1997) Skeletal homologies of echinoderms. *Paleontological Society Papers*, 3, 305–
1104 335.
1105 Mooi, R. & David, B. (1998) Evolution within a bizarre phylum: homologies of the first echinoderms.
1106 *American Zool*ogist, 38, 965–974.
1107 Mooi, R. & David, B. (2008) Radial symmetry, the anterior/posterior axis, and echinoderm Hox genes.
1108 *Annual Review of Ecology, Evolution, and Systematics*, 39, 43–62.
1109 Moore, R.C. (1978) Flexibilia. *In*: Moore R.C (Ed.), *Treatise on Invertebrate Paleontology, Echinodermata*
1110 *2(2)*. Geological Society of America, Boulder & University of Kansas Press, Lawrence, pp. T759–T812.
1111 Moore, R.C. & Plummer, F.B. (1940) Crinoids from the Upper Carboniferous and Permian strata in Texas.
1112 *University of Texas Publications*, 3495, 9–468.
1113 Moore, R.C., Lane, N.G., Strimple, H.L., Sprinkle, J. & Fay, R.O. (1978) Inadunata. *In*: Moore R.C (Ed.),
1114 *Treatise on Invertebrate Paleontology, Echinodermata 2(2)*. Geological Society of America, Boulder &
1115 University of Kansas Press, Lawrence, pp. T520–T759.
1116 Nardin, E., Lefebvre, B., David, B. & Mooi, R. (2009) La radiation des échinodermes au Paléozoïque
1117 inférieur : l'exemple des blastozoaires. *Comptes Rendus Palevol,* 8, 179–188.
1118 Nardin, E., Lefebvre, B., Fatka O., Nohejlová, M., Kašička, L., Šinágl, M. & Szabad M. (2017).
1119 Evolutionary implications of a new transitional blastozoan echinoderm from th*e* mid Cambrian of
1120 Czech Republic. *Journal of Paleontology*, 91, 672–684.
1121 Noailles, F., Lefebvre, B. & Kašička, L. (2014) A probable case of heterochrony in the solutan
1122 *Dendrocystites* Barrande, 1887 (Echinodermata: Blastozoa) from the Upper Ordovician of the Prague
1123 Basin (Czech Republic) and a revision of the family Dendrocystitidae Bassler, 1938. *Bulletin of*
1124 *Geosciences*, 89, 451–476.
1125 Öpik, A.A. (1934) *Ristnacrinus*, a new Ordovician crinoid from Estonia. *Tartu Ülikooli Geoloogia-*
1126 *Instituudi toimetused*, 40, 1–7.
1127 Owen, A.W., Bruton, D.L., Bockelie, J.F. & Bockelie, T.G. (1990) The Ordovician successions of the Oslo
1128 region, Norway. *Norges Geologiske Undersøkelse, Special Publications*, 4, 3–54.
1129 Oxley, P. & Kay, M. (1959) Ordovician Chazyan series of Champlain Valley, New York and Vermont, and
1130 its reefs. *Bulletin of the American Association of Petroleum Geologists*, 43, 817–853.
1131 Pander, C.H. (1830) *Beiträge zur Geognosie des Russischen Reiches*. Karl Kray, Saint-Petersburg, 165 pp.
1132 Parsley, R.L. (1970) Revision of the North American Pleurocystitidae (Rhombifera - Cystoidea). *Bulletins*
1133 *of American Paleontology*, 58, 135–213.
1134 Parsley, R.L. (1991) Review of selected North American mitrate stylophorans (Homalozoa:
1135 Echinodermata). *Bulletins of American Paleontology*, 100, 1–57.
1136 Parsley, R.L. (1998) Community setting and functional morphology of *Echinosphaerites infaustus*
1137 (Fistuliporita: Echinodermata) from the Ordovician of Bohemia. *Věstník Ústředního ústavu*
1138 *geologického*, 73, 253–266.
1139 Parsley, R.L. & Caster, K.E. (1965) North American Soluta (Carpoidea, Echinodermata). *Bulletins of*
1140 *American Paleontology*, 49, 109–174.
1141 Paul, C.R.C. (1967) The functional morphology and mode of life of the cystoid *Pleurocystites* E. Billings,
1142 1854. *In*: Milot, E. (Ed.), *Echinoderm Biology*. *Symposium of the Zoological Society of London*, 20,
1143 105–123.
1144 Pisera, A. (1994) Echinoderms from the Mójcza Limestone. *Palaeontologia Polonica*, 53, 283–307.
1145 Quenstedt, F.A. (1881) *Petrefakten Deutschlands, Part 1, Volume 6. Korallen (Röhren- und Sternkorallen)*.
1146 Fues, Leipzig, 1093 pp.
1147 Rahman, I.A. & Clausen, S. (2009) Re-evaluating the palaeobiology and affinities of the Ctenocystoidea
1148 (Echinodermata). *Journal of Systematic Palaeontology*, 7, 413–426.

1149 Rahman, I.A. & Zamora, S. (2009) The oldest cinctan carpoid (stem-group Echinodermata), and the
1150 evolution of the water-vascular system. *Zoological Journal of the Linnean Society*, 157, 420–432.
1151 Rahman, I.A., Zamora, S. & Geyer, G. (2010) The oldest stylophoran echinoderm: a new *Ceratocystis* from
1152 the Middle Cambrian of Germany. *Paläontologische Zeitschrift*, 84, 227–237.
1153 Rahman, I.A., Waters, J.A., Sumrall, C.D. & Astolfo, A. (2015) Early post-metamorphic, Carboniferous
1154 blastoid reveals the evolution and development of the digestive system in echinoderms. *Biology Letters*,
1155 11, 20150776.
1156 Raymond, P.E. (1905) The faunas of the Chazy Limestone. *American Journal of Science*, 20, 353–382.
1157 Raymond, P.E. (1906) The Chazy Formation and its fauna. *Annals of the Carnegie Museum*, 3, 498–596.
1158 Raymond, P.E. (1913) Ordovician of Montreal and Ottawa. *12th International Geological Congress*
1159 *Canada, Guide book*, 3, 137–162.
1160 Reed, F.R.C. (1899) The Lower Palaeozoic bedded rocks of County Waterford. Quarterly *Journal of the*
1161 *Geological Society of London*, 55, 718–772.
1162 Regnéll, G. (1956) On *Bolboporites*. *Norsk Geologisk Tiddskrift*, 36, 81.
1163 Regnéll, G. (1982) What is *Bolboporites*? *In*: Lawrence, J.M. (Ed.), *International Echinoderms*
1164 *Conference, Tampa Bay*. Balkema, Rotterdam, pp. 97.
1165 Reich, M. (1999) Ordovizische und silurische Holothurien (Echinodermata). *In*: Reich, M. (Ed.),
1166 *Festschrift zum 65. Geburstag von Ekkehard Herrig*. *Greifswalder Geowissenschaftliche Beiträge*, 6,
1167 479–488.
1168 Reich, M. (2001) *Linguaserra*? (Echinodermata: Ophiocistoidea) aus dem Ordovizium Baltoskandiens.
1169 *Greifswalder Geowissenschaftliche Beiträge*, 9, 33–35.
1170 Reich, M. (2010) The oldest synallactid sea cucumber (Echinodermata: Holothuroidea: Aspidochirotida).
1171 *Paläontologische Zeitschrift*, 84, 541–546.
1172 Reich, M. & Smith, A.B. (2009) Origins and biomechanical evolution of teeth in echinoids and their
1173 relatives. *Palaeontology*, 52, 1149–1168.
1174 Roemer, C.F. (1854) Beiträge zur geologischen Kenntnis des nordwestlichen Harzgebirges,
1175 *Palaeontographica*, 3, 1–67.
1176 Roemer, C.F. (1885) Lethaea erratica oder Aufzählung und Beschreibung der in der norddeutschen Ebene
1177 vorkommenden Diluvial-Geschiebe nordischer Sedimentärgesteine. *Paläontologische Abhandlungen*, 2,
1178 250–420.
1179 Rouault, M. (1851) Mémoire sur le terrain paléozoïque des environs de Rennes. *Bulletin de la Société*
1180 *géologique de France*, 8, 358–399.
1181 Rozhnov, S.V. (2002) Morphogenesis and evolution of crinoids and other pelmatozoan echinoderms in the
1182 Early Paleozoic. *Paleontological Journal*, 36, 525–674.
1183 Rozhnov, S.V. (2005) Echinoderms. *In*: Dronov, A., Tolmacheva, T., Raevskaya, E. & Nestell, M. (Eds.),
1184 *Cambrian and Ordovician of St. Petersburg area. Guidebook of the pre-conference field trip*. St.
1185 Petersburg State University, Saint-Petersburg, pp. 23–26.
1186 Rozhnov, S.V. (2009). New data on Ordovician eocrinoids and paracrinoids of the Baltic Region.
1187 *Geophysical Research Abstracts*, 11, EGU2009-3683-1.
1188 Rozhnov, S.V. & Kushlina, V.B. (1994a) A new interpretation of *Bolboporites* (Echinodermata,
1189 ?Eocrinoidea). *Paleontological Journal*, 28, 71–80.
1190 Rozhnov, S.V. & Kushlina, V.B. (1994b) Interpretation of new data on *Bolboporites* Pander, 1830
1191 (Echinodermata; Ordovician). *In*: David, B., Guille, A., Féral, J.P. & Roux, M. (Eds.), *Echinoderms*
1192 *through Time*. Balkema, Rotterdam, pp. 179–180.
1193 Ruedemann, R. (1901) Trenton conglomerate of Rysedorph Hill, Rensselaer County, New York and its
1194 fauna. *New York State Museum Bulletin*, 49, 1–114.
1195 Ruta, M. (1999) A new stylophoran echinoderm, *Juliaecarpus milnerorum*, from the Late Ordovician
1196 Upper Ktaoua Formation of Morocco. *Bulletin of the Natural History Museum, London (Geology)*, 55,
1197 47-79.

1198 Ruta, M. & Bartels, C. (1998) A redescription of the anomalocystitid mitrate *Rhenocystis latipedunculata*
1199 from the Lower Devonian of Germany. *Palaeontology*, 41, 771–806.
1200 Schmidt, H. (1951) Whitehouse's Ur-Echinodermen aus dem Cambrium Australiens. *Paläontologische*
1201 *Zeitschrift*, 24, 142–145.
1202 Seilacher, A. & MacClintock, C. (2005) Crinoid anchoring strategies for soft-bottom dwelling. *Palaios*, 20,
1203 224–240.
1204 Shaw, F.C. & Bolton, T.E. (2011) Ordovician trilobites from the Romaine and Mingan formations
1205 (Ibexian–late Whiterockian), Mingan Islands, Quebec. *Journal of Paleontology*, 85, 406–441.
1206 Smith, A.B. (1980a) Stereom microstructure of the echinoid test. *Special Papers in Palaeontology*, 25, 1–
1207 81.
1208 Smith, A.B. (1980b) The structure and arrangement of echinoid tubercles. *Philosophical Transactions of*
1209 *the Royal Society, B*, 289, 1–46.
1210 Smith, A.B. (1982) The affinities of the Middle Cambrian Haplozoa (Echinodermata). *Alcheringa*, 6, 93–
1211 99.
1212 Smith, A.B. (1988) Patterns of diversification and extinction in Early Palaeozoic echinoderms.
1213 *Palaeontology*, 31, 799–828.
1214 Smith, A.B. & Savill, J.J. (2001) *Bromidechinus*, a new Ordovician echinozoan (Echinodermata), and its
1215 bearing on the early history of echinoids. *Transactions of the Royal Society of Edinburgh (Earth*
1216 *Sciences),* 92, 137–147.
1217 Smith, A.B. & Zamora, S. (2009) Rooting phylogenies of problematic fossil taxa; a case study using
1218 cinctans (stem-group echinoderms). *Palaeontology*, 52, 803–821.
1219 Spencer, W.K. & Wright, C.W. (1966) Asterozoans. *In*: Moore R.C (Ed.), *Treatise on Invertebrate*
1220 *Paleontology, Echinodermata 3(1).* Geological Society of America, Boulder & University of Kansas
1221 Press, Lawrence, pp. U4–U107.
1222 Spjeldnaes, N. & Nitecki, M.H. (1990) *Coelosphaeridium*, an Ordovician alga from Norway. *Institutt for*
1223 *Geologi Universitetet I Oslo, Intern Skriftserie*, 59, 1–53.
1224 Springer, F. (1926) Unusual forms of fossil crinoids. *Proceedings of the United States National Museum*,
1225 67, 1–137.
1226 Sprinkle, J. (1973) Morphology and evolution of blastozoan echinoderms. *Harvard University Museum of*
1227 *Comparative Zoology, Special Publication*, 1–283.
1228 Sprinkle, J. & Guensburg, T.E. (2001) Growing a stalked echinoderm within the Extraxial-Axial Theory.
1229 *In*: Barker, F.K. (Ed.), *Echinoderms 2000*. Swets & Zeitlinger, Lisse, pp. 59–65.
1230 Stanton, R.J., Jr., Lambert, L.L., Webb, G.E. & Lustig, L.D. (2016) *Chaetetes* morphology, environment,
1231 and taxonomy. *Facies*, 62:29, 1–21.
1232 Strimple, H.L. (1961) Late Desmoinesian crinoid faunule from Oklahoma. *Bulletin of the Oklahoma*
1233 *Geological Survey*, 93, 1–189.
1234 Strimple, H.L. & Watkins, W.T. (1969) Carboniferous crinoids of Texas with stratigraphic implications.
1235 *Palaeontographica Americana*, 6, 41–275.
1236 Sumrall, C.D., Sprinkle, J. & Guensburg, T.E. (1997) Systematics and paleoecology of late Cambrian
1237 echinoderms from the western United States. *Journal of Paleontology*, 71, 1091–1109.
1238 Sutton, M.D., Briggs, D.E.G., Siveter, D.J., Siveter, D.J. & Gladwell, D.J. (2005) A starfish with three-
1239 dimensionally preserved soft parts from the Silurian of England. *Proceedings of the Royal Society B*,
1240 272, 1001–1006.
1241 Termier, H. & Termier, G. (1950) Contribution à l'étude des faunes paléozoïques de l'Algérie. *Bulletin du*
1242 *Service de la Carte Géologique de l'Algérie*, 1–83.
1243 Thoral, M. (1935) *Contribution à l'Etude Paléontologique de l'Ordovicien Inférieur de la Montagne Noire*
1244 *et Révision Sommaire de la Faune Cambrienne de la Montagne Noire*. Imprimerie de la Charité,
1245 Montpellier, 362 pp.
1246 Tinn, O. & Ainsaar, L. (2014) Asterozoan pedicellariae and ossicles revealed from the Middle Ordovician
1247 of Baltica. *Acta Palaeontologica Polonica*, 59, 353–358.

Twenhofel, W.H. (1938) Geology and paleontology of the Mingan Islands, Quebec. *Geological Society of America, Special Paper*, 11, 1–132.

Ubaghs, G. (1961) Un échinoderme nouveau de la classe des carpoïdes dans l'Ordovicien inférieur du département de l'Hérault (France). *Comptes-Rendus de l'Académie des Sciences, Paris*, 253, 2565–2567.

Ubaghs, G. (1963) *Rhopalocystis destombesi* n.g., n.sp., éocrinoïde de l'Ordovicien inférieur (Trémadocien supérieur) du Sud marocain. *Notes du Service géologique du Maroc*, 23, 25–44.

Ubaghs, G. (1968a) Stylophora. *In*: Moore, R.C (Ed.), *Treatise on Invertebrate Paleontology, Echinodermata 1(2)*. Geological Society of America, Boulder & University of Kansas Press, Lawrence, pp. S495–S565.

Ubaghs, G. (1968b) *Cymbionites* and *Peridionites* - unclassified Middle Cambrian echinoderms. *In*: Moore, R.C (Ed.), *Treatise on Invertebrate Paleontology, Echinodermata 1(2)*. Geological Society of America, Boulder & University of Kansas Press, Lawrence, pp. S634–S637.

Ubaghs, G. (1969) *Aethocrinus moorei* Ubaghs, n. gen., n. sp., le plus ancien crinoïde dicyclique connu. *University of Kansas Paleontological Contributions*, 38, 1–25.

Ubaghs, G. (1970) *Les Echinodermes "Carpoïdes" de l'Ordovicien Inférieur de la Montagne Noire (France)*. Editions du CNRS, Cahiers de Paléontologie, Paris, 110 pp.

Ubaghs, G. (1972) Le genre *Balantiocystis* Chauvel (Echinodermata, Eocrinoidea) dans l'Ordovicien inférieur de la Montagne Noire (France). *Annales de Paléontologie*, 58, 3–27.

Ubaghs, G. (1978a) Skeletal morphology of fossil crinoids. *In*: Moore R.C (Ed.), *Treatise on Invertebrate Paleontology, Echinodermata 2(1)*. Geological Society of America, Boulder & University of Kansas Press, Lawrence, pp. T58–T216.

Ubaghs, G. (1978b) Camerata. *In*: Moore R.C (Ed.), *Treatise on Invertebrate Paleontology, Echinodermata 2(1)*. Geological Society of America, Boulder & University of Kansas Press, Lawrence, pp. T408–T518.

Ubaghs, G. (1981) Réflexions sur la nature et la fonction de l'appendice articulé des "carpoïdes" Stylophora (Echinodermata). *Annales de Paléontologie*, 67, 33–48.

Ubaghs, G. (1983) Echinodermata. Notes sur les Échinodermes de l'Ordovicien Inférieur de la Montagne Noire (France). *In*: Courtessole, R., Marek, L., Pillet, J., Ubaghs, G. & Vizcaïno, D. (Eds.), *Calymena, Echinodermata et Hyolitha de l'Ordovicien de la Montagne Noire (France Méridionale)*. Société d'Etudes Scientifiques de l'Aude, Carcassonne, p. 33–35.

Wanner, J. (1920) Ueber einige palaeozoische Seeigelstacheln (*Timorocidaris* gen. nov. und *Bolboporites* Pander). *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen*, 22, 696–712.

Wanner, J. (1924) Die permischen Blastoiden von Timor. *Jaarboek van het mijnwezen in Nederlands Oost-Indie*, 51, 163-233.

Webster, G.D. & Kues, B.S. (2006) Pennsylvanian crinoids of New Mexico. *New Mexico Geology*, 28, 3–39.

White, T.G. (1896) Geology of Essex and Willsboro townships, Essex County, New York. *Transactions of the New York Academy of Sciences*, 13, 214–233.

Whitehouse, F.W. (1941) The Cambrian faunas of north-eastern Australia. Part 4: Early Cambrian echinoderms similar to the larval stages of recent forms. *Memoirs of the Queensland Museum*, 12, 1–28.

Wyse Jackson, P.N., Buttler, C.J. & Key, M.M. (2002) Palaeoenvironmental interpretation of the Tramore Limestone Formation (Llandeilo, Ordovician) based on bryozoan colony form. *In*: Wyse Jackson, P.N., Buttler, C.J. & Spencer Jones, M.E. (Eds.), *Bryozoan Studies 2001*. Swets & Zeitlinger, Lisse, pp. 359–365.

Yakovlev, N.N. (1921). *Bolboporites*, its structure and affinity to the Hydrozoa. *Annals of the Russian Paleontological Association*, 3, 1–10 [in Russian, with an abstract in English]

Yeltysheva, R.S. (1955) *Bolboporites*. *Voprosy Paleontologii*, 2, 136–147 [in Russian].

Zamora, S., Alvaro, J.J. & Vizcaïno, D. (2009) Pelmatozoan echinoderms from the Cambrian-Ordovician of the Iberian Chains (NE Spain): early diversification of anchoring strategies. *Swiss Journal of Geosciences*, 102, 43–55.

Zamora, S., Lefebvre, B., Alvaro, J.J., Clausen, S., Elicki, O., Fatka, O., Jell, P., Kouchinsky, A., Lin, J.P., Nardin, E., Parsley, R.L., Rozhnov, S.V., Sprinkle, J., Sumrall, C.D., Vizcaïno, D. & Smith, A.B. (2013) Cambrian echinoderm diversity and palaeobiogeography. *In*: Harper, D.A.T & Servais, T. (Eds.), *Early Palaeozoic Biogeography and Palaeogeography*. *Geological Society, London, Memoirs*, 38, 157–171.

Zittel, K.A. von (1879) *Handbuch der Paläontologie. 1. Band, 1. Protozoa, Coelenterata, Echinodermata und Molluscoidea*. Oldenburg, München & Leipzig, 765 pp.

Figure Captions

FIGURE 1. External morphology of *Bolboporites uncinatus* Pander, 1830, Middle Ordovician (Dapingian), Saint-Petersburg area, Russia; redrawn and modified from Rozhnov & Kushlina (1994a, fig. 3b) and Kushlina (1995, pl. 5 fig. 1b). A: Base of the cone, in front view, showing the two adjoining depressed areas (lunules), and the small orifice corresponding to the outlet of the longitudinal internal canal. B: Lateral view, showing the typical honeycomb ornamentation on lateral sides of the cone.

FIGURE 2. Fourier transform infrared (FT-IR) spectroscopy made on transverse sections of *Bolboporites mitralis* Pander, 1830, included in LR white resin; UCBL-FSL 712510, Middle Ordovician (Dapingian), Saint-Petersburg area, Russia. A: Location of FTIR analyses. B: IF spectrum of LR white resin (control spectrum). C, G: IF spectra of sampling points located on the external margin of the specimen (calcite). D-F: IF spectra of sampling points within the specimen (calcite). Peak values indicated in cm^{-1}.

FIGURE 3. Thin section views of *Bolboporites mitralis* Pander, 1830 (Middle Ordovician (Dapingian), Saint-Petersburg area, Russia; UCBL-FSL 712510) included in the surrounding sedimentary rock (ss) which is a packstone-wackestone with bioclasts and glauconite grains. A: Plane-polarized optical light, showing the conjugated cleavage planes (cp) crosscutting the whole cone. B: Cross-polarized optical light, showing total extinction of the cone. C: Close up

view of A and B (plane polarized light) revealing the tenuous dark-brown patches aligned along cleavage planes, forming a stereom-like structure.

FIGURE 4. EBSD mapping of *Bolboporites mitralis* Pander, 1830; Middle Ordovician (Dapingian), Saint-Petersburg area, sample UCLB-FSL 712510, cross section, perpendicular to the cone axis. A: Periphery of the sample observed by SEM. B: corresponding EBSD map, showing the size heterogeneity of the crystals around the monocrystalline structure; the box on the down right corner gives the color code for crystallographic axes. C: another peripheral zone, at higher magnification. D: corresponding EBSD map. E: central zone of *B. mitralis* visualized by SEM. F and G: two EBSD maps of two areas of the central zone (E) 4 mm apart. Although the structure is fully monocrystalline on the whole cross-section (no grain limit detectable), a slight variation of the colour indicates a tiny crystallographic disorientation (< 4°).

FIGURE 5. Internal structures of *Bolboporites mitralis* Pander, 1830; Middle Ordovician (Dapingian), Saint-Petersburg area, Russia. A-C: Cathodoluminescence view of sectioned *Bolboporites mitralis* Pander, 1830; Middle Ordovician (Dapingian), Saint-Petersburg area, Russia. A: Longitudinal section through specimen UCBL-FSL 712508, with several randomly distributed *Trypanites*-like borings (bor) through the body wall. B-C: Cross section through specimen UCBL-FSL 712510, showing evidence of narrow longitudinal axial canal (int. canal). B: General view of the sectioned specimen. C: Detail showing the central internal canal filled with a syntaxial blocky calcite cement (sbc). The surrounding skeleton reveals a stereom-like structure with aligned luminescent inclusions; specimen UCBL-FSL 712510 D: This SEM view shows the opening part of a tubular boring crosscutting the stereom-like microstructure of a *Bolboporites* cone (UCBL-FSL 712508). Note this stereom-like microstructure is highly cemented near the walls of the boring and at the periphery of the cone, whereas micropores (mp) are presents in the more internal parts. This boring is partially filled with a bioclastic and glauconitic packstone (gp) that also surround the cone. More internal parts of the boring are cemented by dolomite rhombs (dol) and by a syntaxial blocky calcite cement (sbc).


FIGURE 6. CT-scan imagery of *Bolboporites mitralis* Pander, 1830; UCBL-FSL 712509 Middle Ordovician (Dapingian), Saint-Petersburg area, Russia. A: Virtual cross section showing diffuse peripheral rim, made of denser (less porous) stereom than central part of the


cone, as also shown on SEM imagery (Fig 5B). This view is reminiscent, in part, to a section through a cidaroid spine (see, e.g., Donovan 2018). Lateral honeycomb cells appear clearly on external margin of peripheral rim. B: Three-dimensional reconstruction of the specimen, with line indicating location of cross section in A.


FIGURE 7. External aspect and morphological disparity of *Bolboporites*; all specimens from same locality and level, Upper Ordovician (Sandbian), Skien-Langesund area, Norway. A: *Bolboporites* sp., lateral side of the cone showing honeycomb ornamentation, and almost flat base of the cone; PMO 116887. B: *Bolboporites elongatus* Kushlina, 1995, narrow cone in lateral view, with gently convex base; PMO 218238. C-D: *Bolboporites* cf. *mitralis* Pander, 1830; PMO 218249. C: Wide cone in lateral aspect, with convex base. D: Basal surface with two lunules in almost central position, and small orifice at their junction.


FIGURE 8. External morphology of *Bolboporites mitralis* Pander, 1830; Middle Ordovician (Dapingian), Saint-Petersburg area, Russia. A: Cells forming honeycomb ornamentation on lateral sides of the cone; SEM view of specimen UCBL-FSL 713254. B: Lunules and associated small orifice, on the base of the cone; SEM view of specimen UCBL-FSL 713250. Note also the well-preserved stereom-like microstructure that appears on the wall of these lunules.


FIGURE 9. Aboral tubercles of the Recent oreasterid asteroid *Pentaceraster mammilatus* (Audouin, 1826). A: Extracted tubercle in lateral view, with its lateral walls covered by thick, polyplated, granulose aboral membrane; specimen UCBL.2017.01.44. B-E: Specimen UCBL.2017.01.47. B: Extracted tubercle in oblique view, with its basal surface showing several diverging branches, connecting it with surrounding aboral plates. C: Cross-section of extraction site of tubercle shown in (B), on the aboral surface, showing complex articulation of tubercle-bearing plate with surrounding aboral skeletal elements. D: General view of aboral surface. E: Close-up of aboral surface showing area of extraction of the spiny tubercle.



A
B
5 mm
base
apex
orifice
lunules
honeycomb cells



A
B
C
D
E
F
G
5 mm
B
2929
2857
1721
1148
C
D
E
F
G
711
871
1395



A
ss
cp
2 mm
B
ss
2 mm
C
ss
cp
1 mm



A
90 µm
B
100 um
Calcite
0001
$1\bar{1}00$
$0\bar{1}10$
$1\bar{2}10$
C
50 µm
D
60 um
E
100 µm
F
5 µm
G
3 µm


A
B
int. canal
bor
5 mm
C
sbc
0.5 mm
D
gp
dol
sbc
mp
0.25 mm


A
2 mm
B
2 mm


A
5 mm
B
5 mm
C
D
5 mm


A
1 mm
B
1 mm


A
5 mm
B
5 mm
C
5 mm
D
10 mm
E
10 mm