



# Seawater paleotemperature and paleosalinity evolution in neritic environments of the Mediterranean margin: insights from isotope analysis of bivalve shells

Justine Briard, Emmanuelle Pucéat, Emmanuelle Vennin, M. Daëron, V. Chavagnac, R. Jaillet, Didier Merle, Marc de Rafélis

## ► To cite this version:

Justine Briard, Emmanuelle Pucéat, Emmanuelle Vennin, M. Daëron, V. Chavagnac, et al.. Seawater paleotemperature and paleosalinity evolution in neritic environments of the Mediterranean margin: insights from isotope analysis of bivalve shells. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 2020, 543, pp.109582. 10.1016/j.palaeo.2019.109582 . hal-02449079

**HAL Id: hal-02449079**

**<https://hal.science/hal-02449079>**

Submitted on 22 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Seawater paleotemperature and paleosalinity evolution in neritic environments of the Mediterranean margin: insights from isotope analysis of bivalve shells.

J. Briard<sup>a,b,\*</sup>, E. Pucéat<sup>b</sup>, E. Vennin<sup>b</sup>, M. Daëron<sup>c</sup>, V. Chavagnac<sup>a</sup>, R. Jaillet<sup>b</sup>, D. Merle<sup>d</sup>, M. de Rafélis<sup>a</sup>

<sup>a</sup> Géosciences Environnement Toulouse (GET), Université Paul Sabatier Toulouse 3, CNRS UMR 5563, IRD, 14 avenue Edouard Belin F-31400 Toulouse, France.

<sup>b</sup> Biogéosciences, UMR 6282, UBFC/CNRS, Université Bourgogne Franche-Comté, 6 boulevard Gabriel, F-21000 Dijon, France.

<sup>c</sup> Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France.

<sup>d</sup> Muséum National d'Histoire Naturelle, UMR CRNS 7207 (CR2P-MNHN, CNRS, UPMC-Paris 6), Sorbonne Universités, 8 rue Buffon F-75005 Paris, France.

\* E-mail address: [justine.briard@get.omp.eu](mailto:justine.briard@get.omp.eu)

## Abstract

The first step of ice-sheet build-up on Antarctica at the Eocene-Oligocene boundary was followed by a phase of climate instability culminating during the Miocene with a warming event called the Middle Miocene Climate Optimum (MMCO), that ended with a marked cooling phase identified as the Middle Miocene Climate Transition (MMCT). While numerous benthic foraminifera  $\delta^{18}\text{O}$  and Mg/Ca data have been used to capture the global climate evolution during this interval of climate and ice-sheet instability, geochemical records from shallow-water carbonates still remain scarce. Yet such records are crucial to capture the diversity of regional environmental responses to global climate changes, and thus to better



understand the behavior of our climate system during this critical interval. In this work, we test the potential of a multi-proxy approach ( $\delta^{18}\text{O}$ , clumped isotope ( $\Delta_{47}$ ), strontium isotopes ( $^{87}\text{Sr}/^{86}\text{Sr}$ )) applied to bivalves recovered from the Liguro-Provencal Basin and Rhodano-Provencal basin (Northwestern of the Mediterranean Sea) to reconstruct the evolution of shallow seawater temperature and salinity in this region over the latest Oligocene to Middle Miocene interval (~10 Myrs). Our results highlight a local cooling in the northwestern Mediterranean Sea during the MMCO that contrasts with the warming observed in other regions, with seawater temperatures inferred from  $\Delta_{47}$  analyses in the 13-18°C range. These cool seawater temperatures recorded in the studied region during the MMCO are much cooler than those recorded in the open oceans, but are in agreement with the proliferation of bryozoan observed in the Castillon-Du-Gard area during this interval. Low bivalve  $\delta^{18}\text{O}$  values (-3.24 ‰ in average) are recorded during this episode—are interpreted to reflect enhanced freshwater inputs, lowering local salinity and seawater  $\delta^{18}\text{O}$ . Such enhanced freshwater inputs point to a phase of enhanced hydrological cycle in the studied region, possibly linked to the global mid-Miocene warming event. Bivalve  $^{87}\text{Sr}/^{86}\text{Sr}$  data mostly fall within the global seawater Sr isotope reference curve, but remain compatible with enhanced freshwater inputs during the Aquitanian and during the Langhian, as these inputs were probably not important enough to induce a large departure from the seawater curve at a regional scale.

#### **Highlights :**

- Coupled  $\Delta_{47}$  -  $\delta^{18}\text{O}$  -  $^{87}\text{Sr}/^{86}\text{Sr}$  analyses of bivalve shells decipher temperature and salinity.

• Cool seawater temperatures in the northwestern Mediterranean Sea during the MMCO.

• Enhanced hydrological cycle in western Europe during the MMCO.

**Keywords:** Paleoclimate; Middle Miocene Climatic Optimum; Clumped isotopes; Hydrological cycle; Coastal environment

## 1. Introduction

A major feature of the long-term climate evolution during the Cenozoic is the transition from the “greenhouse” conditions of the late Early Eocene (~50 million years ago) to the modern “icehouse” climate mode with its much lower atmospheric CO<sub>2</sub> levels and permanent polar ice-sheets on both poles (Zachos et al., 2008). This transition occurred through a series of steps, beginning with the abrupt global cooling recorded at the Eocene/Oligocene transition (Zachos et al., 1996; Lear et al., 2008; Liu et al., 2009; Galeotti et al., 2016; Cramwinckel et al., 2018) followed by a period of instability of the Antarctic ice-sheet during the Oligocene and part of the Miocene (Billups and Schrag, 2002; Lear et al., 2015; Gasson et al., 2016). This period of instability is marked by several warming and cooling phases associated with waxing and waning of the Antarctica ice-sheet. This includes the Late Oligocene Warming Event, the cooling at the Oligocene-Miocene boundary (~23.03 Ma, Mi-1 event), and the global warming event leading to the Middle Miocene Climatic Optimum (MMCO, ca. 16 Ma to ca. 14.5 Ma) during the latest Early to Middle Miocene interval (e.g. Zachos et al., 2008; Mawbey and Lear, 2013; Beddow et al., 2016, Goldner et al., 2014; Holbourn et al., 2015; Super et al., 2018). This phase of instability ended with the marked cooling of sea surface temperature (SST) defined as the Middle Miocene Climate Transition (MMCT, ca. 14.5 to ca. 13.5 Ma), associated once more with a large-scale

expansion of the Antarctic ice-sheet (Holbourn et al., 2005; Foster et al., 2012). Although the first development of permanent ice-sheets on Antarctica during the Eocene-Oligocene cooling event has been extensively studied in the past decades (Zachos et al., 1996; Ivany et al., 2000; Lear et al., 2008; Liu et al., 2009; Tremblin et al., 2016), little is known about the following interval of ice-sheet instability. For the earliest Oligocene to the middle Miocene interval (~10 million years), most paleoclimatic records available to date are based on isotopic ( $\delta^{18}\text{O}$ ) and elemental (Mg/Ca) composition of benthic foraminifera that record the physico-chemical features of ocean bottom water temperature (Pagani et al., 1999; Miller et al., 1991; Diester-Haass et al., 2009). Marine surface temperature records are scarce, mostly based on various geochemical measurements on planktonic organisms (e.g.  $\delta^{18}\text{O}$ , Mg/Ca,  $\text{TEX}_{86}$ ,  $\text{U}_{37}^{k'}$ ), and are restricted to limited time intervals (Pagani et al., 1999; Super et al., 2018). Palynological data or fauna distribution give further information on paleoenvironmental conditions on the continents. (Bruch et al., 2004; Jimenez and Suc 2007; Pound et al., 2012). However, archives from coastal environments such as bivalves remain understudied for this time period despite their location in a transitional environment between continent and ocean important to characterize as well to obtain a global view of the response of climate to perturbations. Bivalves recovered from shallow-water environments (< 50 m, Stenzel, 1971) have been successfully used to document temperature and/or salinity variations in the past (Klein et al., 1996; Kirby et al., 1998, Steuber et al., 2005; Schöne and Giere 2005; Huyghe et al., 2012, 2015; Bougeois et al., 2014, 2016; Mouchi et al., 2018; de Winter et al., 2018). Bivalves like oyster shells offer key advantages for paleotemperature reconstructions in coastal area, as (1) they are formed of low-magnesian calcite, thus quite resistant to recrystallization (Brand and Veizer, 1981; Al-Aasm and Veizer, 1986; Ulmann and Korte, 2015), (2) they have a large spatial and temporal distribution and (3) they live fixed on a substratum during several years, allowing reconstruction of local seasonal variations of environmental parameters (temperature

and salinity), which cannot be assessed with foraminifera. In coastal environments, reconstruction of shallow water temperatures using bivalve  $\delta^{18}\text{O}$  can however be impeded by local seawater  $\delta^{18}\text{O}$  ( $\delta^{18}\text{O}_{\text{sw}}$ ) variability due to local processes such as evaporation, precipitation, surface runoff and groundwater discharges. One way to overcome this limitation is to combine conventional oxygen isotope ( $\delta^{18}\text{O}$ ) with clumped isotope ( $\Delta_{47}$ ) analyses of bivalves because carbonate clumped isotope ( $\Delta_{47}$ ) composition is solely linked to temperature during mineral formation, and independent from the isotopic composition of the ambient water (Ghosh et al., 2006; Schauble et al., 2006; Eiler et al., 2009; Henkes et al., 2013). Additional information on environmental conditions is provided by the Sr isotopic system applied on bivalves, as the  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios recorded in their umbo has been shown to record the occurrence of large freshwater discharge in their living marine environment (Ingram and Sloan, 1992; Bryant et al., 1995; Winderlund and Anderson, 2006; El Meknassi et al., 2018; Zaky et al., 2018).

In this context, we propose to test the potential of a multi-proxy approach applied on bivalves thriving in coastal environments to reconstruct the evolution of shallow (<50m) seawater temperatures over the latest Oligocene to Middle Miocene interval. The two selected outcrops, i.e. Carry-Le-Rouet and Castillon-Du-Gard located within the Liguro-Provençal and Rhodano-Provençal Basins (SE of the Mediterranean Sea, France), present sediments deposited in shallow (<50m) open marine conditions with abundant bivalve shells. In both localities, a well-constrained chronostratigraphic framework is available (Andreieff et al., 1972; Oudet et al., 2010; Ferrandini et al., 2010; Demory et al., 2011). Our new dataset provides new information on both temperature, salinity and runoff evolution within a still understudied region. The outcomes contribute to better constrain the regional environmental response to global climatic changes during this key interval of Antarctica ice-sheet waning and waxing.

125

## 126 **2. Geological setting**

127       The modern western Mediterranean Sea is structured into four basins (Fig.1A):  
128   Alboran Sea, Adriatic Sea, Tyrrhenian Sea, and Liguro-Provençal basin that extends from the  
129   Gulf of Genoa in the East to the Strait of Gibraltar in the West to Sardinia in the South. These  
130   basins began to form by the end of the Oligocene (~23.03 Ma) and reached their full  
131   extension during the Miocene (23.03 – 5.33 Ma) (Rehault et al., 1984). The oceanization of  
132   the Liguro-Provençal basin corresponds to the opening of a back-arc basin associated with the  
133   eastward retreat of an Adriatic/ionian slab sinking into the mantle (Gueguen et al., 1998;  
134   Faccenna et al., 1997; Gattacceca et al., 2007; van Hinsbergen et al., 2019). During the  
135   Miocene marine transgression, a second basin started to form in the Rhône Valley, called the  
136   Rhodano-Provençal Basin. This basin is located north of the Liguro-Provençal Basin, at the  
137   junction between the perialpine foreland basin to the north and the northern margin of the  
138   western Mediterranean to the south. The Rhodano-Provençal Basin extends to the north as a  
139   channel that follows the alpine chain from Marseille (France) to Vienna (Austria). The pre-  
140   Miocene structuration of the Rhodano-Provençal Basin results in a highly differentiated  
141   topography organized in sub-basins (e.g. Valreas, Sommières, Uzès and Castillon-Du-Gard).  
142   Due to this inheritance, the Rhodano-Provençal sub-basins evolved independently of each  
143   other during the Miocene marine transgression. The two outcrops selected for this study are  
144   located within the Liguro-Provençal (Carry-Le-Rouet) and Rhodano-Provençal (Castillon-Du-  
145   Gard) basins (Fig. 1A).

146

### 147 *2.1. Carry-Le-Rouet outcrops in the Coastal Nerthe area*

148       On the northern part of the Liguro-Provençal Basin (Fig. 1B), a ~100 m succession of  
149   Late Oligocene (Chattian) to Early Miocene (Burdigalian) continental and marine sediments

crops out along the Blue coast (43°19'50.19''N, 5°09'05.67''E), at Carry-Le-Rouet (20 km west of Marseille, France). The sedimentary succession is separated by a major unconformity from the deformed and eroded Mesozoic basement called Nerthe Massif. The Barrémo-Aptian basement (Urgonian facies) is a thrust structure resulting from the Eocene Pyrenean compression (Tempier, 1987). The outcrops are exposed along the coast on 8 km and the sediments are organized in six lithostratigraphic units extending from the Cap de Nautes to the Pointe de Tamaris (Fig. 1B; Andreieff et al., 1972; Maurel-Ferrandini, 1976). The basal Conglomeratic Rouet unit (Unit 1, Chattian) is characterized by a transition from fluvial to deltaic depositional environments. The following formations (Para-reefal Cap de Nautes unit (Unit 2), the Brackish Rousset unit (Unit 3), the Bioclastic Carry unit (Unit 4), the Biodetrital Sausset unit (Unit 5) and the Plan de Sausset unit (Unit 6)) correspond to an alternation of marine to continental-marine transitional depositional environments. This succession benefits from a well-constrained biostratigraphic record using calcareous nannofossils and planktonic foraminifera (Oudet et al., 2010), supplemented with magnetostratigraphy (Demory et al., 2011). Despite the presence of hiatus along the Chattian to Burdigalian succession, the interval sampled in this study did not record any major stratigraphic gaps considering the sedimentary succession continuity and the age model proposed by Demory et al., 2011.

## *2.2. Castillon-Du-Gard outcrops*

The Castillon-Du-Gard Sub-basin (43°58'12.61''N, 4°33'16.05''E) of the Rhodano-Provençal Basin (Fig. 1C) presents marine deposits of lower to middle Miocene in age (Demarcq, 1970). This Sub-basin rests above a thick Mesozoic succession in a corridor and is separated from the main Rhodanian basin by the N040° Nîmes Fault (Reynaud et al., 2012). The sediments are organized into five lithostratigraphic formations (Greenish Molasse, Saint-Siffret Marls, Uzès Molasse, Blue Marls and Castillon Molasse, respectively) ranging from

the Burdigalian to the regional “Helvetian” stage (Langhian-Serravallian equivalent; Fig. 1C). Only “Helvetian” stage is presented here, which is in temporal continuity with the Carry-Le-Rouet outcrops. The Helvetian stage at Castillon-Du-Gard Sub-basin is organized into two distinct formations (Fm.): the blue marls at the base and the Castillon Molasse at the top. The blue marls Fm. varies laterally in thickness from a few meters to 70 m-thick. They consist mainly of lithoclasts composed of calcite, feldspar, glauconite, muscovite and iron oxide grain and planktonic and benthic foraminifera, echinoderms, bryozoans and mollusks (Reynaud et al., 2012). The Castillon Molasse is about 70 m-thick and is composed of a bryozoan and mollusk packstone-rudstone (Bryomol; Reynaud et al., 2012). It is subdivided into two sub-formations: the Vers Unit at the base and the Castillon Unit s.s at the top. The presence of glauconite and oxide of iron in the Vers unit gives a characteristic yellow color to the Bryomol packstone rich in pectenids encrusted by oysters, barnacles and bryozoans. Castillon Unit s.s is a Bryomol floatstone differs from the Vers Unit by the occurrence of green and red algae (Reynaud et al., 2012).

The depositional setting corresponds to an offshore environment for the Blue Marls, protected from currents and waves. The Castillon Molasse is interpreted as being deposited in a shallow tidal bay (Reynaud et al., 2012). The stability and homogeneity of the floro-faunal assemblages throughout the whole sedimentary succession are indicative of open marine conditions without major variation in salinity Reynaud et al., 2012).

Several hundred oyster shells were collected on about thirty different levels of the Carry-Le-Rouet section. All these oysters belong to the *Ostrea* genus and some have been determined to be *Ostrea fimbriata* species. A hundred oyster shell fragments were collected on about twenty different levels for the Castillon-Du-Gard section, unfortunately the sample size did not allow the species identification.

### 3. Materials and methods

#### 3.1. Preservation of the studied material

Each oyster shell was embedded in Araldite ® 2020 resin from Huntsman (Basel, Switzerland), sawed perpendicularly to growth lines along the major growth axis (Fig. 2A), and carefully polished. Polished sections were observed under cathodoluminescence microscopy (CL) to assess the preservation state of each shell. CL analyses were carried out on a luminoscope (ELM – 3R) coupled to LEICA MZ12 microscope surmounted by a camera Lumenera Infinity 3. Luminescent areas with large unorganized calcite crystals (mainly chalky areas that are more sensitive to recrystallization; Kirby et al., 1998; Lartaud et al., 2006; Ulmann et al., 2013; de Winter et al., 2018) and non-luminescent to weakly luminescent areas with identifiable seasonal patterns of growths (Fig. 2B) of each shell section were precisely mapped. The luminescent areas are representative of recrystallized calcite and have not been sampled for isotopic analyses. Only the non-luminescent to weakly luminescent parts of the shells, where a slight luminescence follows the seasonal patterns of growths (induced by natural incorporation of  $Mn^{2+}$  in calcitic structure; Langlet et al., 2006), have been sampled for  $\delta^{18}O$ ,  $\Delta_{47}$  and  $^{87}Sr/^{86}Sr$  analyses, according to recent works (Lartaud et al., 2010; Huyghe et al., 2015; Bougeois et al., 2016). The Carry-le-Rouet and Castillon-Du-Gard outcrops only encountered a shallow burial (<100 m; Reynaud et al., 2006; Hamon et al., 2013; Seibel and James, 2017) as evidenced by the absence of compaction traces and the absence of younger sediment deposition in the region. Such shallow burial depths (Choquette and James, 1990) ensures that the sediments have not encountered large overburden pressure or elevated temperatures. This argues for an absence of solid-state reordering in the analyzed bivalve shells and therefore a preservation of initial  $\Delta_{47}$  signal.

#### 3.2. Oxygen isotope analyses



For this study, 534 samples were recovered from 101 shells. For most of the shell fragments, 1 to 3 different powder sample have been collected, each of them by gathering carbonate powder along a transect perpendicular to the growth line of the individual shell, in order to average the  $\delta^{18}\text{O}$  value over at least part of the oyster life and minimize the variability within and between shells from a single stratigraphic level. For 6 of the largest oyster shells, a microsampling along a transect perpendicular to the growth lines was performed in the ligamental area (Fig. 2A) to assess the variability of isotopic signal during at least part of the bivalve life. For these shells, the  $\delta^{18}\text{O}$  value of each measurement along this transect is reported on Suppl. Table 1 in bold. The recovered powders were analyzed for its oxygen isotope composition : 30 to 40  $\mu\text{g}$  of calcite collected using a scalpel reacted with 100 % phosphoric acid at 70 °C using a KIEL IV Carbonate Device connected to a Delta V Plus Isotope ratio mass spectrometer at the Biogeosciences Laboratory (University of Burgundy, Dijon, France). Oxygen isotope compositions have been corrected from the instrumental drift and are reported in the standard delta notation relative to VPDB (Vienna PeeDee Belemnite) in ‰ by assigning a  $\delta^{18}\text{O}$  value of - 2.20 ‰ to NBS19. External reproducibility was better than  $\pm 0.08$  ‰ (1  $\sigma_{\text{E}}$ ) for oxygen isotopes based that on multiple analyses of NBS19 along the runs. Carbon isotopes (Suppl. Table 1) measured at the same time are also reported in the delta notation relative to VPDB in ‰ by assigning a  $\delta^{13}\text{C}$  value of + 1.95 ‰ to NBS19. Carbon isotope reproducibility is better than  $\pm 0.04$  ‰ (1  $\sigma_{\text{E}}$ ).

### 3.3. Clumped-isotope analyses

Carbonate clumped-isotope thermometry is based on the quantification of statistical anomalies in the abundance of doubly substituted isotopologues of carbonate groups (e.g.,  $^{13}\text{C}^{18}\text{O}^{16}\text{O}^{16}\text{O}^{2-}$ ). For thermodynamic reasons, the equilibrium abundance of  $^{13}\text{C}$ – $^{18}\text{O}$  bonds in a carbonate mineral increases as temperature decreases (e.g. Schauble et al., 2006), and this

isotopic signal may be preserved over geological time scales under favourable conditions (Passey and Henkes, 2012; Stolper and Eiler, 2015). By precisely measuring the abundance of the multiply-substituted, mass-47, isotopologues in the CO<sub>2</sub> produced by acid digestion of a carbonate sample it is thus possible to constrain its original crystallization temperature without making assumptions regarding the composition of parent waters (Ghosh et al., 2006; Eiler et al., 2009).

$\Delta_{47}$  analyses generally require larger amounts of material than conventional, “single-isotope” analyses, and their precision and accuracy are ultimately limited by ion counting statistics and standardization issues. Each analysis reported here required from 2 to 3.5 mg, and two or three replicate analyses per sample were performed (Table 2).

At Carry-Le-Rouet, nine of the larger shells already analyzed for  $\delta^{18}\text{O}$  were selected for  $\Delta_{47}$  measurements. For the largest oyster (CAR17-6-33-b), two sub-sample carbonate powders were collected separately from sections of the shell corresponding to minimum and maximum  $\delta^{18}\text{O}$  values, respectively. For the remaining shells, carbonate powder was collected from the entire preserved oyster section (foliated calcite). At Castillon-Du-Gard, due to the small size of the available bivalve fragments, each sample comprised carbonate from several shell fragments collected from the same stratigraphic level or from adjacent stratigraphic levels. Overall, samples were collected from 3 different stratigraphic levels, corresponding to levels 9–12 m, 13–17 m, and 19.5–20.5 m.

The 13 clumped-isotope samples (considered of 2 or 3 aliquots each) were analyzed at LSCE (Laboratoire des Sciences du Climat et de l'Environnement) using the equipment and procedures described by Peral et al. (2018). Carbonate powders were dissolved in a common, stirred, 104 % phosphoric acid bath at 90 °C for 15 minutes. After cryogenic removal of water, the evolved CO<sub>2</sub> passed through a Porapak Q column (50/80 mesh, 1 m length, 2.1 mm ID) held at –20 °C under helium 6.0 flow (25 mL/min). CO<sub>2</sub> was then quantitatively

recollected by cryogenic trapping, and transferred by gas expansion into an Isoprime 100 dual-inlet mass spectrometer equipped with six Faraday collectors ( $m/z$  44 to 49). Each analysis took about 3 hours, during which sample gas and working reference gas were allowed to flow from matching, 10 mL reservoirs into the source, through a pair of fused silica capillaries (65 cm length, 110  $\mu\text{m}$  ID). Every 20 minutes, gas pressures were adjusted to achieve a mass 44 current of 40 nA, with differences between sample and reference gas generally below 0.1 nA. Background currents were measured in all high-gain collectors ( $m/z$  45 to 49) before and after each pressure adjustment, with gas flowing into the source, and are found to strongly correlate with the mass 44 current. Background-corrected ion current values were processed using the IUPAC  $^{17}\text{O}$ -correction parameters (Brand et al., 2010) to compute  $\delta^{13}\text{C}_{\text{VPDB}}$ ,  $\delta^{18}\text{O}_{\text{VPDB}}$  and “raw”  $\Delta_{47}$  values for each analysis. The isotopic composition ( $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$ ) of our working reference  $\text{CO}_2$  was computed based on nominal  $\delta^{13}\text{C}_{\text{VPDB}}$  and  $\delta^{18}\text{O}_{\text{VPDB}}$  values for carbonate standard ETH-3 ( $\delta^{13}\text{C}_{\text{VPDB}} = 1.71\text{‰}$ ,  $\delta^{18}\text{O}_{\text{VPDB}} = -1.78\text{‰}$ , Bernasconi et al., 2018) and an oxygen-18 acid fractionation factor of 1.008176 (Das Sharma et al., 2002). Three carbonate standards, ETH-1 to ETH-3 (Meckler et al., 2014; Bernasconi et al., 2018), were then used to convert raw  $\Delta_{47}$  to “absolute”  $\Delta_{47}$  values, following the procedure described by Daëron et al. (2016). The nominal  $\Delta_{47}$  values for these standards are those recomputed by Bernasconi et al. (2018) using IUPAC parameters: 0.258 ‰ for ETH-1, 0.256 ‰ for ETH-2 and 0.691 ‰ for ETH-3. The overall external reproducibility (standard deviation) of clumped isotopes measurements for bivalve shells and standard is 14.6 ppm ( $n = 86$ ).

### *3.4. Strontium isotope analyses*

Twelve of the thirteen clumped-isotope samples were also analyzed for  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios. About 10 mg of carbonate powder were dissolved in a savillex beaker with 1 ml solution of 2N  $\text{HNO}_3$  on a hot plate at 70°C. The Sr was isolated from the matrix using

conventional Eichrom Sr-Spec columns. The strontium isotopic composition is measured using a Thermo Triton Plus thermal ionization mass spectrometers at the GET laboratory (Toulouse, France). The  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios were defined as the average of 150 measurements of ion intensities following the static multi-collection mode normalized to  $^{86}\text{Sr}/^{88}\text{Sr} = 0.1194$ . During the measurement period, the standard NBS 987 gave  $^{87}\text{Sr}/^{86}\text{Sr}$  values of  $0.710281 \pm 0.000008$  ( $2\sigma_D$ ,  $n = 4$ ). The accuracy of our measurements is verified against the measured ratios of international standards: 1) Jct-1 Giant Clam *Tridacna gigas* standard with a measured value of  $0.709175 \pm 0.000010$  ( $2\sigma_D$ ;  $n = 7$ ) and a certified value of  $0.709150 \pm 0.000050$  (Ohno and Hirata, 2007); 2) CAL-S carbonate standard with a measured value of  $0.706955 \pm 0.000008$  ( $2\sigma_D$ ;  $n = 4$ ) and a certified value of  $0.706924 \pm 0.000018$  (Yeghicheyan et al., 2014); and 3) IAPSO seawater standard with a measured value of  $0.709181 \pm 0.000003$  ( $2\sigma_D$ ;  $n = 4$ ) and a certified value of  $0.709179 \pm 0.000005$  (Nayar et al., 2014). All  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios measured on our samples were adjusted to the NBS 987 standard reference value of 0.710250.

## **4. Results**

### *4.1. Sedimentary study*

#### 4.1.1. Carry-Le-Rouet outcrops

In the Carry-Le-Rouet section, the transitional continental to shallow marine depositional environments develops following an E-W polarity. Four facies associations are organized in four main depositional environments: (1) continental to coastal plain; (2) upper shoreface; (3) restricted upper shoreface and (4) lower shoreface (Fig. 3A). Facies descriptions are summarized in Table 1 and environmental interpretations are given below.

##### 4.1.1.1. Continental to coastal plain environment

This environment groups facies A1 to A4 (Table 1). The presence of pedogenetic horizons with rhizoliths, rare plant remains, mottling and carbonate nodules indicate widespread subaerial, episodic sedimentation and soil formation. The marly deposits are crosscut by sandy to conglomerate channels that correspond to sediment discharges in a proximal alluvial plain (Miall, 1986). A1 is interpreted as a flood plain (Nury, 1990; Demory et al., 2011). The clast-supported channels (Table 1; Fig. 3B1 and 3B2) and lobe-shape conglomerates (A3 and A4) are interpreted as deltaic distributaries (Demory et al., 2011) or alluvial fan delta (Nemec and Steel, 1988). The composition of the conglomerates indicates several sources with a Nerthe (Barremian stage) carbonate origin for the monogenic ones and Paleozoic basement (Permo-Triassic radiolarites or quartzites) for the polygenic conglomerates (Demory et al., 2011). The presence of oysters encrusting the conglomerate boulders is indicative of at least partly marine conditions during the deposition of the distal alluvial fan system (Holmes, 1965), as oysters can be found in brackish or estuarine conditions.

#### 4.1.1.2. Marine environment: Coastal plain to outer-platform

This environment encompasses facies B1 to D2 (Table 1). B1 consists of bioturbated sandstones with cordon-rich conglomerates organized in planar stratifications. B1 is interpreted as a coastal shallow marine environment in an upper shoreface (Table 1; Fig. 3B3 and 3B4; Nemec and Steel, 1988). The fine to coarse sandstones (B2) with megaripples, Hummocky Cross-stratification (HCS) and convolute beds are interpreted as upper shoreface zone (Table 1; Borgovini, 2003; Demory et al., 2011). Facies C consists of fine siltstone to marls rich in mollusk (bivalves and gastropods), benthic and rare planktonic foraminifera as *Globigerinoides trilobus*, *Globigerinoides altiapertura* and *Catapsydrax dissimilis* and echinoderms, corals and bryozoans. The absence of high-energy sedimentary structures, the presence of marls to fine grain siltstones and the biotic association picture restricted marine

conditions in an upper shoreface environment (Table 1; Fig. 3B7 and 3B8). The carbonated facies D1 is mainly composed of broken and disarticulated bioclats indicating intense reworking and high-energy conditions in shoreface environment (Table 1; Nalin et al., 2010). The D2 is composed of coral carpets and patches surrounded by bioclastic packstones (Table 1; Fig. 3B5 and 3B6; Riegl and Piller, 2000). The coral buildups and the bioclastic limestones are interpreted as deposited in a shallow high-energy platform setting (Demory et al., 2011).

#### 4.1.2. Castillon-Du-Gard outcrop

The sedimentology and stratigraphy of Middle Miocene succession of Castillon-Du-Gard Sub-basin are revisited in Reynaud et al. (2012). In this study, the authors propose that the succession of shallow cool-water carbonates (e.g. Castillon Molasse, Langhian-Serravallian stages) alternates with offshore marls (e.g. Blue Marls, Burdigalian stage). Blue Marls, close to the contact with the Vers unit are grainier and characterized by wave ripples in sands and mollusk and foraminifera packstones (Molechfor; Reynaud et al., 2012). The biotic components composed of molechfor and bryozoans and the wave ripples testify to proximal offshore environment deposited in a low energy bay environment (Reynaud et al., 2012). Vers unit (part of Castillon Molasse sub-unit) shows an erosional surface with the blue marls that indicates decreasing accommodation space. This part of Castillon Molasse is a packstone/rudstone with abundant bivalves and bryozoans (Bryomol; Reynaud et al., 2012). This succession corresponds to shallow tidal deposits in a protected bay environment (Reynaud et al., 2012). Castillon unit (part of Castillon Molasse sub-unit) is a floatstone composed of bryozoan, mollusks and algae (Bryomol; Reynaud et al., 2012), interpreted as subtidal dunes in a sand sheet deposited in deeper marine environments than the Vers Unit. The top of this Castillon unit is a fine-grained packstone with bryozoans and mollusks

(Bryomol; Reynaud et al., 2012) infilling successive channels and is interpreted as a decreasing water-depth with no emersion (Reynaud et al., 2012).

#### 4.1.3. Age Model

The age model proposed for Carry-Le-Rouet is based on a biostratigraphic study (Oudet et al., 2010) and a magnetostratigraphic one (Demory et al., 2011). Based on these two dating methods, we propose ages for each stratigraphic level markers on Carry-Le-Rouet section. Reefal unit (UR1 at 20 meters) is estimated at 23.03 Ma and marks the Chattian-Aquitania boundary. Reefal unit (UR2 at 25 meters; Fig. 3A) is dated at 22.9 Ma. Turritellid-rich level (at 42 meters, Fig. 3) is dated at 22.27 Ma. Finally, two reefal units UR3 (at 61 meters; Fig. 3A) and UR4 (at 66 meters; Fig. 3A) are respectively dated at 21.01 Ma and 21.00 Ma. Tamaris unit has only been dated by biostratigraphy with an age of 18.10 Ma at the base and 16.50 Ma at the top (Fondecave-Wallez et al., 1986). According to the literature, this unit is overlaid by two Langhian bars (Anglada et al., 1988; Arnaud et al., 1988; Oudet et al., 2010). An age of 15.97 Ma (Burdigalian-Langhian boundary) is proposed for the base of these bars.

For Castillon-Du-Gard outcrop, transition between Uzès Molasse and Blue Marls (base of the section) is dated at 15.97 Ma (Burdigalian-Langhian boundary). The Langhian-Serravallian boundary (~ 13.82 Ma) is located within the Castillon Molasse (Reynaud et al., 2012), despite limitation concerning available datation, we propose a position between 24 and 28 meters.

#### *4.2. Bivalve oxygen isotope composition*

Along the sedimentary succession at Carry-Le-Rouet, the bivalve  $\delta^{18}\text{O}$  values display relatively large scattering of about 4 ‰ within the same stratigraphic level (Fig. 4; Suppl.

Table 1). Similar scatter amplitudes are found within single oyster shells (e.g., level 21A). Samples from the Chattian and Aquitanian stages yield  $\delta^{18}\text{O}$  values ranging between about -4 and 0 ‰. By contrast, samples from the Burdigalian stage yield more positive  $\delta^{18}\text{O}$  values ranging between about -3 and 1 ‰. The youngest Late Burdigalian level yields more negative  $\delta^{18}\text{O}$  values than the rest of the Late Burdigalian stage.

Analyses performed on the largest oyster shells reveal different patterns of intra-shell  $\delta^{18}\text{O}$  variability. Two shells present a sinusoidal signal with an amplitude of  $\delta^{18}\text{O}$  variations up to  $\sim 4$  ‰ (Fig. 5D and 5E). Two other shells present variations in  $\delta^{18}\text{O}$  values with an amplitude of about 3 ‰, that could recall a sinusoidal pattern with a truncation of the highest  $\delta^{18}\text{O}$  values (Fig. 5A and 5C). Two shells present a decrease in  $\delta^{18}\text{O}$  values down to a plateau, followed by an increase, with an amplitude of variation of 1 ‰ (Fig. 5F) and 3 ‰ (Fig. 5B). A last shell presents more erratic variations in  $\delta^{18}\text{O}$  values, within a  $\sim 1$  ‰ range (Fig. 5G). For hinges that were sampled with a high spatial resolution, we observe a decrease in the amplitude of the signal between the Aquitanian (CAR17-6-28-a; CAR17-6-31base-a; CAR17-6-33-b and CAR17-6-33-c) and the Burdigalian (CAR17-8-43-h and CAR17-8-43-i), concomitantly with an overall increase in  $\delta^{18}\text{O}$  values.

Similarly to the Carry-Le-Rouet section,  $\delta^{18}\text{O}$  data from the Castillon-Du-Gard section present a large dispersion (Fig. 4B), slightly higher than at Carry-Le-Rouet as it reaches about 5 ‰ in the Castillon Molasse (9 to 17 meters from the base of the section).  $\delta^{18}\text{O}$  values seem to increase at the top of the section, from average values of about -3.16 ‰ at about 20-22 m from the base of the section to average  $\delta^{18}\text{O}$  values around -1.57 ‰ at about 27-30 m from the base of the section.

#### 4.3. Bivalve clumped isotope analyses



The  $\Delta_{47}$  values were measured on 12 bivalves and range between 0.676 and 0.716 ‰ (Table 2). They are not constant over time but fluctuate along the Carry-Le-Rouet and Castillon-Du-Gard sections. The 5 analyzed Aquitanian samples present  $\Delta_{47}$  values comprised between  $0.6852 \pm 0.0093$  ‰ and  $0.705 \pm 0.0095$  ‰. One shell from the Aquitanian was large enough to provide two subsamples (CAR17-6-33-b), one sampled in the part yielding maximum  $\delta^{18}\text{O}$  values from the microsampling approach, and the other in the part yielding minimum  $\delta^{18}\text{O}$  values (Fig. 5D). Both subsamples give quite close  $\Delta_{47}$  values of  $0.6821 \pm 0.01$  ‰ and  $0.6823 \pm 0.0099$  ‰. The 3 analyzed Burdigalian samples yield lower values (p-value of 0.08 based on a T-test), ranging from  $0.6762 \pm 0.0093$  ‰ and  $0.6882 \pm 0.0093$  ‰. The 3 analyzed Langhian samples display the highest values of the studied interval, ranging from  $0.6968 \pm 0.0098$  ‰ and  $0.7137 \pm 0.0099$  ‰ (higher than the Burdigalian samples : p-value of 0.04 based on a T-test; and higher than the Aquitanian samples : p-value of 0.12 based on a T-test).

#### 4.4. $^{87}\text{Sr}/^{86}\text{Sr}$ ratios

$^{87}\text{Sr}/^{86}\text{Sr}$  ratios range between 0.708222 and 0.708816 (Table 3). The strontium ratio average for the Aquitanian is 0.708237, 0.708655 for the Late Burdigalian and 0.708830 for Langhian stage.  $^{87}\text{Sr}/^{86}\text{Sr}$  data increase about 600 ppm over the Aquitanian to Langhian time period.

## 5. Discussion

### 5.1. Intra-shell variability of oyster clumped and oxygen isotope composition

Modern oysters and other bivalves often present growth cessation during part of the year, when environmental conditions become stressful, due to extreme temperature, nutrient availability, or seawater oxygenation (Brown and Hartwick, 1988; Mitchell et al., 2000; Surge

et al., 2001; Schöne 2008; Ulmann et al., 2013; Huygues et al., 2019) and can also occur during reproduction (Clark et al., 1974, Paulet et al., 2006). The growth cessation can result in an incomplete seasonal record of environmental parameters, namely temperature and seawater  $\delta^{18}\text{O}$  ( $\delta^{18}\text{O}_{\text{sw}}$ ), archived in the carbonate  $\delta^{18}\text{O}$  signal of the shells (Ullmann et al., 2010). Growth cessation is apparent in shells CAR17-6-28-a and CAR17-6-31base-a (Fig. 5A and 5C), showing a truncated  $\delta^{18}\text{O}$  signal at maximum  $\delta^{18}\text{O}$  values. Therefore, only minimum amplitude of seasonal variations in environmental parameters can be obtained in these shells, along with temperature and  $\delta^{18}\text{O}_{\text{sw}}$  during summer (represented by minimum carbonate  $\delta^{18}\text{O}$  values, that are not truncated). We estimate that only 4 oysters may provide an estimation of the full amplitude of seasonal temperature and  $\delta^{18}\text{O}_{\text{sw}}$  variations (Fig. 5 B, D, E, F). Other shells either present a truncated signal at maximum  $\delta^{18}\text{O}$  values, or may not have lived long enough to record a complete seasonal cycle. Alternatively, it cannot be excluded that some shells (CAR17-8-43-I; Fig. 5G for instance) had a very low growth rate and that the sampling resolution was not accurate enough to allow to identify the seasonal pattern.

We have first estimated temperature variations using the standard approach based on carbonate  $\delta^{18}\text{O}$  only, using the Anderson and Arthur (1983) equation established for calcitic molluscs with the  $\delta^{18}\text{O}_{\text{sw}}$  reconstructed by Billups and Schrag (2002), of -0.66 ‰ for the Aquitanian, and of -0.94 ‰ for the Burdigalian. We have first considered a constant  $\delta^{18}\text{O}_{\text{sw}}$  throughout the year to calculate the amplitude of seasonal temperature variations ( $\Delta T$ ) from  $\delta^{18}\text{O}$  values for the 4 largest micro-sampled oysters devoid of truncation in the  $\delta^{18}\text{O}$  signal (Fig. 5, B, D, E and F). The obtained  $\Delta T$  with this approach range between 10 and 15°C for the Aquitanian (Fig. 5, B, D and E) and around 5°C for the Late Burdigalian (Fig. 5, F). This approach would thus show a  $\Delta T$  decrease of 5 to 10°C between the Aquitanian and the Late Burdigalian, along with a marked decrease in summer temperatures, of about 15°C. Pollen assemblages indicate a temperate climate characterized by humid winters and drier summers

in Europe during the Late Burdigalian (Jiménez-Moreno et al., 2010; Bruch et al., 2011). Such a climate would likely result in seasonal variations of local  $\delta^{18}\text{O}_{\text{sw}}$  as well, with a higher  $\delta^{18}\text{O}_{\text{sw}}$  during summer and a lower  $\delta^{18}\text{O}_{\text{sw}}$  during winter. Taking these suspected seasonal variations of  $\delta^{18}\text{O}_{\text{sw}}$  into consideration would lead to increased estimates of seasonal temperature amplitude variations compared to those calculated using a constant  $\delta^{18}\text{O}_{\text{sw}}$ .

By comparison, the seasonal temperature contrast calculated from  $\delta^{18}\text{O}$  variations measured within Late Burdigalian oyster shells from Central Europe (Harzhauser et al., 2011), also considering a constant  $\delta^{18}\text{O}_{\text{sw}}$  throughout the year, is much higher ( $\sim 10^\circ\text{C}$ ) as part of another study. Summer temperatures calculated by these authors are also quite high at  $\sim 28^\circ\text{C}$ , but close to the temperatures calculated from our Aquitanian oyster shells. Harzhauser et al. (2011) report winter temperatures at  $\sim 9\text{--}10^\circ\text{C}$  similar those obtained in this study for the Late Burdigalian stage. This seasonal difference between the Aquitanian and the Late Burdigalian may be explained by a microclimate on the Northwestern Mediterranean Sea margin during the Late Burdigalian, or by different seasonal variations in salinity between the two regions, as the oysters collected in Harzhauser et al. (2011) study come from estuarine environments where the salinity is around 23 ‰. The Carry-Le-Rouet and Castillon-Du-Gard latitude changed very little since the early Miocene at  $\sim 43^\circ\text{N}$  (Jolivet et al., 2006). At present, the annual water at ten meters water depth in the Gulf of Lion is around  $18^\circ\text{C}$  fluctuating between winter temperatures of  $13 \pm 1^\circ\text{C}$  and summer temperatures of  $23 \pm 2^\circ\text{C}$  (Data from Levitus94; Levitus and Boyer, 1994). This seasonal temperature variation of  $\sim 10^\circ\text{C}$  is quite similar to our estimates for the Aquitanian stage, although the temperatures calculated for this stage are globally higher. By contrast, the Late Burdigalian seasonal contrast estimates are much weaker compared to current one, with in addition cooler summer ( $\sim 10^\circ\text{C}$ ) and winter ( $\sim 15^\circ\text{C}$ ) temperatures. This differs significantly from increasing temperatures recorded worldwide at

497 this period that marks the initiation of the climate warming leading to the MMCO during the  
498 Langhian (Zachos et al., 2008; Jiménez-Moreno et al., 2010; Harzhauser et al., 2011).

499 An attempt was made to estimate seasonal seawater temperature contrast independently  
500 from variations in  $\delta^{18}\text{O}_{\text{sw}}$ , based on clumped isotopes. Using conventional  $\delta^{18}\text{O}$   
501 measurements, complete seasonal cycles in temperature and  $\delta^{18}\text{O}_{\text{sw}}$  may be identified in 4  
502 oysters (Fig. 5 B, D, E and F) allowing to identify which part of the shells mineralized during  
503 summer (minimum  $\delta^{18}\text{O}$ ) versus winter (maximum  $\delta^{18}\text{O}$ ), and these inferred seasonal extrema  
504 were then sampled for  $\Delta_{47}$  analyses. Unfortunately, due to the amount of material required per  
505  $\Delta_{47}$  analysis and the small size of shells, only one Aquitanian shell (CAR17-6-33-b, Fig. 5D)  
506 could be used, providing two sub-samples corresponding to the  $\delta^{18}\text{O}$  minimum and maximum,  
507 respectively (all other shells provided “bulk” shell material assumed to reflect weighted  
508 average annual temperatures, and the corresponding results are presented in section 5.2.1).

509 There is today a large body of literature pertaining to the calibration of clumped-isotope  
510 thermometry in biogenic and inorganic carbonates. Among other reasons, this is due to the  
511 lingering methodological issues which have so far hindered precise inter-laboratory  
512 comparisons (e.g., Petersen et al., 2019 and references therein). Here we opt for the recent  
513 calibration by Peral et al. (2018), for two main reasons. For one thing, that study was  
514 performed at LSCE using identical methods and instruments as those reported here. In  
515 addition, despite the Peral et al. 2018 calibration being based on foraminifera, published and  
516 unpublished results from LSCE yield a calibration function for modern oysters and pectens,  
517 which is virtually identical to that of Peral et al. (2018).

518 The two “seasonal” sub-samples (CAR17-6-33-b-S1 and -W1) yield  $\delta^{18}\text{O}$  values of -  
519 3.6 ‰ and -1.3 ‰ respectively (Table 2), consistent with those obtained through  
520 microsampling (Fig. 6, Areas 1-2). Somewhat surprisingly, the corresponding  $\Delta_{47}$ -derived  
521 temperatures (Fig. 6) are statistically indistinguishable ( $22.4 \pm 3.1$  °C versus  $22.4 \pm 3.1$  °C, 1

$\sigma_E$ ). If these temperatures are used with the  $\delta^{18}\text{O}$  values of the two shell subsamples and the equation of Anderson and Arthur (1983), the resulting  $\delta^{18}\text{O}_{\text{sw}}$  calculated are of -2.05 ‰ and 0.25 ‰ for the parts of the shell yielding minimum and maximum  $\delta^{18}\text{O}$  values, respectively. The absence of seasonal temperature variation appears unlikely considering the paleolatitude of the Carry-Le-Rouet site during the Aquitanian (~10°C seasonality at this latitude at present time), and considering independent estimates based on pollen data that point to the existence of seasonal variations around 20°C in atmospheric temperature in Europe (Akgün et al., 2007). This oyster shell yields the highest  $\Delta_{47}$ -derived temperatures for the Aquitanian, 3°C higher than the other analyzed shells (Table 2). As summer tend to be the period with the highest growing rate in oysters from subtropical to temperate settings, it is possible that the intra-shell  $\delta^{18}\text{O}$  record defined with the microsampling method actually reflects a much shorter time span of the oyster life than a year. The large variations in  $\delta^{18}\text{O}$  recorded within the oyster shell could then reflect punctual freshwater discharges within the same season. This sample has been recovered from level 33, located at the transition between the Brackish Rousset Unit and the Bioclastic Carry Unit (Demory et al., 2011). The presence of brackish conditions in the lower part of the Bioclastic Carry Unit is supported by the presence of wood fragments and reworked potamids, pointing to the existence of punctual freshwater discharges in the shallow marine deposits of the Bioclastic Rousset unit.

## *5.2. Miocene Climate Change in Mediterranean area*

### 5.2.1. Comparison between bivalve $\delta^{18}\text{O}$ values and benthic foraminifera $\delta^{18}\text{O}$

Bivalve  $\delta^{18}\text{O}$  values present a large dispersion within each stratigraphic level, typically around 4 ‰. As shown by the microsampling realized on sections of the largest oysters (Fig. 5) that presents one seasonality-cycle at most, most of the analyses performed on bivalve fragments likely represent a relatively short time of the bivalve life, shorter than a year and

547 even maybe than a season. Thus, part of this large dispersion is likely to arise from seasonal  
548 or intra-seasonal variations in temperature and  $\delta^{18}\text{O}_{\text{sw}}$ , as a bivalve fragment can have been  
549 built mostly during a winter while another fragment recovered from the same stratigraphic  
550 level may have formed mostly during summer. Due to this large dispersion, we applied a  
551 statistical treatment to the dataset in order to identify significant climatic trends in the signal,  
552 using the SiZer package (Sonderegger and Sonderegger, 2018) in the R software (R Core  
553 Team 2017). The « SiZer » statistical method (based on studying statistical Significance of  
554 ZERo crossings of smoothed estimates; Marron and Chaudhuri, 1998) has been widely used  
555 on paleoclimatic proxies such as carbonate  $\delta^{18}\text{O}$  values (increasing or decreasing) to detect  
556 significant climatic trend at different timescales (Weckstrom et al., 2006; Dera et al., 2011;  
557 Bougeault et al., 2017). This method is based on the construction of curves fitting time series  
558 using different level of smoothing (h). The first derivatives of each curve are simultaneously  
559 computed with their 95 % confidence intervals, allowing the signs of derivative estimated to  
560 be statistically tested. If the sign of the first derivative is different from 0, including entire  
561 95% confidence interval, then the corresponding trend shows a significant increase (for a  
562 positive sign of the first derivative) or a significant decrease (for a negative sign of the first  
563 derivative). These multiple tests are then represented as a map to identify the trend  
564 significance for each size of smoothing. When the result is significant, it is coded in blue if it  
565 is positive and in red if it is negative. On the contrary, it is coded in gray if it does not differ  
566 from 0 and white when data are not numerous enough to identify significant trends ( $n < 5$ ). A  
567 Sizer map can thus be viewed as the summary of a large number (hundreds) of hypothesis test  
568 results, with each pixel of the map representing the result of a hypothesis test where the null  
569 hypothesis corresponds to the 0 slope. More formally, these tests are obtained from  
570 confidence interval calculations using the row-wise method that is detailed in Hannig and  
571 Marron (2006).

In this study, SiZer maps have been generated using a bandwidth  $h$  comprised between 0.1 and 10 Myrs (Fig. 7). This statistical treatment applied on carbonate  $\delta^{18}\text{O}$  data highlights the existence of a long-term increase in  $\delta^{18}\text{O}$  values between 23 and 20 Ma followed by a long-term decrease between 18 and 15 Ma. A second increase of  $\delta^{18}\text{O}$  values is depicted as well during the Serravallian, but over shorter timescales of about 1 to 2 Myrs. Using the equation of Anderson and Arthur (1983) and a  $\delta^{18}\text{O}_{\text{sw}}$  (Table 2) published by Billups and Schrag (2002), as described in Section 5.1., these trends in  $\delta^{18}\text{O}$  values depicted by the SiZer map would correspond to (1) a cooling of about  $10^{\circ}\text{C}$ , from temperatures of about  $25^{\circ}\text{C}$  in the Chattian and earliest Aquitanian to about  $15^{\circ}\text{C}$  in the Late Burdigalian, (2) a warming of about  $12^{\circ}\text{C}$ , with temperatures up to  $27^{\circ}\text{C}$  on average during the Langhian, and (3) a shorter-time cooling of about  $4^{\circ}\text{C}$ , with temperatures down to  $23^{\circ}\text{C}$  during the Serravallian (Fig. 8A). Note that the  $\delta^{18}\text{O}$  signal only depicts the cooling initiation that likely continues further into the Serravallian.

Our new dataset recovered from shallow coastal environments presents some similarities with the evolution of benthic foraminifera  $\delta^{18}\text{O}$  ( $\delta^{18}\text{O}_{\text{bf}}$ ; Zachos et al., 2008; Fig. 7). From the Chattian to the Burdigalian, the  $\delta^{18}\text{O}$  increase highlighted by the SiZer map (Fig. 7) is coherent with the general increasing trend of  $\delta^{18}\text{O}_{\text{bf}}$  values from the latest Oligocene to the Burdigalian, although our dataset is not continuous enough to identify the shorter-time Mi-events as evidenced in former studies (Zachos et al., 2008; Mawbey and Lear, 2014; Beddow et al., 2016). The  $\delta^{18}\text{O}$  minimum identified in our dataset during the Langhian could well correspond to the MMCO (Zachos et al., 2008; Mawbey and Lear, 2013; Beddow et al., 2016, Goldner et al., 2014; Holbourn et al., 2015; Super et al., 2018) detected in the  $\delta^{18}\text{O}_{\text{bf}}$  dataset, while the  $\delta^{18}\text{O}$  increase at the top the Castillon-Du-Gard succession could correspond to the initiation of the MMCT (Zachos et al., 2001; Billups and Schrag, 2002; Shevenell et al., 2004).

In contrast, the amplitude of the  $\delta^{18}\text{O}$  variations is much larger in our bivalves recovered from coastal environments compared to the deep marine environment record. The Early Miocene cooling is expressed by an increase of 0.5 ‰ of the  $\delta^{18}\text{O}_{\text{bf}}$  in deep marine environment (Fig. 7) while the amplitude depicted by bivalve  $\delta^{18}\text{O}$  at Carry-le-Rouet is four times higher. Similarly, for the MMCO, the  $\delta^{18}\text{O}_{\text{bf}}$  decreases by about 1 ‰ but about 2.5 ‰ in our bivalve shells living in proximal environments. This amplitude difference is however expected as shallow marine environments record larger temperature variations than deep-sea environments during climate fluctuations. Both environments records variations in the ocean  $\delta^{18}\text{O}$  linked to the waning and waxing of polar ice-sheets. In shallow coastal environments, an additional source of carbonate  $\delta^{18}\text{O}$  variations is be related to the hydrological cycle, whose variability in time and space can impact local  $\delta^{18}\text{O}_{\text{sw}}$ . Sedimentary facies at Carry-Le-Rouet and Castillon-Du-Gard point to the persistence of an open marine depositional environment, except for the Tamaris section which presents sedimentary facies coherent with a more restricted bay environment (Fig. 3). Yet, even open marine environments are susceptible to be impacted by nearby runoff leading to salinity variations, especially in coastal environments. The sporadic occurrence of wood fragments and the low abundance of the fossils throughout the Tamaris section may point to nearby freshwater inputs. Such an impact of local seawater  $\delta^{18}\text{O}$  fluctuations is further supported by the quite high temperatures of up to 35-40°C (Fig. 8A), calculated from bivalve  $\delta^{18}\text{O}$  using the equation of Anderson and Arthur (1983) and the  $\delta^{18}\text{O}_{\text{sw}}$  of Billups and Schrag (2002) established for the global ocean, that most likely exceeded the tolerance of bivalves (Stenzel, 1971). Thus, we anticipate that part of the large variations recorded in the bivalve  $\delta^{18}\text{O}$  signal may be related to change in the local hydrological cycle in response to global climate change occurring at this period.

#### 5.2.2. Evolution of shallow seawater temperatures inferred from clumped isotope analyses



“Bulk”  $\Delta_{47}$  analyses, representing averaged isotopic composition over at least part of the life span of the organisms (all shells except CAR17-6-33-b) could help decipher whether the bivalve  $\delta^{18}\text{O}$  signal is related to variations in seawater temperature and  $\delta^{18}\text{O}_{\text{sw}}$ . The  $\Delta_{47}$ -derived shallow seawater temperatures “ $T_{47}$ ” (Table 2; Fig. 8A) range from  $15.6 \pm 2.7^\circ\text{C}$  to  $22.4 \pm 3.1^\circ\text{C}$  ( $1 \sigma_{\text{E}}$ ), with an average of  $19.7 \pm 1.1^\circ\text{C}$  ( $1 \sigma_{\text{E}}$ , fully propagated from the analytical errors on individual samples) which is cooler (p-value of 0.003 using a T-test) than those calculated with the  $\delta^{18}\text{O}_{\text{shell}}$  and a  $\delta^{18}\text{O}_{\text{sw}}$  of Billups and Schrag (2002) for the Aquitanian (about  $25^\circ\text{C}$  on average). Previous studies on pollen assemblages in Europe (Akgün et al., 2007) and of coral assemblages in the Mediterranean Sea (Bosellini and Perrin, 2008), argue for atmospheric temperatures between  $16.5$  and  $21.3^\circ\text{C}$  and sea surface temperatures between  $18$  and  $19.3^\circ\text{C}$ , respectively, during the Aquitanian stage. These values are in good agreement with the temperatures calculated from our  $\Delta_{47}$  analyses.

For the Late Burdigalian stage, shallow seawater  $T_{47}$  values are higher (Table 2; Fig. 8A) than those for the Aquitanian (p-value of 0.048 using T-test), ranging between  $20.5 \pm 2.8^\circ\text{C}$  and  $24.3 \pm 2.9^\circ\text{C}$  ( $1 \sigma_{\text{E}}$ ) and with an average of  $22.6 \pm 1.7^\circ\text{C}$  ( $1 \sigma_{\text{E}}$ ). These Late Burdigalian temperatures calculated from  $\Delta_{47}$  data are higher (p-value of 0.021 using a T-test) than those calculated from bivalve  $\delta^{18}\text{O}$  using a  $\delta^{18}\text{O}_{\text{sw}}$  from Billups and Schrag (2002), by about  $15^\circ\text{C}$  on average, but are again in better agreement with those published in the literature (Lecuyer et al., 1996; Goedert et al., 2017). Indeed, with the exception of the slightly cooler temperatures deduced from coral assemblages for the Late Burdigalian ( $18.5$  to  $20.5^\circ\text{C}$ ; Bosellini and Perrin, 2008), our temperature range is quite close to that deduced from combined analyses of marine mammal and fish tooth  $\delta^{18}\text{O}$ , that is comprised between  $20$  and  $28^\circ\text{C}$  (Goedert et al., 2017) in the northern Aquitaine Basin, and between  $18$  and  $22^\circ\text{C}$  in Brittany (Lecuyer et al., 1996).

For the Langhian stage,  $T_{47}$  values range from  $12.5 \pm 2.8^{\circ}\text{C}$  to  $18 \pm 2.9^{\circ}\text{C}$  ( $1 \sigma_E$ ) with an average of  $14.5 \pm 1.6^{\circ}\text{C}$  ( $1 \sigma_E$ ). Again, these Langhian temperatures derived from  $\Delta_{47}$  data are cooler (p-value of 0.026 using a T-test), as for the Aquitanian (Table 2; Fig. 8A), than those calculated from bivalve  $\delta^{18}\text{O}$  using a  $\delta^{18}\text{O}_{\text{sw}}$  from Billups and Schrag (2002), of about  $25^{\circ}\text{C}$  on average. These values might appear cooler than expected, considering that this period encompasses the MMCO for which both terrestrial and marine proxy data suggest globally warmer conditions relative to the Early Miocene, especially at mid- and high latitudes (Mosbrugger et al., 2005; Bruch et al., 2007; Bruch et al., 2011; Shevenell et al., 2008; Zachos et al., 2008; Super et al., 2018). On the other hand, our clumped-isotope results are coherent with atmospheric temperatures inferred from pollen assemblages on the northwestern Mediterranean margin ( $17\text{--}18^{\circ}\text{C}$ , Bruch et al., 2007), and in Germany ( $16\text{--}21^{\circ}\text{C}$ , Böhme et al., 2007). Published sea surface temperatures inferred from combined  $\delta^{18}\text{O}$  analyses of mammals and fish remains, are around  $20^{\circ}\text{C} \pm 2^{\circ}\text{C}$  in Brittany (Lecuyer et al., 1996), slightly higher than our  $\Delta_{47}$ -derived temperatures. Additional estimates, calculated from bivalve  $\delta^{18}\text{O}$ , range between  $18$  and  $28^{\circ}\text{C}$  in the Tethyan domain (Harzhauser et al., 2011) and between  $16.5$  and  $19.5^{\circ}\text{C}$  from coral assemblage (Bosellini and Perrin, 2008). Although the uncertainties associated with the calculation of all these temperatures are difficult to estimate, they all tend to appear quite lower than the  $27\text{--}37^{\circ}\text{C}$  range obtained from  $\text{TEX}_{86}$  analyses at similar latitude in the central Atlantic (Super et al., 2018), similarly to our  $\Delta_{47}$ -derived temperature estimates. Our new temperature data inferred from  $\Delta_{47}$  analyses are thus coherent with the temperatures published in the literature for the Mediterranean and Tethyan regions and support lower temperatures in this general area during the MMCO compared to other sites at similar paleolatitude.

A relatively cool local to regional climate during the MMCO, compared to that inferred from similar paleolatitudes based on  $\text{TEX}_{86}$  data, is also consistent with the high abundance of

bryozoans throughout the Castillon-Du-Gard section. The proliferation of bryozoans, mainly in the form of encrusting cheilostomes, colonized the shallow seawater setting of the Rhodano-Provençal Sub-basins and has been interpreted as deposited in temperate to warm temperate depositional environment (Taylor and James, 2013). Similar bryozoan-rich deposits in the Tethyan domain have been interpreted as cool-water carbonates by Randazzo et al. (1999), based on  $\delta^{18}\text{O}$  analyses and ecological reconstitutions. So far, such lower seawater temperatures in the Mediterranean and Tethyan regions have been tentatively explained by the deep cold-water arrivals coming from development of upwellings (Randazzo et al., 1999; Reuter et al., 2013).

Thus the temperatures obtained from  $\Delta_{47}$  analyses are in relatively good agreement with those published in the literature in the general region, but quite markedly differ from those obtained from bivalve  $\delta^{18}\text{O}$  using the equation of Anderson and Arthur (1983) and the  $\delta^{18}\text{O}_{\text{sw}}$  of the global ocean of Billups and Schrag (2002). This apparent discrepancy between  $\Delta_{47}$ -derived and  $\delta^{18}\text{O}$ -derived temperatures, observed for the different studied periods, can be explained by modifications of the hydrological cycle linked to global climate change, leading to a departure of the local  $\delta^{18}\text{O}_{\text{sw}}$  from that of the global ocean estimated by Billups and Schrag (2002).

### 5.2.3. Variations in the local hydrological cycle

In order to estimate the temporal evolution of the local  $\delta^{18}\text{O}_{\text{sw}}$  we used the seawater temperatures inferred from  $\Delta_{47}$  analyses together with the  $\delta^{18}\text{O}$  values of the oyster shells measured on the same powder sample, and the Anderson and Arthur (1983) equation.

Our calculated values point to low local  $\delta^{18}\text{O}_{\text{sw}}$  values during the Aquitanian (values ranging from -3.9 to -1.06 ‰ with an average of -1.89 ‰), that are likely to reflect enhanced precipitations and a lower salinity of the shallow neritic waters during this period (Table 2; Fig. 8B). For the late Burdigalian, the calculated  $\delta^{18}\text{O}_{\text{sw}}$  are much higher (p-value of 0,021

696 using a T-test), ranging from 0.9 to 1.71 ‰ with an average of 1.27 ‰, that points to a  
697 decrease in precipitation and runoff and/or an increase in evaporation (Table 2; Fig. 8B). This  
698 finding agrees well with the display of a low floro-faunal content in the Late Burdigalian of  
699 the Tamaris succession, an absence of high-energy sedimentary structures and a deposition of  
700 fine-grain sediments, suggesting that the depositional environment was potentially more  
701 restricted. Our calculated  $\delta^{18}\text{O}_{\text{sw}}$  values for the Late Burdigalian are close although slightly  
702 lower than the estimation of about 2 ‰ on average for coastal seawater in the nearby  
703 Aquitaine Basin during the Late Burdigalian by Goedert et al. (2017), based on combined  
704  $\delta^{18}\text{O}$  analyses of marine mammals and fish remains. The local  $\delta^{18}\text{O}_{\text{sw}}$  values for the Langhian  
705 are again lower (ranging from -3.3 to -2.52 ‰, with on average of -3.02 ‰), which would  
706 point to enhanced precipitations and runoff during this period (Table 2; Fig. 8B). These values  
707 are significantly lower than those calculated for the Burdigalian (p-value of 0.0002 using a T-  
708 test). The Castillon-Du-Gard Sub-basin is located on the periphery of the main Rhodano-  
709 Provencal Basin, with the presence of nearshore facies (Bryomol) in the Castillon-Du-Gard  
710 area (Reynaud et al., 2012), indicative of a semi-enclosed basin, and of sediments organized  
711 in a regressive sequence. These conditions would further favor increased freshwater inputs  
712 into this Sub-basin in the case of an enhanced hydrological cycle. In addition, a tectonically-  
713 enhanced uplift of Alps chain occurred between the lower and the middle Miocene, that can  
714 induce changes in the drainage pattern, major fluvial incisions, and increased runoff in the  
715 Castillon-Du-Gard Sub-basin (Reynaud et al., 2012). In any case, following this coupled  $\Delta_{47}$  -  
716  $\delta^{18}\text{O}$  approach, our results would then point to the existence of changes in the local  
717 hydrological cycle during the studied period, from the early to the middle Miocene.  
718 Importantly, changes in local  $\delta^{18}\text{O}_{\text{sw}}$ , likely driven by changes in freshwater inputs, dominate  
719 the recorded changes in bivalve  $\delta^{18}\text{O}$ , while temperature changes appear to have a more minor  
720 influence (Fig. 9A and 9B).

Part of the local  $\delta^{18}\text{O}_{\text{sw}}$  evolution reconstructed from the coupled  $\Delta_{47}$  and  $\delta^{18}\text{O}$  analyses in this study is due to changes in the ocean  $\delta^{18}\text{O}_{\text{sw}}$  at a global scale, linked to variations in ice-sheet volume on Antarctica at that time. In an attempt to estimate first-order variations in local salinity due to changes in the local hydrological cycle only, we subtracted to our calculated local  $\delta^{18}\text{O}_{\text{sw}}$  the ocean  $\delta^{18}\text{O}_{\text{sw}}$  (Fig. 8C) values published in Billups and Schrag (2002), i.e.  $-0.66\text{‰ VPDB}$  (or  $-0.86\text{‰ VSMOW}$ ; see Grossmann (2012) for conversion from the VPDB to the VSMOW scale) for the Aquitanian,  $-0.94\text{‰ VPDB}$  (or  $-1.14\text{‰ VSMOW}$ ) for the Burdigalian, and  $-0.71\text{‰ VPDB}$  (or  $-0.91\text{‰ VSMOW}$ ) for the Langhian. We then used salinity -  $\delta^{18}\text{O}_{\text{sw}}$  relations established in modern environments but still comparable to the sites investigated here. We chose for this rough approach the equation proposed by Railsback and Anderson (1989) based on modern surface North Atlantic Ocean salinity and  $\delta^{18}\text{O}_{\text{sw}}$  data. A salinity -  $\delta^{18}\text{O}_{\text{sw}}$  relation exists for the modern Mediterranean Sea (Pierre, 1999). However, during the Miocene the Mediterranean Sea was still largely open to the Atlantic Ocean on the west, and to the Indian Ocean on the east prior to its closure during the late Miocene (Jolivet et al., 2006; Do Couto et al., 2016). Therefore, we think that this configuration of the Mediterranean Sea during the Miocene, quite different than the modern one, likely impacted the salinity -  $\delta^{18}\text{O}_{\text{sw}}$  relation. To our opinion, it is thus preferable to consider the equation of Railsback and Anderson (1989) based on surface North Atlantic data. Note that if the equation of Pierre (1999) had been used instead, the estimated salinities presented below would be of about 20 to 31‰ for the Aquitanian, about 40 to 43‰ for the Late Burdigalian and about 22 to 25‰ for the Langhian.

For the Aquitanian stage, the reconstructed salinities based on this approach tend to be lower on average than the mean value of 35‰ of the modern open ocean, with an important variability in the 30 and 34‰ range based on bulk shells, i.e an average salinity throughout the bivalve life (Table 2; Fig. 8D). This points to a contribution of freshwater inputs to the

local seawater by runoff, rainfall, or groundwater discharge, although this range remains within the tolerance of the stenohalin organisms (corals, bryozoans, echinoids and foraminifera) observed at Carry le Rouet during the Aquitanian (Oertli, 1964). The Sr isotope composition of bivalve shells can provide additional insights on potential freshwater inputs, if they are large enough to impact the local seawater Sr budget. The strontium isotope composition ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) of bio-carbonates reflects the isotopic composition of the water in which they form their test (Veizer et al., 1999; Palmer and Elderfield, 1985). Therefore, the  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios measured on carbonated organisms enable to reconstruct the isotopic signature of the aqueous environment in which they thrive.

The measured  $^{87}\text{Sr}/^{86}\text{Sr}$  of the bivalves compares well with the Sr isotope signatures of planktonic foraminifera and of mixed calcitic fossils from the Umbria-March Basin (Fig. 8E; Kocsis et al., 2008; Cornacchia et al., 2018). Our Sr isotope record mostly falls within the global seawater Sr isotope reference curve (McArthur and Howarth, 2004), when considering a seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  variability comparable to that recorded in the modern ocean ( $\pm 50$  ppm; El Meknassi et al., 2018). For the Aquitanian, the  $^{87}\text{Sr}/^{86}\text{Sr}$  of our samples tends to fall at the lower limit of this range, and even presents slightly more unradiogenic values than that of the open ocean at that time, as observed from the nearby Umbria-March Basin (Kocsis et al., 2008; Fig. 8E). This slight deviation from the open ocean Sr isotope composition may derive from active volcanism in the region, that can release mantle-derived unradiogenic Sr signature into the local seawater and influence its Sr budget. Sporadic volcanic activity during the Miocene in the western Mediterranean Sea can indeed be generated during the extension of the Tyrrhenan and Algéro-Provencal basins (Frizon de Lamotte et al., 2000; Schettino and Turco, 2011; Bialik et al., 2019). Large eruptions within the 25-19 Ma interval are supported by volcanoclastic sediments in the vicinity of the studied region (from Provence to Sardinia and Corsica and in the Umbria-March Basin ; Montanari et al., 1994; Beccaluva et al., 2004),

and by radiogenic  $\epsilon_{\text{Nd}}$  and unradiogenic  $^{87}\text{Sr}/^{86}\text{Sr}$  values of sediments from the Umbria-March region (Kocsis et al., 2008).

Alternatively, it has been shown that large riverine inputs can also result in a deviation of the Sr isotope composition of neritic seawater from that of the global ocean (Bryant et al., 1995; El Meknassi et al., 2018). At Carry-le Rouet, the Miocene deposits unconformably rest on marine carbonates from the Lower Cretaceous (Barremian to Lower Aptian), that are characterized by low  $^{87}\text{Sr}/^{86}\text{Sr}$  values (typically within the 0.7073 to 0.70745 range; McArthur and Howarth, 2004). Enhanced freshwater input should then result in a decrease of local seawater  $^{87}\text{Sr}/^{86}\text{Sr}$ , if large enough to impact the local Sr budget. Bryant et al. (1995) has shown that a measurable effect occurs at salinities below 12 ‰. Salinities calculated from our coupled  $\delta^{18}\text{O}-\Delta_{47}$  approach lay however in the 30 to 34 ‰ range for this interval at Carry-le-Rouet, and the impact of freshwater inputs on the local seawater Sr isotope composition should then have remained limited. Yet recent studies reported that deviation from the global ocean  $^{87}\text{Sr}/^{86}\text{Sr}$  value can occur at higher salinities, above 30 ‰ (Huang et al., 2011), especially in environments affected by submarine groundwater discharges that can have elevated Sr concentrations when deriving from carbonate-rock aquifer systems (Trezzi et al., 2017; El Meknassi et al., 2018). Thus enhanced riverine inputs may also have contributed to lower the  $^{87}\text{Sr}/^{86}\text{Sr}$  value of the local seawater at Carry-le-Rouet.

Conversely, for the Late Burdigalian, the reconstructed salinities are much higher (p-value of 0.0014 using a T-test), in the 38 to 39 ‰ range (Table 2; Fig. 8D). This would correspond to a hypersaline environment with reduced runoff and high evaporation, in agreement with the higher temperatures inferred from our  $\Delta_{47}$  analyzes in the Late Burdigalian compared to the Aquitanian. Such high salinities are consistent with the disappearance of stenohalin organisms and the restricted environmental conditions recorded at the Tamaris section. High evaporation should not affect  $^{87}\text{Sr}/^{86}\text{Sr}$  values of seawater, which is coherent

with the  $^{87}\text{Sr}/^{86}\text{Sr}$  values recorded by our bivalves, mostly within the expected range of signature for the open ocean at that time.

The calculated salinities for the Langhian stage lay between 30 and 32 ‰ (Table 2; Fig. 8D), significantly lower than those calculated for the Burdigalian (p-value of 0.00002, using a T-test). Combined with the quite cool temperatures deduced from  $\Delta_{47}$  for this period, this suggests again a marine environment subjected to freshwater inputs. A cooler and slightly brackish environment during the MMCO at Castillon-Du-Gard would agree with the proliferation of bryozoans and associated algae in these levels. Similar facies association is described in middle Miocene marine sediments from Austria and considered as deposited in a temperate environment with low salinity (Friebe, 1994). Modern bryozoan ecological reconstructions, occupy a wide variety of environments and can tolerate salinities ranging from 32 to 37 ‰ (Ryland, 1970; Smith, 1995) and temperatures from 10 to 30°C (Domack, 1988), that are within the temperature and salinity ranges inferred from our coupled  $\Delta_{47}$  -  $\delta^{18}\text{O}$  approach at Castillon-Du-Gard.

The Sr isotope composition of Langhian bivalves from the Castillon-du-Gard section fall within the Sr isotope reference curve for the open ocean at that time (Fig. 8E), although our coupled  $\delta^{18}\text{O}$ - $\Delta_{47}$  approach points to salinities comparable to that obtained for the Aquitanian. Deviations toward more radiogenic values at that time were reported from mixed fossils from sites closer to the Alps, as exposure of older continental rocks contributes to imprint river waters with a radiogenic Sr isotope composition (Kocsis et al., 2008; 2009). Surface runoff issued from watersheds in Western Europe exhibits at present quite radiogenic Sr isotope signature (of 0,7093 on average ; Peucker-Ehrenbrink and Fiske, 2019). The Castillon-du-Gard section, located north to the Carry-le-Rouet section within a narrow and incised valley, could have received a larger contribution from river waters draining the Alps watersheds, especially in the context of the tectonically-enhanced uplift of the Alps during the Miocene



(Frisch et al., 2000; Reynaud et al., 2012). Rivers draining old continental material tend to yield much lower Sr concentrations than rivers or groundwaters draining carbonate lithologies. It is thus possible that the enhanced freshwater inputs during the MMCO, inferred from the salinities calculated from the  $\delta^{18}\text{O}$  and  $\Delta_{47}$  values, were not large enough at Castillon-du-Gard to significantly shift the local seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio.

## 6. Conclusions

Our multi-proxy approach based on both  $\delta^{18}\text{O}$  and  $\Delta_{47}$  analyses of bivalves recovered from two outcrops in southern France during the Miocene highlight a warming ( $\sim 3^\circ\text{C}$ ) of the North-West Mediterranean coastal marine environments between the Aquitanian and the Late Burdigalian, accompanied by an increase in local salinity probably linked to more restricted conditions. Surprisingly, low  $\delta^{18}\text{O}$  values recorded in the Langhian, supposedly during the MMCO, are associated with cooler seawater temperatures (between  $13 - 18^\circ\text{C}$ ) inferred from  $\Delta_{47}$  analyzes. This apparent discrepancy can be reconciled if a modification occurred in the local hydrological cycle linked to the overall warmer conditions recorded at a global scale during this event, leading to enhanced discharges of freshwaters in the region during the MMCO. The existence of relatively cool and low salinity local waters would be consistent with the development of an abundant bryozoan fauna observed at that time in the studied section. Local freshwater discharges at the regional scale could also have been promoted by change in the pattern of the drainage system driven by local tectonic uplift and a narrowing of the corridor structure of the Rhodano-Provencal. The Sr isotope analyses performed on the bivalves from the two studied sections are compatible with the existence of freshwater inputs during the Aquitanian and the Langhian in this region. These inputs were however not large enough to result in large deviations of the local seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  composition from that of the global ocean, which is consistent with the reconstructed salinities, laying in the 30-34 ‰

range. Importantly, our results illustrate the risks inherent to paleotemperature reconstructions based solely on  $\delta^{18}\text{O}$  records and highlight the pertinence of coupling several paleoenvironmental and paleoclimatic proxies, geochemical ( $\delta^{18}\text{O} - \Delta_{47}$ ) or sedimentological, to reconstruct the variations of seawater temperature and salinity in shallow coastal environments. The combined  $\delta^{18}\text{O} - \Delta_{47}$  approach thus appears promising to study the local response to global climatic variations in environments that are subjected to salinity variations (lagoonal, mangrove or estuarine environments). Understanding how such environments react to climate change would help to better constrain the behavior of our climate system. Our work highlights the importance of a continuous effort in gathering new data from different regions of the world during MMCO, to capture the diversity of local responses to a global climatic change.

### Acknowledgements

We wish to thank T. COCQUEREZ (Biogeosciences Laboratory, University of Burgundy) for the isotopic analyses. We wish also to thank F. MONNA and R. LAFFONT (Biogeosciences Laboratory, University of Burgundy) for help with statistical treatment of our data. This study was funded by the ANR AMOR, coordinated by Y. DONNADIEU.

### References

Akgün, F., Kayseri, M. S., and Akkiraz, M. S. 2007. Palaeoclimatic evolution and vegetational changes during the Late Oligocene–Miocene period in Western and Central Anatolia (Turkey). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 253(1), 56-90. <https://doi.org/10.1016/j.palaeo.2007.03.034>.

870 Al-Aasm, I. S., and Veizer, J. 1986. Diagenetic stabilization of aragonite and low-Mg calcite;  
871 I, Trace elements in rudists. *Journal of Sedimentary Research*, 56(1), 138-152.  
872 <https://doi.org/10.1306/212F88A5-2B24-11D7-8648000102C1865D>.

873 Anglada, R., Gourinard, Y., Lesueur, J. L., and Rubino, J.-L., 1988. Les séquences de dépôt  
874 du Burdigalien et du Langhien de Basse Provence (Littoral de la Nerthe et région des  
875 Etangs). *Livret Guide d'Excursion Coll. « Paléobathymétrie, eustatisme et séquence*  
876 *de dépôts »*, Marseille, juin 1988: 34p.

877 Anderson, T. F., and Arthur, M. A. 1983. Stable isotopes of oxygen and carbon and their  
878 application to sedimentologic and paleoenvironmental problems. *Unknown Journal*.

879 Andreieff, P., Anglada, R., Carbonnel, G., Catzigras, F., Cavelier, C., Chateauneuf, J. J.,  
880 Colomb, E., Jacob, C., Lai, J., l'Hommer., A., Lezard, L., Lorenz, C., Mercier, H., and  
881 Parfenoff, A. 1972. Contribution à l'étude de Carry-le-Rouet (Bouches-du-Rhône).  
882 Cinquième congrès du Néogène Méditerranéen, *Mémoire du BRGM*, 3, 132.

883 Arnaud, M., Barbaroux, L., Catzigras, F., Colomb, E., Monleau, C., 1988. Le Burdigalien et  
884 la Langhien de Provence occidentale (Bouches-du-Rhône): nouvelles données  
885 stratigraphiques et sédimentologiques. *Comptes Rendus de l'Académie des Sciences*  
886 307(II): 779-784.

887 Beccaluva, L., Coltorti, M., Giunta, G., and Siena, F. 2004. Tethyan vs. Cordilleran  
888 ophiolites: a reappraisal of distinctive tectono-magmatic features of supra-subduction  
889 complexes in relation to the subduction mode. *Tectonophysics*, 393(1-4), 163-174.  
890 <https://doi.org/10.1016/j.tecto.2004.07.034>.

891 Beddow, H. M., Liebrand, D., Sluijs, A., Wade, B. S., and Lourens, L. J. 2016. Global change  
892 across the Oligocene-Miocene transition: High-resolution stable isotope records from  
893 IODP Site U1334 (equatorial Pacific Ocean). *Paleoceanography*, 31(1), 81-97.  
894 <https://doi.org/10.1002/2015PA002820>.

895 Bernasconi, S. M., Müller, I. A., Bergmann, K. D., Breitenbach, S. F., Fernandez, A., Hodell,  
 896 D. A., Jaggi, M., Nele-Meckler, A., Millan, I., and Ziegler, M. 2018. Reducing  
 897 uncertainties in carbonate clumped isotope analysis through consistent carbonate-  
 898 based standardization. *Geochemistry, Geophysics, Geosystems*, 19(9), 2895-2914.  
 899 <https://doi.org/10.1029/2017GC007385>.

900 Bialik, O. M., Frank, M., Betzler, C., Zammit, R., and Waldmann, N. D. 2019. Two-step  
 901 closure of the Miocene Indian Ocean Gateway to the Mediterranean. *Scientific*  
 902 *Reports*, 9(1), 8842.<https://doi.org/10.1038/s41598-019-45308-7>.

903 Billups, K., and Schrag, D. P. 2002. Paleotemperatures and ice volume of the past 27 Myr  
 904 revisited with paired Mg/Ca and  $^{18}\text{O}/^{16}\text{O}$  measurements on benthic foraminifera.  
 905 *Paleoceanography*, 17(1), 3-1. <https://doi.org/10.1029/2000PA000567>.

906 Böhme, M., Bruch, A. A., and Selmeier, A. 2007. The reconstruction of Early and Middle  
 907 Miocene climate and vegetation in Southern Germany as determined from the fossil  
 908 wood flora. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 253(1), 91-114.  
 909 <https://doi.org/10.1016/j.palaeo.2007.03.035>.

910 Borgovini, R. 2003. Stratigraphie et facies de la base du cycle Néogène de Basse Provence  
 911 (littoral de la Nerthe : Carry-Le-Rouet, Sausset-Les-Pins). Rapport Total, Exploration  
 912 and Production, DGEP/GSR/TG/THEM N°03-006 : 134.

913 Bosellini, F. R., and Perrin, C. 2008. Estimating Mediterranean Oligocene–Miocene sea-  
 914 surface temperatures: an approach based on coral taxonomic richness.  
 915 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 258(1-2), 71-88.  
 916 <https://doi.org/10.1016/j.palaeo.2007.10.028>.

917 Bougeault, C., Pellenard, P., Deconinck, J. F., Hesselbo, S. P., Dommergues, J. L., Bruneau,  
 918 L., Cocquerez, T., Laffont, R., Huret, A., and Thibault, N. 2017. Climatic and  
 919 palaeoceanographic changes during the Pliensbachian (Early Jurassic) inferred from

920 clay mineralogy and stable isotope (CO) geochemistry (NW Europe). *Global and*  
921 *Planetary Change*, 149, 139-152. <https://doi.org/10.1016/j.gloplacha.2017.01.005>.

922 Bougeois, L., De Rafélis, M., Reichart, G. J., De Nooijer, L. J., Nicollin, F., and Dupont-  
923 Nivet, G. 2014. A high resolution study of trace elements and stable isotopes in oyster  
924 shells to estimate Central Asian Middle Eocene seasonality. *Chemical Geology*, 363,  
925 200-212. <https://doi.org/10.1016/j.chemgeo.2013.10.037>.

926 Bougeois, L., De Rafélis, M., Reichart, G. J., de Nooijer, L. J., and Dupont-Nivet, G. 2016.  
927 Mg/Ca in fossil oyster shells as palaeotemperature proxy, an example from the  
928 Palaeogene of Central Asia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 441,  
929 611-626. <https://doi.org/10.1016/j.palaeo.2015.09.052>.

930 Brand, U., and Veizer, J. 1981. Chemical diagenesis of a multicomponent carbonate system;  
931 2, Stable isotopes. *Journal of Sedimentary Research*, 51(3), 987-997.  
932 <https://doi.org/10.1306/212F7DF6-2B24-11D7-8648000102C1865D>.

933 Brand, W. A., Assonov, S. S., and Coplen, T. B. 2010. Correction for the  $^{17}\text{O}$  interference in  $\delta$   
934 ( $^{13}\text{C}$ ) measurements when analyzing  $\text{CO}_2$  with stable isotope mass spectrometry  
935 (IUPAC Technical Report). *Pure and Applied Chemistry*, 82(8), 1719-1733.  
936 <https://doi.org/10.1351/PAC-REP-09-01-05>.

937 Brown, J. R., and Hartwick, E. B. 1988. Influences of temperature, salinity and available food  
938 upon suspended culture of the Pacific oyster, *Crassostrea gigas*: I. Absolute and  
939 allometric growth. *Aquaculture*, 70(3), 231-251. [https://doi.org/10.1016/0044-](https://doi.org/10.1016/0044-8486(88)90099-3)  
940 [8486\(88\)90099-3](https://doi.org/10.1016/0044-8486(88)90099-3).

941 Bruch, A. A., Utescher, T., Olivares, C. A., Dolakova, N., Ivanov, D., and Mosbrugger, V.  
942 2004. Middle and Late Miocene spatial temperature patterns and gradients in Europe-  
943 preliminary results based on palaeobotanical climate reconstructions. *Courier-*  
944 *Forschungsinstitut Senckenberg*, 15-28.

945 Bruch, A. A., Uhl, D., and Mosbrugger, V. 2007. Miocene climate in Europe—patterns and  
 946 evolution: a first synthesis of NECLIME. *Palaeogeography, Palaeoclimatology,*  
 947 *Palaeoecology*, 253(1-2), 1-7. <https://doi.org/10.1016/j.palaeo.2007.03.030>.

948 Bruch, A. A., Utescher, T., and Mosbrugger, V. 2011. Precipitation patterns in the Miocene of  
 949 Central Europe and the development of continentality. *Palaeogeography,*  
 950 *Palaeoclimatology,* *Palaeoecology*, 304(3-4), 202-211.  
 951 <https://doi.org/10.1016/j.palaeo.2010.10.002>.

952 Bryant, J. D., Jones, D. S., and Mueller, P. A. 1995. Influence of freshwater flux on  $^{87}\text{Sr}/^{86}\text{Sr}$   
 953 Sr chronostratigraphy in marginal marine environments and dating of vertebrate and  
 954 invertebrate faunas. *Journal of Paleontology*, 69(1), 1-6.  
 955 <https://doi.org/10.1017/S002233600002686X>.

956 Choquette, P. W., and James, N. P. 1990. Limestones—the burial diagenetic environment.  
 957 *Diagenesis: Geoscience Canada, Reprint Series*, 4, 75-111.

958 Clark Jr, R. C., Finley, J. S., and Gibson, G. G. 1974. Acute effects of outboard motor effluent  
 959 on two marine shellfish. *Environmental Science and Technology*, 8(12), 1009-1014.

960 Cornacchia, I., Agostini, S., and Brandano, M. 2018. Miocene Oceanographic Evolution  
 961 Based on the Sr and Nd Isotope Record of the Central Mediterranean.  
 962 *Paleoceanography and Paleoclimatology*, 33(1), 31-47.  
 963 <https://doi.org/10.1002/2017PA003198>.

964 Cramwinckel, M. J., Huber, M., Kocken, I. J., Agnini, C., Bijl, P. K., Bohaty, S. M., Frieling,  
 965 J., Goldner, A., Hilgen, F. J., Kip, E. L., Peterse, F., Van Der Ploeg, R., Röhl, U.,  
 966 Schouten, S., Sluijs, A. 2018. Synchronous tropical and polar temperature evolution in  
 967 the Eocene. *Nature*, 559(7714), 382. <https://doi.org/10.1038/s41586-018-0272-2>.

968 Daëron, M., Blamart, D., Peral, M., and Affek, H. P. 2016. Absolute isotopic abundance ratios  
 969 and the accuracy of  $\Delta_{47}$  measurements. *Chemical Geology*, 442, 83-96.  
 970 <https://doi.org/10.1016/j.chemgeo.2016.08.014>.

971 Das Sharma, S., Patil, D. J., and Gopalan, K. 2002. Temperature dependence of oxygen  
 972 isotope fractionation of CO<sub>2</sub> from magnesite-phosphoric acid reaction. *Geochimica et*  
 973 *Cosmochimica Acta*, 66(4), 589-593. [https://doi.org/10.1016/S0016-7037\(01\)00833-](https://doi.org/10.1016/S0016-7037(01)00833-X)  
 974 X.

975 Demarcq, G. 1970. Étude stratigraphique du Miocène rhodanien. Mém. BRGM, n°61, 257 p.

976 Demory, F., Conesa, G., Oudet, J., Mansouri, H., Münch, P., Borgomano, J., Thouveny, N.,  
 977 Lamarche, J., Gisquet, F., and Marié, L. 2011. Magnetostratigraphy and  
 978 paleoenvironments in shallow-water carbonates: the Oligocene-Miocene sediments of  
 979 the northern margin of the Liguro-Provençal basin (West Marseille, southeastern  
 980 France). *Bulletin de la Société géologique de France*, 182(1), 37-55.  
 981 <https://doi.org/10.2113/gssgfbull.182.1.37>.

982 Dera, G., Brigaud, B., Monna, F., Laffont, R., Pucéat, E., Deconinck, J. F., Pellenard, P.,  
 983 Joachimski, M. M., and Durlet, C. 2011. Climatic ups and downs in a disturbed  
 984 Jurassic world. *Geology*, 39(3), 215-218. <https://doi.org/10.1130/G31579.1>.

985 Diester-Haass, L., Billups, K., Gröcke, D. R., François, L., Lefebvre, V., and Emeis, K. C.  
 986 2009. Mid-Miocene paleoproductivity in the Atlantic Ocean and implications for the  
 987 global carbon cycle. *Paleoceanography and Paleoclimatology*, 24(1).  
 988 <https://doi.org/10.1029/2008PA001605>.

989 Do Couto, D., Gorini, C., Jolivet, L., Lebreton, N., Augier, R., Gumiaux, C., d'Acremont, E.,  
 990 Ammar, A., Jabour, H., and Auxietre, J. L. 2016. Tectonic and stratigraphic evolution  
 991 of the Western Alboran Sea Basin in the last 25 Myrs. *Tectonophysics*, 677, 280-311.  
 992 <https://doi.org/10.1016/j.tecto.2016.03.020>.

993 Domack, E. W. 1988. Biogenic facies in the Antarctic glacimarine environment: basis for a  
 994 polar glacimarine summary. *Palaeogeography, Palaeoclimatology, Palaeoecology*,  
 995 63(4), 357-372. [https://doi.org/10.1016/0031-0182\(88\)90105-8](https://doi.org/10.1016/0031-0182(88)90105-8).

996 Eiler, J. M., Bonifacie, M., and Daëron, M. 2009. 'Clumped isotope' thermometry for high-  
 997 temperature problems. *Geochimica et Cosmochimica Acta*, 73(13), A322-A322.  
 998 <http://dx.doi.org/10.1016/j.gca.2009.05.004>.

999 El Meknassi, S., Dera, G., Cardone, T., De Rafélis, M., Brahmi, C., and Chavagnac, V. 2018.  
 1000 Sr isotope ratios of modern carbonate shells: Good and bad news for  
 1001 chemostratigraphy. *Geology*, 46(11), 1003-1006. <https://doi.org/10.1130/G45380.1>.

1002 Faccenna, C., Mattei, M., Funiciello, R., and Jolivet, L. 1997. Styles of back-arc extension in  
 1003 the central Mediterranean. *Terra Nova*, 9(3), 126-130. <https://doi.org/10.1046/j.1365-3121.1997.d01-12.x>.

1005 Ferrandini, M., BouDagher-Fadel, M. K., Ferrandini, J., Oudet, J., and André, J. P. 2010.  
 1006 Nouvelles observations sur les Miogypsinidés du Miocène inférieur et moyen de  
 1007 Provence et de Corse (France) et de Sardaigne septentrionale (Italie). In *Annales de*  
 1008 *Paléontologie*. Vol. 96, No. 3, pp. 67-94. <https://doi.org/10.1016/j.annpal.2011.04.002>.

1009 Foster, G. L., Lear, C. H., and Rae, J. W. 2012. The evolution of pCO<sub>2</sub>, ice volume and  
 1010 climate during the middle Miocene. *Earth and Planetary Science Letters*, 341, 243-  
 1011 254. <https://doi.org/10.1016/j.epsl.2012.06.007>.

1012 Fondecave-Wallez, M. J., Magné, J., Gourinard, Y., and Demarcq, G. 1986. Nouvelles grade-  
 1013 datations de sédiments burdigaliens des régions rhodaniennes. *Géologie*  
 1014 *Méditerranéenne*, 12(1), 59-63. <https://doi.org/10.3406/geolm.1985.1339>.

1015 Friebe, J. G. 1994. Serpulid-bryozoan-foraminiferal biostromes controlled by temperate  
 1016 climate and reduced salinity: Middle Miocene of the Styrian Basin, Austria. *Facies*,  
 1017 30(1), 51-62. <https://doi.org/10.1007/BF02536889>.



1018 Frisch, W., Dunkl, I., & Kuhleemann, J. 2000. Post-collisional orogen-parallel large-scale  
 1019 extension in the Eastern Alps. *Tectonophysics*, 327(3-4), 239-265.  
 1020 [https://doi.org/10.1016/S0040-1951\(00\)00204-3](https://doi.org/10.1016/S0040-1951(00)00204-3).

1021 Frizon de Lamotte, D., Saint Bezar, B., Bracène, R., and Mercier, E. 2000. The two main  
 1022 steps of the Atlas building and geodynamics of the western Mediterranean. *Tectonics*,  
 1023 19(4), 740-761. <https://doi.org/10.1029/2000TC900003>.

1024 Galeotti, S., DeConto, R., Naish, T., Stocchi, P., Florindo, F., Pagani, M., Barrett, P., Bohaty,  
 1025 S. M., Lanci, L. Pollard, D., Sandroni, S., Talarico, F. M., Zachos, J. C. 2016.  
 1026 Antarctic Ice Sheet variability across the Eocene-Oligocene boundary climate  
 1027 transition. *Science*, 352(6281), 76-80. <https://doi.org/10.1126/science.aab0669>.

1028 Gasson, E., DeConto, R. M., Pollard, D., and Levy, R. H. 2016. Dynamic Antarctic ice sheet  
 1029 during the early to mid-Miocene. *Proceedings of the National Academy of Sciences*,  
 1030 113(13), 3459-3464. <https://doi.org/10.1073/pnas.1516130113>.

1031 Gattacceca, J., Deino, A., Rizzo, R., Jones, D. S., Henry, B., Beaudoin, B., and Vadeboin, F.  
 1032 2007. Miocene rotation of Sardinia: New paleomagnetic and geochronological  
 1033 constraints and geodynamic implications. *Earth and Planetary Science Letters*, 258(3-  
 1034 4), 359-377. <https://doi.org/10.1016/j.epsl.2007.02.003>.

1035 Ghosh, P., Adkins, J., Affek, H., Balta, B., Guo, W., Schauble, E. A., Schrag, D., and Eiler, J.  
 1036 M. 2006.  $^{13}\text{C}$ - $^{18}\text{O}$  bonds in carbonate minerals: a new kind of paleothermometer.  
 1037 *Geochimica et Cosmochimica Acta*, 70(6), 1439-1456.  
 1038 <https://doi.org/10.1016/j.gca.2005.11.014>

1039 Goedert, J., Amiot, R., Arnaud-Godet, F., Cuny, G., Fourel, F., Hernandez, J. A., Pedreira-  
 1040 Segade, U., and Lecuyer, C. 2017. Miocene (Burdigalian) seawater and air  
 1041 temperatures estimated from the geochemistry of fossil remains from the Aquitaine

1042 Basin, France. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 481, 14-28.  
 1043 <https://doi.org/10.1016/j.palaeo.2017.04.024>.  
 1044 Goldner, A., Herold, N., and Huber, M. 2014. The Challenge of Simulating the Warmth of the  
 1045 Mid-Miocene Climatic Optimum in CESM1. *Climate of the Past*, 10, 523-536.  
 1046 <https://doi.org/10.5194/cp-10-523-2014>.  
 1047 Grossman, E. L. 2012. Applying oxygen isotope paleothermometry in deep time. *The*  
 1048 *Paleontological Society Papers*, 18, 39-68.  
 1049 <https://doi.org/10.1017/S1089332600002540>.  
 1050 Gueguen, E., Doglioni, C., and Fernandez, M. 1998. On the post-25 Ma geodynamic  
 1051 evolution of the western Mediterranean. *Tectonophysics*, 298(1-3), 259-269.  
 1052 [https://doi.org/10.1016/S0040-1951\(98\)00189-9](https://doi.org/10.1016/S0040-1951(98)00189-9).  
 1053 Hamon, Y., Santerre, Y., Granjeon, D., Conesa, G., and Borgomano, J. 2013. Early diagenesis  
 1054 in meteoric versus brackish environments: Example of the Late Oligocene-Early  
 1055 Miocene, littoral, mixed sedimentary succession of Carry-Le-Rouet (southeastern  
 1056 France). *Bulletin de la Société géologique de France*, 184(6), 601-620.  
 1057 <https://doi.org/10.2113/gssgfbull.184.6.601>.  
 1058 Hannig, J., and Marron, J. S. 2006. Advanced distribution theory for SiZer. *Journal of the*  
 1059 *American Statistical Association*, 101(474), 484-499.  
 1060 Harzhauser, M., Piller, W. E., Müllegger, S., Grunert, P., and Micheels, A. 2011. Changing  
 1061 seasonality patterns in Central Europe from Miocene Climate Optimum to Miocene  
 1062 Climate Transition deduced from the *Crassostrea* isotope archive. *Global and*  
 1063 *Planetary Change*, 76(1-2), 77-84. <https://doi.org/10.1016/j.gloplacha.2010.12.003>.  
 1064 Henkes, G. A., Passey, B. H., Wanamaker Jr, A. D., Grossman, E. L., Ambrose Jr, W. G., and  
 1065 Carroll, M. L. 2013. Carbonate clumped isotope compositions of modern marine

1066 mollusk and brachiopod shells. *Geochimica et Cosmochimica Acta*, 106, 307-325.  
1067 <https://doi.org/10.1016/j.gca.2012.12.020>.

1068 van Hinsbergen, D. J., Torsvik, T. H., Schmid, S. M., Matenco, L. C., Maffione, M., Vissers,  
1069 R. L., Güreş, D., and Spakman, W. 2019. Orogenic architecture of the Mediterranean  
1070 region and kinematic reconstruction of its tectonic evolution since the Triassic.  
1071 *Gondwana Research*. <https://doi.org/10.1016/j.gr.2019.07.009>.

1072 Holbourn, A., Kuhnt, W., Schulz, M., and Erlenkeuser, H. 2005. Impacts of orbital forcing  
1073 and atmospheric carbon dioxide on Miocene ice-sheet expansion. *Nature*, 438(7067),  
1074 483. <https://doi.org/10.1038/nature04123>.

1075 Holbourn, A., Kuhnt, W., Kochhann, K. G., Andersen, N., and Sebastian Meier, K. J. 2015.  
1076 Global perturbation of the carbon cycle at the onset of the Miocene Climatic  
1077 Optimum. *Geology*, 43(2), 123-126. <https://doi.org/10.1130/G36317.1>.

1078 Holmes, A. 1965. *Principles of Physical Geology*. 2<sup>nd</sup> edn., Thomas Nelson, London, 1288 pp.

1079 Huang, K. F., You, C. F., Chung, C. H., and Lin, I. T. 2011. Nonhomogeneous seawater Sr  
1080 isotopic composition in the coastal oceans: A novel tool for tracing water masses and  
1081 submarine groundwater discharge. *Geochemistry, Geophysics, Geosystems*, 12(5).  
1082 <https://doi.org/10.1029/2010GC003372>.

1083 Huyghe, D., Mouthereau, F., and Emmanuel, L. 2012. Oxygen isotopes of marine mollusc  
1084 shells record Eocene elevation change in the Pyrenees. *Earth and Planetary Science*  
1085 *Letters*, 345, 131-141. <https://doi.org/10.1016/j.epsl.2012.06.035>.

1086 Huyghe, D., Lartaud, F., Emmanuel, L., Merle, D., and Renard, M. 2015. Palaeogene climate  
1087 evolution in the Paris Basin from oxygen stable isotope ( $\delta^{18}\text{O}$ ) compositions of marine  
1088 molluscs. *Journal of the Geological Society*, 172(5), 576-587.  
1089 <https://doi.org/10.1144/jgs2015-016>.

1090 Huyghe, D., de Rafélis, M., Ropert, M., Mouchi, V., Emmanuel, L., Renard, M., and Lartaud,  
 1091 F. 2019. New insights into oyster high-resolution hinge growth patterns. *Marine*  
 1092 *biology*, 166(4), 48. <https://doi.org/10.1007/s00227-019-3496-2>.  
 1093 Ingram, B. L., and Sloan, D. 1992. Strontium isotopic composition of estuarine sediments as  
 1094 paleosalinity-paleoclimate indicator. *Science*, 255(5040), 68-72.  
 1095 <https://doi.org/10.1126/science.255.5040.68>.  
 1096 Ivany, L. C., Patterson, W. P., and Lohmann, K. C. 2000. Cooler winters as a possible cause  
 1097 of mass extinctions at the Eocene/Oligocene boundary. *Nature*, 407(6806), 887.  
 1098 <https://doi.org/10.1038/35038044>.  
 1099 Jiménez-Moreno, G., and Suc, J. P. 2007. Middle Miocene latitudinal climatic gradient in  
 1100 Western Europe: evidence from pollen records. *Palaeogeography, Palaeoclimatology,*  
 1101 *Palaeoecology*, 253(1-2), 208-225. <https://doi.org/10.1016/j.palaeo.2007.03.040>.  
 1102 Jiménez-Moreno, G., Fauquette, S., and Suc, J. P. 2010. Miocene to Pliocene vegetation  
 1103 reconstruction and climate estimates in the Iberian Peninsula from pollen data. *Review*  
 1104 *of Palaeobotany and Palynology*, 162(3), 403-415.  
 1105 <https://doi.org/10.1016/j.revpalbo.2009.08.001>.  
 1106 Jolivet, L., Augier, R., Robin, C., Suc, J. P., and Rouchy, J. M. 2006. Lithospheric-scale  
 1107 geodynamic context of the Messinian salinity crisis. *Sedimentary geology*, 188, 9-33.  
 1108 <https://doi.org/10.1016/j.sedgeo.2006.02.004>.  
 1109 Kirby, M. X., Soniat, T. M., and Spero, H. J. 1998. Stable isotope sclerochronology of  
 1110 Pleistocene and Recent oyster shells (*Crassostrea virginica*). *Palaios*, 13(6), 560-569.  
 1111 <https://doi.org/10.2307/3515347>.  
 1112 Klein, R. T., K. C. Lohmann, and C. W. Thayer. 1996. Sr/Ca and  $^{13}\text{C}/^{12}\text{C}$  ratios in skeletal  
 1113 calcite of *Mytilus trossulus*: Covariation with metabolic rate, salinity, and carbon

1114 isotopic composition of seawater, *Geochim. Cosmochim. Acta*, 60, 4207– 4221.  
 1115 [https://doi.org/10.1016/S0016-7037\(96\)00232-3](https://doi.org/10.1016/S0016-7037(96)00232-3).  
 1116 Kocsis, L., Vennemann, T. W., Fontignie, D., Baumgartner, C., and Montanari, A. 2008.  
 1117 Oceanographic and climatic evolution of the Miocene Mediterranean deduced from  
 1118 Nd, Sr, C, and O isotope compositions of marine fossils and sediments.  
 1119 *Paleoceanography*, 23, PA4211. <https://doi.org/10.1029/2007PA001540>.  
 1120 Kocsis, L., Vennemann, T. W., Hegner, E., Fontignie, D., and Tütken, T. 2009. Constraints on  
 1121 Miocene oceanography and climate in the Western and Central Paratethys: O-, Sr-,  
 1122 and Nd-isotope compositions of marine fish and mammal remains. *Palaeogeography*,  
 1123 *Palaeoclimatology*, *Palaeoecology*, 271(1-2), 117-129.  
 1124 <https://doi.org/10.1016/j.palaeo.2008.10.003>.  
 1125 Lartaud, F., De Rafélis, M., Ropert, M., Emmanuel, L., Geairon, P., and Renard, M. 2010. Mn  
 1126 labelling of living oysters: artificial and natural cathodoluminescence analyses as a  
 1127 tool for age and growth rate determination of *C. gigas* (Thunberg, 1793) shells.  
 1128 *Aquaculture*, 300(1-4), 206-217. <https://doi.org/10.1007/s10347-009-0196-2>.  
 1129 Langlet, D., Alunno-Bruscia, M., Rafélis, M., Renard, M., Roux, M., Schein, E., and Buestel,  
 1130 D. 2006. Experimental and natural cathodoluminescence in the shell of *Crassostrea*  
 1131 *gigas* from Thau lagoon (France): ecological and environmental implications. *Marine*  
 1132 *Ecology Progress Series*, 317, 143-156. <https://doi.org/10.3354/meps317143>.  
 1133 Lartaud, F., Langlet, D., De Rafelis, M., Emmanuel, L., and Renard, M. 2006. Mise en  
 1134 évidence de rythmicité saisonnière dans la coquille des huîtres fossiles *Crassostrea*  
 1135 *aginensis* Tournouer, 1914 (Aquitaniens) et *Ostrea bellovacina* Lamarck, 1806  
 1136 (Thanétien). Approche par cathodoluminescence et par sclérochronologie. *Geobios*,  
 1137 39(6), 845-852. <https://doi.org/10.1016/j.geobios.2005.11.001>.

1138 Lear, C. H., Bailey, T. R., Pearson, P. N., Coxall, H. K., and Rosenthal, Y. 2008. Cooling and  
 1139 ice growth across the Eocene-Oligocene transition. *Geology*, 36(3), 251-254.  
 1140 <https://doi.org/10.1130/G24584A.1>.

1141 Lear, C. H., Coxall, H. K., Foster, G. L., Lunt, D. J., Mawbey, E. M., Rosenthal, Y., Sosdian,  
 1142 S. M., Thomas, E., and Wilson, P. A. 2015. Neogene ice volume and ocean  
 1143 temperatures: Insights from infaunal foraminiferal Mg/Ca paleothermometry.  
 1144 *Paleoceanography*, 30(11), 1437-1454. <https://doi.org/10.1002/2015PA002833>.

1145 Lecuyer, C., Grandjean, P., Paris, F., Robardet, M., and Robineau, D. 1996. Deciphering  
 1146 “temperature” and “salinity” from biogenic phosphates: the  $\delta^{18}\text{O}$  of coexisting fishes  
 1147 and mammals of the Middle Miocene sea of western France. *Palaeogeography*,  
 1148 *Palaeoclimatology*, *Palaeoecology*, 126(1-2), 61-74. [https://doi.org/10.1016/S0031-](https://doi.org/10.1016/S0031-0182(96)00070-3)  
 1149 [0182\(96\)00070-3](https://doi.org/10.1016/S0031-0182(96)00070-3).

1150 Levitus, S., and Boyer, T. P. 1994. World ocean atlas 1994. volume 4. temperature (No. PB-  
 1151 95-270112/XAB; NESDIS-4). National Environmental Satellite, Data, and  
 1152 Information Service, Washington, DC (United States).

1153 Liu, Z., Pagani, M., Zinniker, D., DeConto, R., Huber, M., Brinkhuis, H., Shah, S. R., Leckie,  
 1154 M., and Pearson, A. 2009. Global cooling during the Eocene-Oligocene climate  
 1155 transition. *Science*, 323(5918), 1187-1190. <https://doi.org/10.1126/science.1166368>.

1156 Marron, J. S., and Chaudhuri, P. 1998. When is a feature really there: the SiZer approach. In  
 1157 Automatic Target Recognition VIII (Vol. 3371, pp. 306-313). International Society for  
 1158 Optics and Photonics. <https://doi.org/10.1117/12.323850>.

1159 Maurel-ferrandini M. 1976. Reconstitution paléogéographique du Burdigalien du littoral de la  
 1160 chaîne de la Nerthe et de la région des étangs (Bouches-du-Rhône, France). – PhD  
 1161 Thesis, University of Provence, Marseille, 87 p.

1162 Mawbey, E. M., and Lear, C. H. 2013. Carbon cycle feedbacks during the Oligocene-Miocene  
 1163 transient glaciation. *Geology*, 41(9), 963-966. <https://doi.org/10.1130/G34422.1>.

1164 McArthur, J., and Howarth, R. J. 2004. Strontium isotope stratigraphy. In F. Gradstein, J.  
 1165 Ogg, and A. Smith (Eds.), *A geologic time scale* (pp. 96–105). Cambridge: Cambridge  
 1166 University Press.

1167 Meckler, A. N., Ziegler, M., Millán, M. I., Breitenbach, S. F., and Bernasconi, S. M. 2014.  
 1168 Long-term performance of the Kiel carbonate device with a new correction scheme for  
 1169 clumped isotope measurements. *Rapid Communications in Mass Spectrometry*,  
 1170 28(15), 1705-1715. <https://doi.org/10.1002/rcm.6949>.

1171 Miall, A. D. 1986. Eustatic sea level changes interpreted from seismic stratigraphy: a critique  
 1172 of methodology with particular reference to the North Sea Jurassic record. *AAPG*  
 1173 *Bull.* 70, 131-137.

1174 Miller, K. G., Wright, J. D., and Fairbanks, R. G. 1991. Unlocking the ice house: Oligocene-  
 1175 Miocene oxygen isotopes, eustasy, and margin erosion. *Journal of Geophysical*  
 1176 *Research: Solid Earth*, 96(B4), 6829-6848. <https://doi.org/10.1029/90JB02015>.

1177 Mitchell, I. M., Crawford, C. M., and Rushton, M. J. 2000. Flat oyster (*Ostrea angasi*) growth  
 1178 and survival rates at Georges Bay, Tasmania (Australia). *Aquaculture*, 191(4), 309-  
 1179 321. [https://doi.org/10.1016/S0044-8486\(00\)00441-5](https://doi.org/10.1016/S0044-8486(00)00441-5).

1180 Montanari, A., Carey, S., Coccioni, R., and Deino, A. 1994. Early Miocene tephra in the  
 1181 Apennine pelagic sequence: an inferred Sardinian provenance and implications for  
 1182 western Mediterranean tectonics. *Tectonics*, 13(5), 1120-1134.  
 1183 <https://doi.org/10.1029/94TC00295>.

1184 Mosbrugger, V., Utescher, T., and Dilcher, D. L. 2005. Cenozoic continental climatic  
 1185 evolution of Central Europe. *PNAS*, 102(42), 14964-14969.  
 1186 <https://doi.org/10.1073/pnas.0505267102>.

1187 Mouchi, V., Briard, J., Gaillot, S., Argant, T., Forest, V., and Emmanuel, L. 2018.  
 1188 Reconstructing environments of collection sites from archaeological bivalve shells:  
 1189 Case study from oysters (Lyon, France). *Journal of Archaeological Science: Reports*,  
 1190 21, 1225-1235. <https://doi.org/10.1016/j.jasrep.2017.10.025>.  
 1191 Nalin, R., Ghinassi, M., and Basso, D. 2010. Onset of temperate carbonate sedimentation  
 1192 during transgression in a low-energy siliciclastic embayment (Pliocene of the Val  
 1193 d'Orcia Basin, Tuscany, Italy). *Facies*, 56(3), 353-368.  
 1194 <https://doi.org/10.1007/s10347-010-0211-7>.  
 1195 Nayar, K. G., Panchanathan, D., McKinley, G. H., and Lienhard, J. H. 2014. Surface tension  
 1196 of seawater. *Journal of Physical and Chemical Reference Data*, 43(4), 043103.  
 1197 <https://doi.org/10.1063/1.4899037>.  
 1198 Nemec, W., and Steel, R. J. 1988. What is a fan delta and how do we recognize it. *Fan Deltas:*  
 1199 *sedimentology and tectonic settings*, 3-13.  
 1200 Nury, D. 1990. L'Oligocène de Provence méridionale: stratigraphie, dynamique sédimentaire,  
 1201 reconstitutions paléogéographiques. PhD Thesis, University of Provence, Marseille,  
 1202 411 p.  
 1203 Oertli, H. J. 1964. The Venice system for the classification of marine waters according to  
 1204 salinity. *Publicazioni della Stazione Zoologica di Napoli*, 33, Supplement, p.611.  
 1205 Ogg, J. G., Ogg, G., and Gradstein, F. M. 2016. A concise geologic time scale: 2016. Elsevier.  
 1206 <https://doi.org/10.1016/C2009-0-64442-1>.  
 1207 Ohno, T., and Hirata, T. 2007. Simultaneous determination of mass-dependent isotopic  
 1208 fractionation and radiogenic isotope variation of strontium in geochemical samples by  
 1209 multiple collector-ICP-mass spectrometry. *Analytical Sciences*, 23(11), 1275-1280.  
 1210 <https://doi.org/10.2116/analsci.23.1275>.



1211 Oudet, J., Münch, P., Borgomano, J., Quillevere, F., Melinte-Dobrinescu, M. C., Demory, F.,  
 1212 Viseur, S., and Cornee, J. J. 2010. Land and sea study of the northeastern golfe du  
 1213 Lion rifted margin: the Oligocene–Miocene of southern Provence (Nerthe area, SE  
 1214 France). *Bulletin de la Société géologique de France*, 181(6), 591-607.  
 1215 <https://doi.org/10.2113/gssgfbull.181.6.591>.  
 1216 Pagani, M., Arthur, M. A., and Freeman, K. H. 1999. Miocene evolution of atmospheric  
 1217 carbon dioxide. *Paleoceanography*, 14(3), 273-292.  
 1218 <https://doi.org/10.1029/1999PA900006>.  
 1219 Palmer, M. R., and Elderfield, H. 1985. Sr isotope composition of sea water over the past 75  
 1220 Myr. *Nature*, 314(6011), 526. <https://doi.org/10.1038/314526a0>.  
 1221 Passey, B. H., and Henkes, G. A. 2012. Carbonate clumped isotope bond reordering and  
 1222 geospeedometry. *Earth and Planetary Science Letters*, 351, 223-236.  
 1223 <https://doi.org/10.1016/j.epsl.2012.07.021>.  
 1224 Paulet, Y. M., Lorrain, A., Richard, J., and Pouvreau, S. 2006. Experimental shift in diet  $\delta^{13}\text{C}$ :  
 1225 a potential tool for ecophysiological studies in marine bivalves. *Organic*  
 1226 *Geochemistry*, 37(10), 1359-1370. <https://doi.org/10.1016/j.orggeochem.2006.01.008>.  
 1227 Peral, M., Daëron, M., Blamart, D., Bassinot, F., Dewilde, F., Smialkowski, N., Isguder, G.,  
 1228 Bonnin, J., Jorissen, F., Kissel, C., Michel, E., Vasquez Riveiros, N., and  
 1229 Waelbroeck, C. 2018. Updated calibration of the clumped isotope thermometer in  
 1230 planktonic and benthic foraminifera. *Geochimica et Cosmochimica Acta*, 239, 1-16.  
 1231 <https://doi.org/10.1016/j.gca.2018.07.016>.  
 1232 Petersen, S. V., Defliese, W. F., Saenger, C., Daëron, M., Huntington, K. W., John, C. M.,  
 1233 Kelson, J. R., Bernasconi, S. M., Colman, A. S., Klug, T., Olack, G. A., Schauer, A. J.,  
 1234 Bajnai, D., Bonifacie, M., Breitenbach, S. F. M., Fiebig, J., Fernandez, A. B., Henkes,  
 1235 G. A., Hodell, D., Katz, A., Kele, S., Lohmann, K. C., Passey, B. H., Peral, M. Y.,

1236 Petrizzo, D. A., Rosenheim, B. E., Tripathi, A., Venturelli, R., Young, E. D., and  
 1237 Winkelstern, I. Z. 2019. Effects of Improved  $^{17}\text{O}$  Correction on Inter-Laboratory  
 1238 Agreement in Clumped Isotope Calibrations, Estimates of Mineral-Specific Offsets,  
 1239 and Temperature Dependence of Acid Digestion Fractionation. *Geochemistry,*  
 1240 *Geophysics, Geosystems.* 20(7), 3495-3519. <https://doi.org/10.1029/2018GC008127>.  
 1241 Peucker-Ehrenbrink, B., and Fiske, G. J. 2019. A continental perspective of the seawater  
 1242  $^{87}\text{Sr}/^{86}\text{Sr}$  record: A review. *Chemical Geology,* 510, 140-165.  
 1243 <https://doi.org/10.1016/j.chemgeo.2019.01.017>.  
 1244 Pierre, C. 1999. The oxygen and carbon isotope distribution in the Mediterranean water  
 1245 masses. *Marine Geology,* 153, 41–55. [https://doi.org/10.1016/S0025-3227\(98\)00090-](https://doi.org/10.1016/S0025-3227(98)00090-5)  
 1246 5.  
 1247 Pound, M. J., Haywood, A. M., Salzmann, U., and Riding, J. B. 2012. Global vegetation  
 1248 dynamics and latitudinal temperature gradients during the Mid to Late Miocene  
 1249 (15.97–5.33 Ma). *Earth-Science Reviews,* 112(1-2), 1-22.  
 1250 <https://doi.org/10.1016/j.earscirev.2012.02.005>.  
 1251 R Core Team 2017. R: A language and environment for statistical computing. R Foundation  
 1252 for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.  
 1253 Railsback, L. B., and Anderson T.F. 1989. Paleoceanographic modeling of temperature-  
 1254 salinity profiles from stable isotopic data. *Paleoceanography,* v. 4, 801-814.  
 1255 <https://doi.org/10.1029/PA004i005p00585>.  
 1256 Randazzo, A. F., Muller, P., Lelkes, G., Juhasz, E., and Hamor, T. 1999. Cool-water  
 1257 limestones of the Pannonian basinal system, Middle Miocene, Hungary. *Journal of*  
 1258 *Sedimentary Research,* 69(1), 283-293. <https://doi.org/10.2110/jsr.69.283>.

1259 Rehault, J. P., Boillot, G., and Mauffret, A. 1984. The western Mediterranean basin geological  
 1260 evolution. *Marine Geology*, 55(3-4), 447-477. [https://doi.org/10.1016/0025-](https://doi.org/10.1016/0025-3227(84)90081-1)  
 1261 3227(84)90081-1.

1262 Reuter, M., Piller, W. E., Brandano, M., and Harzhauser, M. 2013. Correlating Mediterranean  
 1263 shallow water deposits with global Oligocene–Miocene stratigraphy and oceanic  
 1264 events. *Global and planetary change*, 111, 226-236.  
 1265 <https://doi.org/10.1016/j.gloplacha.2013.09.018>.

1266 Reynaud, J. Y., Dalrymple, R. W., Vennin, E., Parize, O., Besson, D., and Rubino, J. L. 2006.  
 1267 Topographic controls on production and deposition of tidal cool-water carbonates,  
 1268 Uzes Basin, SE France. *Journal of Sedimentary Research*, 76(1), 117-130.  
 1269 <https://doi.org/10.2110/jsr.2006.07>.

1270 Reynaud, J. Y., Vennin, E., Parize, O., Rubino, J. L., and Bourdillon, C. 2012. Incised valleys  
 1271 and tidal seaways: the example of the Miocene Uzès-Castillon basin, SE France.  
 1272 *Bulletin de la Société Géologique de France*, 183(5), 471-486.  
 1273 <https://doi.org/10.2113/gssgfbull.183.5.471>.

1274 Riegl, B., and Piller, W. E. 2000. Biostromal coral facies—a Miocene example from the  
 1275 Leitha Limestone (Austria) and its actualistic interpretation. *Palaios*, 15(5), 399-413.  
 1276 [https://doi.org/10.1669/0883-1351\(2000\)015<0399:BCFAME>2.0.CO;2](https://doi.org/10.1669/0883-1351(2000)015<0399:BCFAME>2.0.CO;2).

1277 Ryland, J. S. 1970. *Bryozoans*. Hutchinson Univ. Library, London, 175.

1278 Schauble, E. A., Ghosh, P., and Eiler, J. M. 2006. Preferential formation of  $^{13}\text{C}$ – $^{18}\text{O}$  bonds in  
 1279 carbonate minerals, estimated using first-principles lattice dynamics. *Geochimica et*  
 1280 *Cosmochimica Acta*, 70(10), 2510-2529. <https://doi.org/10.1016/j.gca.2006.02.011>.

1281 Schettino, A., and Turco, E. 2011. Tectonic history of the western Tethys since the Late  
 1282 Triassic. *Bulletin*, 123(1-2), 89-105. <https://doi.org/10.1130/B30064.1>.

1283 Schöne, B. R., and Giere, O. 2005. Growth increments and stable isotope variation in shells of  
 1284 the deep-sea hydrothermal vent bivalve mollusk *Bathymodiolus brevior* from the  
 1285 North Fiji Basin, Pacific Ocean. *Deep Sea Research Part I: Oceanographic Research*  
 1286 *Papers*, 52(10), 1896-1910. <https://doi.org/10.1016/j.dsr.2005.06.003>.

1287 Schöne, B. R. 2008. The curse of physiology—challenges and opportunities in the  
 1288 interpretation of geochemical data from mollusk shells. *Geo-Marine Letters*, 28(5-6),  
 1289 269-285. <https://doi.org/10.1007/s00367-008-0114-6>.

1290 Seibel, M. J., and James, N. P. 2017. Diagenesis of Miocene, incised valley-filling limestones;  
 1291 Provence, Southern France. *Sedimentary geology*, 347, 21-35.  
 1292 <https://doi.org/10.1016/j.sedgeo.2016.09.006>.

1293 Shevenell, A. E., Kennett, J. P., and Lea, D. W. 2004. Middle Miocene southern ocean  
 1294 cooling and Antarctic cryosphere expansion. *Science*, 305(5691), 1766-1770.  
 1295 <https://doi.org/10.1126/science.1100061>.

1296 Shevenell, A. E., Kennett, J. P., and Lea, D. W. 2008. Middle Miocene ice sheet dynamics,  
 1297 deep-sea temperatures, and carbon cycling: A Southern Ocean perspective.  
 1298 *Geochemistry, Geophysics, Geosystems*, 9(2).  
 1299 <https://doi.org/10.1029/2007GC001736>.

1300 Smith, A. M. 1995. Palaeoenvironmental interpretation using bryozoans: a review. *Geological*  
 1301 *Society, London, Special Publications*, 83(1), 231-243.  
 1302 <https://doi.org/10.1144/GSL.SP.1995.083.01.11>.

1303 Sonderegger, D., and Sonderegger, M. D. 2018. Package ‘SiZer’.

1304 Stenzel, H. B. 1971. Oysters. *Treatise on Invertebrate Paleontology, Part N, Bivalvia 3*, N953-  
 1305 N1224.

1306 Steuber, T., Rauch, M., Masse, J. P., Graaf, J., and Malkoč, M. 2005. Low-latitude seasonality  
 1307 of Cretaceous temperatures in warm and cold episodes. *Nature*, 437(7063), 1341.  
 1308 <https://doi.org/10.1038/nature04096>.

1309 Stolper, D. A., and Eiler, J. M. 2015. The kinetics of solid-state isotope-exchange reactions  
 1310 for clumped isotopes: A study of inorganic calcites and apatites from natural and  
 1311 experimental samples. *American Journal of Science*, 315(5), 363-411.  
 1312 <https://doi.org/10.2475/05.2015.01>.

1313 Super, J. R., Thomas, E., Pagani, M., Huber, M., O'Brien, C., and Hull, P. M. 2018. North  
 1314 Atlantic temperature and pCO<sub>2</sub> coupling in the early-middle Miocene. *Geology*, 46(6),  
 1315 519-522. <https://doi.org/10.1130/G40228.1>.

1316 Surge, D., Lohmann, K. C., and Dettman, D. L. 2001. Controls on isotopic chemistry of the  
 1317 American oyster, *Crassostrea virginica*: implications for growth patterns.  
 1318 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 172(3-4), 283-296.  
 1319 [https://doi.org/10.1016/S0031-0182\(01\)00303-0](https://doi.org/10.1016/S0031-0182(01)00303-0).

1320 Taylor, P. D., and James, N. P. 2013. Secular changes in colony-forms and bryozoan  
 1321 carbonate sediments through geological history. *Sedimentology*, 60(5), 1184-1212.  
 1322 <https://doi.org/10.1111/sed.12032>.

1323 Tempier, C., 1987. Modèle nouveau de mise en place des structures provençales. *Bulletin*  
 1324 *Société Géologique France* 8 (3), 533–540. <https://doi.org/10.2113/gssgfbull.III.3.533>.

1325 Tremblin, M., Hermoso, M., and Minoletti, F. 2016. Equatorial heat accumulation as a long-  
 1326 term trigger of permanent Antarctic ice sheets during the Cenozoic. *Proceedings of the*  
 1327 *National Academy of Sciences*, 113(42), 11782-11787.  
 1328 <https://doi.org/10.1073/pnas.1608100113>.

1329 Trezzi, G., Garcia-Orellana, J., Rodellas, V., Masqué, P., Garcia-Solsona, E., and Andersson,  
 1330 P. S. 2017. Assessing the role of submarine groundwater discharge as a source of Sr to

1331 the Mediterranean Sea. *Geochimica et Cosmochimica Acta*, 200, 42-54.  
1332 <https://doi.org/10.1016/j.gca.2016.12.005>.

1333 Ullmann, C. V., Wiechert, U., and Korte, C. 2010. Oxygen isotope fluctuations in a modern  
1334 North Sea oyster (*Crassostrea gigas*) compared with annual variations in seawater  
1335 temperature: Implications for palaeoclimate studies. *Chemical Geology*, 277(1-2),  
1336 160-166. <https://doi.org/10.1016/j.chemgeo.2010.07.019>.

1337 Ullmann, C. V., Campbell, H. J., Frei, R., Hesselbo, S. P., von Strandmann, P. A. P., and  
1338 Korte, C. 2013. Partial diagenetic overprint of Late Jurassic belemnites from New  
1339 Zealand: Implications for the preservation potential of  $\delta^7\text{Li}$  values in calcite fossils.  
1340 *Geochimica et Cosmochimica Acta*, 120, 80-96.  
1341 <https://doi.org/10.1016/j.gca.2013.06.029>.

1342 Ullmann, C. V., and Korte, C. 2015. Diagenetic alteration in low-Mg calcite from  
1343 macrofossils: a review. *Geological Quarterly*, 59(1), 3-20.  
1344 <https://doi.org/10.7306/gq.1217>.

1345 Veizer, J., Ala, D., Azmy, K., Bruckschen, P., Buhl, D., Bruhn, F., Carden, G., Diener, A.,  
1346 Godderis, Y., Jasper, T., Korte, C., Pawellek, F., Podlaha, O., and Strauss, H. 1999.  
1347  $^{87}\text{Sr}/^{86}\text{Sr}$ ,  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  evolution of Phanerozoic seawater. *Chemical geology*,  
1348 161(1-3), 59-88. [https://doi.org/10.1016/S0009-2541\(99\)00081-9](https://doi.org/10.1016/S0009-2541(99)00081-9).

1349 Weckström, J., Korhola, A., Erästö, P., and Holmström, L. 2006. Temperature patterns over  
1350 the past eight centuries in Northern Fennoscandia inferred from sedimentary diatoms.  
1351 *Quaternary Research*, 66(1), 78-86. <https://doi.org/10.1016/j.yqres.2006.01.005>.

1352 Widerlund, A., and Andersson, P. S. 2006. Strontium isotopic composition of modern and  
1353 Holocene mollusc shells as a palaeosalinity indicator for the Baltic Sea. *Chemical*  
1354 *Geology*, 232(1-2), 54-66. <https://doi.org/10.1016/j.chemgeo.2006.02.010>.

1355 de Winter, N. J. D., Vellekoop, J., Vorsselmans, R., Golreihan, A., Soete, J., Petersen, S. V.,  
 1356 Meyer, K. W., Casadio, S., Speijer, R. P., Claeys, P. 2018. An assessment of latest  
 1357 Cretaceous *Pycnodonte vesicularis* (Lamarck, 1806) shells as records for  
 1358 palaeoseasonality: a multi-proxy investigation. *Climate of the Past*, 14(6), 725-749.  
 1359 <https://doi.org/10.5194/cp-14-725-2018>.

1360 Yeghicheyan, D., Bossy, C., Le Coz, M. B., Douchet, C., Granier, G., Heimburger, A., Lacan,  
 1361 F., Lanzanova, A., Rousseau, T., Seidel, J., Tharaud, M., Candaudap, F., Chmeleff, J.,  
 1362 Cloquet, C., Delpoux, S., Labatut, M., Losno, R., Pradoux, C., Sivry, Y., and Sonke, J.  
 1363 E. 2013. A compilation of silicon, rare earth element and twenty-one other trace  
 1364 element concentrations in the natural river water reference material SLRS-5 (NRC-  
 1365 CNRC). *Geostandards and Geoanalytical Research*, 37(4), 449-467.  
 1366 <https://doi.org/10.1111/j.1751-908X.2013.00232.x>.

1367 Zachos, J. C., Quinn, T. M., and Salamy, K. A. 1996. High-resolution (104 years) deep-sea  
 1368 foraminiferal stable isotope records of the Eocene-Oligocene climate transition.  
 1369 *Paleoceanography*, 11(3), 251-266. <https://doi.org/10.1029/96PA00571>.

1370 Zachos, J., Pagani, M., Sloan, L., Thomas, E., and Billups, K. 2001. Trends, rhythms, and  
 1371 aberrations in global climate 65 Ma to present. *Science*, 292(5517), 686-693.  
 1372 <https://doi.org/10.1126/science.1059412>.

1373 Zachos, J. C., Dickens, G. R., and Zeebe, R. E. 2008. An early Cenozoic perspective on  
 1374 greenhouse warming and carbon-cycle dynamics. *Nature*, 451(7176), 279.  
 1375 <https://doi.org/10.1038/nature06588>.

1376 Zaky, A. H., Brand, U., Buhl, D., Blamey, N., Bitner, M. A., Logan, A., Gaspard, D., and  
 1377 Popov, A. 2018. Strontium isotope geochemistry of modern and ancient archives:  
 1378 tracer of secular change in ocean chemistry. *Canadian Journal of Earth Sciences*,  
 1379 56(3), 245-264. <https://doi.org/10.1139/cjes-2018-0085>.

1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397

TABLE CAPTION

**Table 1** Facies and depositional environments of Carry-Le-Rouet section.

**Table 2**  $\Delta_{47}$  data for all analyzed samples and conversion to temperature according to Peral et al. (2018) equation. Analytical errors reported here are derived from the external reproducibility of carbonate standards (N = 47) and samples (N = 39) within each analytical session, and conservatively account for the uncertainties in raw  $\Delta_{47}$  measurements as well as those associated with the conversion to the “absolute”  $\Delta_{47}$  reference frame. Local  $\delta^{18}\text{O}_{\text{sw}}$  values were calculated using  $\Delta_{47}$ -derived temperatures and  $\delta^{18}\text{O}_{\text{shell}}$  with Anderson and Arthur (1983) equation. Salinity estimations derives from calculated local  $\delta^{18}\text{O}_{\text{sw}}$  values, from which the  $\delta^{18}\text{O}_{\text{sw}}$  of the global ocean (from Billups and Schrag; 2002) has been subtracted, and Railsback and Anderson (1989) equation based on North Atlantic Ocean.

**Table 3**  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios of bulk carbonate and their error bars.



FIGURE CAPTION

**Fig. 1.** Location of the studied outcrops. Absolute ages presented on the left are from Ogg et al. (2016), along with the correspondence with magnetostratigraphic frame and foraminifera zonation (C6C to C5A). A. Location of the outcrops on a modern map of the North Western Mediterranean area. B. Geological map of outcropping latest Chattian to latest Burdigalian (Oligo-Miocene) formations in the Coastal Nerthe area and their sedimentary units (modified from Andreieff et al., 1972; Oudet et al., 2010; Demory et al., 2011). The numbers 1 to 8 correspond to logs positions used to build a composite log of over than 80 meters, that is used in this study. C. Geological map of Burdigalian to earliest Serravallian (Miocene) formations in the Castillon-Du-Gard sub-basin and their sedimentary units (modified by from Reynaud et al., 2012). The numbers 1 to 5 correspond to logs positions.

**Fig. 2.** A. Picture representing the location of ligamental area on an oyster shell (modified from Stenzel, 1971). B. Sections of oyster shells under natural light (LN) and cathodoluminescence (CL), showing luminescent part in orange (mainly chalky areas) and non-luminescent to weakly luminescent parts with identifiable seasonal patterns of growths. Only the non-luminescent to weakly luminescent parts with identifiable seasonal patterns of growths parts have been sampled for isotopic analyses.

**Fig. 3.** A. Composite log of the Carry-Le-Rouet sedimentary succession with the different units identified along the log (after Andreieff et al., 1972; Oudet et al., 2010; Demory et al., 2011), lithologies and marker beds (UR1 to UR4 in red). The position of logs 1 to 8, used to build this synthetic log, is reported on Fig. 1. B. Outcrop pictures: 1. Transition from continental conglomerate to marine sandstone in the Chattian Conglomeratic Rouet unit; 2. Detail of the conglomerate beds eroding sandstones with oyster debris in the Conglomerate

1423 Rouet Unit. 3. Bioclastic carbonate deposits with abundant oyster shells at the transition  
1424 between the Brackish Rousset and the Bioclastic Carry units; 4. *Ostrea fimbriata* localized at  
1425 the base of the Bioclastic Carry unit; 5. Reefal Unit 4 (UR4) outcropping the Bioclastic Carry  
1426 unit in Carry harbor; 6. Detail of the surface topping the UR4 and composed of dense  
1427 Poritidae corals preserved in life position; 7. Marls of Tamaris section covered by the  
1428 Langhian carbonate deposits; 8. Detail of marls with a low fossil record.

1429

1430 **Fig. 4.** Evolution of bivalve  $\delta^{18}\text{O}$  throughout the section of Carry-Le-Rouet (A, Black plain  
1431 circles) and Castillon-Du-Gard (B, Grey plain circles). Several microsamplings were realized  
1432 per shells (levels where a microsampled oyster has been analyzed are identified by the name  
1433 of the shell newt to the relevant data) and several shells were analyzed per stratigraphic levels.  
1434 Each plotted data represents  $\delta^{18}\text{O}$  data acquired in this work, including the multiple  $\delta^{18}\text{O}$   
1435 measurements realized on some shells, appears on the figure. Analytical uncertainties are  
1436 lower than the size of the circles ( $\pm 0.08\text{‰}$ , see Method section) and cannot be reported for  
1437 each data point. Plain squares correspond to  $\delta^{18}\text{O}_{\text{shell}}$  measured with clumped isotopes (White  
1438 squares correspond to bulk of shell; Red square corresponds to summer part of shell CAR17-  
1439 6-33-b; Blue square corresponds to winter part of shell CAR17-6-33). Vertical error bars  
1440 reported on two Langhian samples, representing an uncertainty on the stratigraphic position of  
1441 the samples, correspond to bulk shells that were gathered from stratigraphic levels newt to  
1442 each other into a single analysis.

1443

1444 **Fig. 5.** Intra-shells  $\delta^{18}\text{O}$  evolution from bivalves recovered from the Carry-Le-Rouet outcrop.  
1445 A. CAR17-6-28-a; B. CAR17-6-30top-e; C. CAR17-6-31base-a; D. CAR17-6-33-b; E.  
1446 CAR17-6-33-c; F. CAR17-8-43-h; G. CAR17-8-43-i. Corresponding temperatures have been  
1447 calculated using the equation of Anderson and Arthur (1983) and a  $\delta^{18}\text{O}_{\text{sw}}$  of  $-0.66\text{‰}_{\text{VPDB}}$

(Billups and Schrag, 2002) for the Aquitanian (A to E) and of  $-0.94\text{‰}$   $\text{VPDB}$  (Billups and Schrag, 2002) for the Late Burdigalian (F and G). The error bar that applies to  $\delta^{18}\text{O}$  measurements corresponds to the analytical reproducibility specified in the Method part, of  $\pm 0.08\text{‰}$ .

**Fig. 6.** Distribution of  $\delta^{18}\text{O}$  values along CAR17-6-33-b umbo obtained from microsampling (black plain circles). The dotted red and blue lines represent the  $\delta^{18}\text{O}$  values of the two samples recovered from this oyster for  $\Delta_{47}$  analyses, over the length of the shell corresponding to minimum  $\delta^{18}\text{O}$  values and maximum  $\delta^{18}\text{O}$  values, respectively, as described in the text (Part 3.3). The temperature calculated from  $\Delta_{47}$  ( $T(\Delta_{47})$ ) is reported for these two samples. The error bar that applies to  $\delta^{18}\text{O}$  measurements corresponds to the analytical reproducibility specified in the Method part, of  $\pm 0.08\text{‰}$ .

**Fig. 7.** Comparison of geochemical data from bivalve shells obtained in this work, with the results of the applied statistical treatment (SiZer Map:  $0.1 < h < 10$  Myrs ; Blue: significant increase; Red: significant decrease; Grey: No significant; White: No data), with the benthic foraminifera  $\delta^{18}\text{O}$  global record (Zachos et al., 2008). Black plain circles correspond to bivalve  $\delta^{18}\text{O}$  values for the Carry-Le-Rouet section and Grey plain circles correspond to bivalve  $\delta^{18}\text{O}$  values from Castillon-Du-Gard. Smoothed black curves and their grey envelope on foraminifera  $\delta^{18}\text{O}$  global record and on our bivalve  $\delta^{18}\text{O}$  values are realized using *locally.weighted.polynomial* function (the confidence intervals are created using the row-wise method of Hannig and Marron, 2006) of SiZer Package (R software) with a bandwidth “h” at 1 Ma for both curves.

**Fig. 8.** A. Comparison between  $\delta^{18}\text{O}$ -derived temperatures and  $\Delta_{47}$ -derived Temperatures.  $\delta^{18}\text{O}$ -derived temperatures are calculated using the  $\delta^{18}\text{O}_{\text{sw}}$  from Billups and Scharg (2002) and the equation of Anderson and Arthur (1983) as described in the text, Sections 5.1 and 5.2.1. Black plain circles correspond to Carry-Le-Rouet section; Grey plain circles correspond to Castillon-Du-Gard Section. Vertical error bars reported on 2 bulk shells, representing an uncertainty on the stratigraphic position of the samples, correspond to bulk shells that were gathered from stratigraphic levels new to each other into a single analysis. Black smoothed curve and associated grey interval have been realized using the *locally.weighted.polynomial* function with a bandwidth “h” at 1 Ma.  $\Delta_{47}$ -derived mean annual temperatures are estimated from clumped isotope analyses bulk shells using the equation of Peral et al. (2018): Orange squares: mean annual temperatures; Red square: summer seawater CAR17-6-33-b); Blue square: winter seawater temperatures (CAR17-6-33-b). Vertical error bars as for  $\delta^{18}\text{O}$ -derived temperatures. B. Evolution of local  $\delta^{18}\text{O}_{\text{sw}}$  calculated from  $\delta^{18}\text{O}_{\text{shell}}$  and  $\Delta_{47}$ -derived temperatures using Anderson and Arthur (1983) equation (see section 5.2.3 for further detail) C. Global  $\delta^{18}\text{O}_{\text{sw}}$  curve, is from Billups and Schrag (2002). The shaded areas reported on this curve highlight the portions of the curve selected to calculate the average  $\delta^{18}\text{O}_{\text{sw}}$  value of the global ocean for the three periods for which coupled  $\Delta_{47}$ - $\delta^{18}\text{O}$  analyses have been performed on bivalves, with a minimum of 3 data of the curve gathered for the calculation. D. Salinity estimations are calculated using Railsback and Anderson (1989) equation based on North Atlantic Ocean and local  $\delta^{18}\text{O}_{\text{sw}}$  values from which global ocean  $\delta^{18}\text{O}_{\text{sw}}$  value has been subtracted (see section 5.2.3 for further detail) E. Bivalve  $^{87}\text{Sr}/^{86}\text{Sr}$  evolution (white squares) compared to the  $^{87}\text{Sr}/^{86}\text{Sr}$  of the global ocean (black line; McArthur and Howarth, 2004) with the two red dashed lines showing the variability of  $\pm 50$  ppm recorded in the modern ocean (El Meknassi et al., 2018), and compared to published planktonic foraminifera (Cornacchia et

1497 al., 2018) and mixed fossils (Kocsis et al., 2008)  $^{87}\text{Sr}/^{86}\text{Sr}$  from the North Western  
1498 Mediterranean Sea.

1499

1500 **Fig. 9.** A.  $\delta^{18}\text{O}_{\text{shell}}$  values versus  $\Delta_{47}$ -derived temperatures measured with clumped isotopes.

1501 B. Local  $\delta^{18}\text{O}_{\text{sw}}$  values calculated with Anderson and Arthur (1983) equation versus  $\delta^{18}\text{O}_{\text{shell}}$   
1502 values measured with clumped isotopes.

1503

**Figure 1** Geographical and stratigraphic context of the study area. **A** Map of Europe showing the location of the Rhodano-provençal basin (Castillon-Du-Gard) in southern France, between the Atlantic Ocean and the Mediterranean Sea. **B** Detailed map of the Carry-Le-Rouet area, showing the Mediterranean Sea, the Gard River, and various geological features like the Phare de Carry-le-Rouet and Cap de Nautes. **C** Map of the Castillon-Du-Gard area, highlighting the Vers and Castillon-Du-Gard sites. A vertical scale on the left indicates age in millions of years (m.y.) from 12 to 23, with corresponding geological stages: Serravallian, Langhian, Burdigalian, and Aquitanian. Two sedimentary sections are detailed: Carry-Le-Rouet and Castillon-Du-Gard. The Carry-Le-Rouet section includes Quaternary alluvions, Budigalian-Langhian deposits, Biodebtal Sausset-Les-Pins Unit, Bioclastic Carry Unit, Brackish Rousset Unit, Parareefal Cap de Nautes Unit, Conglomeratic Rouet Unit, and Basement (Jurassic-Cretaceous). The Castillon-Du-Gard section includes Quaternary, Pliocene, Castillon Molasse, Blue Marls, Uzès Molasse, Greenish Molasse, and Lower Cretaceous. Regional stages 'Helvetian' and 'Burdigalian' are also indicated.

Figure 2






Figure 3






Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Table 1

Facies	Non-biotic elements		Biotic components	Matrix	Sedimentary structures	Environments
A1. Marls to fine siltstones	Rhizo-concretion, root traces			Clayey silts		Palaeosoils, Floodplain
A2. Conglomerates	Monogenic	Clasts source: Rudist Limestone (Barremian of the Nerthe)		Heterogeneous and clast-supported conglomerates with a sandstone matrix	Erosional base of decimetric to meter large channels	Fluviatile
	Poligenic	Clasts source : Rudist Limestones (Barremian of the Nerthe) and Paleozoic rocks				
A3. Conglomerates	Polygenic conglomerate with clasts composed of Rudist Limestones (Barremian of the Nerthe) and Paleozoic rocks		Rare bivalves and gastropods			Proximal deltaic domain
A4. Sandstones	Quartz grains, feldspath		Rare bivalves and gastropods	Calcareous sandstone	Continuous cm-thick beds intercalated with oyster and bioturbated sandstones	Distal deltaic lobes, transitional continental to marine
B1.Sandstones with conglomerates	Sandstones with quartz		Common bivalves, gastropods and barnacles	Fine grain sandstones	Planar stratification Bioturbation	Upper shoreface
B2. Fine to coarse sandstones	Quartz grains		Abundant bivalves Common gastropods Rare wood fragments and bryozoans	Fine grain sandstones	Megaripples, HCS (Hummocky cross stratification), Convolute beds, Bioturbation	Lower shoreface
C. Siltstones to marls	Quartz grains		Common wood fragments and foraminifera Rare bivalves	Silt and Marls	Bioturbation ( <i>Thalassinoides</i> / <i>Planolites</i> ), HCS	Lagoon, protected environments
D1. Bioclastic packstones	Quartz grains		Abundant bivalves, gastropods and corals Common bryozoans and echinoids	Broken bioclastic grains and micritic matrix	Bioturbation	Upper to lower shoreface
D2. Coral-rich floatstones to framestones	Floatstones	Reworked corals, and fragmented bioclasts	Abundant corals Common bivalves, gastropods and bryozoans		Bioturbation, bioerosion, lenticular bioclastic accumulation and sigmoid reefal units	Lower shoreface
	Framestones	In situ corals, embedded in a bioclastic matrix				

Table 2.

Age (m.y)	Sample	N	Seasonality	$\delta^{18}\text{O}_{\text{shell}}$ (‰VPDB)	$\Delta_{47}$ (‰, $\pm 1 \sigma_E$ )	T (°C)	$\delta^{18}\text{O}_{\text{sw}}$ calculated (‰VSMOW)	$\delta^{18}\text{O}_{\text{sw}}$ (Billups and Schrag, 2002) (‰VPDB)	Salinity (‰)
22.9	CAR17_4_21A_a_B1	3	Bulk	-2.02	$0.6903 \pm 0.0093$	$19.9 \pm 2.8$	$-1.06 \pm 0.68$	-0.66	$34.3 \pm 1.1$
22.3	CAR17_6_28_a_B1	3	Bulk	-3.79	$0.705 \pm 0.0095$	$15.6 \pm 2.7$	$-3.9 \pm 0.65$	-0.66	$29.6 \pm 1.1$
22.2	CAR17_6_30TOP_B1	2	Bulk	-2.37	$0.6934 \pm 0.0097$	$19 \pm 2.9$	$-1.68 \pm 0.7$	-0.66	$33.3 \pm 1.2$
22.2	CAR17_6_31BASE_a_B1	3	Bulk	-2.96	$0.6852 \pm 0.0093$	$21.5 \pm 2.9$	$-1.67 \pm 0.7$	-0.66	$33.3 \pm 1.2$
22.1	CAR17_6_33_b_S1	3	Summer	-3.58	$0.6821 \pm 0.01$	$22.4 \pm 3.1$	$-2.05 \pm 0.75$	-0.66	$32.7 \pm 1.3$
22.1	CAR17_6_33_b_W1	3	Winter	-1.3	$0.6823 \pm 0.0099$	$22.4 \pm 3.1$	$0.25 \pm 0.75$	-0.66	$36.5 \pm 1.3$
22.1	CAR17_6_33_c_B1	3	Bulk	-1.53	$0.6986 \pm 0.01$	$17.5 \pm 3$	$-1.14 \pm 0.72$	-0.66	$34.2 \pm 1.2$
18.1	CAR17_8_43_h_B1	3	Bulk	0	$0.6801 \pm 0.0093$	$23.1 \pm 2.9$	$1.71 \pm 0.70$	-0.94	$39.4 \pm 1.2$
18.1	CAR17_8_43_i_B1	3	Bulk	0.12	$0.6882 \pm 0.0093$	$20.5 \pm 2.8$	$1.19 \pm 0.68$	-0.94	$38.5 \pm 1.1$
17.7	CAR17_8_45_a_B1	3	Bulk	-1.13	$0.6762 \pm 0.0093$	$24.3 \pm 2.9$	$0.9 \pm 0.70$	-0.94	$38.1 \pm 1.2$
14.9	CDG18_3&4_BASE_B1	3	Bulk	-2.37	$0.7157 \pm 0.0099$	$12.5 \pm 2.8$	$-3.25 \pm 0.68$	-0.71	$30.8 \pm 1.1$
14.6	CDG18_3&4_TOP_B1	3	Bulk	-2.59	$0.7137 \pm 0.0099$	$13.1 \pm 2.8$	$-3.3 \pm 0.68$	-0.71	$30.7 \pm 1.1$
14.2	CDG18_5_1_B1	3	Bulk	-2.98	$0.6968 \pm 0.0098$	$18 \pm 2.9$	$-2.52 \pm 0.70$	-0.71	$32 \pm 1.2$



Table 3

Age (m.y)	Sample	Nature	$^{87}\text{Sr}/^{86}\text{Sr}$	+ 2 $\sigma_E$
22.9	CAR17_4_21A_a_B1	Bulk	0.708255	0.000005
22.3	CAR17_6_28_a_B1	Bulk	0.708299	0.000007
22.2	CAR17_6_30TOP_B1	Bulk	0.708289	0.000005
22.2	CAR17_6_31BASE_a_B1	Bulk	0.708266	0.000005
22.1	CAR17_6_33_b_B1	Bulk	0.708228	0.000005
22.1	CAR17_6_33_c_B1	Bulk	0.708271	0.000009
18.1	CAR17_8_43_h_B1	Bulk	0.708696	0.000005
18.1	CAR17_8_43_i_B1	Bulk	0.708673	0.000006
17.7	CAR17_8_45_a_B1	Bulk	0.708695	0.000005
14.9	CDG18_3&4_BASE_B1	Bulk	0.708873	0.000005
14.6	CDG18_3&4_TOP_B1	Bulk	0.708868	0.000006
14.2	CDG18_5_1_B1	Bulk	0.708849	0.000005