High-Quality Draft Genome Sequence of Xanthomonas alfalfae subsp. alfalfae Strain CFBP 3836.

Marie Agnes Jacques, Stéphanie S. Bolot, Erika Charbit, Armelle Darrasse, Martial Briand, Matthieu Arlat, Lionel Gagnevin, R Koebnik, Laurent Noel, Perrine Portier, et al.

▶ **To cite this version:**

Marie Agnes Jacques, Stéphanie S. Bolot, Erika Charbit, Armelle Darrasse, Martial Briand, et al.. High-Quality Draft Genome Sequence of Xanthomonas alfalfae subsp. alfalfae Strain CFBP 3836.. Genome Announcements, 2013, 1 (6), 10.1128/genomeA.01035-13 . hal-02649259

HAL Id: hal-02649259
https://hal.inrae.fr/hal-02649259

Submitted on 9 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution 4.0 International License

genomeA
Journals.ASM.org

High-Quality Draft Genome Sequence of *Xanthomonas alfalfae* subsp. *alfalfae* Strain CFBP 3836

M. A. Jacques,[a,b,c] S. Bolot,[d,e] E. Charbit,[a,b,c] A. Darrasse,[a,b,c] M. Briand,[a,b,c] M. Arlat,[d,e,f] L. Gagnevin,[g] R. Koebnik,[h] L. D. Noël,[d,e] P. Portier,[a,b,c] S. Carrère,[d,e] T. Boureau[a,b,c]

INRA, Institut de Recherche en Horticulture et Semences (IRHS), UMR 1345 SFR 4207 QUASAV, Beaucouzé, France[a]; Université d'Angers, Institut de Recherche en Horticulture et Semences (IRHS), UMR 1345 SFR 4207 QUASAV, Beaucouzé, France[b]; Agrocampus Ouest, Institut de Recherche en Horticulture et Semences (IRHS), UMR 1345 SFR 4207 QUASAV, Beaucouzé, France[c]; INRA, Laboratoire des Interactions Plantes Micro-Organismes (LIPM), UMR 441, Castanet-Tolosan, France[d]; CNRS, Laboratoire des Interactions Plantes Micro-Organismes (LIPM), UMR 2594, Castanet-Tolosan, France[e]; Université de Toulouse, Université Paul Sabatier, Toulouse, France[f]; UMR "Peuplements Végétaux et Bioagresseurs en Milieu Tropical" (PVBMT), CIRAD, Saint-Pierre, La Réunion, France[g]; UMR 186 IRD-Cirad-Université Montpellier 2 "Résistance des Plantes aux Bioaggresseurs," Montpellier, France[h]

M.A.J. and S.B. contributed equally to this study.

We report the high-quality draft genome sequence of *Xanthomonas alfalfae* subsp. *alfalfae* strain CFBP 3836, the causal agent of bacterial leaf and stem spot in lucerne (*Medicago sativa*). Comparative genomics will help to decipher the mechanisms provoking disease and triggering the defense responses of this pathogen of the model legume *Medicago truncatula*.

Received 2 November 2013 **Accepted** 11 November 2013 **Published** 12 December 2013

Citation Jacques MA, Bolot S, Charbit E, Darrasse A, Briand M, Arlat M, Gagnevin L, Koebnik R, Noël LD, Portier P, Carrère S, Boureau T. 2013. High-quality draft genome sequence of *Xanthomonas alfalfae* subsp. *alfalfae* strain CFBP 3836. Genome Announc. 1(6):e01035-13. doi:10.1128/genomeA.01035-13.

Copyright © 2013 Jacques et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

Address correspondence to M. A. Jacques, marie-agnes.jacques@angers.inra.fr.

Xanthomonas alfalfae subsp. *alfalfae* (formerly *Xanthomonas axonopodis* pv. alfalfae) is the causal agent of bacterial leaf and stem spot of lucerne (*Medicago sativa*) (1, 2). While lucerne is the main host of *X. alfalfae* subsp. *alfalfae*, other hosts, such as soybean (*Glycine max*), clover (*Trifolium* spp.), and vetch (*Vicia* spp.), have been reported (3). *X. alfalfae* subsp. *alfalfae* is also pathogenic on the model legume *Medicago truncatula* (4). Lesions first appear on the leaf as water-soaked spots surrounded by a diffuse chlorotic area, which then turns dry, yellow-brown, and papery. Severe defoliation is a common result of leaf infection. Postemergence damping off, stunting of seedlings, and damage of lucerne stands are other symptoms. This seed-borne disease may cause severe losses in hot, moist environments (3). To better understand the molecular basis of legume-bacterium interactions, we sequenced the genome of the *X. alfalfae* subsp. *alfalfae* strain CFBP 3836. This strain is the pathotype strain of the former pathovar *X. axonopodis* pv. alfalfae (5). This strain originates from the Sudan (http://www.angers-nantes.inra.fr/cfbp/resultnum_e.php?r0=3836).

This genome was sequenced using the Illumina HiSeq 2000 platform (GATC Biotech, Germany). Shotgun sequencing yielded 116,576,062 read pairs (37,799,928 100-bp paired-end reads with an insert size of ca. 250 bp and 78,776,134 bp mate-pair reads with an insert size of approximately 3 kb). A combination of Velvet (6), SOAP*denovo*, and SOAP Gapcloser (7) yielded 22 contigs >500 bp (N_{50}, 763,181 bp), with the largest contig being 2,433,808 bp, for a total assembly size of 5,077,532 bp. Multilocus sequence analyses of seven housekeeping genes described earlier for xanthomonads confirmed that strain CFBP 3836 belongs to *X. alfalfae* subsp. *alfalfae* (8).

The genome carries the core characteristics shared by most plant-pathogenic xanthomonads, including chemotaxis- and flagellum-encoding genes, many genes encoding two-component systems, and TonB-dependent transporters. This genome sequence displays the genes coding for the six secretion systems (from T1SS to T6SS) that have been identified so far in Gram-negative bacteria and multiple effectors. A minimum repertoire of 23 type III effectors is predicted. It comprises two putative transcription activator-like effectors, the highly repetitive sequence of which could not be completely resolved with this sequencing strategy. This genome sequence will be a valuable tool for transcriptomic and genetic studies to allow for a better understanding of disease development and plant defense triggering in legume-bacterium interactions.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited in GenBank under the accession no. AUWN00000000. The version described in this paper is the first version, AUWN01000000.

ACKNOWLEDGMENTS

This work was supported by the French Agence Nationale de la Recherche (no. ANR-2010-GENM-013) and the LABEX TULIP (no. ANR-10-LABX-41).

REFERENCES

1. **Riker AJ, Jones FR, Davis MC.** 1935. Bacterial leaf spot of alfalfa. J. Agric. Res. **51**:177–182.
2. **Esnault R, Buffard D, Breda C, Sallaud C, el Turk J, Kondorosi A.** 1993. Pathological and molecular characterizations of alfalfa interactions with compatible and incompatible bacteria, *Xanthomonas campestris* pv. *alfalfae* and *Pseudomonas syringae* pv. *pisi*. Mol. Plant Microbe Interact. **6**:655–664.
3. **Bradbury JF.** 1981. *Xanthomonas campestris* pv. *alfalfae*. IMI Descrip. Fungi Bact. **70**:698. http://www.cabi.org/dfb/?loadmodule=review&page=4048&reviewid=9807&site=159.
4. **Horning JL, Stuteville DL, White FF.** 2002. Stunting of 10 accessions of

Medicago truncatula by *Xanthomonas axonopodis alfalfae*. Phytopathology **92:**S138. doi:10.1094/PHYTO.2002.92.6.S137.

5. **Bull CT, De Boer SH, Denny TP, Firrao G, Fischer-Le Saux M, Saddler GS, Scortichini M, Stead DE, Takikawa Y.** 2010. Comprehensive list of names of plant pathogenic bacteria, 1980–2007. J. Plant Pathol. **92:** 551–592.
6. **Zerbino DR, Birney E.** 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. Genome Res. **18:**821–829.
7. **Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu SM, Peng S, Xiaoqian Z, Liu G, Liao X, Li Y, Yang H, Wang J.** 2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. Gigascience **1:**18. doi:10.1186/2047-217X-1-18.
8. **Mhedbi-hajri N, Hajri A, Boureau T, Darrasse A, Durand K, Brin C, Fischer-Le Saux M, Manceau C, Poussier S, Pruvost O, Lemaire C, Jacques MA.** 2013. Evolutionary history of the plant pathogenic bacterium *Xanthomonas axonopodis*. PLoS One **8:**e58474. doi:10.1371/journal.pone.0058474.