

Behavioural traits modulate the use of heterospecific social information for nest site selection: experimental evidence from a wild bird population

Jennifer Morinay, Jukka Forsman, Marion Germain, Blandine Doligez

► To cite this version:

Jennifer Morinay, Jukka Forsman, Marion Germain, Blandine Doligez. Behavioural traits modulate the use of heterospecific social information for nest site selection: experimental evidence from a wild bird population. Proceedings of the Royal Society B: Biological Sciences, 2020, 287 (1925), pp.20200265. 10.1098/rspb.2020.0265. hal-02990528

HAL Id: hal-02990528 https://cnrs.hal.science/hal-02990528v1

Submitted on 24 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Behavioural traits modulate the use of heterospecific social information for nest
2	site selection: experimental evidence from a wild bird population
3	
4	
5	Jennifer Morinay ^{1,2,*} , Jukka T. Forsman ^{3,4} , Marion Germain ^{1,2,5} and Blandine Doligez ¹
6	
7	¹ Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie
8	Evolutive UMR 5558, France
9	² Department of Ecology and Evolution, Animal Ecology, Evolutionary Biology Centre,
10	Uppsala University, Sweden
11	³ Department of Ecology and Genetics, University of Oulu, Finland
12	⁴ Natural Resources Institute Finland (Oulu), University of Oulu, Finland
13	⁵ Department of Animal and Plant Sciences, University of Sheffield, United Kingdom
14	
15	* Correspondance: LBBE, CNRS UMR5558, Université Claude Bernard Lyon 1, Bâtiment
16	Gregor Mendel, 43 boulevard du 11 novembre 1918, 69622 Villeurbanne Cedex, France.
17	Email: jennifer.morinay@gmail.com
18	
19	
20	
21	Keywords: collared flycatcher, copying, personality, boldness, aggressiveness, neophobia

22 Abstract

23

The use of social information for making decisions is common but can be constrained by 24 behavioural traits via e.g. the ability to gather information. Such constrained information use 25 has been described in foraging habitat selection; yet it remains unexplored in the breeding 26 habitat selection context, despite potentially strong fitness consequences. We experimentally 27 tested whether three behavioural traits (aggressiveness, boldness, neophobia) affected the use 28 29 of heterospecific social information for nest site selection in wild collared flycatchers Ficedula albicollis. Flycatchers have previously been found to copy or reject an artificial 30 apparent preference of tits (their main competitors) for a nest site feature: they preferred nest 31 boxes with the same or a different feature, depending on tit early reproductive investment. 32 Here, we confirmed this result and showed that, shy individuals and less aggressive old males 33 34 (i.e. 2 years old or older) copied tit apparent preference, while more aggressive old males rejected the tit preference. Aggressiveness and boldness may allow males to access more 35 36 information sources or affect males' interactions with dominant tits when selecting a nest site. Our study highlights the links between variation in behaviours and social information use for 37 breeding habitat selection and calls for further work to explore underlying mechanisms. 38

39 Introduction

40

In spatio-temporally variable environments, individuals can use a great variety of information 41 42 to make decisions. In particular, they can use personal information (derived from their own knowledge about - or experience with - the environment) and/or social information (derived 43 from observing other individuals' actions in the environment; [1,2]). Depending on the 44 relative reliability and availability of these two types of information, individuals can flexibly 45 use personal and/or social information [e.g. 3,4]. Social information use is known to depend 46 on environmental conditions [e.g. population size, spatio-temporal predictability; 5,6] but also 47 48 on individual factors, such as age [7] or personality traits [8]. Personality traits may constrain the use of social information by affecting either the propensity to acquire information or the 49 decisions made once information is acquired. Personality traits may in particular shape 50 individual's willingness to prospect in general (activity), and more specifically in new or risky 51 environments (exploration and boldness), or when prospecting involves social interactions 52 53 (aggressiveness and sociality); thereby, they may affect individuals' overall knowledge of the 54 environment.

Thus far, 24 published studies (to our knowledge) have investigated the links between 55 social information use and personality traits, mostly in the context of foraging decisions 56 (Table 1; see Table S1 for full details). Among these studies, the investigation of the 57 exploration / neophobia axis [8] was predominant (19 over 24 studies, i.e. 79%). Higher 58 neophobia level was frequently associated with higher social information use (in 7 59 relationships over 11; Table 1). Conversely, no overall pattern was found for the links 60 between social information use and other personality traits, either because most relationships 61 62 were non-significant (for exploration and boldness) or very few studies (or even none) investigated these links (for activity, sociality and in particular aggressiveness; Table 1, Table 63

64 S1). Furthermore, testing the causality of links between personality traits and social 65 information use requires experimentally manipulating information sources. Yet, such 66 manipulations have only been conducted in captivity thus far. Therefore, the extent to which 67 different personality traits, but also more flexible behavioural traits in general may favour or 68 constrain the use of social information for decision-making in the wild remains poorly 69 understood.

Social information use for breeding habitat selection and dispersal decisions is well 70 documented [e.g. 5,9,10]. In parallel, dispersal syndromes involving behavioural traits have 71 been well studied [e.g. 11,12]. However, to our knowledge, no study has directly investigated 72 73 the link between behavioural traits and social information use for breeding site choice (Table S1). Yet, prospecting to gather social information on potential breeding sites can be costly in 74 terms of time, energy and increased agonistic interactions with competitors [13] and only 75 76 individuals displaying specific behaviours may be able to face these costs. For example, more aggressive, bold and/or explorative individuals may have access to more and/or larger-scale 77 78 social information sources. Social information use itself may also increase intra- and interspecific competition when individuals spatially aggregate because of con- or hetero-79 specific attraction or because they use the same information [6,9]. Therefore, the realised 80 breeding site choices may notably depend on aggressiveness allowing individuals to acquire 81 and defend the chosen site/territory against competitors. Assessing to what extent behavioural 82 traits shape social information use for breeding site choice is needed to understand how 83 selective pressures act on behaviour over different decision-making contexts. 84

Here, we tested whether difference in the use of an experimentally manipulated source of social information for nest site selection was related to three main behavioural traits (aggressiveness, boldness and neophobia), previously shown to be partly repeatable [14], in a natural population of a small passerine bird, the collared flycatcher *Ficedula albicollis*.

Collared and pied flycatchers F. hypoleuca (a sister species) have been repeatedly shown to 89 use social information from con- and heterospecific (titmice) competitors when choosing a 90 nest site [5,10,15–19]. However, this social information use shows high between-individual 91 variability, only partly explained by sex [5], age [20] or dispersal status [15], and depends on 92 years [21] and environmental conditions (e.g. clutch size [20], or titmice density [24]). We 93 tested here whether differences in the three behavioural traits investigated could explain part 94 of the observed variability in social information use. Using an experimental design already 95 successfully implemented in our study species, we created an apparent local preference of 96 dominant tutors (here tits) for a specific nest box feature observable from a distance 97 (geometric symbols) [10,23]. We then recorded whether flycatchers copied or rejected this 98 preference by settling in boxes displaying the same feature. After settlement, we measured 99 levels of aggressiveness, boldness and neophobia of the experimental birds to test the link 100 101 between these behavioural traits and the probability of copying tit apparent preference. Due to potential risks of collecting information at the vicinity of tit territories, we expected 102 103 aggressive individuals to be more likely to copy tutors' preference than less aggressive ones. 104 Furthermore, shyness (lack of boldness) and/or neophobia could restrain access to other conspecific or heterospecific information sources besides tit apparent preference (e.g. if they 105 affect the gathering of information available at a large-scale, for neophobia [5], or risky to 106 107 acquire, for boldness [9]), and thus shyer and/or more neophobic individuals could be expected to be more likely to copy tutors' preference than less neophobic and/or bolder ones. 108

109

110

111 Methods

112

113 Species and study site

The experiment was conducted in spring 2012 and 2013 in a wild breeding population of 114 collared flycatchers on the island of Gotland (Baltic Sea, Sweden). Collared flycatchers are 115 sexually dimorphic migratory hole-nesting passerine birds that readily breed in artificial nest 116 boxes provided in the forest patches of the study area. Breeding flycatchers were captured in 117 boxes (during incubation for females and chick rearing for males) as part of the long-term 118 monitoring of the population. Caught individuals were measured and aged based on plumage 119 criteria (yearling vs. older individuals. In this population, collared flycatchers compete for 120 121 nest boxes with great tits Parus major and blue tits Cyanistes caeruleus (Gustafsson 1988), which are resident passerine species, are competitively dominant over flycatchers and 122 typically start laying on average two weeks before flycatchers' arrival (but see Table S2 and 123 [21]). 124

125

126 *Heterospecific preference copying: experimental design*

In 12 (in 2012) and 17 (in 2013) experimental forest patches (2,048 nest boxes in total over 127 128 the two years), we created an apparent preference of tits for a specific nest box feature to measure flycatchers' subsequent copying behaviour by attaching around the entrance of boxes 129 one of two geometric symbols (white plastic shapes; either a triangle or a circle) depending on 130 the species occupying the box [21]. Before flycatchers' arrival (i.e. in the first two weeks of 131 April), we attached on all boxes occupied by great and blue tits (and the few coal tits 132 Periparus ater) in a given forest patch the same symbol (shape alternated between patches, 133 see Figure S1 for more details). At the same time, we randomly attached a triangle on half of 134 the remaining (empty) boxes, i.e. boxes available for newcomers' settlement, and a circle on 135 the other half. We attached no symbol on the few boxes occupied by other species (nuthatches 136 Sitta europaea, sparrows Passer domesticus and P. montanus and wrynecks Jynx torquilla; 25 137 boxes in total over the two years). Therefore, when flycatchers arrived from migration (late 138

April to mid-May), they had the choice between copying tit preference by settling in a box 139 with the same symbol as on tit boxes, or rejecting it by settling in a box with the opposite 140 symbol. When a flycatcher pair had settled in a box, as shown by the presence of new nest 141 material in the box, we removed the symbol on this box. This avoided providing conspecific 142 information via the symbol chosen to later arriving flycatchers. We checked empty boxes 143 every other day to detect newly started nest building and removed (for new flycatcher nests) 144 or changed if needed (for new tit nests) the symbol accordingly. At the same time, we 145 adjusted the number of triangles and circles on empty boxes within a forest patch to keep an 146 equal proportion of available boxes displaying each symbol, and thus an equal probability for 147 newcomers to choose a symbol at random. Because this equal proportion of both symbols 148 could not always be met (e.g. when an odd number of empty boxes remained in a patch), we 149 controlled for the deviation from random (0.5) of the proportion of empty boxes matching the 150 151 tit apparent preference within a plot on the day of choice for each flycatcher pair [see 21 for more details]. Because we can assume that flycatchers naïve to the experimental design have 152 153 no previous experience with geometric symbols, this design minimises genetic and ecological effects on nest site choice and is a powerful method to reveal factors affecting individuals' 154 choices [10]. 155

156

157 Measuring behavioural traits

The three behavioural traits of interest here, namely aggressiveness, boldness and neophobia, were measured as described in a former study on the same population (see [14] for detailed methods). In this former study based on a larger sample [14], all three traits were found to be weakly repeatable between years (R=0.2, 0.1 and 0.4 for aggressiveness, boldness and neophobia respectively) and weakly phenotypically correlated (-0.2 for aggressivenessneophobia, -0.3 for boldness-neophobia) but they did not associate in behavioural syndromes (i.e. no between-individual covariance between them [14]). In the present study, we refer to these traits as behavioural rather than personality traits, because we could not separate the effect of the repeatable vs. flexible part of the traits on the use of social information; indeed the copying behaviour was measured only once (i.e. in naïve birds).

We measured aggressiveness through the agonistic response of a focal pair to a 168 simulated intrusion by competitors on the nest box during nest building stage, i.e. when the 169 risk of losing a nest site is highest [as in 14]. We used both conspecific and heterospecific 170 (great tit) decoys (in successive tests) because flycatchers respond aggressively to both 171 species [24,25]. A total of 2 to 4 tests were conducted for each focal pair (1 or 2 tests per 172 173 stimuli species, depending on field constraints), with one test maximum per day and 2 days maximum in a row to avoid habituation. The decoy species was randomized for the first test 174 and alternated between subsequent tests. At the start of a test, an observer attached on the box 175 decoys of either a flycatcher pair or a male great tit, randomly chosen among 10 different sets 176 for each species, as well as a loudspeaker broadcasting songs of the same species as the 177 178 decoy(s), randomly chosen among 5 different song tracks per species. The observer then hid under a camouflage net approximately 8-10 meters away from the box and recorded all 179 behaviours performed by each member of the focal pair during 15 minutes on average (mean 180 15.12 minutes \pm 0.96 SD): movements around- and distance from- the box, flights and attacks 181 towards a decoy or live birds attracted by the stimulus. To account for differences in the 182 latency to respond between individuals, each behavioural variable recorded was converted 183 into frequency per minute using the time interval between the first observation of the 184 individual during the test and the end of the test. We then estimated an aggressiveness score 185 for each individual and for each test as the sum of the frequencies of (i) movements within 2 186 meters from the box, (ii) attacks or stationary flights towards a decoy and (iii) chases towards 187 live intruders [similarly to 14]. We excluded from the datasets individuals that were observed 188

189 less than 5 minutes. In total, we used 1168 behavioural responses of both sexes, performed 190 during 790 aggressiveness tests on 224 males and 271 females and 313 reproductive events 191 over the two years of the experiment. The final individual aggressiveness score was calculated 192 as the average of the scores measured for each individual within one season.

We measured boldness through the reaction to the presence of a human observer near 193 the box and neophobia through the reaction to the presence of a novel object attached on the 194 box (i.e. in a familiar environment) [as in 14]. We conducted one combined boldness / 195 neophobia test per breeding pair per year, when chicks were 5 or 6 days old. The test 196 consisted of two consecutive periods lasting one hour each, during which the provisioning 197 behaviour of both parents was video-recorded from a distance (6-8m). In the first period, an 198 observer settled a recorder and opened the box to check chick satiety before leaving the area. 199 In the second period, the observer came back to the box, checked chick satiety again, attached 200 201 a novel object (here a coloured figurine approximately 7 cm high) near the entrance of the box and left again for one hour. Chick satiety was checked in order to avoid performing 202 203 behavioural tests if chicks' condition was too poor. We estimated a boldness score for each parent based on the latency to enter the box after the observer's departure in the first period of 204 the test, i.e. without the novel object. To obtain meaningful boldness scores (i.e. increasing 205 boldness for decreasing latency), we subtracted this latency from the maximum latency 206 observed in our data set [as in 14]. We estimated a neophobia score for each parent based on 207 the latency to enter the box after the departure of the observer in the second period of the test, 208 i.e. in the presence of the novel object. Among the 318 individuals that entered the box in the 209 210 first period, 38% did not enter in the second period and thus had no latency available. To take into account those highly neophobic individuals, we discretized neophobia as a 5-level score, 211 212 with the first four levels corresponding to latency quartiles and the last level assigned to these non-returning individuals [as in 14]. Results however remained quantitatively unchanged (not 213

detailed here) when considering neophobia as a continuous latency instead of a score by attributing a maximum latency to non-returning individuals (here 4,000 seconds, the maximum observed latency plus one minute).

217

218 *Statistical analyses*

We analysed the probability for flycatchers to copy the apparent preference of tits (binary 219 response variable: copy vs. reject) in the two years of the experiment (2012 and 2013). In the 220 221 second year, we retained only individuals naïve to the symbol experiment, i.e. which had not been caught as breeders in the first year. The overall lower number of males in the sample and 222 223 differences in sample sizes between models were mostly due to early breeding failures (before the boldness/neophobia test and/or male capture). Because aggressiveness, boldness and 224 neophobia are slightly phenotypically correlated within individuals [14], we fitted separate 225 models for each trait. Furthermore, because nest site choice is a joint decision by both pair 226 members, the most appropriate model to estimate the effect of individual behavioural traits on 227 228 the joint copying decision would include both male and female trait estimates simultaneously. However, retaining only nests where both pair members have been captured and aged, are 229 naïve to the symbols and have responded to behavioural tests strongly reduced sample size 230 (by up to 33%). Therefore, we first fitted sex-specific models. Second, we fitted models with 231 both male and female estimates of the behavioural trait and age, the same other main effects 232 as above, the two-way interaction between male and female behavioural trait estimates, and 233 the interactions that were significant in the sex-specific models. In total, we fitted 6 sex- and 234 behaviour-specific models (see Table S3 for the full models) and 3 behaviour-specific models 235 with both male and female trait estimates simultaneously (see Table S4 for the full and final 236 models). 237

Models included as fixed effects the individual's behavioural trait estimate considered 238 (aggressiveness, boldness or neophobia score), individual's age (yearling vs. older), tit density 239 and tit early reproductive investment within the forest patch on the day of flycatcher 240 settlement, and the potential bias in the proportion of empty boxes with each symbol in the 241 patch on the same day. These latter variables have indeed been found to influence the 242 probability of copying tit apparent symbol preference in this and other populations 243 [9,16,21,23]. Tit density was estimated as the proportion of boxes occupied by great tits (i.e. 244 with tit nest material) within the forest patch on the day of flycatcher choice. Tit early 245 investment was measured as the average great tit clutch (or possibly brood for the earliest 246 247 great tit nests) size within the forest patch on the day of flycatcher choice. The bias in the proportion of empty boxes with each symbol was calculated as the proportion of boxes 248 bearing the symbol associated to the tit preference on the day of flycatcher choice minus 0.5. 249 250 To account for age-specific behavioural effects, we included in the models the two-way interaction between age and the behavioural trait estimate considered. We also included the 251 252 two-way interactions of age or the behavioural trait estimate with tit density and tit early reproductive investment. This gave a total of 5 main effects and 5 two-way interactions for 253 each sex- and behaviour-specific model; the maximum number of fixed effects for the models 254 with both male and female trait estimates was 7 main effects and 3 two-way interactions (see 255 full models output in Tables S3-S4). Prior to analyses, all continuous fixed effects were 256 scaled. Finally, we included forest patch and year as random factors to control for potential 257 spatio-temporal effects on social information use. 258

We fitted generalized linear mixed-effects models (GLMM) in R [26] with the *glmer* function (*lme4* R package [27]) and 'binomial' family and selected our fixed effects using a stepwise backward selection procedure. Because the stepwise approach can increase the risk of type-I error [28,29], we checked that the significant effects retained in the final models were significant in the full models too (see Tables S3-S4 for the full models output). Overall, results remained similar when using a model averaging approach (AIC-based selection of subset models with $\Delta AIC < 2$ using the '*MuMIn*' R package [30]; results not detailed). The fit of final models was assessed based on ROC curves, and Areas Under the Curves (AUC), estimated using *pROC* R package [31].

- 268
- 269
- 270 **Results**
- 271

The probability for flycatchers to copy tit preference was affected by male aggressiveness score differently between yearlings and older males (interaction aggressiveness by age; Table 2). Among older males, less aggressive ones significantly copied tit preference, whereas more aggressive ones rejected it (Figure 1a); conversely, there was no relation between the probability of copying tit preference and aggressiveness in yearling males (Figure 1a). Female aggressiveness did not affect the probability of copying tit preference (z-value= -0.47, pvalue= 0.64; Table S3).

In addition, the probability of copying tit preference was affected by boldness score, again differently between yearlings and older individuals, but this time both in males and females (interaction boldness by age; Table 2). Among older individuals of both sexes, shyer ones were more likely to copy tit preference than bolder ones, while the reverse was observed in yearlings, even though in females, 95% Confidence Intervals largely overlapped a random choice, i.e. a probability of copying of 0.5; Figure 1b&c).

Finally, in females, the interaction between neophobia score and tit clutch/brood size seemed to affect the probability of copying tit preference (N= 173, z-value= -2.33, p-value= 0.020; Table 2): for the most neophobic females (neophobia score of 5, i.e. non-returning

females in the presence of the novel object), tit clutch/brood size had no effect on copying, 288 while high tit clutch/brood size was associated with higher probability of copying in other 289 females (neophobia category 1 to 4; Figure S2). However, this interaction was not strongly 290 291 supported in a model averaging approach (relative importance = 0.78) and when the most neophobic females were excluded, no effect of neophobia remained among females with score 292 1 to 4 (z-value= 0.154, p-value= 0.877 for the interaction between female neophobia and tit 293 clutch/brood size; z-value= 0.268, p-value= 0.788 for the simple neophobia effect). This 294 295 suggests that the effect of neophobia was not strong. Male neophobia did not affect the probability of copying tit preference (z-value= 1.34, p-value= 0.18; Table S3). 296

As found previously, both male and female flycatchers were more likely to copy (resp. 297 reject) tit apparent preference when tit clutch / brood size was high (resp. low) in the forest 298 patch on the day of settlement (z-value> 2.73 and p-value ≤ 0.01 over all models; Tables 2, S3; 299 300 Figure S3). The probability of copying tit preference also increased with the bias in the proportion of empty boxes with the symbol associated to tits in the model with female 301 302 aggressiveness (z-value= 2.59, p-value= 0.01 in the final model; see Table S3) but not in other 303 models (Table S3). Tit density did not affect the probability of copying tit preference (Table S3). 304

305 Variances associated to forest patch and year were negligible in all models (not 306 detailed here). Including both male and female behavioural trait estimates simultaneously in 307 models led to similar results (Table S4).

308

309

310 Discussion

We have experimentally shown in our wild bird population that the use of heterospecific 312 social information for nest site selection depended not only on external factors (here, the early 313 reproductive investment of the heterospecific tutors) but also on individual factors, and more 314 particularly on behavioural traits (here, male aggressiveness and both parents' boldness). 315 Among old males, the probability of copying heterospecific competitors' preference 316 decreased with increasing male aggressiveness. In addition, both parents' boldness score 317 modulated the probability of copying tit apparent preference depending on age: pairs with old 318 319 and shy individuals, on the one hand, and young and bold individuals, on the other hand, were more likely to copy tit preference compared to other pairs. Finally, we found no strong effect 320 of female or male neophobia, even though increased exploration and decreased neophobia [8] 321 could be expected to favour prospecting and thereby large-scale (social and non-social) 322 information gathering and use. This was in contrast with former studies in the foraging 323 context, which usually found neophobia to promote conspecific attraction or scrounging 324 strategies, i.e. foraging strategies based on social information (e.g. [32-38], but see [39]). The 325 326 joint copying behaviour of the pair was therefore affected by different behavioural traits that 327 may in particular impact information access and thus availability but also the ability to cope with the consequences of information use. Flycatcher pairs were besides also more likely to 328 copy apparent preference of tits when average tit clutch / brood size in the patch was high at 329 the time of nest site choice. This is in line with previous results [9,16,17,21,23] and suggests 330 that flycatchers adjusted the use of this heterospecific social information source depending on 331 the quality and/or decisions of tit tutors, on top of their own behavioural traits. 332

333

334 Social information use and male aggressiveness

335 Our results provide clear evidence that aggressiveness, i.e. the agonistic reaction towards 336 competitors, can shape the use of heterospecific social information, with different effects depending on age. Aggressive individuals (especially those high in the dominance hierarchy, e.g. older individuals) could be more likely to acquire social information than less aggressive ones when this involves engaging in agonistic interactions with others, including heterospecifics. Here, however, less aggressive old males copied tit apparent preference and more aggressive ones rejected it, suggesting that all old males could have access to information about tit preference independently from their aggressiveness level.

More aggressive individuals could be expected to be more prone to copy competitors' 343 decisions because they would benefit from competitors' experience [40] while at the same 344 time being able to cope with potentially increased competitive costs. Contrary to this 345 expectation, aggressive males avoided competitors' apparent preference. One possible 346 explanation may be that more aggressive individuals pay a greater cost from competition with 347 tits compared to less aggressive ones, because they engage more in territorial defence. More 348 349 aggressive males may thus reject tit apparent preference to avoid costs of heterospecific agonistic interactions with dominant competitors. Conversely, less aggressive flycatchers may 350 351 engage less in agonistic interactions with tits and thus benefit more from using information from tits. Indeed, even though tits are dominant over flycatchers, they tolerate flycatchers' 352 settlement in the vicinity of their nest (Doligez, pers. obs.). In line with this prediction, house 353 crickets Acheta domesticus with a high Resource Holding Potential were more likely to win 354 355 contests, but if losing, they ended the contest sooner [41]. Assessing whether increased aggressiveness increases the risk of heterospecific agonistic interactions and thus potential 356 costs for flycatchers would be needed to confirm this explanation. 357

358

359 Social information use and boldness

Boldness also affected the probability of copying tit preference: especially in males, copyingprobability was higher for old and shy individuals, as well as young and bold ones, compared

to old and bold, and young and shy ones, respectively (Figure 1). Former studies reported 362 highly contrasting results regarding the link between boldness and social information use 363 (Tables 1 and S1). In some studies, shy individuals were more likely to shoal and follow 364 others [42–44], as a result of higher attention paid to, and higher probability to rely on, others' 365 decisions. Other studies however found that bold individuals paid more attention to others 366 [45], or even no support for boldness to affect the propensity to use three different types of 367 social information [46]. The effect of boldness on social information use may thus be strongly 368 dependent on the context and in particular on social organisation and individual's experience. 369 Here, the link between boldness and social information use depended on age, which likely 370 shapes both competitive ability and experience [47,48] and thereby information access and 371 use. Yearlings may have a restrained access to information, but this effect may be 372 compensated for by boldness. Furthermore, old and bold individuals may have access to 373 374 additional information sources such that only old and shy individuals may rely on tit apparent preference (that can be obtained from a distance with limited risks) over other sources. 375 376 Constraints on the access to social information imposed by the behavioural trait considered 377 may shape the link between this trait and information use.

378

379 *Modulation of social information use or of the response to our behavioural tests?*

The three behavioural traits considered here are only weakly repeatable [14] and thus mostly plastic. Therefore, we cannot exclude that flycatchers adjusted their behavioural responses to our behavioural tests depending on whether they copied tit apparent preference for nest box choice rather than adjusting their copying behaviour depending on their behavioural traits. Yet, our experiment was designed so that choosing a given symbol had no subsequent reproductive consequence for flycatchers, because symbols were randomized in space and thus independent from intrinsic site quality [21]. Post-settlement adjustment of behavioural

responses to our tests would require different levels (or expectance) of competition level or 387 predation risk depending on the symbol chosen. For example, for this mechanism to explain 388 the observed patterns in aggressiveness, old males that rejected tit preference would have had 389 to expect, or to be exposed to, higher competition level by settling in a box displaying the 390 opposite symbol than the one associated with tits, and thus increased their aggressiveness 391 response to defend their nest box. We consider as unlikely such age-specific difference in 392 competition level due to the presence of an artificial nest feature that was removed days (for 393 aggressiveness tests) or weeks (for boldness and neophobia tests) before. Many social factors 394 after settlement are likely to affect flycatchers' behaviour, making the alternative explanation 395 396 of a post-settlement adjustment of behavioural responses unlikely.

397

398 Modulation of heterospecific social information use based on tutors' investment

399 The increase in the probability of copying tit apparent preference with increasing tit clutch / brood size in the patch on the day of choice implies that flycatchers can estimate average tit 400 401 reproductive investment at the patch scale when they settle and use it for modulating nest site 402 choice according to tit preference. This is in line with former experimental results at a smaller scale, showing that pied flycatchers use tit clutch size as social information (i) to choose 403 between two close-by boxes according to the feature (symbol) associated to tit nest [9,16,23] 404 but also (ii) to adjust breeding investment later on ([49], see also [17] for an experimental test 405 of patch choice according to tit phenology). Overall, our results provide clear evidence that 406 flycatchers modulated their use of heterospecific social information obtained from tit apparent 407 preference for nest site features based on other information sources (here, tit early 408 reproductive investment). This modulation did not depend on their behavioural traits, but 409 410 more work is needed to investigate whether behavioural traits can in general affect the relative use of different social information sources. 411

Our study extends the importance of behavioural traits in shaping the use of social information reported in previous studies to the context of breeding habitat selection in the

412

413

information reported in previous studies to the context of breeding habitat selection in the 414 415 wild, using a powerful experimental manipulation of social information. The joint copying behaviour of the pair for nest site selection was likely constrained both by access to social 416 information, explaining the age-dependent link with boldness, and by competitive costs 417 related to the use of social information after gathering it, explaining the link with age-418 419 dependent male aggressiveness. More generally, how behavioural traits affect access to social information and resulting decision-making based on this information may be a prevalent issue 420 421 in explaining among-individual variation in social information use over contexts. Such constraints may have evolutionary consequences through the costs / benefits balance of the 422 use of social information, which may favour functional integration between certain types of 423 424 personality traits and social information use depending on the relative availability and 425 reliability of these and other information sources. The evolution of such trait associations 426 however relies on genetic bases for both social information use and behavioural traits, which 427 was not the case in our system [14,21,50]. Yet, whether the same behavioural traits may be expected to shape social information use in different contexts and/or the use of different types 428 of social information remains to be explored. Theoretical approaches could prove useful in 429 this respect to explore whether features of the decisions to be made (e.g. spatio-temporal 430 scales) may lead to associations between certain behavioural traits and social information use 431 432 across contexts.

433

434

435 Acknowledgements

We thank Lars Gustafsson for providing access to the field site, the landowners from Gotland to allow us collecting data on their properties, all fieldworkers who collected the data, especially Laure Cauchard, Gregory Daniel, Sami M. Kivelä, Uéline Courcoux-Caro and Louise Riotte-Lambert. We are grateful to all the students who contributed to extracting behaviours from the recordings. We also thank Jelmer Samplonius, an anonymous reviewer and the associate editor Sarah Brosnan for constructive comments on a previous version of the manuscript.

443

444 Funding

This work was funded by the Ministère de l'Enseignement Supérieur et de la Recherche and the Department of Ecology and Genetics from Uppsala University (PhD grants to JM and MG), by research grants from Uppsala University (Stiftelsen för Zoologisk Forskning to JM and MG), by the Région Auvergne Rhone-Alpes (Explora'Doc grants to JM) and by the University of Lyon (IDEX mobility grant to JM). JTF was funded by the Kone Foundation, and BD by the Centre National pour la Recherche Scientifique and the Région Auvergne Rhone-Alpes (CIBLE programme).

452

453 Ethics

Permission for catching and ringing adult and young birds was granted every year by the Ringing Centre from the Museum of Natural History in Stockholm (licence nb. 471: M015 to B.D.). Personality tests only required observing individuals from a distance from below a camouflage net or video-recording their behaviour, without catching them, and special care was taken to avoid any detrimental effect of behavioural test on the health of the nestlings (see the Methods section).

461 Data, code and materials

462 The datasets supporting this article have been uploaded as part of the supplementary material

463 (Table S5).

464 **Competing interests**

465 The authors declare no competing interests.

466 Authors' contributions

467 BD designed the study; MG, BD, JTF and many assistants carried out the fieldwork; JM

468 analysed the data, and drafted the manuscript. JM, JTF, MG, and BD critically revised the

469 manuscript. All authors gave approval for publication and agree to be held accountable for the

- 470 work performed therein.
- 471
- 472

473 **References**

- Danchin E, Giraldeau L-A, Valone TJ, Wagner RH. 2004 Public information: from nosy neighbors to cultural evolution. *Science* (80-.). 305, 487–491. (doi:10.1126/science.1098254)
- 477 2. Dall SRX, Giraldeau L-A, Olsson O, McNamara JM, Stephens DW. 2005 Information
 478 and its use by animals in evolutionary ecology. *Trends Ecol. Evol.* 20, 187–193.
 479 (doi:http://dx.doi.org/10.1016/j.tree.2005.01.010)
- 480 3. van Bergen Y, Coolen I, Laland KN. 2004 Nine-spined sticklebacks exploit the most
 481 reliable source when public and private information conflict. *Proc. Biol. Sci.* 271, 957–
 482 62. (doi:10.1098/rspb.2004.2684)
- 483 4. Templeton JJ, Giraldeau L-A. 1996 Vicarious sampling: the use of personal and public
 484 information by starlings foraging in a simple patchy environment. *Behav. Ecol.*485 Sociobiol. 38, 105–114. (doi:10.1007/s002650050223)
- 5. Doligez B, Danchin E, Clobert J, Gustafsson L. 1999 The use of conspecific reproductive success for breeding habitat selection in a non-colonial, hole-nesting species, the collared flycatcher. J. Anim. Ecol. 68, 1193–1206. (doi:10.1046/j.1365-2656.1999.00362.x)
- 490 6. Doligez B, Cadet C, Danchin E, Boulinier T. 2003 When to use public information for
 491 breeding habitat selection? The role of environmental predictability and density
 492 dependence. *Anim. Behav.* 66, 973–988. (doi:10.1006/anbe.2002.2270)
- 493 7. Dugatkin LA, Godin JGJ. 1993 Female mate copying in the guppy (*Poecilia reticulata*): Age-dependent effects. *Behav. Ecol.* 4, 289–292.
 495 (doi:10.1093/beheco/4.4.289)
- 496 8. Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. 2007 Integrating animal
 497 temperament within ecology and evolution. *Biol. Rev.* 82, 291–318.

- 498 (doi:10.1111/j.1469-185X.2007.00010.x)
- 499 9. Loukola OJ, Seppänen J-T, Krams I, Torvinen SS, Forsman JT. 2013 Observed fitness
 500 may affect niche overlap in competing species via selective social information use. *Am.*501 *Nat.* 182, 474–483. (doi:10.1086/671815)
- Seppänen J-T, Forsman JT. 2007 Interspecific social learning: novel preference can be
 acquired from a competing species. *Curr. Biol.* 17, 1248–1252.
 (doi:10.1016/j.cub.2007.06.034)
- 505 11. Dingemanse NJ, Both C, van Noordwijk AJ, Rutten AL, Drent PJ. 2003 Natal dispersal and personalities in great tits (*Parus major*). *Proceedings. Biol. Sci.* 270, 741–7. (doi:10.1098/rspb.2002.2300)
- Cote J, Clobert J, Brodin T, Fogarty S, Sih A. 2010 Personality-dependent dispersal: 508 12. characterization, ontogeny and consequences for spatially structured populations. 509 Lond. Biol. Sci. 4065-4076. 510 Philos. Trans. *R*. Soc. В. 365. (doi:10.1098/rstb.2010.0176) 511
- 512 13. Forsman JT, Thomson RL. 2008 Evidence of information collection from heterospecifics in cavity-nesting birds. *Ibis (Lond. 1859).* 150, 409–412.
 514 (doi:10.1111/j.1474-919X.2007.00773.x)
- Morinay J, Daniel G, Gustafsson L, Doligez B. 2019 No evidence for behavioural syndrome and genetic basis for three personality traits in a wild bird population. *Anim. Behav.* 153, 69–82. (doi:10.1016/j.anbehav.2019.05.001)
- 518 15. Kivelä SM, Seppänen J-T, Ovaskainen O, Doligez B, Gustafsson L, Mönkkönen M,
 519 Forsman JT. 2014 The past and the present in decision-making: the use of conspecific
 520 and heterospecific cues in nest site selection. *Ecology* 95, 3428–3439. (doi:10.1890/13521 2103.1)
- Forsman JT, Seppänen J-T. 2011 Learning what (not) to do: testing rejection and copying of simulated heterospecific behavioural traits. *Anim. Behav.* 81, 879–883. (doi:10.1016/j.anbehav.2011.01.029)
- 525 17. Samplonius JM, Both C. 2017 Competitor phenology as a social cue in breeding site selection. J. Anim. Ecol. 86, 615–623. (doi:10.1111/1365-2656.12640)
- Jaakkonen T, Kivelä SM, Meier CM, Forsman JT. 2015 The use and relative importance of intraspecific and interspecific social information in a bird community. *Behav. Ecol.* 26, 55–64. (doi:10.1093/beheco/aru144)
- Samplonius JM, Kromhout Van Der Meer IM, Both C. 2017 Nest site preference
 depends on the relative density of conspecifics and heterospecifics in wild birds. *Front. Zool.* 14. (doi:10.1186/s12983-017-0246-5)
- Doligez B, Pärt T, Danchin E, Clobert J, Gustafsson L. 2004 Availability and use of
 public information and conspecific density for settlement decisions in the collared
 flycatcher. J. Anim. Ecol. 73, 75–87. (doi:10.1111/j.1365-2656.2004.00782.x)
- Morinay J, Forsman JT, Kivelä SM, Gustafsson L, Doligez B. 2018 Heterospecific nest site copying behavior in a wild bird: assessing the influence of genetics and past experience on a joint breeding phenotype. *Front. Ecol. Evol.* 5, 167. (doi:10.3389/fevo.2017.00167)
- Forsman JT, Hjernquist MB, Taipale J, Gustafsson L. 2008 Competitor density cues for
 habitat quality facilitating habitat selection and investment decisions. *Behav. Ecol.* 19,
 539–545. (doi:10.1093/beheco/arn005)
- Seppänen J-T, Forsman JT, Mönkkönen M, Krams I, Salmi T. 2011 New behavioural trait adopted or rejected by observing heterospecific tutor fitness. *Proc. R. Soc. B Biol. Sci.* 278, 1736–1741. (doi:10.1098/rspb.2010.1610)
- 546 24. Gustafsson L. 1988 Interspecific and intraspecific competition for nest holes in a 547 population of the collared flycatcher Ficedula albicollis. *Ibis (Lond. 1859)*. **130**, 11–16.

- 548 (doi:10.1111/j.1474-919X.1988.tb00951.x)
- 549 25. Král M, Bícík V. 1992 Nest defence by the collared flycatcher (*Ficedula albicollis*)
 550 against the great tit (Parus major). *Folia zooogica* 41, 263–269.
- 551 26. R Core Team. 2016 R: a language and environment for statistical computing.
- 552 27. Bates D, Maechler M, Bolker BM, Walker S. 2015 Fitting linear mixed-effects models
 553 using lme4. J. Stat. Softw. 67, 1–48. (doi:10.18637/jss.v067.i01.)
- Mundry R, Nunn CL. 2009 Stepwise model fitting and statistical inference: turning noise into signal pollution. *Am. Nat.* 173, 119–123. (doi:10.1086/593303)
- Forstmeier W, Wagenmakers E, Parker TH. 2017 Detecting and avoiding likely falsepositive findings a practical guide. *Biol. Rev.* 92, 1941–1968.
 (doi:10.1111/brv.12315)
- 30. Barton K. 2016 MuMIn: Multi-Model Inference. R package version 1.15.6.
- 31. Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J-C, Müller M. 2011
 pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* 12, 77. (doi:10.1186/1471-2105-12-77)
- 32. Michelena P, Sibbald AM, Erhard HW, McLeod JE. 2009 Effects of group size and personality on social foraging: the distribution of sheep across patches. *Behav. Ecol.*20, 145–152. (doi:10.1093/beheco/arn126)
- 566 33. Kurvers RHJM, Van Oers K, Nolet BA, Jonker RM, Van Wieren SE, Prins HHT,
 567 Ydenberg RC. 2010 Personality predicts the use of social information. *Ecol. Lett.* 13,
 568 829–837. (doi:10.1111/j.1461-0248.2010.01473.x)
- 569 34. Kurvers RHJM, Prins HHT, van Wieren SE, van Oers K, Nolet BA, Ydenberg RC.
 570 2010 The effect of personality on social foraging: shy barnacle geese scrounge more.
 571 *Proc. R. Soc. B Biol. Sci.* 277, 601–608. (doi:10.1098/rspb.2009.1474)
- 572 35. Kurvers RHJM, Adamczyk VMAP, van Wieren SE, Prins HHT. 2011 The effect of boldness on decision-making in barnacle geese is group-size-dependent. *Proc. R. Soc.*574 *B Biol. Sci.* 278, 2018–2024. (doi:10.1098/rspb.2010.2266)
- S75 36. Carter AJ, Marshall HH, Heinsohn R, Cowlishaw G. 2013 Personality predicts decision making only when information is unreliable. *Anim. Behav.* 86, 633–639. (doi:10.1016/J.ANBEHAV.2013.07.009)
- Trompf L, Brown C. 2014 Personality affects learning and trade-offs between private and social information in guppies, *Poecilia reticulata*. *Anim. Behav.* 88, 99–106. (doi:10.1016/j.anbehav.2013.11.022)
- 38. Snijders L, Naguib M, van Oers K. 2017 Dominance rank and boldness predict social attraction in great tits. *Behav. Ecol.* 28, 398–406. (doi:10.1093/beheco/arw158)
- Jolles JW, Ostojić L, Clayton NS. 2013 Dominance, pair bonds and boldness determine
 social-foraging tactics in rooks, *Corvus frugilegus. Anim. Behav.* 85, 1261–1269.
 (doi:10.1016/j.anbehav.2013.03.013)
- Forsman JT, Seppänen J-T, Mönkkönen M. 2002 Positive fitness consequences of
 interspecific interaction with a potential competitor. *Proc. R. Soc. B* 269, 1619–1623.
 (doi:10.1098/rspb.2002.2065)
- 589 41. Briffa M. 2008 Decisions during fights in the house cricket, Acheta domesticus: mutual
 590 or self assessment of energy, weapons and size? *Anim. Behav.* 75, 1053–1062.
 591 (doi:10.1016/j.anbehav.2007.08.016)
- 42. Ward AJW, Thomas P, Hart PJB, Krause J. 2004 Correlates of boldness in three-spined sticklebacks (Gasterosteus aculeatus). *Behav. Ecol. Sociobiol.* 55, 561–568. (doi:10.1007/s00265-003-0751-8)
- 595 43. Dyer JRG, Croft DP, Morrell LJ, Krause J. 2008 Shoal composition determines
 596 foraging success in the guppy. *Behav. Ecol.* 20, 165–171. (doi:10.1093/beheco/arn129)
- 597 44. Croft DP, Krause J, Darden SK, Ramnarine IW, Faria JJ, James R. 2009 Behavioural

- trait assortment in a social network: patterns and implications. *Behav. Ecol. Sociobiol.* **63**, 1495–1503. (doi:10.1007/s00265-009-0802-x)
- Carter AJ, Marshall HH, Heinsohn R, Cowlishaw G. 2014 Personality predicts the propensity for social learning in a wild primate. *PeerJ* 2, e283. (doi:10.7717/peerj.283)
- 46. Harcourt JL, Biau S, Johnstone R, Manica A. 2010 Boldness and information use in three-spined sticklebacks. *Ethology* 116, 440–447. (doi:10.1111/j.1439-0310.2010.01757.x)
- 47. Qvarnström A. 1997 Experimentally increased badge size increases male competition and reduces male parental care in the collared flycatcher. *Proc. R. Soc. B Biol. Sci.* 264, 1225–1231. (doi:10.1098/rspb.1997.0169)
- 48. Pärt T. 1995 The importance of local familiarity and search costs for age- and sexbiased philopatry in the collared flycatcher. *Anim. Behav.* 49, 1029–1038.
 (doi:10.1006/anbe.1995.0132)
- 49. Forsman JT, Seppänen J-T, Nykänen IL. 2012 Observed heterospecific clutch size can affect offspring investment decisions. *Biol. Lett.* 8, 341–343. (doi:10.1098/rsbl.2011.0970)
- 50. Tolvanen J, Kivelä SM, Doligez B, Morinay J, Gustafsson L, Bijma P, Pakanen V,
 Forsman JT. In press. Quantitative genetics of the use of conspecific and heterospecific
 social cues for breeding site choice. *Submitted*
- Budaev SV, Zworykin DD. 1998 Difference in shoaling behaviour between ocellated 51. 617 (Symphodus ocellatus) and long-striped (S. tinca) wrasses and its relation with other 618 patterns. 619 behavioural Mar. Freshw. Behav. Physiol. 31. 115-121. (doi:10.1080/10236249809387067) 620
- 52. Marchetti C, Drent P. 2000 Individual differences in the use of social information in
 foraging by captive great tits. *Anim. Behav.* 60, 131–140.
 (doi:10.1006/anbe.2000.1443)
- 53. Webster MM, Hart AJW, Ward PJB. 2007 Boldness is influenced by social context in threespine sticklebacks (Gasterosteus aculeatus). *Behaviour* 144, 351–371. (doi:10.1163/156853907780425721)
- 54. Nomakuchi S, Park PJ, Bell MA. 2009 Correlation between exploration activity and use of social information in three-spined sticklebacks. *Behav. Ecol.* 20, 340–345. (doi:10.1093/beheco/arp001)
- 55. Sibbald AM, Erhard HW, McLeod JE, Hooper RJ. 2009 Individual personality and the
 spatial distribution of groups of grazing animals: An example with sheep. *Behav. Processes* 82, 319–326. (doi:10.1016/j.beproc.2009.07.011)
- 56. David M, Cézilly F, Giraldeau LA. 2011 Personality affects zebra finch feeding
 success in a producer-scrounger game. *Anim. Behav.* 82, 61–67.
 (doi:10.1016/j.anbehav.2011.03.025)
- Aplin LM, Farine DR, Mann RP, Sheldon BC. 2014 Individual-level personality
 influences social foraging and collective behaviour in wild birds. *Proc. R. Soc. B Biol. Sci.* 281, 20141016–20141016. (doi:10.1098/rspb.2014.1016)
- 58. Webster MM, Laland KN. 2015 Space-use and sociability are not related to publicinformation use in ninespine sticklebacks. *Behav. Ecol. Sociobiol.*(doi:10.1007/s00265-015-1901-5)
- 59. Smit JAH, Oers K Van. 2019 Personality types vary in their personal and social information use. *Anim. Behav.*, 3–11. (doi:10.1016/j.anbehav.2019.02.002)
- 644 60. Carter AJ, Ticó MT, Cowlishaw G. 2016 Sequential phenotypic constraints on social 645 information use in wild baboons. *Elife* **5**, 1–21. (doi:10.7554/eLife.13125)
- 646 61. Aplin LM, Farine DR, Morand-Ferron J, Sheldon BC. 2012 Social networks predict
 647 patch discovery in a wild population of songbirds. *Proc. R. Soc. B Biol. Sci.* 279, 4199–

- 4205. (doi:10.1098/rspb.2012.1591) Evans JC, Jones TB, Morand-ferron J. 2018 Dominance and the initiation of group feeding events: the modifying effect of sociality. *Behav. Ecol.* **29**, 448–458. (doi:10.1093/beheco/arx194) 62.



Figure 1. Probability for flycatchers to copy tit apparent preference depending on (a)
male aggressiveness and age, (b) male boldness and age, and (c) female boldness and age
(yearlings: light grey; older: dark grey). Data points show actual choices (copy = 1 / reject =
0). The predicted means (lines) and corresponding 95%CI (shaded areas) were derived from
the final model for an averaged value of tit clutch / brood size.

Table 1. Summary of the results of studies investigating relations between personality traits and social information use: for each personality trait, number of studies that found a positive (+), negative (-), or non-significant (NS) relation. Full details on each study and measured traits are given in Table S1. Note that neophobia and exploration were often referred to as 'boldness' in articles, but we follow here the definitions from [8] and refer to boldness as the reaction in a risky situation (presence of potential predators, including humans).

Definitions	Nb. and signs of the links with social information use			References	
	-	0	+		
Reaction in a known environment	0	1	1	[49]	
Agonistic reaction towards others	0	0	0		
Reaction in a risky situation	2	5	1	[34, 40, 41, 43, 44]	
Reaction in a novel environment	3	4	2	[36, 50-57]	
Reaction towards a novel object	2	2	7	[30-35, 37, 43, 57,58]	
Non-agonistic reaction towards others	1	1	2	[35, 56, 59, 60]	

665

[1] Budaev and Zworykin [51]; [2] Ward et al. [42]; [3] Dyer et al. [43]; [4] Carter et al. [36];
[5] Carter et al. [45]; [6] Harcourt et al. [46]; [7] Marchetti and Drent [52]; [8] Webster et al.
[53]; [9] Nomakuchi et al. [54], [10] Sibbald et al. [55]; [11] David et al. [56]; [12] Aplin et al. [57]; [13] Webster and Laland [58]; [14] Snijders et al. [38]; [15] Smit and van Oers [59];
[16] Michelena et al. [32]; [17] Kurvers et al. [34]; [18] Kurvers et al. [33]; [19] Kurvers et al.
[35]; [20] Jolles et al. [39]; [21] Trompf and Brown [37]; [22] Carter et al. [60]; [23] Aplin et

672 al. [61]; [24] Evans et al. [62].

673	Table 2. Influence of male aggressiveness and boldness scores, female neophobia score
674	and tit clutch / brood 'c/b' size on the probability of copying tit apparent preference of
675	nest box artificial features. Aggressiveness has been log-transformed and all continuous
676	traits have been standardized prior analysis. Age estimates are given for yearling individuals
677	(older individuals being the reference). P-values below the risk α of 0.05 are highlighted in bold.

	$Estimate \pm SE$			z-value	p-value			
Final model with male aggressiveness score, N=224								
Intercept	-0.37	±	0.19	-1.95	0.05			
$Age_{\hat{c}}$	0.46	±	0.33	1.36	0.17			
Aggressiveness	-0.83	±	0.30	-2.79	0.01			
Tit clutch / brood (c/b) size	0.68	±	0.15	4.48	< 10 ⁻⁵			
$\operatorname{Aggressiveness}_{\widehat{\operatorname{C}}} : \operatorname{Age}_{\widehat{\operatorname{C}}}$	0.93	±	0.36	2.59	0.01			
Final model with male boldness score, N=142								
Intercept	0.05	±	0.22	0.22	0.83			
Age∂	0.35	±	0.41	0.85	0.39			
Boldness♂	-0.53	±	0.27	-1.95	0.05			
Tit c/b size	0.75	±	0.20	3.77	< 10 ⁻³			
$\mathbf{Boldness}_{\hat{c}}: \mathbf{Age}_{\hat{c}}$	0.98	±	0.45	2.21	0.03			
Final model with female bold	Final model with female boldness score. $N=173$							
Intercept	-0.20	±	0.20	-1.01	0.31			
$\mathrm{Age}_{\mathbb{Q}}$	0.54	±	0.35	1.54	0.12			
Boldnessç	-0.25	±	0.21	-1.23	0.22			
Tit c/b size	0.64	±	0.17	3.78	< 10 ⁻³			
$\mathbf{Boldness}_{\mathbb{Q}}: \mathbf{Age}_{\mathbb{Q}}$	0.71	±	0.35	2.03	0.04			
Final model with female neophobia score, N=173								
Intercept	0.00	±	0.16	-0.01	0.99			
Neophobia _♀	-0.02	±	0.17	-0.10	0.92			
Tit c/b size	0.68	±	0.17	3.92	< 10 ⁻⁴			
Neophobia _♀ : Tit c/b size	-0.47	±	0.20	-2.33	0.02			