

Seeds of Life in Space (SOLIS). IX. Chemical Segregation of SO 2 and SO toward the Low-mass Protostellar Shocked Region of L1157

Siyi Feng, Claudio Codella, C. Ceccarelli, Paola Caselli, Ana Lopez-Sepulcre, Roberto Neri, Francesco Fontani, L Podio, B. Lefloch, H B Liu, et al.

▶ **To cite this version:**

Siyi Feng, Claudio Codella, C. Ceccarelli, Paola Caselli, Ana Lopez-Sepulcre, et al.. Seeds of Life in Space (SOLIS). IX. Chemical Segregation of SO 2 and SO toward the Low-mass Protostellar Shocked Region of L1157. The Astrophysical Journal, 2020, 896 (1), pp.37. 10.3847/1538-4357/ab8813 . hal-02999347

HAL Id: hal-02999347
https://hal.science/hal-02999347

Submitted on 13 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Draft version May 15, 2020
Typeset using LaTeX **twocolumn** style in AASTeX62

arXiv:2005.04629v2 [astro-ph.GA] 14 May 2020

Seeds of Life in Space (*SOLIS*). IX. Chemical segregation of SO_2 and SO toward the low-mass protostellar shocked region of L1157*

S. Feng,[1,2,3] C. Codella,[4,5] C. Ceccarelli,[5] P. Caselli,[3] A. Lopez-Sepulcre,[5,6] R. Neri,[6] F. Fontani,[3,4] L. Podio,[4] B. Lefloch,[5] H. B. Liu,[7] R. Bachiller,[8] and S. Viti[9]

[1]*National Astronomical Observatory of China, Datun Road 20, Chaoyang, Beijing, 100012, P. R. China*
[2]*CAS Key Laboratory of FAST, NAOC, Chinese Academy of Sciences, P. R. China*
[3]*Max-Planck-Institut für Extraterrestrische Physik, Gießenbachstraße 1, D-85748, Garching bei München, Germany*
[4]*INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125, Florence, Italy*
[5]*Univ. Grenoble Alpes, CNRS, IPAG, F-38000 Grenoble, France*
[6]*Institut de Radioastronomie Millimétrique, 300 rue de la Piscine, Domaine Universitaire de Grenoble, F-38406 Saint-Martin d'Hères, France*
[7]*Academia Sinica Institute of Astronomy and Astrophysics, No.1, Sec. 4, Roosevelt Road, Taipei 10617, Taiwan, Republic of China*
[8]*Observatorio Astronomico Nacional (OAN, IGN), Calle Alfonso XII, 3, E-28014 Madrid, Spain*
[9] *Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK*

(Received; Revised; Accepted 2020 Apr 7)

Submitted to ApJ

ABSTRACT

We present observations of SO and SO_2 lines toward the shocked regions along the L1157 chemically rich outflow, taken in the context of the Seeds Of Life In Space IRAM-NOrthern Extended Millimeter Array Large Program, and supported by data from Submillimeter Array and IRAM-30 m telescope at 1.1–3.6 mm wavelengths. We simultaneously analyze, for the first time, all of the brightest shocks in the blueshifted lobe, namely, B0, B1, and B2. We found the following. (1) SO and SO_2 may trace different gas, given that the large(-scale) velocity gradient analysis indicates for SO_2 a volume density (10^5–10^6 cm^{-3}) denser than that of the gas emitting in SO by a factor up to an order of magnitude. (2) Investigating the 0.1 pc scale field of view, we note a tentative gradient along the path of the precessing jet. More specifically, $\chi(SO/SO_2)$ decreases from the B0-B1 shocks to the older B2. (3) At a linear resolution of 500–1400 au, a tentative spatial displacement between the two emitting molecules is detected, with the SO peak closer (with respect to SO_2) to the position where the recent jet is impinging on the B1 cavity wall. Our astrochemical modeling shows that the SO and SO_2 abundances evolve on timescales less than about 1000 years. Furthermore, the modeling requires high abundances (2×10^{-6}) of both H_2S/H and S/H injected in the gas phase due to the shock occurrence, so pre-frozen OCS only is not enough to reproduce our new observations.

Keywords: Stars: formation; Stars: low-mass; ISM: lines and bands; Submillimeter: ISM

1. INTRODUCTION

In the earliest stage of star formation, supersonic jets generated by protostars impact on the natal dense parent cloud and create shocks (e.g., Frank et al. 2014). Shocks compress and heat the surrounding gas, driving complex chemistry in a short timescale, including endothermic chemical reactions, ice mantle sublimation, and sputtering. These chemical processes are infeasible toward the pre-shock gas (e.g. Viti et al. 2011). As a result, gas-phase abundances of species such as sulfur (S)-bearing molecules are enhanced by several orders of magnitude compared with the prestellar cores and protostellar envelopes (e.g. Tieftrunk et al. 1994; Kaufman & Neufeld 1996; Bachiller et al. 2001; Jiménez-Serra et al. 2005; Jørgensen et al. 2007).

1.1. *S-bearing chemistry*

S-bearing molecules are highly reactive in the gas phase and are very sensitive to the thermal and kinetic properties of the gas. It has long been proposed that the relative abundance ratio between H_2S, SO_2, and SO can be used to date

Corresponding author: Siyi Feng
siyi.s.feng@gmail.com

*

the evolutionary stages of star-formation (i.e., constraining the age of the outflow and associated shocks; Charnley et al. 1997; Hatchell et al. 1998; Buckle & Fuller 2003; Wakelam et al. 2004a). For example, in agreement with theoretic predictions, observations revealed that the SO_2/SO ratio is enhanced in young ($< 10^3$ yr) shocked region compared with the cloud, due to gaseous endothermic reactions; while this ratio decreases for the conversion of SO_2 to atomic sulfur in the old shocks ($> 3 \times 10^4$ yr, e.g., Pineau des Forets et al. 1993; Hatchell et al. 1998; Wakelam et al. 2004a; Feng et al. 2015). However, the feasibility of taking the relative abundances of S-bearing species as a star formation dating tool depends on the physical condition of the source and a proper assumption of the S grain reservoir.

Moreover, understanding of the S-bearing chemistry is challenged by a collection of mysteries, the major one being the "S-depletion problem" (e.g, Tieftrunk et al. 1994); i.e., although S is known to be significantly depleted in molecular dark clouds, its depleted form (the main reservoir on grain mantles) is far from being established. Several species have been proposed to be the main carriers of S on dust grains, such as H_2S (e.g., Li et al. 2015; Holdship et al. 2016) and OCS (e.g., Hatchell et al. 1998; Wakelam et al. 2004a; Liu et al. 2012; Podio et al. 2014; Minh et al. 2016; Holdship et al. 2019). Historically, H_2S has never been definitively detected on ices, which calls for further and more comprehensive S-chemistry study using telescopes at higher sensitivity. Recent observations and models have shown that, at the very least, H_2S is not the only S-bearing species on grains (e.g., Wakelam et al. 2004b; Codella et al. 2005). By far, OCS is detected (e.g., Geballe et al. 1985; Palumbo et al. 1995; Podio et al. 2014; Boogert et al. 2015), and SO_2 (e.g., Boogert et al. 1997) is tentatively detected in interstellar ices. Therefore, the main reservoir of S on dust is still unclear.

1.2. *The blueshifted outflow lobe of L1157*

Regions containing successive young ($\lesssim 10^3$ yr) shocks driven by mass losses from protostars are ideal laboratories for studying time-dependent S-chemistry, as the short timescales of the shocks provide the possibility to either directly observe the material coming off grains, or to trace the chemistry back to the original content of interstellar ices.

This well applies to the target of the present study, the shocks associated with the L1157 outflow region. The Class 0 protostar L1157-mm ($\simeq 3\ L_\odot$, Tobin et al. 2010), located at d ~352 pc (Zucker et al. 2019), drives an episodic and precessing jet (e.g. Gueth et al. 1996, 1998; Podio et al. 2016), creating a bipolar outflow (e.g., Bachiller et al. 2001; Nisini et al. 2010). The impacts between the jet and the high-density (10^4–10^5 cm^{-3}, Lefloch et al. 2012; Benedettini et al. 2013; Gómez-Ruiz et al. 2015) cavity walls triggered several shocks in the last $\sim 2 \times 10^3$ yr (e.g., Lefloch et al. 2010).

The targets of the present work are the B0, B1, and B2 shocked regions, which are the brightest shocks associated with the southern blueshifted outflow lobe, located within 0.1 pc from the protostar in the plane of the sky. Specifically, the B1 region has been studied in detail by using either single-dish telescopes (e.g., *Herschel-CHESS*, Ceccarelli et al. 2010; *30 m-ASAI*, Lefloch et al. 2018) or interferometers (SMA, Very Large Array, NOEMA, Tafalla & Bachiller 1995; Codella et al. 2009; Lefloch et al. 2012; Benedettini et al. 2013), revealing a clumpy bow-like structure associated with several chemical processes, such as the sputtering of dust mantles, the disruption of the dust refractory cores, and warm gas-chemistry reactions. The abundance of molecular species, from diatomic–such as CO, CS–to molecular ions–such as $HOCO^+$, SO^+, and HCS^+ Podio et al. (2014)–and more complex molecules such as CH_3OH, CH_3CHO, and NH_2CHO (e.g., Bachiller et al. 2001; Benedettini et al. 2007; Arce et al. 2008; Codella et al. 2010, 2015, 2017; Busquet et al. 2014; Fontani et al. 2014a; Lefloch et al. 2017, 2018) are consequently enhanced. Specifically, S-bearing species (e.g., CS, SO, SO_2, H_2S, OCS) have been detected toward this region (see, e.g., Bachiller et al. 2001; Benedettini et al. 2013; Podio et al. 2014; Holdship et al. 2016).

Kinematically, B2 is older than B1 (e.g., Bachiller et al. 2001; Podio et al. 2016); B1 mainly consists of three shock layers produced by episodes of the precessing jet, with the oldest layer located the farthest from the protostar, i.e. in the southernmost edge of B1 (Codella et al. 2015). Observing the jet radial velocity in the inner knots, the precessing model (Podio et al. 2016) indicates that[1] B0 is ~1340 yr old, B2 is ~2530 yr old, and the kinematic age of B1 knots ranges from 1550 yr at the north edge to 1760 yr at the southern edge. Therefore, the entire B0-B1-B2 region provides us one of the best laboratories to study the time-dependent S-chemistry.

In this paper, we present the data in Section 2, which are obtained from the "Seeds Of Life In Space"(*SOLIS*[2]*; Ceccarelli et al. 2017)* large program and supplementary Submillimeter Array (SMA) observations. Our results are presented in Section 3 and analyzed in Section 4. We study the map of relative abundance ratio between SO and SO_2 in Section 5.1, and discuss the possible origin of SO and SO_2 by using our chemical model to fit the observation measurement in Section 5.2. We conclude in Section 6.

2. OBSERVATIONS

[1] Podio et al. (2016) estimated the kinematic ages of each knot by using d ~250 pc. The kinematic ages of these knots in this work are updated by using d ~352 pc (Zucker et al. 2019).

[2] The "Seeds Of Life In Space" large programme is aiming at investigating molecular formation and processes during the early stages of the star formation process, by using IRAM NOrthern Extended Millimeter Array (NOEMA).

Using the NOrthern Extended Millimeter Array (NOEMA) and the SMA, we carried out a line-imaging survey at 1.1–3.6 mm toward our targets. Given that the antennae of NOEMA and SMA have different diameters, and that the molecular lines we observed are at different wavelengths, the primary beam attenuation is different. Therefore, the B0, B1, and B2 shocked regions have been imaged by using one point or two mosaic points with NOEMA or SMA at different wavelengths.

2.1. *NOEMA*

Using NOEMA, we observed B0-B1 at 3.6 mm, 3.1 mm, and 1.5 mm in its A (7 antennae), C (6-8 antennae), and D (5-8 antennae) configurations from 2015 July to 2017 January. With the shortest projected baseline of 24 m, the observations are sensitive to structures up to 19″ (at 3.6 mm), 16″ (at 3.1 mm), and 8″ (at 1.5 mm), respectively. For all observations, a common phase center at $20^h39^m10^s.20$, $68°01'10''.50$ (J2000) and the systemic velocity $V_{sys} \sim 2.7\,km\,s^{-1}$ was used. The precipitable water vapor (PWV) varied between 1 and 40 mm during the observations. Standard interferometric calibrations were performed during the observations. The bandpass, gain (phase/amplitude), and flux calibrators used during the observations are recorded in Table A1. The uncertainty of absolute flux scale is estimated to be correct to within ~ 5%.

With the wide-band receiver (WIDEX), three spectral setups cover the frequency ranges of 80.790–84.407, 95.794-99.499, and 203.962-207.632 GHz, at a frequency resolution of 1.95 MHz. We detected one SO_2 line and three SO (^{34}SO) lines (with $E_u/k_B < 40$ K). From our *ASAI*-30 m single-point observations at a frequency resolution of 0.195 MHz (Lefloch et al. 2018), these lines have full-width half-maximum (FWHM) linewidth of $> 8\,km\,s^{-1}$ toward the B1 shocked region (Mendoza et al. 2014), so the velocity resolution from WIDEX observations ($\sim 7\,km\,s^{-1}$ at 83.688 GHz, see Table 1 and Figure 1) is sufficient for our chemical analysis. Moreover, we use the narrow-band correlator and set four windows in each spectral setup (80 MHz backends with a spectral resolution of 156 kHz) to target on several complex organics lines (see the spectral setups in Ceccarelli et al. 2017; Codella et al. 2017, for details). Other lines detected toward this region will be presented in the follow-up papers (e.g., CS/CCS/OCS/NS/SiS/H_2S/H_2CS, cyanopolyynes; S. Feng et al. in prep.).

Furthermore, aiming at a robust analysis on the possible chemical differentiations between SO_2 and SO toward different time-dependent shock layers of B1, we also include one ^{34}SO and four SO_2 ($^{34}SO_2$) lines from our previous NOEMA-WIDEX observations (obsID: W068, Fontani et al. 2014a; obsID: X058, Podio et al. 2017; see observation details therein). The spectroscopic parameters of all these lines are taken from the Cologne Database for Molecular Spectroscopy (CDMS, Müller et al. 2002, 2005) and listed in Table 1.

Calibration and imaging were performed by using the standard procedures with the CLIC and MAPPING softwares of the GILDAS[3] package. After testing different weighting algorithms, we used the natural weighting to achieve a better signal-to-noise ratio (S/N). However, our NOEMA data recover only 40–60% of the emission when compared with our *ASAI*-30 m pointing observations, indicating that these SO_2 and SO lines should have extended emissions toward our target.

Averaging the line-free channels of the WIDEX band, we did not detect significant ($> 5\sigma$ rms) continuum emission at 3.6 mm with a synthesized beam of $5.08'' \times 4.69''$ (σ=0.016 mJy beam^{-1}), at 3.1 mm with a synthesized beam of $3.12'' \times 2.30''$ (σ=0.012 mJy beam^{-1}), or 1.5 mm with a synthesized beam of $1.20'' \times 0.85''$ (σ=0.036 mJy beam^{-1}).

From our NOEMA observations, the detected SO isotopologue lines have low-J levels (E_u/k_B=9–18 K), while the SO_2 isotopologue lines have mid-J or high-J levels (E_u/k_B=16–102 K). They may selectively trace gas with different temperatures and therefore show different spatial distributions. With the aim of investigating any chemical difference from the spatial distribution maps of both species, it is necessary to observe more SO_2 and SO lines, to cover the same range of upper energy level (E_u/k_B).

2.2. *SMA*

To minimize the excitation effect on the spatial distribution differentiation between SO and SO_2, we carried out four-track observations toward our source with the SMA at 230/255 GHz (1.14-1.40 mm), by using the compact configuration. From 2017 June to July, we observed B1 and B2 with a two-point mosaic per track. The phase centers of B1 and B2 are at $20^h39^m09^s.500$, $68°01'10''.500$ and $20^h39^m11^s.358$, $68°00'51''.00$ (J2000), respectively. For all of the observations, the phase and amplitude calibrations were performed via frequent switch (every 20 minutes) on quasars 1849+670 and 1927+739.

The baselines range from 16 m to 76 m with 7 or 8 antennae at different dates. B1 and B2 share the bandpass (3C273 or 3C454.3) and flux (Callisto and MWC349a) calibrators. The zenith opacities, measured with water vapor monitors mounted on the James Clerk Maxwell Telescope (JCMT), were satisfactory during all tracks with τ(225 GHz) ~ 0.05–0.3. The system temperature was 100–200 K at 230 GHz and 200–400 K at 255 GHz. Further technical descriptions of the SMA and its calibration schemes can be found in Ho et al. (2004).

[3] http://www.iram.fr/IRAMFR/GILDAS

We employed the dual-RX, single polarization and double-sideband observing mode. Our frequency coverages (214.935–221.030, 229.004–235.015, 240.738–246.828, 254.804–260.890 GHz) were sampled using the SWARM correlator with a 0.140 MHz frequency resolution (~0.182 km s^{-1} at 230.538 GHz). We tuned 230.0 GHz and 255.8 GHz to be at the center of the spectral chunk 1 in the upper sideband of the receivers Rx230 and Rx240, respectively. To increase the S/N ratio per channel, we smooth the frequency resolution to 4.469 MHz (a velocity resolution of 5.812 km s^{-1} at 230.538 GHz).

The flagging and calibration was done with the MIR package (Scoville et al. 1993; Qi 2003)[4]. The mosaic imaging was conducted with MIRIAD package by using the natural weighting (Sault et al. 1995).

We detected four SO lines in the entire frequency coverage (E_u/k_B=35–56 K), while no SO_2 line is detected with $> 3\sigma$ emission (σ=42.7 mJy beam^{-1} at an angular resolution of ~ 3.4″). Their spectroscopic parameters taken from CDMS are listed in Table 1 as well. In addition, we also detected lines from SiO, HNCO, CO, CS, H_2CO in these observations.

Averaging the line-free channels, the sensitivity of continuum emission at 1.1–1.4 mm achieves σ=0.17 mJy beam^{-1} with a synthesized beam of 3.37″ ×3.13″. At such sensitivity, we have marginally detected 3–4 σ of continuum emission toward B1-B2 region (see H. B. Liu in prep. for a detailed continuum study).

2.3. *IRAM-30 m Telescope*

To compensate the missing flux, we use IRAM-30 m telescope and performed the observations in the on-the-fly (OTF) mode from 2017 May to November. We mapped a 2.5′ × 2.5′ area (at least twice that of the primary beam of the interferometers for successful combination, according to Pety et al. 2013, covering both B1 and B2), centered at $20^h39^m09^s.50$, 68°01′10″.5 (J2000). By superpositioning different spectral tunings, the broad bandpass of EMIR (16 GHz bandwidth simultaneously for each spectral tuning) covers the frequency range of 79.302–87.084, 91.821–102.764, 106.902–114.683, 130.703–138.485, 161.102–169.882, 203.925–211.703, 213.422–227.385, 229.105–236.882, 239.420–247.204, and 255.103–262.880 GHz, with a frequency resolution of 0.195 MHz (by using the FTS200 backend). The FWHM beam of the 30 m telescope at 83.688 GHz is ~ 30.86″.

To combine the emissions of SO_2 and SO isotopologue lines obtained from NOEMA or SMA at high spatial resolution with their large-scale emission data obtained from IRAM-30 m, we aligned the phase center and smoothed the velocity resolution of lines obtained from IRAM-30 m to those from NOEMA or SMA. Afterwards, we use the task "UV-SHORT" in the MAPPING software for NOEMA-30 m combination; and use the MIRIAD package for SMA-30 m combination. The synthesized beam and the 1σ rms value of the data cube of each line, after NOEMA-30 m or SMA-30 m combination, are listed in Table 1.

3. MOLECULAR SPATIAL DISTRIBUTION

For the first time, the interferometric data of L1157 B0-B1-B2 are complemented with the IRAM-30 m mapping, providing us images with full spatial scale coverage, as well as high angular resolutions. Moreover, the wide frequency coverage allows us to study five SO_2 lines (including one $^{34}SO_2$ line) and eight SO lines (including two ^{34}SO line), with E_u/k_B in the range of 16–102 K. At a linear resolution of < 1000 au, we will be able to search for chemical differentiations from the systematic differentiations in the spatial distribution of both species.

All lines show a significant blueshifted emission toward B0-B1-B2, and the S/N > 3 part of each line is in the velocity range of −10 to +5 km s^{-1} (Figure 1 and A1). Specifically, the peaks of all SO_2 lines are close to the V_{sys}, when observing at a velocity resolution of 3.6–7.0 km s^{-1}. Observed at higher velocity resolutions (2.8–5.9 km s^{-1}), some SO lines seem to peak at ~ −2 km s^{-1} toward positions such as B0e, B1f, and B1a. However, such a velocity offset (~ 5 km s^{-1}) only corresponds to 1–2 channel(s), and it smears out when these SO lines are binned to a coarser velocity resolution (7.0 km s^{-1}). Therefore, whether these SO and SO_2 lines trace different gas calls for further observations at a higher-velocity resolution.

We imaged the spatial distribution of each line by integrating their emissions over the velocity range[5] of −10 to +5 km s^{-1}. Specifically, Figure 2 show the spatial distributions of SO_2 ($8_{1,7}-8_{0,8}$) and four SO lines, i.e., 6_5-5_4, 5_5-4_4, 7_6-6_5, and 6_6-5_5, which are sensitive to a comparably larger field than the rest of the lines (see their primary beams in Table 1). Overlaid with CO (1–0) emission contours (Gueth et al. 1996), we found that all of these SO and SO_2 line emissions trace the extended structure from the eastern cavity wall B0 to two cavities associated with B1 and B2 (denoted as C2 and C1, respectively in Gueth et al. 1998). These lines show the same shock knots in B0 and B1 as those in SiO (5–4) from the same SMA-30 m observations[6] (Figure 2) as well

[4] The MIR package was originally developed for the Owens Valley Radio Observatory, and is now adapted for the SMA, http://cfa-www.harvard.edu/~cqi/mircook.html.

[5] We compared the integrated intensity of each line over the velocity range of −10 to +5 km s^{-1} (S/N > 3) with −20 to +15 km s^{-1} (line emission down to zero), and found the difference between these integrations is < 5%, which is less than the systematic uncertainty from observations and data reduction.

[6] The combination of two-point mosaic of SMA observations and IRAM-30 m observations recover extended emission of SiO (5–4), which that was

Table 1. Target lines in this work with their spectroscopic and observation parameters

Mol.	Freq.[a] (GHz)	Transition[a]	$S\mu^2$ (D^2)	E_u/k_B[a] (K)	ΔV (km s^{-1})	$\Delta\theta$ (P.A.) [b] ($''\times''$, $^\circ$)	rms[b] (mJy beam^{-1} ch^{-1})	Primary beam ($''$)	Interferometric Observations
SO_2	83.688	$8_{1,7}-8_{0,8}$	17.01	37	7.01	$5.30''\times4.86''$ (-94°)	0.7	61	*SOLIS*-NOEMA
SO_2	134.004	$8_{2,6}-8_{1,7}$	15.18	43	4.37	$2.64''\times2.43''$ (-89°)	1.1	38	NOEMA[c]
SO_2	135.696	$5_{1,5}-4_{0,4}$	8.35	16	4.32	$2.65''\times2.51''$ (75°)	1.3	38	NOEMA[c]
SO_2	163.606	$14_{1,13}-14_{0,14}$	17.10	102	3.58	$2.01''\times1.87''$ (43°)	1.4	31	NOEMA[d]
$^{34}SO_2$	162.776	$7_{1,7}-6_{0,6}$	11.86	26	3.60	$1.98''\times1.86''$ (47°)	0.9	31	NOEMA[d]
SO	99.230	3_2-2_1	6.91	9	5.90	$2.95''\times2.20''$ (25°)	0.4	52	*SOLIS*-NOEMA
SO	206.176	4_5-3_4	8.91	39	2.84	$1.15''\times0.80''$ (-171°)	1.3	25	*SOLIS*-NOEMA
SO	215.221	5_5-4_4	11.31	44	6.23	$3.91''\times3.66''$ (-2°)	21.9	55	SMA
SO	219.949	6_5-5_4	14.01	35	6.09	$3.82''\times3.58''$ (-3°)	32.1	53	SMA
SO	258.256	6_6-5_5	13.74	56	5.19	$3.23''\times3.04''$ (-4°)	22.2	45	SMA
SO	261.844	7_6-6_5	16.38	48	5.12	$3.18''\times3.00''$ (-4°)	73.1	45	SMA
^{34}SO	97.715	3_2-2_1	6.92	9	5.99	$2.98''\times2.33''$ (25°)	0.4	52	*SOLIS*-NOEMA
^{34}SO	135.776	4_3-3_2	9.28	16	4.31	$2.64''\times2.50''$ (75°)	1.2	38	NOEMA[c]
SiO	217.105	$5-4$	48.0	32	6.17	$3.99''\times3.75''$ (-3°)	22.7	54	SMA

Note. *a*. Spectroscopic parameters are taken from the CDMS.
b. Data from NOEMA-30 m or SMA-30 m combination, without angular resolution smoothing, measured with the referred beam size ("beam") and channel width ("ch").
c. Unpublished data was taken from NOEMA observations W068 reported by Fontani et al. (2014a).
d. Unpublished data was taken from NOEMA observations X058 reported by Podio et al. (2017).

as SiO (2–1) observed by NOEMA-30 m (Gueth et al. 1998), which is consistent with the prediction of a precessing jet model (Podio et al. 2016).

Aside from the above five lines, the rest SO and SO_2 isotopologue lines in this work are only sensitive to the B0-B1 region. Nevertheless, they offer images observed at higher angular resolutions. At a spatial resolution of 500–1400 au, six clumpy substructures appear toward the B0-B1 region in Figure 3. These substructures are already identified from previous observations (e.g., by using line emissions from CH_3CN, HCN, CH_3OH, SiO, CS, H_2CO, HCO^+, and HNCO in Benedettini et al. 2007; Codella et al. 2009; Gómez-Ruiz et al. 2013; Burkhardt et al. 2016). In this work, we use the consistent names of B0e, B1a, B1b, B1c, and B1f with those defined by Benedettini et al. (2007); Codella et al. (2009); Burkhardt et al. (2016). Moreover, we denote a position B1i, which was noted previously but not labeled as a clumpy substructure from emission of other species. For any particular substructure traced by more than one molecule, the difference in the measurement of its absolute coordinate by different molecular tracers is about $1''$–$2''$. Based on Figure 3, the absolute coordinates of these clumpy substructures are listed in Table 2.

Morphologically, B0e outlines the eastern wall of the northernmost blueshifted bow-shock cavity (e.g., named as "C2" in Gueth et al. 1998), showing higher abundance of species such as SiO and HCN than the western wall and the connecting ridge B1f (Gómez-Ruiz et al. 2013) . B0e and B1a are suggested as knots where episodic ejection impacting against the cavity wall (Podio et al. 2016), and high-velocity SiO and CS emissions (Benedettini et al. 2007; Gómez-Ruiz et al. 2013) are detected toward them. A U-shape structure connecting B1a-B1c-B1b pins down the shock front, and the apex B1i is coincident with the "finger" structure detected on the maps of CS and low-velocity SiO, which is suggested as the magnetic precursor of the shock (Gueth et al. 1998).

According to the spatial distributions of detected complex organic molecules (COMs), several chemical layers are defined in B1. One chemical layer is composed of B0e-B1a-B1f-B1b, where the detected COMs include HDCO (Fontani et al. 2014b), CH_3CHO (Codella et al. 2017), and CH_3OH (Codella et al. 2017), indicating the place where the current dust mantle and core are released by sputtering. To the southern shock front, B1c and B1i denote another chemical layer where NH_2CHO is the main COM emitter (Codella et al. 2017), and this layer contains the gas older than B1a-B1f-B1b (Gueth et al. 1998; Podio et al. 2016).

To compare the line emissions along the path of the processing jet, we also denote a position B2a, where SO and SO_2 lines show the emission local maximum within B2.

All of the SO and SO_2 lines show the strongest emission toward B1a. More specifically, ^{34}SO, SO_2, and $^{34}SO_2$ lines show the U-shape structure toward the wall and southern rim

missed toward B2 in previous SMA-only observations (Gómez-Ruiz et al. 2013).




Figure 1. Profiles of two identified SO and SO_2 lines with the same E_u/k_B and similar beam attenuation. The profiles of all identified SO and SO_2 lines are shown in Appendix Figure A1. Lines are averaged from a beam-sized region with the center toward each clumpy substructure in the plane of the sky (Table 2). All lines are extracted from images smoothed to the same pixel size and the same angular resolution (5.1″). The line profile of SO (6_5–5_4) shown in black or dashed blue has a velocity resolution of 6.09 km s^{-1} (original spectral resolution) or 7.01 km s^{-1} (the same as that of SO_2 $8_{1,7}$–$8_{0,8}$), respectively. In each panel, two red vertical lines at −10.0 km s^{-1} and +5.0 km s^{-1} indicate the velocity range for which we integrate the intensity; the gray dashed vertical line indicate the V_{sys} = 2.7 km s^{-1} of the cloud. The horizontal cyan line indicates the baseline (T_B=0 K).

of B1, linking B0e-B1a- B1c- B1b. In contrast, SO lines suggest an anticorrelated spatial distributions, having a bright emission toward the eastern B1a. This finding could either reflect a chemical segregation or be due to optical depth effects, and it will be investigated in the next sections.

4. ANALYSIS

To quantify the possible chemical differentiations between the SO and SO_2 molecules by excluding excitation effects, we need to constrain their column densities over the entire targeted region. Uncertainty in the molecular column density measurement is mainly from two parts, whether the optical depth of each line can or has to be corrected, and whether these lines can be assumed in the local thermal equilibrium (LTE) condition.

Table 2. Positions corresponding to the clumpy structures shown on the maps in Figure 3.

	R.A. (J2000)	Dec (J2000)
B0e	$20^h39^m10^s.365$	68°01′23.80″
B1a	$20^h39^m10^s.301$	68°01′13.51″
B1b	$20^h39^m08^s.734$	68°01′10.54″
B1c	$20^h39^m09^s.677$	68°01′06.45″
B1f	$20^h39^m09^s.444$	68°01′16.70″
B1i	$20^h39^m10^s.000$	68°01′01.11″
B2a	$20^h39^m12^s.322$	68°00′43.85″

4.1. *Optical depths*

In the shocked region, we cannot exclude the possibility that some low-J SO and SO_2 lines may be optically thick. We note that $^{34}SO\,(3_2-2_1)$ and $^{32}SO\,(3_2-2_1)$ are a pair of lines that have similar upper-level energies ($E_u/k_B \sim 9$ K) and Einstein coefficients ($A_{ij} \sim 10^{-6}$ s^{-1}). Therefore, their brightness temperature ratio, $T_{B,^{34}SO(3_2-2_1)}/T_{B,^{32}SO(3_2-2_1)}$, can be used to measure the optical depth τ of $^{32}SO\,(3_2-2_1)$ line (Myers et al. 1983)[7].

Figure 4 shows that such ratio is in the range of 0.1–0.15 toward the pixels where both lines show > 5σ emissions. Considering the 10% systematic uncertainty per pixel, $T_{B,^{32}SO(3_2-2_1)}/T_{B,^{34}SO(3_2-2_1)}$, toward pixels within B0-B1 can be treated as uniform. Therefore, if we assume that $^{34}SO(3_2-2_1)$ is optically thin, we can reasonably assume that $^{32}SO(3_2-2_1)$ has uniform optical depths toward all valid pixels over the entire B0-B1 region.

Quantitatively, using the standard isotopic ratio $\mathcal{R}_{^{32}S/^{34}S} \sim 22$ in the solar system (e.g., Wilson & Rood 1994; Chin et al. 1996; Lodders 2003), τ of the $^{32}SO\,(3_2-2_1)$ line in B0-B1 can be estimated. Solving numerically the Myers et al. (1983) approximation (see footnote 7), we present the range of τ on Figure 4 as well. In general, τ of the $^{32}SO\,(3_2-2_1)$ line is not negligible ($\leq$ 1.5 toward B0e and B1a, as well as $\leq$ 3 toward B1f, B1c, and B1i, and $\leq$ 4 toward the western edge), but it has small dynamic range in B0-B1 region. As the absolute value of τ is an approximation and depends on the assumption of $\mathcal{R}_{^{32}S/^{34}S}$, τ toward individual pixels is not im-

[7] Assuming that fractionation (isotopic exchange reaction) of $^{34}S \Leftrightarrow {}^{32}S$ is stable, according to Myers et al. (1983), we can estimate the optical depth at the line center of a given line $\tau_{\alpha,0}$ by measuring the ratio between the main beam brightness temperature of the main line $T_{mb,\,\alpha,0}$ and its isotopologue $T_{mb,\,\beta,0}$ as $\frac{1-\exp(-\tau_{\alpha,0}/\mathcal{R}_\alpha)}{1-\exp(-\tau_{\alpha,0})} \approx \frac{T_{mb,\,\beta,0}}{T_{mb,\,\alpha,0}}$. Here, $\mathcal{R}_\alpha$ is the intrinsic abundance of the main isotope (e.g., ^{32}S) compared to its rare isotope (e.g., ^{34}S) in the ISM. A numerical solver to this approximation indicates that $\tau_{\alpha,0}$ is appropriate to $\sqrt{\ln(\frac{T_{mb,\,\beta,0}}{T_{mb,\,\alpha,0}})}$ at each pixel.




Figure 2. Outline of the southern (blueshifted) outflow lobe from the low-mass protostar L1157-mm. Color maps show the intensity maps of several shock tracer lines integrated over the velocity range of −10 to +5 km s^{-1} , which are sensitive to the extended B0-B1-B2 region. All of the maps are obtained from the combination of NOEMA (or SMA) and IRAM-30 m, and line emissions with < 5σ rms are blanked. The red contours, starting from 8σ (σ = 0.27 Jy beam^{-1}) and increasing with the step of 8σ, show the CO (1-0) emission, obtained from NOEMA and IRAM-30 m combination (Gueth et al. 1996). Gray circle(s) in each panel indicates the primary beam of the NOEMA/SMA at corresponding line rest frequency, and the angular resolution after IRAM-30 m combination is plotted at the bottom left in each panel. The black curve guidelines the path of the precessing jet from L1157-mm as modeled by Podio et al. (2016). The black squares indicate the driving source, L1157-mm, and the shock knots in B0, B1, and B2.

portant in this work. Instead, the relative gradient of τ from B0e to B1a and B1i is ∼1, indicating that the relative gradient of SO column density in B0-B1 does not change significantly before and after line optical depth correction.

We are not able to estimate the τ of any $^{32}SO_2$ line, due to the lack of detections on their $^{34}SO_2$ line counterparts (with the same transition level). Nevertheless, the relative abundance ratio between this isotopologue pair is comparable with the standard Solar $\mathcal{R}_{^{32}S/^{34}S}$ (see Section 4.2), which indicates that the optical depths of the SO_2 lines may not lead to an underestimation of the SO_2 column density in this work.

4.2. *Rotation temperature and molecular column density from Rotational Diagram (RD)*

Although the line emissions are clumpy, their spatial distributions toward the positions list in Table 2 are more extended than their synthesized beams, so we do not take the beam filling factor into account in the following estimates. As the first-order approximation, we assume that all of the detected SO and SO_2 lines are optically thin and under LTE. Smoothing the line-intensity maps to the same angular resolution (i.e., 5.30″×4.86″) and the same pixel size (i.e., 0.5″), we use the RD method, and fit the relative population distribution of all the detected energy levels by assuming a single component toward each clumpy substructure (see RD fit toward B1a in Figure 5 as an example). The least-square solutions of the molecular rotational temperature T_{rot}, as well as total column density N_{rot} from fittings are list Table 3.

When taking the optical depth of the SO $(3_2 - 2_1)$ lines into account, the SO column density in pixels within B0-B1 increases by a factor of 1.5–2.5 (Table 3). Taking into account the uncertainties, it seems that the difference between the SO column density throughout the B0-B1 structure is little, ranging from (3.9 ± 1.7) × 10^{14} cm^{-2} (B1a) to (1.5 ± 1.0) × 10^{14} cm^{-2} (B1i). Also, the SO_2 column densities, as derived from RD, looks quite uniform, being around 1 × 10^{14} cm^{-2}. SO seems to have uniform T_{rot} toward the B0-B1 region (10–14 K), and the optical depth correction for the SO $(3_2 - 2_1)$ line does not bring in significant change in the T_{rot} estimates. In contrast, precise T_{rot} estimation of SO_2 depends on whether the high-J line (with E_u/k_B >100 K) is included, i.e., fitting to the mid-J SO_2 lines yields the same T_{rot} as that from fitting the low-J and mid-J SO lines (with E_u/k_B <70 K). A possible reason could be that the high-J SO_2 may trace a warmer component.

The upper energy level of the detected ^{34}SO lines are too close to derive the molecular rotational temperature map, and we are not able to obtain the rotational temperature of $^{34}SO_2$ from only one detected line ($7_{1,7}$−$6_{0,6}$). With the LTE and optically thin assumptions, we estimate the column densities of ^{34}SO and $^{34}SO_2$ by using the T_{rot} of their ^{32}S- isotopologues toward the clumpy substructures in B0-B1. The relative abundance ratio between each pair of ^{32}S- and ^{34}S- isotopologues is uniform (with a small standard deviation among pixels), especially toward B1. The ratio of $\chi(^{32}SO/^{34}SO)$ is 9 ± 2 over the entire B0-B1 (∼ 7 toward B0e and 15 ± 4 toward B1 after SO $(3_2 - 2_1)$ optical depth correction), while




Figure 3. Color maps show the intensity maps of all of the detected SO_2 and SO lines integrated over the velocity range of −10 to +5 km s^{-1}, zoomed in toward the central 1′ × 1′ region. All the maps are obtained from the combination of NOEMA (or SMA) and IRAM-30 m, and the emission < 5σ rms are blanked. Seven clumpy substructures labeled as B1a, B1b, B1c, B0e, B1f, B1i, and B2a show strong line emissions. Gray circle(s) in each panel indicates the primary beam of the NOEMA/SMA at corresponding line rest frequency, and the angular resolution after IRAM-30 m combination is plotted at the bottom left in each panel.

the ratio of $\chi(^{32}SO_2/^{34}SO_2)$ is ∼ 6 toward B0e and ∼ 15 toward B1a-B1c-B1b (assuming $\tau \leq 1$ of all of the SO_2 lines detected in this work). These ratios are consistent with the $\chi(C^{32}S/C^{34}S) \sim 15$ measured toward B1 from our previous single-dish observations (Bachiller & Pérez Gutiérrez 1997)[8], and is comparable with the $\mathcal{R}_{^{32}S/^{34}S} \sim 22$ in the solar system. Therefore, the optically thin assumption for the SO_2 lines can be used to measure the SO_2 column density in this work.

[8] The column density of $C^{34}S$ given in Bachiller & Pérez Gutiérrez (1997) was a typo, and it is corrected as "1.8(13)".

4.3. *Kinetic temperature and molecular column density from Large Velocity Gradient (LVG)*

In the temperature range of 10–100 K, the critical densitie of all of the detected SO and SO_2 isotopologue lines are in the range of $10^5 - 10^6$ cm^{-3} (estimated by using the Einstein coefficient and the collision rate given in Leiden Atomic and Molecular Database, Schöier et al. 2005, which are consistent with the effective critical density given by, e.g., Reiter et al. 2011). According to previous studies, such as Benedettini et al. (2013); Gómez-Ruiz et al. (2015), the number density toward the entire B1 region is 10^5–10^6 cm^{-3}. Therefore, excitation condition of SO and SO_2 gas in this shocked region need a LVG consideration.

Using the collision coefficients of SO-(para) H_2 and SO_2-(para and ortho) H_2 obtained from the BASECOL database




Figure 4. Map of the integrated intensity ratio $T_{B,^{34}SO_{(3_2-2_1)}}/T_{B,^{32}SO_{(3_2-2_1)}}$ (top widget) as well as the optical depth of $^{32}SO\,(3_2-2_1)$ (right widget) toward each pixel. The gray dashed contours, starting from 8σ ($\sigma = 0.27$ Jy beam^{-1}) and increasing with the step of 8σ, show the CO (1-0) emission (Gueth et al. 1996). The pixels where $^{34}SO\,(3_2-2_1)$ shows $< 6\sigma$ emissions are blanked. Six clumpy structures toward B1 are labeled.

(Dubernet et al. 2013), and applying the code of LVG approximation to non-LTE analysis described in Ceccarelli et al. (2002), we run grids of parameters and find the least-square (χ^2) best-fitting solution toward four positions along the precessing path, i.e., B0e, B1a (as well as B1b as a double-check), and B1i. B2a is not fit because less than four lines show $> 5\sigma$ emission. In the fittings, we adopted a FWHM linewidth of 7 km s^{-1} toward each position. Then, we vary three parameters in a relatively large range: H_2 number density n (10^4 cm^{-3}–10^8 cm^{-3}), gas kinetic temperature T_{kin} (30 K–300 K), molecular column density N_T (10^{13} cm^{-2}–10^{18} cm^{-2}). Note that the FWHM linewidth we adopt in this work is a compromise made to obtain the best fit by including all SO and SO_2 lines observed at a coarse velocity resolution (6–7 km s^{-1}). According to the previous analysis based on CO observations at a velocity resolution of 0.1–0.5 km s^{-1} (Lefloch et al. 2012), at least three components overlap in the velocity range from -30 to 5 km s^{-1}, with their intensity peaks close to the V_{sys}. Before we resolve them with observations at higher-velocity resolution, the adopted FWHM linewidth in the range of 5–10 km s^{-1} provides results that represent the average behavior of the different kinematical components within this velocity range.

Comparing the best-fit results for the column density by using LVG and RD methods (Table 3), we have a good agreement. In addition, we find that the SO column density toward B1i is slightly lower than those measured toward the rest of the clumps, even after taking into account the uncertainties (see Table 3). The kinetic temperatures are in the range of 30–200 K, being substantially in agreement with the kinetic temperatures found by Lefloch et al. (2012) in the different gas components coexisting in the B1 shocked gas. On the other hand, the gas volume density n from LVG fit of SO lines is well constrained, from $(3-5)\times10^4$ cm^{-3} (B0e) to 10^5 cm^{-3} (B1i), while n as traced by SO_2 (10^5–10^6 cm^{-3}) is denser than that traced by SO by a factor up to an order of magnitude (see Table 3). The higher values of n are in agreement with those derived by Gómez-Ruiz et al. (2015) using a CS multiline analysis. Therefore, SO and SO_2 may be tracing different gas portions.

5. DISCUSSION

5.1. *Map of relative abundance ratio between SO and* SO_2

Provided that the column densities as derived by the LVG analysis are in agreement with those derived from the LTE-RD analysis (after correction due to the optical depths), to investigate the possible chemical differentiation between SO and SO_2 along the path of the precessing jet in the entire B0-B1-B2 (in a time frame of thousand years), we fit the one-component RD toward pixels in B0-B1-B2 region.

When investigating the 0.1 pc scale field of view, we note that there is a tentative gradient along the path of the precessing jet in the $\chi(SO/SO_2)$ map before optical depth correction[9] (Figure 6): $\chi(SO/SO_2) < 1$ in B2, while $1.5 < \chi(SO/SO_2) < 3$ from the younger B0e. Unfortunately, only two SO_2 lines show $> 5\sigma$ emission toward B2, so SO_2 column density toward B2a is only an upper limit, i.e., $\chi(SO/SO_2)$ is a lower limit there. Only further SO_2 observations toward B2 will allow us to verify the occurrence of such a spatial trend.

To test whether the tentative gradient of $\chi(SO/SO_2)$ is due to optical depth effects, we zoom into the region where SO column density can be corrected by considering the optical depth of $SO\,(3_2-2_1)$ line. Even after the correction for optical depth effects, Figure 7 allows us to speculate a possible trend for the $\chi(SO/SO_2)$ ratio decreasing moving from north (younger gas) to south (older gas). Again, the present results can only suggest the occurrence of such a spatial trend and only future multiline observations, possibly on smaller spatial scales, will verify this speculation.

[9] Because of the line image sensitivity, the optical depth correction only works for the $SO\,(3_2-2_1)$ line toward the B0-B1 region. As a compromise for inspecting the map of column density ratio between SO and SO_2 in a larger field, we assume that all lines are optically thin and under LTE in Figure 6.




Figure 5. Rotation diagrams of SO (*left*) and SO_2 (*right*) toward B1a as an example. The black dots correspond to the line emission integrated in the velocity range of −10 to +5 km s^{-1}. The error bar of each data point comes from the uncertainty in observations and data reduction (calibration, combination, and intensity integration). The black line is fitted with the assumption that lines from the same species are optically thin, under LTE, and trace a single component toward B1a. The gray dashed line and gray dot in the *left* panel is fitted by taking the optical depth of SO ($3_2 - 2_1$) into account. The gray dashed line in the *right* panel is fitted by only taking the low-J and mid-J lines with $E_u/k_B < 70$ K into account.

Table 3. Best-fit results by using RD and LVG methods

		RD				LVG			
Location	Species	N_{rot} (10^{14} cm^{-2})	T_{rot} (K)	$N_{rot,\tau}$[a] (10^{14} cm^{-2})	$T_{rot,c}$[b] (K)	N_T (10^{14} cm^{-2})	T_{kin} (K)	n (10^5 cm^{-3})	χ^2
B0e	SO	2.1 ± 0.8	13 ± 1	1.8 ± 1.3	...	2.0 ± 0.2	70–200	0.3–0.5	0.94
	SO_2	1.0 ± 0.8	28 ± 9	...	12 ± 3	––[c]	––[c]	––[c]	––[c]
B1a	SO	2.4 ± 0.7	14 ± 1	3.9 ± 1.7	...	2.5 ± 0.5	40–70	0.8–0.9	0.6
	SO_2	1.2 ± 1.0	31 ± 8	...	15 ± 4	1.4 ± 0.4	50 ± 10	3 ± 2	0.5
B1b	SO	2.3 ± 1.4	13 ± 1	3.5 ± 2.0	...	2.0 ± 0.2	30–100	0.8–1	1.2
	SO_2	1.2 ± 1.0	25 ± 5	...	15 ± 4	1.0 ± 0.2	30 ± 20	10 ± 5	0.7
B1i	SO	1.4 ± 0.8	13 ± 1	1.5 ± 1.0	...	1.4 ± 0.2	30–100	0.8–1	0.5
	SO_2	1.4 ± 0.7	28 ± 9	...	12 ± 3	1.2 ± 0.2	50 ± 5	3 ± 1	0.5
B2a	SO	0.8 ± 0.5	19 ± 3	...	...	––[c]	––[c]	––[c]	––[c]
	SO_2	< 1.7[d]	...	...	< 64[d]	––[c]	––[c]	––[c]	––[c]

Note. *a*. Column density estimated by taking the optical depth of SO ($3_2 - 2_1$) into account.

b. Rotation temperature derived by only taking the lines with $E_u/k_B < 70$ K into account.

c. Less than four lines show emission with S/N > 5 in the primary beam.

d. Upper limit is given by using two lines which show emission with S/N > 5 .

5.2. *Chemical modeling*

Our previous chemical study, especially the observations of the ions and COMs, unveiled at least three chemically different layers with gas-grain components from the north (B1a) to south (B1i) toward B1 (Codella et al. 2017, 2020). To trace back the possible forming path of SO and SO_2 from the $\chi(SO/SO_2)$ gradient through B0-B1-B2, we apply the same time-dependent gas-phase chemical model as that introduced by Codella et al. (2017). Briefly, we used the MyNahoon code, which is based on the Nahoon code originally developed by Wakelam et al. (2012). The code follows the gaseous abundances evolution in time. Note that it does not consider the reactions possibly occurring on the grain surfaces, except the formation of H_2 molecules, as our goal is to study a subsequent phase: namely, when the grain mantles are already formed, in conditions and timescales where the surface reactions are completely negligible (see later). In short, we simulate the physical effects caused by the shock passage by suddenly increasing the H_2 gas number density n from 10^4 to 10^5 cm^{-3}, and the temperature from 10 to 70 K.

In addition, because of the sputtering and/or grain-grain collisions caused by the shock, a number of species previously frozen on the grain mantles are injected into the gas phase. In our model, we assumed the injected species and their abundance as those in Codella et al. 2017 (their Table B.1), which is based on several previous studies of the same source (Tafalla & Bachiller 1995; Benedettini et al. 2013; Podio et al. 2014; Codella et al. 2015) or known properties of




Figure 6. SO (*left*) and SO_2 (*middle*) column density maps and their relative abundance ratio map (*right*) derived by using the RD method, with the assumption that all lines are optically thin and under LTE. In the *left* panel, the pixels where less than three SO lines show $> 5\sigma$ emission are blanked. In the *middle* panel, the pixels where less than two SO_2 lines show $> 5\sigma$ emission are blanked, and the black contour denotes the region within which at least three lines show $> 5\sigma$ emission. In the *right* panel, the pixels where less than two SO_2 lines show $> 5\sigma$ emission are blanked, and the gray line indicates the path of the precessing jet (Podio et al. 2016). These maps are derived by smoothing all lines to the same angular resolution, i.e., $5.30'' \times 4.86''$, shown as the synthesised beam in the bottom left of each panel. The gray dashed contours and the labeled clumpy structures are the same as shown in Figure 4.

the ice mantle compositions (Boogert et al. 2015; Busquet et al. 2014).

In particular, with respect to the S-bearing frozen species, Podio et al. (2014) could only roughly constrain them, based on the S-bearing ions SO^+ and HCS^+. These authors found that OCS is injected in the gas phase with an abundance of 6×10^{-6} (with respect to the H nuclei) but could not constrain the injected abundances of H_2S and S, two species that are expected to be also present in the grain mantles (see e.g., Wakelam et al. 2004a; Vidal & Wakelam 2018). Therefore, we ran models with different injected abundances of H_2S and S and tried to constrain them based on the new observations presented in this work.

The results of the modeling are shown in Figure 8. In the figure, we report the model predicted SO/SO_2 abundance ratio as a function of the time, namely the age of the shocked gas and for different values (2×10^{-6} and 1×10^{-8}) of the injected H_2S/H and S/H abundances. In the same figure, we also report the values measured in the previous sections (the uncertainty is given based on RD optically thin assumption, RD optical correction for the low-J SO line, and LVG) toward B1a, B1b and B1i and their ages estimated by Gueth et al. (1996) and Podio et al. (2016). Finally, we also show the model predictions for values slightly different of the density and cosmic ray ionization rate to have a rough "sensitivity" study and understand how much the predictions also depend on these, only roughly constrained, quantities.

The figure shows that the model predictions compare rather well with the observations and, remarkably, with the increasing ages of B1a (B1b) to B1i. Based on the predictions, the injected H_2S/H and S/H abundances are equal to 2×10^{-6}: injecting pre-frozen OCS only is not enough to reproduce our new observations, an additional injection of previously frozen S-bearing species in form of H_2S and S in approximately the same amount is also needed (a similar result was also found by Holdship et al. 2016). Second, the cosmic ray ionization rate is around $3 \times 10^{-16}\,s^{-1}$, namely slightly (a factor 2) lower than that used in previous studies. This may mean that it slightly decreases going outward from the central object from which the outflow/jets emanate, which is consistent with the hypothesis that cosmic rays like particles are accelerated by the dense shocks at the jet base (Padovani et al. 2016; Ivlev et al. 2019).

Of course our model, although so far is successful in reproducing several species (see, for example, Codella et al. 2017, 2020), is rather simplified; it does not take into account, for




Figure 7. SO column density map (*upper*) and map of its relative abundance ratio with respect to SO_2 (*lower*) derived by using the RD method, with the LTE assumption and optical depth correction for the SO ($3_2 - 2_1$) line. The pixels where ^{34}SO ($3_2 - 2_1$) shows $< 6\sigma$ emissions are blanked. In the *right* panel, the gray line indicates the path of the precessing jet (Podio et al. 2016). These maps are derived by smoothing all lines to the same angular resolution, i.e., $5.30'' \times 4.86''$, shown as the synthesised beam in the bottom left of each panel. The gray dashed contours and the labelled clumpy structures are the same as shown in Figure 4.




Figure 8. SO/SO_2 abundance ratio predicted by our astrochemical model (see text) as a function of the time for different H_2S/H and S/H abundances, 2×10^{-6} (red curves) and 1×10^{-8} (green curves). The model predictions are also obtained for two values of the cosmic rays ionization rates, 6×10^{-16} s^{-1} (solid curves) and 3×10^{-16} s^{-1} (dashed curves), and a smaller H density, 2×10^4 cm^{-3} (dotted-dashed curve). The results from the analysis of the observations are reported in the dashed areas toward three positions along the B1 region: B1a, B1b, and B1i, respectively.

example, the short-lived hot/warm phase of the post-shocked gas, which, in the case of SO_2 could have an impact on the predictions. Having said that, it is encouraging to see that such a simplified model succeeds to grab the basic gaseous chemical composition changes caused by the shock.

6. CONCLUSIONS

We simultaneously analyze the chemical properties of SO and SO_2 toward all of the brightest shocked regions along the L1157 chemically rich blueshifted outflow, namely, B0, B1, and B2. Based on the data taken in the context of the SOLIS IRAM-NOEMA Large Program, and the supplementary observations from SMA interferometer as well as IRAM-30 m telescope at 1.1–3.6 mm wavelengths, our conclusions are as follows:

1. At an angular resolution of 1″–3″ (500–1400 au at the source distance of 352 pc), SO and SO_2 may trace different gas toward B1. This is indicated from the different spatial distributions of five SO_2 lines (including one $^{34}SO_2$ line) and eight SO lines (including two ^{34}SO lines) which we detected. The emission peak of the SO lines are in general toward northeast, where the gas is impacted by the recent jet impinging on the B1 cavity wall; while the emission peak of the SO_2 lines are all toward the southern edge of the older shock front. Moreover, LVG constraints on gas volume density indicates that, the gas traced by SO_2 has a volume density of 10^5–10^6 cm^{-3}, which is denser than that traced by SO by a factor up to an order of magnitude.

2. Investigating the 0.1 pc scale field of view, there is a tentative gradient in the relative abundance ratio of $\chi(SO/SO_2)$ along the path of the associated precessing jet, showing smooth decrease from B0-B1a (> 2) to B1i-B2 (< 1). These findings call for further analysis at higher spatial resolutions to verify the tentative trends.

3. Our astrochemical modeling shows that the SO and SO_2 abundances evolve on timescales less than about 1000 yr. The computed values of timescales toward individual pixels are consistent with the ages estimated in L1157 B0-B1 by previous studies. Furthermore, the modeling requires high abundances of both H_2S/H and S/H (2×10^{-6}) injected in the gas phase due to the shock occurrence, so pre-frozen OCS only is not enough to reproduce our new observations.

We thank the NOEMA, SMA, and IRAM-30 m staff for their helpful support during the observations and data reduction of the *SOLIS* project. We also thank the referee for constructive suggestions.

S.F. acknowledges the support of National Natural Science Foundation of China No. 11988101, the support of the CAS

International Partnership Program No.114A11KYSB20160008, and the support of the EACOA fellowship from the East Asia Core Observatories Association (EACOA). EACOA consists of the National Astronomical Observatory of China, the National Astronomical Observatory of Japan, the Academia Sinica Institute of Astronomy and Astrophysics, and the Korea Astronomy and Space Science Institute.

This work was also supported by (i) the French program "Physique et Chimie du Milieu Interstellaire" (PCMI) funded by the Conseil National de la Recherche Scientifique (CNRS) and Centre National d'Etudes Spatiales (CNES), (ii) by the Italian project PRIN-INAF 2016 The Cradle of Life - GENESIS-SKA (General Conditions in Early Planetary Systems for the rise of life with SKA), (iii) by the program PRIN-MIUR 2015 STARS in the CAOS - Simulation Tools for Astrochemical Reactivity and Spectroscopy in the Cyberinfrastructure for Astrochemical Organic Species (2015F59J3R, MIUR Ministero dell'Istruzione, dell'Università della Ricerca e della Scuola Normale Superiore), (iv) by the French Agence Nationale de la Recherche (ANR), under reference ANR-12-JS05-0005, and (v) by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme, for the Project "The Dawn of Organic Chemistry" (DOC), grant agreement No. 741002, (vi) the Ministry of Science and Technology (MoST) of Taiwan (grant No. 108-2112-M-001-002-MY3). This research made use of NASA's Astrophysics Data System.

Software: GILDAS/CLASS (Pety 2005), MIR (Scoville et al. 1993), MIRIAD (Sault et al. 1995)

REFERENCES

Arce, H. G., Santiago-García, J., Jørgensen, J. K., Tafalla, M., & Bachiller, R. 2008, ApJL, 681, L21

Bachiller, R. & Pérez Gutiérrez, M. 1997, ApJL, 487, L93

Bachiller, R., Pérez Gutiérrez, M., Kumar, M. S. N., & Tafalla, M. 2001, A&A, 372, 899

Benedettini, M., Viti, S., Codella, C., et al. 2007, MNRAS, 381, 1127

Benedettini, M., Viti, S., Codella, C., et al. 2013, MNRAS, 436, 179

Boogert, A. C. A., Gerakines, P. A., & Whittet, D. C. B. 2015, ARA&A, 53, 541

Boogert, A. C. A., Schutte, W. A., Helmich, F. P., Tielens, A. G. G. M., & Wooden, D. H. 1997, A&A, 317, 929

Buckle, J. V. & Fuller, G. A. 2003, A&A, 399, 567

Burkhardt, A. M., Dollhopf, N. M., Corby, J. F., et al. 2016, ApJ, 827, 21

Busquet, G., Lefloch, B., Benedettini, M., et al. 2014, A&A, 561, A120

Ceccarelli, C., Bacmann, A., Boogert, A., et al. 2010, A&A, 521, L22

Ceccarelli, C., Baluteau, J. P., Walmsley, M., et al. 2002, A&A, 383, 603

Ceccarelli, C., Caselli, P., Fontani, F., et al. 2017, ApJ, 850, 176

Charnley, S. B., Tielens, A. G. G. M., & Rodgers, S. D. 1997, ApJL, 482, L203

Chin, Y.-N., Henkel, C., Whiteoak, J. B., Langer, N., & Churchwell, E. B. 1996, A&A, 305, 960

Codella, C., Bachiller, R., Benedettini, M., et al. 2005, MNRAS, 361, 244

Codella, C., Benedettini, M., Beltrán, M. T., et al. 2009, A&A, 507, L25

Codella, C., Ceccarelli, C., Bianchi, E., et al. 2020, A&A, 635, A17

Codella, C., Ceccarelli, C., Caselli, P., et al. 2017, A&A, 605, L3

Codella, C., Fontani, F., Ceccarelli, C., et al. 2015, MNRAS, 449, L11

Codella, C., Lefloch, B., Ceccarelli, C., et al. 2010, A&A, 518, L112

Dubernet, M. L., Alexander, M. H., Ba, Y. A., et al. 2013, A&A, 553, A50

Feng, S., Beuther, H., Henning, T., et al. 2015, A&A, 581, A71

Fontani, F., Codella, C., Ceccarelli, C., et al. 2014a, ApJL, 788, L43

Fontani, F., Sakai, T., Furuya, K., et al. 2014b, MNRAS, 440, 448

Frank, A., Ray, T. P., Cabrit, S., et al. 2014, in Protostars and Planets VI, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 451

Geballe, T. R., Baas, F., Greenberg, J. M., & Schutte, W. 1985, A&A, 146, L6

Gómez-Ruiz, A. I., Codella, C., Lefloch, B., et al. 2015, MNRAS, 446, 3346

Gómez-Ruiz, A. I., Hirano, N., Leurini, S., & Liu, S.-Y. 2013, A&A, 558, A94

Gueth, F., Guilloteau, S., & Bachiller, R. 1996, A&A, 307, 891

Gueth, F., Guilloteau, S., & Bachiller, R. 1998, A&A, 333, 287

Hatchell, J., Thompson, M. A., Millar, T. J., & MacDonald, G. H. 1998, A&A, 338, 713

Ho, P. T. P., Moran, J. M., & Lo, K. Y. 2004, ApJL, 616, L1

Holdship, J., Jimenez-Serra, I., Viti, S., et al. 2019, ApJ, 878, 64

Holdship, J., Viti, S., Jimenez-Serra, I., et al. 2016, MNRAS, 463, 802

Ivlev, A. V., Silsbee, K., Sipilä, O., & Caselli, P. 2019, ApJ, 884, 176

Jiménez-Serra, I., Martín-Pintado, J., Rodríguez-Franco, A., & Martín, S. 2005, ApJL, 627, L121

Jørgensen, J. K., Bourke, T. L., Myers, P. C., et al. 2007, ApJ, 659, 479
Kaufman, M. J. & Neufeld, D. A. 1996, ApJ, 456, 250
Lefloch, B., Bachiller, R., Ceccarelli, C., et al. 2018, MNRAS, 477, 4792
Lefloch, B., Cabrit, S., Busquet, G., et al. 2012, ApJL, 757, L25
Lefloch, B., Cabrit, S., Codella, C., et al. 2010, A&A, 518, L113
Lefloch, B., Ceccarelli, C., Codella, C., et al. 2017, MNRAS, 469, L73
Li, J., Wang, J., Zhu, Q., Zhang, J., & Li, D. 2015, ApJ, 802, 40
Liu, H. B., Jiménez-Serra, I., Ho, P. T. P., et al. 2012, ApJ, 756, 10
Lodders, K. 2003, ApJ, 591, 1220
Mendoza, E., Lefloch, B., López-Sepulcre, A., et al. 2014, MNRAS, 445, 151
Minh, Y. C., Liu, H. B., & Galvań-Madrid, R. 2016, ApJ, 824, 99
Müller, H. S. P., Schlöder, F., Stutzki, J., & Winnewisser, G. 2005, Journal of Molecular Structure, 742, 215
Müller, K. E., Shirley, Y. L., Evans, II, N. J., & Jacobson, H. R. 2002, ApJS, 143, 469
Myers, P. C., Linke, R. A., & Benson, P. J. 1983, ApJ, 264, 517
Nisini, B., Giannini, T., Neufeld, D. A., et al. 2010, ApJ, 724, 69
Padovani, M., Marcowith, A., Hennebelle, P., & Ferrière, K. 2016, A&A, 590, A8
Palumbo, M. E., Tielens, A. G. G. M., & Tokunaga, A. T. 1995, ApJ, 449, 674
Pety, J. 2005, in SF2A-2005: Semaine de l'Astrophysique Francaise, ed. F. Casoli, T. Contini, J. M. Hameury, & L. Pagani, 721
Pety, J., Schinnerer, E., Leroy, A. K., et al. 2013, ApJ, 779, 43
Pineau des Forets, G., Roueff, E., Schilke, P., & Flower, D. R. 1993, MNRAS, 262, 915
Podio, L., Codella, C., Gueth, F., et al. 2016, A&A, 593, L4
Podio, L., Codella, C., Lefloch, B., et al. 2017, MNRAS, 470, L16
Podio, L., Lefloch, B., Ceccarelli, C., Codella, C., & Bachiller, R. 2014, A&A, 565, A64
Qi, C. 2003, in SFChem 2002: Chemistry as a Diagnostic of Star Formation, ed. C. L. Curry & M. Fich, 393
Reiter, M., Shirley, Y. L., Wu, J., et al. 2011, ApJS, 195, 1
Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in Astronomical Society of the Pacific Conference Series, Vol. 77, Astronomical Data Analysis Software and Systems IV, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes, 433
Schöier, F. L., van der Tak, F. F. S., van Dishoeck, E. F., & Black, J. H. 2005, A&A, 432, 369
Scoville, N. Z., Carlstrom, J. E., Chandler, C. J., et al. 1993, PASP, 105, 1482
Tafalla, M. & Bachiller, R. 1995, ApJL, 443, L37
Tieftrunk, A., Pineau des Forets, G., Schilke, P., & Walmsley, C. M. 1994, A&A, 289, 579
Tobin, J. J., Hartmann, L., Looney, L. W., & Chiang, H.-F. 2010, ApJ, 712, 1010
Vidal, T. H. G. & Wakelam, V. 2018, MNRAS, 474, 5575
Viti, S., Jimenez-Serra, I., Yates, J. A., et al. 2011, ApJL, 740, L3
Wakelam, V., Caselli, P., Ceccarelli, C., Herbst, E., & Castets, A. 2004a, A&A, 422, 159
Wakelam, V., Castets, A., Ceccarelli, C., et al. 2004b, A&A, 413, 609
Wakelam, V., Herbst, E., Loison, J.-C., et al. 2012, ApJS, 199, 21
Wilson, T. L. & Rood, R. 1994, ARA&A, 32, 191
Zucker, C., Speagle, J. S., Schlafly, E. F., et al. 2019, ApJ, 879, 125

APPENDIX

Figure A1 shows the profile of the SO and SO_2 isotopologue lines we use to measure the molecular column densities toward seven positions.

Table A1 lists the NOEMA observation dates, spectral setup, configurations, calibrators, and weather conditions.




Figure A1. Profiles of identified SO and SO_2 isotopologue lines, averaged from a beam-sized region with the center toward each clumpy substructure in the plane of the sky (Table 2). All lines are extracted from images smoothed to the same pixel size and angular resolution (5.1″). Their velocity resolutions are listed in Table 1. In each panel, two red vertical lines at $-10\,\mathrm{km\,s^{-1}}$ and $+5\,\mathrm{km\,s^{-1}}$ indicate the velocity range for which we integrate the intensity; the gray dashed vertical line indicate the V_{sys} = $2.7\,\mathrm{km\,s^{-1}}$ of the cloud. The horizontal cyan line indicates the baseline (T_B=0 K).




Figure A1. (continued)

Table A1. NOEMA observation log toward L1157-B1

Setup	WIDEX (GHz)	Configuration	Observation Date	Calibrators (Bandpass)	(Phase/Amp)	(Flux)	T_{sys} (K)	PWV (mm)
1	80.80-84.40	D (6ant, 24-97m)	2015-Jul-15	1928+738	1928+738	MWC349	100-150	10-30
			2015-Jul-23	1749+096	1928+738	MWC349	90-160	10-15
			2015-Jul-24	3C84	1928+738	MWC349	100-300	20-40
		D (5ant, 32-97m)	2015-Jul-26	3C454.3	1928+738	MWC349	60-130	4-8
		C (7ant, 24-240m)	2015-Oct-25	3C454.3	1928+738	MWC349	90-120	4-6
			2015-Oct-27	3C84	1928+738	LKHA101	200-400	10-40
			2015-Nov-04	3C345	1928+738	MWC349	100-300	10-40
			2015-Nov-04	1928+738	1928+738	LKHA101	110-130	4-8
2	95.85-99.45	C (7ant, 48-304m)	2016-Oct-26	0923+392	2007+659	MWC349	100-250	6-20
			2016-Oct-27	1418+546	2007+659	MWC349	50-100	2-5
			2016-Oct-28	2013+370	2007+659	MWC349	50-90	2-5
		C (8ant, 24-304m)	2016-Oct-22	3C454.3	2007+659	MWC349	80-250	4-5
		D (8ant, 24-176m)	2017-Jan-22	2200+420	2007+659	MWC349	60-100	4-5
			2017-Jan-24	2013+370	2007+659	MWC349	50-90	1-3
3	204.00-207.60	A (7ant, 96-760m)	2016-Jan-29	2013+370	1928+738	MWC349	200-300	4-6
			2016-Feb-06	2013+370	1928+738	MWC349	160-300	3-6
			2016-Feb-11	3C454.3	1928+738	MWC349	200-400	3-10
			2016-Feb-14		1928+738	LKHA101	130-250	1-3
			2016-Feb-16	3C84, 1055+018	1928+738	LKHA101	200-400	3-4
		C (6ant, 24-240m)	2016-Apr-20	1949+096	1928+738	MWC349	100-250	1-4
		C (7ant, 24-192m)	2016-May-06	3C454.3	1928+738	MWC349	110-180	1-2
		C (7ant, 48-304m)	2016-Oct-15	3C454.3	1928+738	MWC349	100-150	2-3