



Link between steroidogenesis, the cell cycle, and PKA in adrenocortical tumor cells

Marthe Rizk-Rabin, Sabrina Chaoui-Ibadioune, Anna Vaczlavik, Christopher Ribes, Michel Polak, Bruno Ragazzon, Jérôme Bertherat

► To cite this version:

Marthe Rizk-Rabin, Sabrina Chaoui-Ibadioune, Anna Vaczlavik, Christopher Ribes, Michel Polak, et al.. Link between steroidogenesis, the cell cycle, and PKA in adrenocortical tumor cells. *Molecular and Cellular Endocrinology*, 2020, 500, 10.1016/j.mce.2019.110636 . hal-03014038

HAL Id: hal-03014038

<https://cnrs.hal.science/hal-03014038>

Submitted on 19 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Link between steroidogenesis, the cell cycle, and PKA in adrenocortical tumor cells

2
3 Marthe Rizk-Rabin¹, Sabrina Chaoui- Ibadioune¹, Anna Vaczlavik¹, Christopher Ribes¹,
4 Michel Polak^{1,3}, Bruno Ragazzon¹, and Jérôme Bertherat^{1,2}.

5
6 ¹Institut Cochin, U1016, CNRS (UMR 8104), Université Paris Descartes, Paris, France,

7 ²Hôpital Cochin, Department of Endocrinology. Center for Rare Adrenal Diseases, Paris,
8 France.

9 ³Hopital Necker Enfants maladies , Department of Endocrinology , Paris France.

10
11 Corresponding author: Marthe Rizk-Rabin, marthe.rizk@inserm.fr

12 Institut Cochin, Faculté Cochin Département Endocrinologie Métabolisme et Diabète. 24 Rue
13 du Fbg St Jacques. 75014 Paris.

14 Tel: 01 53 73 27 31, Fax: 01 44 41 24 21

15 Short title: Cell cycle and PKA control steroidogenesis

16 Key words: cell-cycle phases, PKA activity, MAP kinase, steroidogenesis genes, Star, PKA
17 R1A and CA.

Abstract

Adrenocortical tumors (ACTs) frequently cause steroid excess and present cell-cycle dysregulation. cAMP/PKA signaling is involved in steroid synthesis and play a role in cell-cycle regulation. We investigated, by cell synchronization in the different phases of the cell-cycle, the control of steroidogenesis and the contribution of PKA in adrenocortical cells (H295R and culture of primary pigmented nodular adrenocortical disease cells). Cells showed increased steroidogenesis and a maximal PKA activity at G2 phase, and a reduction at G1 phase. *PRKACA* overexpression, or cAMP stimulation, enhanced PKA activity and induced steroidogenesis in all synchronized groups but is not sufficient to drive cell-cycle progression. *PRKARIA* inactivation enhanced PKA activity and induced *STAR* gene expression, only in cells in G1, and triggered cell-cycle progression in all groups. These findings provide evidence for a tight association between steroidogenesis and cell-cycle in ACTs. Moreover, *PRKARIA* is essential for mediating the function of PKA activity on both steroidogenesis and cell-cycle progression in adrenocortical cells.

1. Introduction

Adrenocortical tumors (ACTs) are often associated with increased steroid secretion, which contributes to morbidity. This has been observed for benign tumors, such as those of primary pigmented nodular adrenocortical disease (PPNAD) (Espiard et al., 2014), macronodular hyperplasia (Louiset et al., 2013), adrenocortical adenomas (Bassett et al., 2005), and malignant adrenal cortical carcinoma (ACC) (Nakamura et al., 2015; Abiven et al., 2006). Even ACCs, which are classified as endocrine-inactive based on routine hormonal assessment, are often responsible for abnormal steroid secretion, as demonstrated by mass spectrometry analysis (Arlt et al., 2011). ACT cells also frequently display cell-cycle control defects (Ragazzon et al., 2011; Stratakis, 2009; Assie et al., 2014). Consistent with this observation, amplifications and deletions of genes encoding key regulators of the cell cycle (CDK4, CCNE1, RB1, CDKN2A) have been described in several studies and are more frequent in ACCs (Assie et al., 2014; De Martino et al., 2013; Ross et al., 2014; Pereira et al., 2018).

It is well established that the effects of steroid hormones, such as glucocorticoids, androgens, and estrogens, on cell proliferation and differentiation are mediated by the regulation of key events in the control of cell-cycle progression or arrest (Kullmann et al., 2013; Heisler et al., 1997; Amanatullah et al., 2002), but little is known concerning steroid-hormone secretion during the cell cycle in the adrenal gland. The adrenolytic drug mitotane (op'DDD), used in ACC treatment, decreases cortisol secretion and induces the destruction of the zona fasciculata and zona reticularis of the adrenal cortex (Cerquetti et al., 2008). Its use has been shown to be associated with cell-cycle arrest at the G2/M phase and apoptosis (Doghman et al., 2013; Hescot et al., 2013).

Genetic alterations leading to constitutive activation of the cAMP/PKA pathway in various types of ACTs, such as cortisol secreting adenomas (ACA) or PPNAD, result in steroid over-secretion (Lodish et al., 2016). The best examples are inactivating mutations of the regulatory subunit R1A of PKA (*PRKAR1A*), which is mutated in the germline DNA of patients with PPNAD and Carney complex or tumor DNA in a subset of secreting ACAs and ACCs (Kirschner et al., 2000; Bertherat et al., 2003; Zheng et al., 2016). In benign adrenal tumors associated with Cushing syndrome, *PRKACA* somatic mutations also result in constitutive PKA activity by disrupting the interaction between *PRKACA* and the regulatory subunits of PKA, including *PRKAR1A* (Beuschlein et al., 2014; Goh et al., 2014; Calebiro et al., 2014; Cao et al., 2014; Sato et al., 2014). Changes in cAMP signal transduction also play a role in regulating the cell cycle (Massimi et al., 2017; Gupta et al., 2017; Sun et al., 2009). Activation of the cAMP/PKA pathway in many cell types inhibits proliferation by preventing cells from entering the S phase and arresting cells in G1 (Duncan et al., 2006; Hewer et al., 2011). In contrast, stimulation of the cAMP/PKA pathway may facilitate cell proliferation, such as in thyroid and pituitary tumors (Medri et al., 1994; Spada et al., 1992).

Cyclic AMP exerts its actions primarily through the activation of protein kinase A (PKA). PKA is composed of two types of regulatory and four types of catalytic subunits. The PKA regulatory subunits have been studied for their role in tumor formation (Taylor et al., 2013). *In vivo* and *in vitro* studies have shown that inactivating *PRKAR1A* leads to tumorigenesis (Kirschner et al., 2005; Nadella et al., 2005). These studies showed that the lack of *Prkar1a* protein in primary mouse cells *in vitro* causes constitutive PKA activation and immortalization of primary mouse embryonic fibroblasts (MEFs). At the molecular level, knocking out *Prkar1a* led to the upregulation of D-type cyclins (Kirschner et al., 2005). Similarly, we showed that inactivation of *PRKAR1A* confers resistance to apoptosis and enhances proliferation in the adrenocortical cell line H295R (Basso et al., 2014). *PRKAR1A*

99 inactivation disrupted the cell-cycle checkpoint and cells accumulated at the G2 phase with an
100 increase in cyclin D and PKA CA activity. These results suggested that there may be a link
101 between the increase in steroid secretion and cell-cycle dysregulation, which are
102 concomitantly observed in adrenal tumors.

103 Here, we investigated the cell-cycle-dependent control of adrenal-steroid oversecretion and
104 the contribution of cAMP/PKA in two cell lines synchronized by pharmacological drugs:
105 H295R, a human adrenal cortex cell line derived from an ACC (Gazdar et al., 1990) that does
106 not harbor any mutations in the cAMP/PKA pathway, and primary cultured PPNAD cells,
107 with inactivating mutations of *PKARIA*. We mimicked the constitutive activation of PKA that
108 is observed in adrenal tumors with *PRKARIA* inactivating (Kirschner et al., 2000; Bertherat et
109 al., 2003) or *PRKACA* activating (Beuschlein et al., 2014; Calebiro et al., 2014; Weigand et al.,
110 2017) mutations using either cAMP treatment, transient silencing of *PRKARIA* by
111 transfection with an siRNA, or overexpression of one of the catalytic subunits of PKA,
112 concomitant with the pharmacological drugs.

113 Overall, our data show that the hypersecretion of steroids by the H295R ACC cell line and
114 PPNAD is controlled by the phase of the cell cycle and that PKA activity plays a role in such
115 coordination.

116

2. Materials and methods

2.1 Cell culture

Human H295R adrenocortical carcinoma cells, purchased from the ATCC, were grown as previously described (Bouizar et al., 2010; Ragazzon et al., 2009; Groussin et al., 2000), in DMEM Ham/F12 medium (Sigma, St. Quentin Fallavier France) supplemented with 50 units/ml penicillin, 50 mg/ml streptomycin, 2 mmol/L glutamine, 2% Ultrosor G2 (Biosepra, Fremont, CA), and ITS (5 µg/ml insulin, 5 µg/ml transferrin, and 5 ng/ml selenium) (Sigma) at 37°C in an atmosphere of 5% carbon dioxide/95% air. Cells were allowed to attach for 24 h before use.

2.2 Primary cell culture of PPNAD

Human adrenals were obtained after informed consent from two CNC patients undergoing surgery for PPNAD. Adrenal tissue collection was approved by the ethics committee of Cochin Hospital. PPNAD adrenal tissues were obtained during surgery and immediately immersed in culture medium. Briefly, adrenocortical explants were enzymatically dispersed as previously described (Cazabat et al., 2014; Bram et al., 2016). Adrenocortical cells were cultured at 37°C in 5% CO₂ and allowed to attach for four days before use. Cells were incubated with various pharmacological inhibitors to synchronize the cell-cycle for 24 h with fresh DMEM medium, as for the H295R cell line.

2.3 Western blotting

Whole-cell protein lysates were obtained by extraction with RIPA lysis buffer containing 50 mM Tris-HCl, pH 7.5, 1 mM EDTA, 150 mM NaCl, 0.1% v/v Nonidet P-40 (NP40), anti-phosphatase, and anti-protease (Roche, Boulogne Billancourt, France) as previously described (Basso et al., 2014). Western blotting was performed with primary mouse antibodies for R1A (1/500, BD Laboratories, Rungis, France), R2B (1/1000), PKA catalytic subunits (1/1000)

(Santa Cruz Biotechnology, Nanterre, France), and Star (1/1000) (Cell Signaling, St Quentin en Yvelines, France). We used anti- β -actin (1/2500) (Sigma) or GAPDH antibodies (1/1000) (Santa Cruz Biotechnology) as loading controls. The antigen-antibody complexes were visualized with appropriate secondary antibodies (Santa Cruz Biotechnology) by chemiluminescence detection (ECLTM kit, Thermo scientific, Saint Herblan France). The signals were digitized with the Syngene/Gene Tool analysis system. (Ozyme, Montigny Le Bretonneux, France)

2.4 Cell-cycle distribution

Cell cultures were harvested, rinsed with PBS, fixed in 500 μ l 70% ethanol, and frozen (-20°C). Fixed cells were centrifuged, rinsed in PBS, suspended in 500 μ l PBS containing 50 μ g/ml propidium iodide (Sigma) and 100 μ g/ml RNase A (Sigma), and examined by flow cytometry (FACscan, EPICS XL Coulter, Villepinte France). Data were analyzed using Multicycle software. All experiments were carried out three times. Each histogram was constructed from at least 10,000 cells (Rizk-Rabin et al., 2008).

2.5 Cell-cycle synchronization

We used three different inhibitors, each specific for a cell-cycle check point: L mimosine for G1 (Vackova et al., 2003), aphidicolin for G1/S (Sorscher et al., 1991), and nocodazole for G2 (Ho et al., 2001) (Sigma). The cell-cycle phases of H295R cells arrested at a specific point in the cell cycle were well defined in a previous report (Bouizar et al., 2010). Cells were cultured in 12-well plates at a density of 150×10^3 cells/well and then treated for 24 h with L-mimosine (400 μ M) or aphidicolin (10 μ M) or for 24 h or 48 h with nocodazole (10 μ M). For cell-cycle release of the H295R cells, following 24 or 48 h of synchronization, cells were washed three times with PBS and cultured in medium without drugs for 6, 12, 24, 36, or 48 h. For PKA activation or inhibition, cells were transiently transfected with an siRNA for

PRKARIA, the control scramble RNA (siS), *PKACA* (in the PCDNA3+ vector, a gift from Dr D. Calebiro), or the empty vector, or incubated with the PKA inhibitor PKI 14-22 (Calbiochem, Fontenay sous-Bois France) and cultured with the different drugs for 24 h.

2.6 Transfection and co-transfection

H295R cells cultured in six-well plates were transfected with 100 pmol siRNA *PRKARIA* (UGAAUGGGCAACCAGUGUdTdT), the siRNA negative control (siS) (CAGUCGCGUUUGCGACUGGdTdT), 500 ng *PKACA* or PCDNA3+, as empty vector, using the Effectene reagent (Promega, Charbonnières les Bains, France). Cells were analyzed 24 h after transfection. For luciferase assays, 24-well plates were used. Cells were co-transfected with the siRNAs or vectors and a luciferase reporter gene driven by Star promoters. Firefly and Renilla luciferase activities (Promega) were measured as previously described (Basso et al., 2014; Ragazzon et al., 2009).

2.7 Steroid measurement.

Cortisol and aldosterone levels in cell medium were measured by competitive immunoassays and revealed by chemiluminescence assays (Laboratory of Hormonology at Cochin Hospital, France).

2.8 Protein kinase A assay

The PepTag nonradioactive protein kinase assay kit (Promega) was used to measure PKA activity, according to the manufacturer's instructions, as described in (Rizk-Rabin et al., 2008)

2.9 Analysis of RNA by quantitative PCR

Total RNA, extracted from the cell lines, was treated with DNase and further purified using the RNeasy Mini kit and RNase-free DNase Set (Promega), according to the manufacturer's instructions. Purified RNA was reverse transcribed with high-capacity cDNA reverse

transcriptase (Applied Biosystems, Illkirch France) and expression levels of the target genes analyzed by quantitative PCR using a LightCycler Fast Start SyberGreen kit (Roche Diagnostics, Meylan France), according to the manufacturer's instructions. The PCR conditions for all target steroidogenic and cyclin genes are shown in Table 1. Relative quantification of target cDNA was determined by calculating the difference in cross-threshold (CT) values after normalization to PPIA (CYCLO) signals (DDCT method).

2.10 Statistical analysis

Data were analysed by ANOVA, Student's t-test, or regression analysis with Fisher's projected least significant difference or Scheffe's F test for comparison of the means (StatView 5.0, SAS Institute, Cary, NC, USA). Data are expressed as the means \pm S.E.M. Values of $P = 0.05$ were considered significant. The number of experiments is indicated in each figure legend. Results of all data for which statistical analysis was performed represent the quantification of at least three duplicate experiments. Control conditions were set as 1. Significance was set at $p < 0.05$ (*), $p < 0.01$ (**), and $p < 0.001$ (***).

3. Results

3.1 Cell-cycle synchronization of H295R and PPNAD cells

We synchronized the cells by treatment with L-mimosine, aphidicolin, or nocodazole, assessed by propidium iodide staining and cytometry analysis (Figure 1). We obtained similar results for both cell types. Asynchronous cells treated with vehicle (DMSO) displayed a typical profile, with a higher percentage of cells in G1 (Figure 1A I,II) . Treatment of the cells with L-mimosine resulted in the accumulation of cells in G1 and decreased the percentage of cells in G2 (Figure 1B I,II). Aphidicolin treatment resulted in the accumulation of cells in late G1, with fewer cells in S and G2 than for the control (Figure 1C I,II) as demonstrated in (Bouizar et al., 2010). Nocodazole treatment resulted in a decrease in the number of cells in G1 and the accumulation of cells in G2 (Figure 1D I,II) as assessed in (Bouizar et al., 2010). All phases of the cell cycle were verified by analyzing the expression of various cyclins to distinguish between the different check points of the cell cycle (Supplemental Figure 1). Cyclin D1 mRNA levels increase substantially in cells blocked with L-mimosine and this increase serves as a marker of initiation of the G1 phase, the level of Cyclin E increases in Aphidicolin-treated cells and serves as a marker of late G1/S, and increases in cyclin B and D levels serve as markers of the G2 Phase (Pereira et al., 2018).

3.2 Steroidogenesis is higher in the G2 phase in synchronized H295R and PPNAD cells

H295R cells secrete both glucocorticoids and mineralocorticoids, whereas PPNAD secretes only glucocorticoids. The analysis of steroid secretion showed significantly greater cortisol secretion by both H295R and PPNAD cells at G2 (Figure 1D III, IV). In H295R cells, aldosterone secretion was also higher at G2 (Figure 1D III). In contrast, cortisol and aldosterone secretion was lower in H295R at G1 ($p < 0.01$), whereas there was no difference for PPNAD (Figure 1A III, IV). Finally, we observed no differences for either cell type at G1/S (Figure 1C III, IV).

We analyzed the impact of cell-cycle synchronization on the expression of genes responsible for steroidogenesis by real-time RT-PCR. Cells synchronized with nocodazole accumulated at G2 and showed a significant increase in mRNA levels for all studied steroidogenic enzyme genes, both in H295R and PPNAD, except for *CYP11A1* in H295R and *CYP17A* in PPNAD (Figure 2C I, II). In contrast, we only observed a significant increase in *CYP17A* ($p < 0.05$) and decrease in *CYP21A2* mRNA levels in H295R in cells synchronized in G1 (Figure 2B I), whereas we observed a decrease in *CYP11A1* and *CYP21A2* mRNA levels in PPNAD cells (Figure 2A II). H295R cells at the early G1/S transition also showed a significant decrease in *CYP11A1* mRNA levels (Figure 2B I). Star protein expression only increased in cells at G2 in H295R (Figure 2 III). The increase in Star protein levels at G2 correlated with an increase in mRNA expression in cultured the H295R and PPNAD cells (Supplemental Figure 2).

We verified that the changes in steroidogenesis during the cell cycle resulted from physiological regulation and were not due to side effects of the drugs by performing similar analyses on H295R cells that were released from synchronization and cultured in normal medium to follow their progression through the cell cycle from 0 to 6, 12, 24, 36, and 48 h after release. The release of cells synchronized at G1 (53.2% G1, 37% S, and 9.7% G2) by L-mimosine treatment, resulted in an increase in the percentage of cells in S at 12 h after release, followed by an increase in cells in G2 at 36 and 48 h (Figure 3A I). The expression of steroidogenic enzyme genes by RT/PCR at various times after release is presented as the fold change with respect to asynchronous cells at time 0, set to one. There was a progressive increase of *CYP11B1*, *CYP11B2*, *CYP11A1*, *NR5A1* (*SF1*), and *STAR* mRNA levels from 6 to 12 h, followed by a peak of expression after 36 h, concomitant with entry into G2 (Figure 3B I and C I).

The release of cells blocked in G2 (11.7% G1, 16.8% S, and 71.5% G2) with nocodazole (48 h of treatment) resulted in their progression through the cell cycle (Figure 3A, right). They then re-entered G2 phase from 36 to 48 h after release (Figure 3A II). There was a progressive decrease in *CYP11B2* mRNA levels from 0 to 24 h, coinciding with the decrease in the percentage of cells in G2, followed by a distinct increase from 36 h that reached the initial level observed at time 0 of the release (Figure 3B II). The expression of *CYP11B1* and *CYP11A1* mRNA levels did not change from 0 to 24 h and then increased (Figure 3B II). *NR5A1* (*SF1*) and *STAR* mRNA levels increased at 36 h (Figure 3C II).

These data confirm that steroidogenic enzyme mRNA levels change during cell-cycle progression.

3.3 PKA and MAP kinase activity in synchronized H295R cells

Cell-cycle progression is often associated with the modulation of PKA and MAP kinase activity. Thus, we next assessed PKA and MAP kinase activity in synchronized cells. ERK1/2 phosphorylation, reflecting MAP kinase pathway activity, was significantly higher in all synchronized groups than in asynchronous cells (Figure 4A-C I and Supplemental Figure 3). MEK phosphorylation was also significantly higher in cells in G2 (Figure 4C I). Moreover, there was significantly less PKA activity in cells in G1 (Figure 4A II), whereas it was significantly higher in cells in G2 (Figure 4C II) than in asynchronous cells.

Under the same conditions, PKA R1A protein levels were markedly lower in cells at G2 (Figure 4C III), whereas those of PKA R2B and PKA CA were not significantly altered in any of the studied groups (Figure 4A-C).

3.4 The effect of PKA activation on the control of the cell cycle and steroidogenesis in synchronized H295R cells

The overexpression of steroidogenic enzyme genes at G2 is associated with an increase in the level of PKA activity. We next investigated whether increased PKA activity effects the phases of the cell cycle and steroidogenesis in synchronized cells. We thus increased PKA activity by overexpressing *PRKACA* (Ca) or silencing *PRKARIA* (Figure 5 I-II). PKA Ca was overexpressed in transfected cells of the asynchronous control and all synchronized groups. However, it was less highly overexpressed in the nocodazole group, for which the basal level was much higher than that of the control asynchronous cells (Supplemental Figure 4A). The expression of the PKA RI protein was reduced in all depleted *PRKARIA* synchronized groups (Supplemental Figure 4 B). However, the magnitude of the increase in Ca protein levels and decrease in RIa protein levels depended on the phase of arrest.

Overexpression of *PRKACA* (Ca) for 24h in H295R cells had no effect on the cell cycle (Figure 5A I) but increased both the activity of PKA and a luciferase reporter construct driven by the *STAR* promoter (*STAR*-Luc) in all groups, independently of the phase of cell-cycle arrest (Figure 5B and C I). *STAR*-Luc activity was the highest in cells in G2 (Figure 5C I). Similarly, PKA activation by cAMP had no effect on the cell cycle (Supplemental Figure 5).

Transient depletion of *PRKARIA* for 24 h was sufficient to increase the percentage/number of cells in G2 in the asynchronous group and decreased the percentage of cells in G1 relative to the control (siS, asynchronous cells; Figure 5A II). In L-mimosine treated cells, the invalidation of *PRKARIA* shifted the cells toward the S and G2 phases and decreased the percentage of cells in G1 ($p \leq 0.02$ siR1A vs siS). Simultaneous transfection with siR1A and aphidicolin treatment for 24 h decreased the percentage/number of cells in G1 ($p < 0.001$ siR1A vs siS) and subsequently increased those in S ($p < 0.05$ siR1A vs siS). Simultaneous nocodazole treatment and R1A silencing increased the percentage of cells in S and decreased the percentage of those in G2 ($p \leq 0.05$) (Figure 5A II). As expected, *PRKARIA* silencing

enhanced PKA activity in all groups (Figure 5B II). Under the same conditions, *PRKARIA* silencing led to increased activity of the *STAR*-Luc reporter in the asynchronous group ($p < 0.05$) and L-mimosine treated cells ($p < 0.05$), whereas its activity remained unchanged in the aphidicolin and nocodazole-treated cells (Figure 5C II).

3.5 Effect of PKA inhibition on control of the cell cycle in synchronized H295R cells

We confirmed the above data with the pharmacological inhibition of PKA activity by treatment with the cell-permeable specific inhibitor Myr-PKI (14-22), which contains a PKA substrate consensus sequence with the serine replaced by alanine (Wiley et al., 1999). The major effect of PKA inhibition was observed in nocodazole treated cells (G2), for which there was a decrease in the percentage of cells in G2 ($p < 0.001$) and an increase in the percentage of those in S ($p < 0.02$) (Figure 6A). This effect was paralleled by a marked and significant decrease in *STAR*-Luc activity in the nocodazole-treated cells ($p < 0.001$) (Figure 6B). Overall, our results show that the PKA R1A subunit mediates the effect of PKA activity on cell-cycle progression and the activation of steroidogenesis, depending on the phase of the cell cycle.

4. Discussion.

Excess adrenal-steroid secretion can be explained by mutations of various key components of the cAMP pathway that are also likely to be involved in adrenal tumorigenesis (Bertherat et al., 2003;Beuschlein et al., 2014;Calebiro et al., 2014;Horvath et al., 2006;Horvath et al., 2008). Here, we demonstrate that steroidogenesis is modulated according to specific cell-cycle phases in both an established human adrenal cortical carcinoma cell line, H295R, and a primary cell culture of PPNAD (harboring *PKARIA* inactivating mutations) from patients with cortisol excess.

Major actors of steroidogenesis (NR5A, Star, and enzymes) were preferentially induced during G2 (Figures 1, 2 and 3), leading to peak steroidogenesis during the same phase.

It is well known that cAMP/PKA and the MAP kinase activity vary during the cell cycle and that MAP kinase activity is higher during the G2 phase in most cells (Philipova et al., 1998). However, in most cells, cAMP levels and PKA activity are low during G2 and mitosis and then become high in G1 (Bolte et al., 2003;Kurokawa et al., 1998;Yu et al., 2008). This variation is cell-type specific, as the elevation of intracellular cyclic AMP (cAMP) is considered to be a potent mitogenic signal for a number of cell types, including Swiss 3T3 cells, thyroid epithelial cells, and somatotroph cells of the anterior pituitary (Withers, 1997).

Here, we demonstrated that only the level of PKA activity varied notably according to the stage of the cell cycle in the adrenocortical cell line H295R. PKA activity was high in G2 (Figure 4C II) and lower in G1 (Figure 4A II). Thus, both PKA and MAP kinase were activated in G2 (Figure 4C, I, II). The change of PKA activity in H295R cells at G2 is a molecular mechanism in which the cAMP/PKA pathway is specifically involved in the control of mitosis, as it phosphorylates several substrates involved in its progression. The in vivo dynamics of PKA activity during mitosis was highlighted by the study of Vandame P et

al. (Vandame et al., 2014) using FRET-based biosensors in single living HeLa cells. They showed that PKA activity is required for the maintenance of chromosome condensation and that it may also be required during mitosis, because it phosphorylates histone H1.4, resulting in the dissociation of this protein from the chromatin during mitosis. Moreover, Aurora A kinase, a key protein for the progression of mitosis, is also a target of PKA. The high PKA activity in G2 corresponded with a decrease in the level of the regulatory subunit R1A of PKA (Figure 4C III). One explanation for the decrease in the level of PKA R1A protein is that the increase in PKA activity at G2 liberates the PKA R1A protein from PKA CA, making it more accessible to degradation. Another possibility is the consequence of the association of PKA R1A with the cytoskeleton or microtubules through either low-affinity or highly dynamic interactions, as shown in BW1J hepatoma cells (Imaizumi-Scherrer et al., 2001).

Mutations of PKA signaling components (*GNAS*, *PRKARIA*, *PRKACA*, and phosphodiesterase genes) in adrenal tumors activate the pathway (Nakajima et al., 2016; Mete et al., 2018; Drougat et al., 2015). Although these mutations are associated with the pathogenesis of benign tumors, we tested whether PKA may regulate the cell cycle and steroidogenesis in a coordinated fashion by activating PKA in the adrenocortical cell line H295R through overexpression of the PKA catalytic subunit alpha, stimulating the cells with Bu2cAMP, or silencing *PRKARIA* in combination with cell synchronization. The responses of the cells to the activation of PKA were different depending on the PKA subunits used. *PRKACA* overexpression stimulated the activity of a PKA-dependent construct (*STAR-Luc*) in all groups but did not alter the phase of the cell cycle (Figure 5 A and C, I). The activation of PKA by wild type *PRKACA* overexpression may not mimic the effect of *PRKACA* mutations, as Bathon et al (Bathon et al., 2019) recently showed that transfection of HEK293 cells with various *PRKACA* mutant constructs for 48 h highly increased PKA activity over that of wild type levels and increased the phosphorylation of the specific PKA substrate histone H1.4,

involved in proliferation, relative to that by wild type *PRKACA*. Our result represents the early effect of PKA activation (24-h transfection of H295R) by wild type *PRKACA*, which indiscriminately activates substrates and not selected ones. Similarly, 24-h activation of PKA by Bu2 cAMP did not alter the cell cycle profile, neither of asynchronous nor synchronized cells. Our results are consistent with those of the study of Liu et al (Liu et al., 2004), in which the effect of Bu2cAMP on proliferation and apoptosis was not observed at an early time point, one to two days of incubation, but later, after three days to one week of incubation.

As expected and described by Basso *et al.* (Basso et al., 2014), inactivation of *PRKARIA* in asynchronous cells for 24 h increased both PKA and *STAR*-Luc reporter activity (Figure 5B and C II) and led to the accumulation of the cells in G2 and a decrease in the number of cells in G1 (Figure 5A II). In synchronized cells, PKA activity increased after *PRKARIA* depletion. The increase was greater in cells treated for G1 arrest than those treated for G1/S and G2 phase arrest. In contrast, the effects of the increase in PKA activity on steroidogenesis were restricted to the cells treated for G1 arrest. In addition, the depletion of *PRKARIA* induced cell-cycle progression in all synchronized groups, as demonstrated by a significant decrease in the percentage/number of cells in G1 reflecting progression to the S and then G2 phases (Figure 5A II).

Moreover, PKA controlled progression of the cells through the cell cycle, as the inhibition of PKA activity by PKI slowed the progression of cells with a high level of PKA activation and those in G2 (Figure 6).

Thus, in our model, the essential and the early determinant of PKA activity as a regulator of cell-cycle progression is the level of PKA R1A. However, our findings on malignant adrenocortical cells that harbor different mutations (Tissier et al., 2005; Ragazzon et al., 2010)

might not apply to normal adrenocortical cells or benign adrenocortical tumors carrying alterations in PKA signaling (Bathon et al., 2019).

PKA R1A can also interact directly with other specific proteins and modulates a number of other signaling pathways. It has been demonstrated that PKA and PKA R1A are important for regulation of the meiotic spindle in mammalian oocytes, in which increased PKA activity is responsible for maintaining meiotic arrest. On the other hand, the knockdown of *PRKAR1A*, as expected, increased PKA activity but the meiotic arrest of mouse oocytes was not maintained (Duncan et al., 2006). PKA R1A is also associated with the second subunit of replication factor C (RFC40) and functions as a nuclear transporter for RFC40, causing G1 arrest of MCF7 cells (Gupte et al., 2005). In the absence of PKA R1A, RFC40 may remain in the cytoplasm and cells progress toward the G2 and M phases.

The role of PKA R1A on the control of cell-cycle progression can be explained by the spatial regulation of PKA activity by A-kinase anchor proteins (AKAPs). AKAPs serve as scaffold proteins to tether PKA in close proximity to its substrate, thereby locally regulating its activity and PKA subunit specificity (Pidoux et al., 2010; Day et al., 2011). Recently, R1A was shown to be associated with AKAPs, which form a complex with the ribosomal S6 protein kinase RSK1 kinase (Chaturvedi et al., 2009; Chaturvedi et al., 2006). *Prkar1a* silencing in mouse lung fibroblasts (B82L) decreased the nuclear accumulation of active RSK1 and increased its cytoplasmic localization. The resulting increase in active cytosolic RSK1 levels is accompanied by increased phosphorylation of its cytosolic substrate tuberous sclerosis complex 2 (TSC-2) and the anti-apoptotic factor BAD. As RSK1 shares common phosphorylation sites with PKA on certain substrates, such as CREB, BAD, and Nur77, part of the cell response in the absence of PKA R1A may not be due to increased PKA activity (Chaturvedi et al., 2009)

412 In conclusion, we have provided evidence of a tight association between the phase of the cell
413 cycle and steroidogenesis in *two* adrenal tumors, *PPNAD* and the malignant *H295R* cell line.
414 Enhancing PKA activity clearly appears to be critical for steroid excess, whereas the PKA
415 R1A subunit is essential for triggering the cell cycle.

416 **Declaration of interest:** there is no conflict of interest that could be perceived to affect the
417 impartiality of the research reported. All authors declare that there is no conflict of interest.

418 **Funding:** this work was supported by the National Research Agency: grants ANR 14 CE12-
419 000-02 and ANR 15 CE 12-0017

420 **Acknowledgements:** We thank the members of the FACS Core Facility CYBIO of the
421 Institute Cochin for their cooperation. The English of the manuscript has been corrected by a
422 native English-speaking scientific editor of Alex Edelman and Associates.

423

References

- [1] Espiard, S., Ragazzon, B. and Bertherat, J., 2014. Protein kinase A alterations in adrenocortical tumors, *Horm Metab Res.* 46, 869-75.
- [2] Louiset, E., Duparc, C., Young, J., Renouf, S., Tetsi Nomigni, M., Boutelet, I., Libe, R., Bram, Z., Groussin, L., Caron, P., Tabarin, A., Grunenberger, F., Christin-Maitre, S., Bertagna, X., Kuhn, J.M., Anouar, Y., Bertherat, J. and Lefebvre, H., 2013. Intraadrenal corticotropin in bilateral macronodular adrenal hyperplasia, *N Engl J Med.* 369, 2115-25.
- [3] Bassett, M.H., Mayhew, B., Rehman, K., White, P.C., Mantero, F., Arnaldi, G., Stewart, P.M., Bujalska, I. and Rainey, W.E., 2005. Expression profiles for steroidogenic enzymes in adrenocortical disease, *J Clin Endocrinol Metab.* 90, 5446-55.
- [4] Nakamura, Y., Yamazaki, Y., Felizola, S.J., Ise, K., Morimoto, R., Satoh, F., Arai, Y. and Sasano, H., 2015. Adrenocortical carcinoma: review of the pathologic features, production of adrenal steroids, and molecular pathogenesis, *Endocrinol Metab Clin North Am.* 44, 399-410.
- [5] Abiven, G., Coste, J., Groussin, L., Anract, P., Tissier, F., Legmann, P., Dousset, B., Bertagna, X. and Bertherat, J., 2006. Clinical and biological features in the prognosis of adrenocortical cancer: poor outcome of cortisol-secreting tumors in a series of 202 consecutive patients, *J Clin Endocrinol Metab.* 91, 2650-5.
- [6] Arlt, W., Biehl, M., Taylor, A.E., Hahner, S., Libe, R., Hughes, B.A., Schneider, P., Smith, D.J., Stiekema, H., Krone, N., Porfiri, E., Opocher, G., Bertherat, J., Mantero, F., Allolio, B., Terzolo, M., Nightingale, P., Shackleton, C.H., Bertagna, X., Fassnacht, M. and Stewart, P.M., 2011. Urine steroid metabolomics as a biomarker tool for detecting malignancy in adrenal tumors, *J Clin Endocrinol Metab.* 96, 3775-84.
- [7] Ragazzon, B., Assie, G. and Bertherat, J., 2011. Transcriptome analysis of adrenocortical cancers: from molecular classification to the identification of new treatments, *Endocr Relat Cancer.* 18, R15-27.
- [8] Stratakis, C.A., 2009. New genes and/or molecular pathways associated with adrenal hyperplasias and related adrenocortical tumors, *Mol Cell Endocrinol.* 300, 152-7.
- [9] Assie, G., Letouze, E., Fassnacht, M., Jouinot, A., Luscap, W., Barreau, O., Omeiri, H., Rodriguez, S., Perlemoine, K., Rene-Corail, F., Elarouci, N., Sbiera, S., Kroiss, M., Allolio, B., Waldmann, J., Quinkler, M., Mannelli, M., Mantero, F., Papathomas, T., De Krijger, R., Tabarin, A., Kerlan, V., Baudin, E., Tissier, F., Dousset, B., Groussin, L., Amar, L., Clauser, E., Bertagna, X., Ragazzon, B., Beuschlein, F., Libe, R., de Reynies, A. and Bertherat, J., 2014. Integrated genomic characterization of adrenocortical carcinoma, *Nat Genet.* 46, 607-12.
- [10] De Martino, M.C., Al Ghuzlan, A., Aubert, S., Assie, G., Scoazec, J.Y., Leboulleux, S., Do Cao, C., Libe, R., Nozieres, C., Lombres, M., Pattou, F., Borson-Chazot, F., Hescot, S., Mazoyer, C., Young, J., Borget, I., Colao, A., Pivonello, R., Soria, J.C., Bertherat, J., Schlumberger, M., Lacroix, L. and Baudin, E., 2013. Molecular screening for a personalized treatment approach in advanced adrenocortical cancer, *J Clin Endocrinol Metab.* 98, 4080-8.
- [11] Ross, J.S., Wang, K., Rand, J.V., Gay, L., Presta, M.J., Sheehan, C.E., Ali, S.M., Elvin, J.A., Labrecque, E., Hiemstra, C., Buell, J., Otto, G.A., Yelensky, R., Lipson, D., Morosini, D., Chmielecki, J., Miller, V.A. and Stephens, P.J., 2014. Next-generation sequencing of adrenocortical carcinoma reveals new routes to targeted therapies, *J Clin Pathol.* 67, 968-73.

- 474 [12] Pereira, S.S., Monteiro, M.P., Bourdeau, I., Lacroix, A. and Pignatelli, D., 2018.
475 MECHANISMS OF ENDOCRINOLOGY: Cell cycle regulation in adrenocortical
476 carcinoma, *Eur J Endocrinol.* 179, R95-R110.
- 477 [13] Kullmann, M.K., Grubbauer, C., Goetsch, K., Jakel, H., Podmirseg, S.R.,
478 Trockenbacher, A., Ploner, C., Cato, A.C., Weiss, C., Kofler, R. and Hengst, L., 2013.
479 The p27-Skp2 axis mediates glucocorticoid-induced cell cycle arrest in T-lymphoma
480 cells, *Cell Cycle.* 12, 2625-35.
- 481 [14] Heisler, L.E., Evangelou, A., Lew, A.M., Trachtenberg, J., Elsholtz, H.P. and Brown,
482 T.J., 1997. Androgen-dependent cell cycle arrest and apoptotic death in PC-3 prostatic
483 cell cultures expressing a full-length human androgen receptor, *Mol Cell Endocrinol.*
484 126, 59-73.
- 485 [15] Amanatullah, D.F., Zafonte, B.T. and Pestell, R.G., 2002. The cell cycle in steroid
486 hormone regulated proliferation and differentiation, *Minerva Endocrinol.* 27, 7-20.
- 487 [16] Cerquetti, L., Bucci, B., Marchese, R., Misiti, S., De Paula, U., Miceli, R., Muleti, A.,
488 Amendola, D., Piergrossi, P., Brunetti, E., Toscano, V. and Stigliano, A., 2008.
489 Mitotane increases the radiotherapy inhibitory effect and induces G2-arrest in
490 combined treatment on both H295R and SW13 adrenocortical cell lines, *Endocr Relat*
491 *Cancer.* 15, 623-34.
- 492 [17] Doghman, M. and Lalli, E., 2013. Lack of long-lasting effects of mitotane adjuvant
493 therapy in a mouse xenograft model of adrenocortical carcinoma, *Mol Cell*
494 *Endocrinol.* 381, 66-9.
- 495 [18] Hescot, S., Slama, A., Lombes, A., Paci, A., Remy, H., Leboulleux, S., Chadarevian,
496 R., Trabado, S., Amazit, L., Young, J., Baudin, E. and Lombes, M., 2013. Mitotane
497 alters mitochondrial respiratory chain activity by inducing cytochrome c oxidase
498 defect in human adrenocortical cells, *Endocr Relat Cancer.* 20, 371-81.
- 499 [19] Lodish, M. and Stratakis, C.A., 2016. A genetic and molecular update on
500 adrenocortical causes of Cushing syndrome, *Nat Rev Endocrinol.* 12, 255-62.
- 501 [20] Kirschner, L.S., Sandrini, F., Monbo, J., Lin, J.P., Carney, J.A. and Stratakis, C.A.,
502 2000. Genetic heterogeneity and spectrum of mutations of the PRKAR1A gene in
503 patients with the carney complex, *Hum Mol Genet.* 9, 3037-46.
- 504 [21] Bertherat, J., Groussin, L., Sandrini, F., Matyakhina, L., Bei, T., Stergiopoulos, S.,
505 Papageorgiou, T., Bourdeau, I., Kirschner, L.S., Vincent-Dejean, C., Perlempine, K.,
506 Gicquel, C., Bertagna, X. and Stratakis, C.A., 2003. Molecular and functional analysis
507 of PRKAR1A and its locus (17q22-24) in sporadic adrenocortical tumors: 17q losses,
508 somatic mutations, and protein kinase A expression and activity, *Cancer Res.* 63,
509 5308-19.
- 510 [22] Zheng, S., Cherniack, A.D., Dewal, N., Moffitt, R.A., Danilova, L., Murray, B.A.,
511 Lerario, A.M., Else, T., Knijnenburg, T.A., Ciriello, G., Kim, S., Assie, G., Morozova,
512 O., Akbani, R., Shih, J., Hoadley, K.A., Choueiri, T.K., Waldmann, J., Mete, O.,
513 Robertson, A.G., Wu, H.T., Raphael, B.J., Shao, L., Meyerson, M., Demeure, M.J.,
514 Beuschlein, F., Gill, A.J., Sidhu, S.B., Almeida, M.Q., Fragoso, M., Cope, L.M.,
515 Kebebew, E., Habra, M.A., Whitsett, T.G., Bussey, K.J., Rainey, W.E., Asa, S.L.,
516 Bertherat, J., Fassnacht, M., Wheeler, D.A., Hammer, G.D., Giordano, T.J. and
517 Verhaak, R.G.W., 2016. Comprehensive Pan-Genomic Characterization of
518 Adrenocortical Carcinoma, *Cancer Cell.* 29, 723-736.
- 519 [23] Beuschlein, F., Fassnacht, M., Assie, G., Calebiro, D., Stratakis, C.A., Osswald, A.,
520 Ronchi, C.L., Wieland, T., Sbiera, S., Faucz, F.R., Schaak, K., Schmittfull, A.,
521 Schwarzmayer, T., Barreau, O., Vezzosi, D., Rizk-Rabin, M., Zabel, U., Szarek, E.,
522 Salpea, P., Forlino, A., Vetro, A., Zuffardi, O., Kisker, C., Diener, S., Meitinger, T.,
523 Lohse, M.J., Reincke, M., Bertherat, J., Strom, T.M. and Allolio, B., 2014.

- Constitutive activation of PKA catalytic subunit in adrenal Cushing's syndrome, *N Engl J Med.* 370, 1019-28.
- [24] Goh, G., Scholl, U.I., Healy, J.M., Choi, M., Prasad, M.L., Nelson-Williams, C., Kunstman, J.W., Korah, R., Suttorp, A.C., Dietrich, D., Haase, M., Willenberg, H.S., Stalberg, P., Hellman, P., Akerstrom, G., Bjorklund, P., Carling, T. and Lifton, R.P., 2014. Recurrent activating mutation in PRKACA in cortisol-producing adrenal tumors, *Nat Genet.* 46, 613-7.
- [25] Calebiro, D., Hannawacker, A., Lyga, S., Bathon, K., Zabel, U., Ronchi, C., Beuschlein, F., Reincke, M., Lorenz, K., Allolio, B., Kisker, C., Fassnacht, M. and Lohse, M.J., 2014. PKA catalytic subunit mutations in adrenocortical Cushing's adenoma impair association with the regulatory subunit, *Nat Commun.* 5, 5680.
- [26] Cao, Y., He, M., Gao, Z., Peng, Y., Li, Y., Li, L., Zhou, W., Li, X., Zhong, X., Lei, Y., Su, T., Wang, H., Jiang, Y., Yang, L., Wei, W., Yang, X., Jiang, X., Liu, L., He, J., Ye, J., Wei, Q., Li, Y., Wang, W., Wang, J. and Ning, G., 2014. Activating hotspot L205R mutation in PRKACA and adrenal Cushing's syndrome, *Science.* 344, 913-7.
- [27] Sato, Y., Maekawa, S., Ishii, R., Sanada, M., Morikawa, T., Shiraishi, Y., Yoshida, K., Nagata, Y., Sato-Otsubo, A., Yoshizato, T., Suzuki, H., Shiozawa, Y., Kataoka, K., Kon, A., Aoki, K., Chiba, K., Tanaka, H., Kume, H., Miyano, S., Fukayama, M., Nureki, O., Homma, Y. and Ogawa, S., 2014. Recurrent somatic mutations underlie corticotropin-independent Cushing's syndrome, *Science.* 344, 917-20.
- [28] Massimi, M., Cardarelli, S., Galli, F., Giardi, M.F., Ragusa, F., Panera, N., Cinque, B., Cifone, M.G., Biagioni, S. and Giorgi, M., 2017. Increase of Intracellular Cyclic AMP by PDE4 Inhibitors Affects HepG2 Cell Cycle Progression and Survival, *J Cell Biochem.* 118, 1401-1411.
- [29] Gupta, A., Tiwari, M., Prasad, S. and Chaube, S.K., 2017. Role of Cyclic Nucleotide Phosphodiesterases During Meiotic Resumption From Diplotene Arrest in Mammalian Oocytes, *J Cell Biochem.* 118, 446-452.
- [30] Sun, Q.Y., Miao, Y.L. and Schatten, H., 2009. Towards a new understanding on the regulation of mammalian oocyte meiosis resumption, *Cell Cycle.* 8, 2741-7.
- [31] Duncan, F.E., Moss, S.B. and Williams, C.J., 2006. Knockdown of the cAMP-dependent protein kinase (PKA) Type Ialpha regulatory subunit in mouse oocytes disrupts meiotic arrest and results in meiotic spindle defects, *Dev Dyn.* 235, 2961-8.
- [32] Hewer, R.C., Sala-Newby, G.B., Wu, Y.J., Newby, A.C. and Bond, M., 2011. PKA and Epac synergistically inhibit smooth muscle cell proliferation, *J Mol Cell Cardiol.* 50, 87-98.
- [33] Medri, G., Sergi, I., Papandreou, M.J., Beck-Peccoz, P., Verrier, B. and Ronin, C., 1994. Dual activity of human pituitary thyrotrophin isoforms on thyroid cell growth, *J Mol Endocrinol.* 13, 187-98.
- [34] Spada, A., Vallar, L. and Faglia, G., 1992. G protein oncogenes in pituitary tumors, *Trends Endocrinol Metab.* 3, 355-60.
- [35] Taylor, S.S., Zhang, P., Steichen, J.M., Keshwani, M.M. and Kornev, A.P., 2013. PKA: lessons learned after twenty years, *Biochim Biophys Acta.* 1834, 1271-8.
- [36] Kirschner, L.S., Kusewitt, D.F., Matyakhina, L., Towns, W.H., 2nd, Carney, J.A., Westphal, H. and Stratakis, C.A., 2005. A mouse model for the Carney complex tumor syndrome develops neoplasia in cyclic AMP-responsive tissues, *Cancer Res.* 65, 4506-14.
- [37] Nadella, K.S. and Kirschner, L.S., 2005. Disruption of protein kinase a regulation causes immortalization and dysregulation of D-type cyclins, *Cancer Res.* 65, 10307-15.

- [38] Basso, F., Rocchetti, F., Rodriguez, S., Nesterova, M., Cormier, F., Stratakis, C.A., Ragazzon, B., Bertherat, J. and Rizk-Rabin, M., 2014. Comparison of the effects of PRKAR1A and PRKAR2B depletion on signaling pathways, cell growth, and cell cycle control of adrenocortical cells, *Horm Metab Res.* 46, 883-8.
- [39] Gazdar, A.F., Oie, H.K., Shackleton, C.H., Chen, T.R., Triche, T.J., Myers, C.E., Chrousos, G.P., Brennan, M.F., Stein, C.A. and La Rocca, R.V., 1990. Establishment and characterization of a human adrenocortical carcinoma cell line that expresses multiple pathways of steroid biosynthesis, *Cancer Res.* 50, 5488-96.
- [40] Weigand, I., Ronchi, C.L., Rizk-Rabin, M., Dalmazi, G.D., Wild, V., Bathon, K., Rubin, B., Calebiro, D., Beuschlein, F., Bertherat, J., Fassnacht, M. and Sbiera, S., 2017. Differential expression of the protein kinase A subunits in normal adrenal glands and adrenocortical adenomas, *Sci Rep.* 7, 49.
- [41] Bouizar, Z., Ragazzon, B., Viou, L., Hortane, M., Bertherat, J. and Rizk-Rabin, M., 2010. 8Cl-cAMP modifies the balance between PKAR1 and PKAR2 and modulates the cell cycle, growth and apoptosis in human adrenocortical H295R cells, *J Mol Endocrinol.* 44, 331-47.
- [42] Ragazzon, B., Cazabat, L., Rizk-Rabin, M., Assie, G., Groussin, L., Fierrard, H., Perlemoine, K., Martinez, A. and Bertherat, J., 2009. Inactivation of the Carney complex gene 1 (protein kinase A regulatory subunit 1A) inhibits SMAD3 expression and TGF beta-stimulated apoptosis in adrenocortical cells, *Cancer Res.* 69, 7278-84.
- [43] Groussin, L., Massias, J.F., Bertagna, X. and Bertherat, J., 2000. Loss of expression of the ubiquitous transcription factor cAMP response element-binding protein (CREB) and compensatory overexpression of the activator CREMtau in the human adrenocortical cancer cell line H295R, *J Clin Endocrinol Metab.* 85, 345-54.
- [44] Cazabat, L., Ragazzon, B., Varin, A., Potier-Cartereau, M., Vandier, C., Vezzosi, D., Risk-Rabin, M., Guellich, A., Schittl, J., Lechene, P., Richter, W., Nikolaev, V.O., Zhang, J., Bertherat, J. and Vandecasteele, G., 2014. Inactivation of the Carney complex gene 1 (PRKAR1A) alters spatiotemporal regulation of cAMP and cAMP-dependent protein kinase: a study using genetically encoded FRET-based reporters, *Hum Mol Genet.* 23, 1163-74.
- [45] Bram, Z., Louiset, E., Ragazzon, B., Renouf, S., Wils, J., Duparc, C., Boutelet, I., Rizk-Rabin, M., Libe, R., Young, J., Carson, D., Vantyghem, M.C., Szarek, E., Martinez, A., Stratakis, C.A., Bertherat, J. and Lefebvre, H., 2016. PKA regulatory subunit 1A inactivating mutation induces serotonin signaling in primary pigmented nodular adrenal disease, *JCI Insight.* 1, e87958.
- [46] Rizk-Rabin, M., Assie, G., Rene-Corail, F., Perlemoine, K., Hamzaoui, H., Tissier, F., Lieberherr, M., Bertagna, X., Bertherat, J. and Bouizar, Z., 2008. Differential expression of parathyroid hormone-related protein in adrenocortical tumors: autocrine/paracrine effects on the growth and signaling pathways in H295R cells, *Cancer Epidemiol Biomarkers Prev.* 17, 2275-85.
- [47] Vackova, I., Engelova, M., Marinov, I. and Tomanek, M., 2003. Cell cycle synchronization of porcine granulosa cells in G1 stage with mimosine, *Anim Reprod Sci.* 77, 235-45.
- [48] Sorscher, D.H. and Cordeiro-Stone, M., 1991. Gene replication in the presence of aphidicolin, *Biochemistry.* 30, 1086-90.
- [49] Ho, Y.S., Duh, J.S., Jeng, J.H., Wang, Y.J., Liang, Y.C., Lin, C.H., Tseng, C.J., Yu, C.F., Chen, R.J. and Lin, J.K., 2001. Griseofulvin potentiates antitumorigenesis effects of nocodazole through induction of apoptosis and G2/M cell cycle arrest in human colorectal cancer cells, *Int J Cancer.* 91, 393-401.

- [50] Wiley, J.C., Wailes, L.A., Idzerda, R.L. and McKnight, G.S., 1999. Role of regulatory subunits and protein kinase inhibitor (PKI) in determining nuclear localization and activity of the catalytic subunit of protein kinase A, *J Biol Chem.* 274, 6381-7.
- [51] Horvath, A., Boikos, S., Giatzakis, C., Robinson-White, A., Groussin, L., Griffin, K.J., Stein, E., Levine, E., Delimpasi, G., Hsiao, H.P., Keil, M., Heyerdahl, S., Matyakhina, L., Libe, R., Fratticci, A., Kirschner, L.S., Cramer, K., Gaillard, R.C., Bertagna, X., Carney, J.A., Bertherat, J., Bossis, I. and Stratakis, C.A., 2006. A genome-wide scan identifies mutations in the gene encoding phosphodiesterase 11A4 (PDE11A) in individuals with adrenocortical hyperplasia, *Nat Genet.* 38, 794-800.
- [52] Horvath, A., Giatzakis, C., Tsang, K., Greene, E., Osorio, P., Boikos, S., Libe, R., Patronas, Y., Robinson-White, A., Remmers, E., Bertherat, J., Nesterova, M. and Stratakis, C.A., 2008. A cAMP-specific phosphodiesterase (PDE8B) that is mutated in adrenal hyperplasia is expressed widely in human and mouse tissues: a novel PDE8B isoform in human adrenal cortex, *Eur J Hum Genet.* 16, 1245-53.
- [53] Philipova, R. and Whitaker, M., 1998. MAP kinase activity increases during mitosis in early sea urchin embryos, *J Cell Sci.* 111 (Pt 17), 2497-505.
- [54] Bolte, M., Dieckhoff, P., Krause, C., Braus, G.H. and Irniger, S., 2003. Synergistic inhibition of APC/C by glucose and activated Ras proteins can be mediated by each of the Tpk1-3 proteins in *Saccharomyces cerevisiae*, *Microbiology.* 149, 1205-16.
- [55] Kurokawa, K. and Kato, J., 1998. Cyclic AMP delays G2 progression and prevents efficient accumulation of cyclin B1 proteins in mouse macrophage cells, *Cell Struct Funct.* 23, 357-65.
- [56] Yu, A., Zhang, Z., Bi, Q., Sun, B., Su, W., Guan, Y., Mu, R., Miao, C., Zhang, J. and Yu, B., 2008. Regulation of cAMP on the first mitotic cell cycle of mouse embryos, *Mol Reprod Dev.* 75, 489-95.
- [57] Withers, D.J., 1997. Signalling pathways involved in the mitogenic effects of cAMP, *Clin Sci (Lond).* 92, 445-51.
- [58] Vandame, P., Spriet, C., Trinel, D., Gelaude, A., Caillau, K., Bompard, C., Biondi, E. and Bodart, J.F., 2014. The spatio-temporal dynamics of PKA activity profile during mitosis and its correlation to chromosome segregation, *Cell Cycle.* 13, 3232-40.
- [59] Imaizumi-Scherrer, T., Faust, D.M., Barradeau, S., Hellio, R. and Weiss, M.C., 2001. Type I protein kinase a is localized to interphase microtubules and strongly associated with the mitotic spindle, *Exp Cell Res.* 264, 250-65.
- [60] Nakajima, Y., Okamura, T., Horiguchi, K., Gohko, T., Miyamoto, T., Satoh, T., Ozawa, A., Ishii, S., Yamada, E., Hashimoto, K., Okada, S., Takata, D., Horiguchi, J. and Yamada, M., 2016. GNAS mutations in adrenal aldosterone-producing adenomas, *Endocr J.* 63, 199-204.
- [61] Mete, O. and Duan, K., 2018. The Many Faces of Primary Aldosteronism and Cushing Syndrome: A Reflection of Adrenocortical Tumor Heterogeneity, *Front Med (Lausanne).* 5, 54.
- [62] Drougat, L., Omeiri, H., Lefevre, L. and Ragazzon, B., 2015. Novel Insights into the Genetics and Pathophysiology of Adrenocortical Tumors, *Front Endocrinol (Lausanne).* 6, 96.
- [63] Bathon, K., Weigand, I., Vanselow, J.T., Ronchi, C.L., Sbiera, S., Schlosser, A., Fassnacht, M. and Calebiro, D., 2019. Alterations in Protein Kinase A Substrate Specificity as a Potential Cause of Cushing Syndrome, *Endocrinology.* 160, 447-459.
- [64] Liu, J., Li, X.D., Ora, A., Heikkila, P., Vaheri, A. and Voutilainen, R., 2004. cAMP-dependent protein kinase activation inhibits proliferation and enhances apoptotic effect of tumor necrosis factor-alpha in NCI-H295R adrenocortical cells, *J Mol Endocrinol.* 33, 511-22.

- [65] Tissier, F., Cavard, C., Groussin, L., Perlemoine, K., Fumey, G., Hagnere, A.M., Rene-Corail, F., Jullian, E., Gicquel, C., Bertagna, X., Vacher-Lavenu, M.C., Perret, C. and Bertherat, J., 2005. Mutations of beta-catenin in adrenocortical tumors: activation of the Wnt signaling pathway is a frequent event in both benign and malignant adrenocortical tumors, *Cancer Res.* 65, 7622-7.
- [66] Ragazzon, B., Libe, R., Gaujoux, S., Assie, G., Fratticci, A., Launay, P., Clauser, E., Bertagna, X., Tissier, F., de Reynies, A. and Bertherat, J., 2010. Transcriptome analysis reveals that p53 and {beta}-catenin alterations occur in a group of aggressive adrenocortical cancers, *Cancer Res.* 70, 8276-81.
- [67] Gupte, R.S., Pozarowski, P., Grabarek, J., Traganos, F., Darzynkiewicz, Z. and Lee, M.Y., 2005. RIalpha influences cellular proliferation in cancer cells by transporting RFC40 into the nucleus, *Cancer Biol Ther.* 4, 429-37.
- [68] Pidoux, G. and Tasken, K., 2010. Specificity and spatial dynamics of protein kinase A signaling organized by A-kinase-anchoring proteins, *J Mol Endocrinol.* 44, 271-84.
- [69] Day, M.E., Gaietta, G.M., Sastri, M., Koller, A., Mackey, M.R., Scott, J.D., Perkins, G.A., Ellisman, M.H. and Taylor, S.S., 2011. Isoform-specific targeting of PKA to multivesicular bodies, *J Cell Biol.* 193, 347-63.
- [70] Chaturvedi, D., Cohen, M.S., Taunton, J. and Patel, T.B., 2009. The PKARIalpha subunit of protein kinase A modulates the activation of p90RSK1 and its function, *J Biol Chem.* 284, 23670-81.
- [71] Chaturvedi, D., Poppleton, H.M., Stringfield, T., Barbier, A. and Patel, T.B., 2006. Subcellular localization and biological actions of activated RSK1 are determined by its interactions with subunits of cyclic AMP-dependent protein kinase, *Mol Cell Biol.* 26, 4586-600.

Figure legends

Figure 1. The effect of cell-cycle synchronization of the H295R cell line and primary PPNAD cell culture for 24 h on the cell-cycle profile and steroid secretion.

Flowcytometry analysis of DNA content by propidium iodide incorporation in synchronized H295R cells and PPNAD primary culture for **A)** control cultures without treatment and cultures treated with **B)** L-mimosine, **C)** aphidicolin, or **D)** nocodazole. Histograms of cortisol and aldosterone content secreted into the cell medium after cell-cycle synchronization are shown to the right. The results are presented as the means \pm SEM of three duplicate independent experiments for H295R and one representative example of two duplicate experiments for PPNAD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 2. The effect of cell synchronization on steroidogenic enzyme and Star protein expression, mRNA expression by RT/PCR of steroidogenic enzymes in H295R cells and PPNAD primary cultures in the **A)** G1, **B)** G1/S, and **C)** G2 phases of the cell cycle. Each studied steroidogenic enzyme is normalized to the corresponding results in the control asynchronous cells, which is set to one. For Star protein expression, the results are normalized against those of control cells and a histogram of the western blot in synchronized H295R cells and PPNAD primary cultures are shown on the right. The results are presented as the means \pm SEM of three duplicate independent experiments for H295R and one representative example of two duplicate experiments for PPNAD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 3. Cell-cycle profile and the kinetics of steroidogenic enzyme expression after cell-cycle arrest and release of H295R cells. 0 h represents the profile at the end of synchronization after 24 h of treatment with L-mimosine (I) or 48 h with nocodazole (II). Each time point is normalized to asynchronous control cells at 0 h of release. **A)** Histogram showing the percentage of cells in each phase of the cell cycle at each time point, analyzed by

flow cytometry, showing the progression in the cell cycle. RT/PCR analysis of **B)** steroidogenic enzyme and **C)** SF-1 and Star mRNA levels for each time point. The results are presented as the means \pm SEM of three duplicate independent experiments. The level of significance is indicated at the top of each figure.

Figure 4. Effect of the different drugs on cell signaling pathways and PKA subunit regulation in H295R cells. The effects are shown for **A)** L-mimosine, **B)** aphidicolin, and **C)** nocodazole. MAP Kinase activity is shown by histograms of western blots of pERK/total ERK, and pMEK/total MEK protein levels normalized to the control, set to one. We spliced the original full gel images of ERK and MEK, presented in Supplementary Figure 3, to re-order the samples in a more logical manner. PKA activity was measured by a nonradioactive PKA assay and is presented as the ratio of free PKA/total PKA. Protein levels of the PKA subunits R1A, R2B, and Ca in cells treated with the different drugs are shown by histograms of western blots. The results of the histograms are expressed as the percentage relative band density/the band density of GaPDH, normalized to the corresponding controls (arbitrarily set to one) in H295R-treated cells. The results are presented as the means \pm SEM of three duplicate independent experiments. * $p < 0.05$, ** $p < 0.01$.

Figure 5. Effect of PKA activation, by the over expression of *PRKACA* or *PRKARIA* depletion, on the regulation of the cell cycle, PKA activity, and steroidogenic response (Star/luc-reporter gene activity) simultaneously with the synchronization of H295R cells. **A)** Cell-cycle distribution analyzed by flow cytometry. **B)** PKA activity measured by a non-radioactive assay and presented as the ratio of free phosphorylated PKA/total PKA. **C)** *STAR* reporter gene activity measured by Luciferase/Renilla. The results are presented as the means \pm SEM of three duplicate independent experiments normalized to the control of each group,

set to one. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ for the over expression of PKA CA.

Figure 6. Inhibition of PKA activity by the PKA inhibitor PKI (5 μ M) throughout cell cycle synchronization (24 h) showing that mostly the nocodazole-treated cells were affected. **A)** Cell-cycle phase distribution. **B)** Star/luc reporter gene activity. The measurement of the PKA activity is not presented because PKA activity was inhibited and thus no PKA phosphorylation. The results are presented as the means \pm SEM of three duplicate independent experiments. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Supplementary Figures:

Supplementary Figure 1. Distribution of cyclin mRNA expression in H295R cells arrested at various phases of the cell cycle. **A)** Arrested in early G1 by L-mimosine treatment. **B)** Arrested in late G1/S by aphidicolin treatment. **C)** Arrested in G2 by nocodazole treatment. The results are presented as the means \pm SEM of three duplicate independent experiments normalized to the control of each group, set to one. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.


Supplementary Figure 2. Star mRNA expression by RT/PCR in H295R cells and PPNAD primary cultures in the G1, G1/S, and G2 phases of the cell cycle. The results are presented as the means \pm SEM of three duplicate independent experiments for H295R and one representative example of two duplicate experiments for PPNAD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Supplementary Figure 3. Image of the full western blot of MAPKinase cell signaling used for Figure 4. **A)** Image of the full western blot. **B)** The corresponding Histogram, as in Figure 4.

780 Supplementary Figure 4. PKA CA and PKA R1A protein expression in simultaneous A)
781 overexpression of wildtype *PRKACA* or B) depleted *PRKARIA* and cell-cycle
782 synchronization. Results are normalized to the control vector or sis, set to one. The
783 comparison is inter group (Vector vs Ca or SiS vs siRI) for each group. The results are
784 presented as the means \pm SEM of three duplicate independent experiments normalized to the
785 control of each group, set to one. *p < 0.05, **p < 0.01, ***p < 0.001.

786 Supplementary Figure 5. Effects of cAMP stimulation on cell-cycle synchronization. The
787 distribution of the phases resulting from cell-cycle arrest were not affected by stimulation
788 with the cAMP analogue (dibutyryl adenosine 3'-5' cyclic monophosphate). The
789 measurement of PKA activity and Star Luc reporter gene activity are not presented because
790 they are equally fully saturated in all of the synchronized groups.

Cell cycle synchronization



Steroids secretion




Figure 1






Figure 2

Release from G1 L -mimosine synchronization



Release from G2 nocodazole synchronization

A



Steroidogenic enzymes

B



Star and NR5A1 (SF-1)

C





Figure 3

MAPKinase Cell signaling

PKA activity

PKA subunits regulation




Figure 4




Figure 5






Figure 6







Supplementary Figure 5




Supplementary Figure 1

MAPKinase Cell signaling: full western blot image





Expression of Star mRNA in synchronized cells



Supplementary Figure 2

Link between steroidogenesis, the cell cycle, and PKA in adrenocortical tumor cells

Marthe Rizk-Rabin¹, Sabrina Chaoui- Ibadioune¹, Anna Vaczlavik¹, Christopher Ribes¹, Michel Polak^{1,3}, Bruno Ragazzon¹, and Jérôme Bertherat^{1,2}.

Declaration of interest: there is no conflict of interest that could be perceived to affect the Impartiality of the research reported. All authors declare that there is no conflict of interest.

Highlights

Modulation of steroidogenesis by specific cell cycle phases in adrenal cell tumors.

PKA activity varied notably according to the stage of cell cycle in the H295R cells.

Early effect of enhancing PKA activity appears to be critical for steroid excess.

PRKARIA mediated the function of PKA activity on both steroidogenesis and cell cycle.