

Chiral catalysts derived from biomass: design, synthesis and applications in asymmetric catalysis

Thi Thuy Duong Ngo, Khanh Duy Huynh,
Houssein Ibrahim, Thi Huong Nguyen, Chloée
Bournaud, Martial Toffano and Giang Vo-Thanh*

*Institut de Chimie Moléculaire et des Matériaux d'Orsay,
ICMMO. UMR 8182. Laboratoire de Catalyse
Moléculaire. Université Paris-Sud, Université Paris
Saclay, 91405 Orsay Cedex, France.*

Abstract: The development of new and more efficient catalytic systems is the subject of increasing attention from both academic and industrial research. Moreover, biomass is an endless supply of hydrocarbon materials that can be used as renewable raw materials for the development of new organic compounds. In this context, many research groups have devoted their works to products with a natural chirality source, non-toxic, biodegradable and usually cost effective for the development of new chiral catalysts.

Some new chiral molecules derived from biomass such as carbohydrates and natural amino acids have been synthesized and used as chiral ligands or organocatalysts for asymmetric transformations leading to the formation of expected products in good yields with high enantioselectivities

Key-words: Asymmetric transfer hydrogenation, asymmetric allylic substitution, [4+2] annulation reaction, isosorbide, isomannide, L-proline.

Introduction

Nowadays, science and technology are growing strongly and serve better for human life. In the midst of scientific disciplines, chemistry contributes a considerable part in the development of human. However, besides the great achievements, chemists have to face with a lot of new challenges, health issues, food safety, environment... Hence, sustainable development is really a target to be achieved for the chemists today.

In 1998, Paul Anastas and John C. Warner published a set of 12 principles to guide the practice of green chemistry. The twelve principles give a range of ways to reduce the environmental and health impacts of chemical production, and also indicate research priorities for the development of green chemistry technologies.^[1] Remarkable among the principles is the catalysis. Indeed, catalysts could reduce the energy consumption which presents an economic and environmental interest, increase yields by limiting the formation of by-products or could improve selectivity of reaction.

Ongoing use of petroleum for chemical feedstocks and energy is not sustainable. It is imperative that new catalysts for existing processes and new products are developed based on renewable sources.

The development of available, inexpensive, modular and innovative ligands and catalysts from biomass products such as carbohydrates and natural amino acids is expected to be one of the key procedures for expanding the reaction scope and the synthetic potential of metal-catalyzed enantioselective catalysis. Besides metal complexes and biocatalysis, small organic molecules may promote chemical transformations. Organocatalysis could accelerate the reactions with a sub-stoichiometric amount of organic molecules, which do not contain any metal element. Advantages of organocatalysts include their lack of sensitivity to moisture and oxygen, their ready availability, low cost, and low toxicity, which confers a huge direct benefit in the production of pharmaceutical intermediates when compared with (transition) metal catalysts. Organocatalysis is a 'metal-free method' and therefore the metal pollution can be prevented. Due to all these features, organocatalysis appears as a very attractive area in modern organic synthesis.

The conversion of biomass and its derivatives to chemicals has been the subject of intense research efforts during the past decade.^[2] Many chemical transformations on biomass such as carbohydrates,^[3] glycerol,^[4] triglycerides,^[5] cellulose,^[6] lignin^[7], etc into high-added value products have been reported so far. However, the design and synthesis of novel chiral catalysts starting from biomass still presents a challenge and an opportunity to researchers.

This mini-review focuses on our works dealing with the synthesis of chiral ligands and organocatalysts derived from carbohydrates and natural amino acids and their applications in asymmetric catalysis.

Design and synthesis of novel chiral ligands derived from carbohydrates. Applications to asymmetric metal catalysis

We report herein the synthesis of a new class of chiral β -amino alcohol, diamine and monophosphine ligands starting from isosorbide and isomannide as chiral renewable sources. The efficiency of these ligands has been evaluated for the ruthenium-catalyzed enantioselective reduction of aromatic ketones by transfer hydrogenation as well as for the rhodium-catalyzed enantioselective hydrogenation of olefins.

Isosorbide **1** and its diastereoisomer, isomannide **2**, also known as (3*R*,3*aR*,6*S*,6*aR*)-hexahydrofuro[3,2-*b*]furan-3,6-diol and (3*R*,3*aR*,6*R*,6*aR*)-hexahydrofuro[3,2-*b*]furan-3,6-diol, are renewable, and commercially available chiral carbohydrates. Isosorbide is basically giving a wedge-shaped molecule with two fused tetrahydrofuran rings having the *cis*-arrangement at the ring junction.^[8] The compound possesses two hydroxyl groups, one at C6 having the *exo*-orientation with respect to the wedge-shaped molecule, and the other at C3 having the *endo*-orientation, which

makes possible the intramolecular hydrogen bonding with the oxygen atom of the neighboring tetrahydrofuran ring. In the case of isomannide, the two hydroxyl groups at C6 and C3 have both the *endo*-orientation (Figure 1).





Figure 1: Structure of isosorbide 1 and isomannide 2



Scheme 1: Synthesis of amino alcohols 5a-h.

The complete inversion of configuration in the substitution reaction of 4 with an amine was confirmed by ¹H NMR analysis based on the observation of the H_a, H_b coupling constant of products 5a-h. For compounds 4, the proton H_a appears as doublet corresponding to *trans*-coupling $J_{H_a-H_b} = 0$ Hz. The signal turns to a doublet of doublet for compounds 5a-h corresponding to a *cis*-coupling $J_{H_a-H_b} = 5$ Hz (Figure).





Figure 2: ¹H NMR studies for 3 and 4a-h

Using the similar reaction conditions previously described for the synthesis of amino alcohol ligands 5, sulfonate 4 afforded diamine 6 and *N*-tosylated amine 7 with yields of 82% and 60%, respectively (Scheme 2).



Scheme 2: Synthesis of diamine 6 and *N*-tosylated amine 7.

Asymmetric reduction of aromatic ketones by transfer hydrogenation reaction.

The reduction of carbonyl compounds to the corresponding alcohols is one of the most fundamental and useful reactions that are important in the pharmaceutical and chemical industry.

Isosorbide and isomannide, industrially obtained by dehydration of D-sorbitol and D-mannitol, represent commercially available and low cost chiral starting materials for the synthesis of sophisticated molecules including chiral ionic liquids,^[9] phase-transfer catalysts^[10] and ligands (amino alcohols, amines, diphosphines, diphosphites, bis diaminophosphites, diamidophosphites).^[11,12]


Synthesis of chiral amino alcohols and diamines

Starting from isosorbide 1, a series of eight amino alcohols 5 was synthesized and isolated in moderate to good overall yields using classical organic transformations (Scheme 1).

The asymmetric transfer hydrogenation (ATH) of ketones has recently emerged as a highly efficient method for the synthesis of enantiomerically enriched secondary alcohols that are key intermediates for the manufacture of a variety of molecules and molecular scaffolds of biological and therapeutic interest.^[13] Transition metal-catalyzed asymmetric transfer hydrogenation of prochiral ketones provides a powerful alternative to asymmetric hydrogenation owing to its ease of handling, the easy availability of hydrogen sources, lower cost and safety.^[14] Many chiral ligands and transition metals have been developed and applied to this reaction. The common catalysts are rhodium, ruthenium and iridium complexes with chiral phosphine and chiral diamine ligands.^[15]

ATH reaction offers an interesting and common tool to evaluate the activity of these new chiral ligands. At this point of our investigation, we decided to study the reduction of acetophenone by ATH as a model reaction to explore the catalytic behavior of our complexes.

After optimization of reaction conditions by screening many parameters including hydride sources, catalyst precursors, base nature, base/ligand/metal ratio, solvent, reaction temperature, reaction time, substrate concentration, and structure of ligands, the reduction of aromatic ketones by ATH reaction with ligand 5f led to the formation of expected product in excellent yields and with good enantioselectivities up to 91% ee as shown in Scheme 3.^[16]




Scheme 3: Asymmetric transfer hydrogenation of aromatic ketones with ligand **5f**.

These results clearly indicated the efficiency of the isosorbide structure for the preparation of chiral ligands and encouraged us to evaluate our amino alcohol ligands **5a-h** in the addition of diethylzinc to aldehydes. This reaction has been extensively studied and offers an excellent tool for evaluating of new chiral ligands.^[17]

Asymmetric addition reaction of diethylzinc to aldehydes with amino alcohol ligands.

Based on these chiral amino alcohols **5a-h** as well as the ATH reaction of aromatic ketones, we decided to study the zinc-catalyzed ethylation of aliphatic and aromatic aldehydes. In the same way, after examining many reaction parameters, the addition of diethylzinc to a selection of aldehydes bearing electron-withdrawing or -donating groups, under optimized reaction conditions, afforded the corresponding products in very good yield and with high enantioselectivity.^[18]




Scheme 4: Asymmetric addition reaction of diethylzinc to aldehyde with ligand **5f**.

In all cases, amino alcohol **5f** was found to be the best ligand in all two studied reactions. The presence of phenyl group at the α position of the hydroxyl function was crucial for improving the enantioselectivity. The use of diamine ligands **6** or **7** led to the formation of the expected product in good yield but with low enantioselective excess.

Synthesis of chiral monophosphine ligands derived from isomannide and isosorbide.

Starting from isomannide **2**, two monophosphine-borane complexes **10a** and **10b** were respectively synthesized from bromide derivatives **9a** and **9b** by nucleophilic substitution with diphenylphosphine anion, followed by a protection with borane dimethylsulfide complex (Scheme 5). It should be noted that the substitution of sulfonate **4** with the same diphenylphosphine anion did not work at all, and produced only the alcohol **3** (see Scheme 1).



Scheme 5: Synthesis of chiral monophosphine-borane **10a** and **10b**.

Surprisingly, X-ray single-crystal diffraction analysis of **10b** and **10a** confirms an *exo* position of phosphine group due to a total retention of configuration during the substitution step (Figure 3).





Figure 3: ORTEP drawing of **10b**. Ellipsoids are drawn at the 50 % probability level


On the other hand, hydrophosphination of the olefin **11**, easily obtained in some steps from isosorbide **1**, conducted to monophosphine-borane **10b** in 47% yield using microwave heating (Scheme 6). Almost no conversion was observed when performing hydrophosphination using classical heating conditions even for several days.

Treatment of the phosphine-borane **10b** with an excess of tetrafluoroboric acid dimethylether complex resulted in a quantitatively formation of the phosphines **12** (Scheme 6).



Scheme 6: Synthesis of chiral monophosphine **12**.

Complexes formed *in situ* from $\text{Rh}[(\text{COD})_2]\text{BF}_4$ and ligand **12** were examined as catalysts for the enantioselective hydrogenation of itaconic acid derivatives, leading to the formation of expected products in excellent yield and enantioselectivity up to 96% ee (Scheme 7).^[19]



Scheme 7: Hydrogenation of itaconic acid derivatives.


These results indicated clearly the efficiency of the isosorbide and isomannide structure for the synthesis of new chiral ligands for asymmetric catalysis applications. Development of new chiral compounds derived from this carbohydrate family as ligands or organocatalysts and their applications in asymmetric catalysis is currently underway in our laboratory.

Design and synthesis of bifunctional chiral organocatalysts. Applications to asymmetric organocatalysis.

The development of suitable, inexpensive, modular chiral organocatalysts has attracted considerable interest in recent years. Many chiral thiourea-phosphine compounds derived from natural or unnatural products have been reported in literature.^[20,21] Moreover, the potentiality and effectiveness of L-proline derivatives as organocatalysts have demonstrated very broad applications.^[22] Thus, in the course of our study to promote the use of natural and inexpensive compounds to elaborate catalytic systems, we focus on the use of simple and well-known L-proline as a chiral renewable resource for the synthesis of a new class of chiral thiourea-phosphines. The original structure of the chiral framework offers an interesting potential to the construction of bifunctional organocatalysts for asymmetric transformations.


Starting from L-proline, a new family of bifunctional chiral thiourea-phosphine organocatalysts was designed and synthesized. Our design of modular catalyst involves: 1) a urea function for interaction between catalyst and substrates; 2) a phosphorus center that is responsible for catalytic activity; 3) a stereogenic center that is ensured by the natural chirality of L-

proline; 4) a chiral carbon framework which offers potential modularity of a catalyst (Scheme 8).



Scheme 8: Retrosynthetic pathway to thiourea-phosphines **13**, **14**, **15**.

Thus, a series of 5 chiral thiourea-phosphine organocatalysts derived from L-proline was isolated in good overall yields.^[23] The synthesis is quite short, flexible and could be carried out in large gram scale. An example of the synthesis of **15** was presented in Scheme 9.



Scheme 9: Synthesis of thiourea-phosphine **15** from CBz-proline.

Thiourea-phosphine-organocatalyzed asymmetric C-N and C-S bond formation reaction


After achieving the synthesis of these chiral thiourea-phosphines, we were interested in testing their potential as organocatalysts for asymmetric organocatalyzed transformations.

Asymmetric allylic substitution reactions catalyzed by a chiral amine or phosphine have been powerful and versatile tools for the C-C, C-N, C-O and C-P bond construction.^[24] However, if several organocatalytic procedures exist for the C-S bond formation,^[25] examples including the allylic substitution of modified Morita-Baylis-Hillman (MBH) adducts are very scarce.^[26] To the best of our knowledge, only one example of asymmetric allylic substitution of alkyl thiols with MBH carbonate adducts catalyzed by a chiral amine has been described.^[27] No example of phosphine-catalyzed allylic

substitution of MBH acetate or carbonate adducts for C-S bond creation has been reported so far.

We report hereafter the use of chiral thiourea-phosphine as organocatalysts in the asymmetric allylic substitution of modified MBH adducts with phthalimide or alkyl thiols for the C-N or C-S bond formation.

After screening many reaction parameters including solvent, catalyst structure, additive, reaction time and temperature, the asymmetric allylic substitution reaction of tert-butoxycarbonyloxy-MBH adduct **16** with phthalimide using thiourea-phosphine based-organocatalysts derived from L-proline worked smoothly leading to the formation of expected product in very good yields and moderate to good enantioselectivities. We have also described the first example of thiourea-phosphine organocatalyzed C-S bond formation affording enantioenriched α -methylene- β -mercapto esters in excellent yields and with very high enantiomeric excess up to 93% ee (Scheme 10).^[28]



Scheme 10: Asymmetric allylic substitution of *tert*-butoxycarbonyloxy-MBH adduct **16**.

Reaction of **16a** (Ar = *p*-NO₂C₆H₄) with trityl thiol conducted to optically pure **17a** (R = Ph₃C) as a single crystals after recrystallization. Its absolute configuration was determined by X-ray single-crystal diffraction analysis (Figure 4).




Figure 4: ORTEP drawing representation of the molecular structure of **17a**. Ellipsoids are drawn at the 30 % probability level.

Synthesis of functionalized 2*H*- and 4*H*-pyran derivatives via organocatalyzed [4+2] annulation of tetrasubstituted alkenes with allenates.

Allenoates are interesting substrates for the Lewis base-catalyzed reactions because of their high reactivity to give






Figure 5: Some bioactive 4*H*-pyran and 2*H*-pyran compounds.

We have recently reported the first amine-organocatalyzed [4+2] annulation of allenolate **19** and all-carbon tetrasubstituted alkenes **18** for the synthesis of highly functionalized 2*H*- and 4*H*-pyran **20** and **21** derivatives. Different chiral amines including quinidine, cinchonidine, quinine, α -ICPN were tested for this reaction (Figure 6). Only β -ICD, conducted to the formation of expected enantioenriched 2*H*-pyran **20** bearing a chiral all-



access to the structurally divergent molecules. The addition of a Lewis base (LB) to the *sp*-hybridized β -carbon of allenolate generates the zwitterionic intermediates (Scheme 11) which participate in various annulation reactions for the synthesis of carbocyclic and heterocyclic compounds.^[29]



Scheme 11: Lewis base (LB)-catalyzed reactions of allenates with electrophiles (E).

Among them, the [4+2] and [2+2] annulations of allenates catalyzed by a tertiary amine bring potentialities for preparation of six and four membered ring structures towards the synthesis of natural products and biologically active molecules (Figure 5).^[30,31] However, to the best of our knowledge, the annulation of all-carbon tetrasubstituted alkenes with allenates has not been reported so far.

carbon quaternary center with enantiomeric excess up to 71% (Scheme 12).^[32] The results obtained will open the opportunities in synthesis of bioactive heterocyclic products and other pharmaceutical products.

Scheme 12: β -ICD catalyzed [4+2] annulation reaction of allenoate **19** with alkene **18**




Figure 6: Structure of chiral amine catalysts for [4+2] annulation reaction.

Based on these results obtained, the design of novel multifunctional organocatalysts, which are expected to afford higher levels of enantioselectivities, is currently underway in our laboratory.

Conclusion

We have designed and synthesized a new family of chiral amino alcohols, diamines and monophosphines derived from isosorbide and isomannide. These chiral ligands were obtained in moderate to good overall yields. The procedures for the preparation of these compounds were quite simple and practical and could be performed in large scale thanks to the inexpensive available starting materials. These ligands were successfully used in the metal-catalyzed enantioselective hydrogenation of ketones or activated olefins leading to the formation of expected products in excellent yields and enantioselectivities. We have recently reported the synthesis of new chiral thiourea-phosphines starting from L-proline. These molecules catalyzed efficiently the asymmetric allylic substitution of modified MBH adducts. Moreover, the synthesis of enantioenriched 2*H*-pyran derivatives bearing a chiral all-carbon quaternary center was also described, for the first time, using β -ICD catalyzed [4+2] annulation reaction.

Although, a number of chiral organocatalysts were designed and synthesized, their applications in asymmetric organocatalysis are still limited in catalyst loading and in asymmetric reaction scope. The concern of finding out an advanced and 'perfect' organocatalytic system always inspires chemists. We hope that this work would bring a promising way in the construction of chiral molecules with high enantiomeric purity which will be used as precursors for the synthesis of molecules or molecular scaffolds of biological and therapeutic interest.

Acknowledgements. The authors warmly thank MENSUR, Université Paris Sud, CNRS (UMR 8182) and Charm3At LabEx (ANR-11-LABEX-0039) for financial support. Our research has been also funded with the support of Vietnamese Government with 911 program

REFERENCES

1. P. T. Anastas, J. C. Warner, *Green Chemistry: Theory and practice*, Oxford University Press, New York, **1998**, 30
2. (a) A. Corma, S. Iborra, A. Velty. Chemical routes for the transformation of biomass into chemicals, *Chem. Rev.* **2007**, 107, 2411-2502. (b) P. Mäki-Arvela, B. Holmbom, T. Salmi, D. Y. Murzin. Recent progress in synthesis of fine and specialty chemicals from wood and other biomass by heterogeneous catalytic processes, *Catal. Rev. - Sci. Eng.* **2007**, 49, 197-340. (c) P. Gallezot. Catalytic routes from renewables to fine chemicals, *Catal. Today*, **2007**, 121, 76-91. (d) J. C. Serrano-Ruiz, R. Luque, A. Sepulveda-Escribano. Transformations of biomass-derived platform molecules: from high added-value chemicals to fuels via aqueous-phase processing, *Chem. Soc. Rev.*, **2011**, 40, 5266-5281. (e) P. Gallezot. Conversion of biomass to selected chemical products, *Chem. Soc. Rev.*, **2012**, 41, 1538-1558. (f) D. M. Alonso, S. G. Wettstein, J. A. Dumesic. Bimetallic catalysts for upgrading of biomass to fuels and chemicals, *Chem. Soc. Rev.*, **2012**, 41, 8075-8098.
3. (a) D. De Oliveira-Vigier, F. Jérôme. Heterogeneously-catalyzed conversion of carbohydrates, *Top. Curr. Chem.*, **2010**, 295, 63-92. (b) M. J. Climent, A. Corma, S. Iborra. Converting carbohydrates to bulk chemicals and fine chemicals over heterogeneous catalysts, *Green Chem.*, **2011**, 13, 520-540.
4. (a) C. H. Zhou, J. N. Beltramini, Y. X. Fan, G. Q. Lu. Chemoselective catalytic conversion of glycerol as a biorenewable source to valuable commodity chemicals, *Chem. Soc. Rev.*, **2008**, 37, 527-549. (b) J. Barrault, F. Jérôme. [Design of new solid catalysts for the selective conversion of glycerol](#), *Eur. J. Lipid Sci. Technol.*, **2008**, 110, 825-830. (c) A. Brandner, K. Lehnert, A. Bienholz, M. Lucas, P. Claus. Production of biomass-derived chemicals and energy: Chemocatalytic conversions of glycerol, *Top. Catal.*, **2009**, 52, 278-287. (d) B. Katryniok, H. Kimura, E. Skrzynsk, J. S. Girardon, P. Fongarland, M. Capron, R. Ducoulombier, N. Mimura, S. Paul, F. Dumeignil. Selective catalytic oxidation of glycerol: perspectives for high value chemicals, *Green Chem.*, **2011**, 13, 1960-1979.
5. (a) M. A. R. Meier, J. O. Metzger, U. S. Schubert. Plant oil renewable resources as green alternatives in polymer science, *Chem. Soc. Rev.*, **2007**, 36, 1788-1802. (b) A. Behr, J. Perez-Gomes. [The refinement of renewable resources: New important derivatives of fatty acids and glycerol](#), *Eur. J. Lipid Sci. Technol.*, **2010**, 112, 31-50.
6. (a) R. Palkovits, K. Tajvidi, J. Procelewska, R. Rinaldi, A. Ruppert. *Green Chem.* **2010**, 12, 972-978. (b) H. Kobayashi, H. Ohta, A. Fukuoka. Conversion of lignocellulose into renewable chemicals by heterogeneous catalysis. *Catal. Sci. Technol.* **2012**, 2, 869-883.
7. (a) J. Zakzeski, P. C. A. Bruijninx, A. L. Jongerius, B. M. Weckhuysen. The catalytic valorization of lignin for the production of renewable chemicals, *Chem. Rev.* **2010**, 110, 3552-3599. (b) M. P. Pandey, C. S. Kim. [Lignin depolymerization and conversion: A review of thermochemical methods](#), *Chem. Eng. Technol.*, **2011**, 34, 29-41.
8. H. B. Kagan, O. Riant. Catalytic asymmetric Diels Alder reactions, *Chem. Rev.*, **1992**, 92(5), 1007-1019.
9. (a) Nguyen Van Buu O., A. Aupoix, Doan T. H. N., Vo-Thanh. G. Chiral ionic liquids derived from isosorbide: synthesis, properties and applications in asymmetric synthesis *New J. Chem.*, **2009**, 33, 2060-2072; (b) Nguyen Van Buu O., A. Aupoix, Vo-Thanh G. Synthesis of novel chiral imidazolium based ionic liquids derived from isosorbide and their applications in asymmetric aza Diels-Alder reaction, *Tetrahedron*, **2009**, 65, 2260-2265; (c) V. Kumar, C. E. Olsen, S. J. C. Schaffer, V. S. Parmar and S. V. Malhotra. Synthesis and applications of novel bis(ammonium) chiral ionic liquids derived from isomannide, *Org. Lett.*, **2007**, 9(5), 3905-3908; (d) Truong, T. K. T.; Nguyen Van Buu, O.; Aupoix, A.; Pégot, B.; Vo-Thanh, G. *Curr. Org. Synth.* **2012**, 1, 53.

10. S. Kumar, U. Ramachandran. The synthesis and applications of asymmetric phase-transfer catalysts derived from isomannide and isosorbide, *Tetrahedron*, **2005**, 61, 4141.
11. For amino-alcohol and diamine ligands, see: (a) S. Guillarme, T. X. M. Nguyen. New chiral ligands from isosorbide: application in asymmetric transfer hydrogenation, *Tetrahedron: Asymmetry*, **2008**, 19, 1450-1454; (b) C. Paolucci, G. Rosini. Approach to a better understanding and modeling of (S)-dihydrofuran-2-yl, (S)-tetrahydrofuran-2-yl-, and furan-2-yl- β -dialkylaminoethanol ligands for enantioselective alkylation, *Tetrahedron: Asymmetry*, **2007**, 18, 2923-2946; (c) F. W. Lui, L. Yan, J. Y. Zhang, H. M. Liu. Highly stereoselective synthesis and structural characterization of new amino sugar derivatives, *Carbohydr. Res.*, **2006**, 314(3), 332-336; (d) G. De Coster, K. Vandyck, E. Van der Eycken, J. Van der Eycken, M. Elseviers, H. Röper. [D-Isomannide in synthesis: asymmetric Diels–Alder reactions with novel homochiral bis-imine Cu²⁺-catalysts](#), *Tetrahedron: Asymmetry*, **2002**, 13, 1673-1679.
12. For phosphorus ligands, see: (a) M. Diéguez, O. Pàmies, C. Claver. [Palladium-diphosphite catalysts for the asymmetric allylic substitution reactions](#), *J. Org. Chem.*, **2005**, 70, 3363-3368; (b) K. N. Gavrilov, S. V. Zheglov, P. A. Vologzhanin, M. G. Maksimova, A. S. Safronov, S. E. Lyubimov, V. A. Davankov, B. Schäffner, A. Börner. A P⁺-chiral bisdiamidophosphite ligand with a 1,4:3,6-dianhydro-d-mannite backbone and its application in asymmetric catalysis, *Tetrahedron Lett.*, **2008**, 49, 3120-3123; (c) M. T. Reetz, T. Neugebauer. [New diphosphite ligands for catalytic asymmetric hydrogenation: The crucial role of conformationally enantiomeric diols](#), *Angew. Chem. Int. Ed.*, **1999**, 38, 179-181; (d) H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu, H. Wang. [Asymmetric hydroesterification of styrene by PdCl₂-CuCl₂-chiral phosphine catalyst systems](#), *J. Organomet. Chem.*, **1997**, 543, 227-228; (e) M. T. Reetz, G. Mehler. Mixtures of configurationally stable and fluxional atropisomeric monodentate P ligands in asymmetric Rh-catalyzed olefin hydrogenation, *Angew. Chem. Int. Ed.*, **2000**, 44, 2959-2962; (f) K. N. Gavrilov, S. V. Zheglov, P. A. Vologzhanin, E. A. Rastorguev, A. A. Shiryayev, M. G. Maximova, S. E. Lyubimov, E. B. Benetsky, A. S. Safronov, P. V. Petrovskii, V. A. Davankov, B. Schäffner, A. Börner. [Diastereomeric P⁺-mono- and P⁺, P⁺-bidentate diamidophosphite ligands based on 1,4:3,6-dianhydro-d-mannitol in asymmetric metallocomplex catalysis](#), *Russ. Chem. Bull. Int. Ed.* **2008**, 57, 2311-2319; (g) R. K. Sharma, A. G. Samuelson. [Asymmetric allylation of aldehydes with chiral platinum phosphinite complexes](#), *Tetrahedron: Asymmetry* **2007**, 18, 2387.
13. For selected examples, see: (a) A. Schlatter, M. K. Kundu, W. D. Woggon. Enantioselective reduction of aromatic and aliphatic ketones catalyzed by ruthenium complexes attached to β -cyclodextrin, *Angew. Chem. Int. Ed.* **2004**, 43, 6731-6734; (b) D. Xue, Y.-C. Chen, X. Cui, Q.-W. Wang, J. Zhu, J.-G. Deng. Transfer hydrogenation of activated C=C bonds catalyzed by ruthenium amido complexes: Reaction scope, limitation, and enantioselectivity, *J. Org. Chem.*, **2005**, 70, 3584-3591; (c) X. Wu, X. Li, F. King, J. Xiao. Insight into and practical application of pH-controlled asymmetric transfer hydrogenation of aromatic ketones in water, *Angew. Chem. Int. Ed.*, **2005**, 44, 3407-3411; (d) A. M. Hayes, D. J. Morris, G. J. Clarkson, M. Wills. A class of ruthenium(II) catalyst for asymmetric transfer hydrogenations of ketones, *J. Am. Chem. Soc.*, **2005**, 127, 7318-7319; (e) W. Baratta, G. Chelucci, S. Gladiali, K. Siega, M. Toniutti, M. Zanette, E. Zangrando, P. Rigo. Ruthenium(II) terdentate CNN complexes: superlative catalysts for the hydrogen-transfer reduction of ketones by reversible insertion of a carbonyl group into the Ru-H bond, *Angew. Chem. Int. Ed.* **2005**, 44, 6214-6219; (f) S. Enthaler, R. Jackstell, B. Hagemann, K. Junge, G. Erre, M. Beller. [Efficient transfer hydrogenation of ketones in the presence of ruthenium N-heterocyclic carbene catalysts](#), *J. Organomet. Chem.* **2006**, 691, 4652-4659; (g) M. T. Reetz, X. Li. J. An efficient catalyst system for the asymmetric transfer hydrogenation of ketones: Remarkably broad substrate scope, *Am. Chem. Soc.*, **2006**, 128, 1044-1045.
14. ¹(a) T. Ohkuma, R. Noyori. *Comprehensive Asymmetric Catalysis Supplement* (Eds.: E. N. Jacobsen, A. Pfaltz, H. Yamamoto). Springer:Berlin, **2004**, Vol. 1, p. 43. (b) S. Gladiali, E. Alberico. Asymmetric transfer hydrogenation: chiral ligands and applications, *Chem. Soc. Rev.* **2006**, 35, 226-236. (c) T. Ikariya, A. J. Blacker. Asymmetric transfer hydrogenation of ketones with bifunctional transition metal-based molecular catalysts, *Acc. Chem. Res.* **2007**, 40, 1300-1308. (d) X. Wu, J. Xiao. Aqueous-phase asymmetric transfer hydrogenation of ketones – a greener approach to chiral alcohols, *Chem. Commun.*, **2007**, 24 2449-2466. (e) C. Wang, X. Wu, J. Xiao. [Broader, greener, and more efficient: Recent advances in asymmetric transfer hydrogenation](#), *Chem. Asian J.* **2008**, 3, 1750-1770. (f) P. Paradez, J. Diez, M. P. Gamasa. Synthesis of enantiopure iridium(I) and iridium(III) pybox complexes and their application in the asymmetric transfer hydrogenation of ketones, *Organometallics*, **2008**, 27, 2597-2607.
15. (a) M. Kitamura, R. Noyori. *Ruthenium in Organic Synthesis* (Ed.: S.-I. Murahashi). Wiley-VCH: Weinheim, **2004**, p. 3. (b) W. He, P. Liu, B. L. Zhang, X. L. Sun, S. Y. Zhang. Efficient iridium and rhodium-catalyzed asymmetric transfer hydrogenation using 9-amino(9-deoxy) cinchona alkaloids as chiral ligands, *Appl. Organometal. Chem.* **2006**, 20, 328-334.
16. (a) Huynh, K. D., H. Ibrahim, M. Toffano, Vo-Thanh G. [New class of chiral ligands derived from isosorbide: first application in asymmetric transfer hydrogenation](#), *Tetrahedron: Asymmetry*, **2010**, 21, 1542-1548. (b) Huynh K.D., H. Ibrahim, E. Kolodziej, M. Toffano, Vo-Thanh G. Synthesis of a new class of ligands derived from isosorbide and their application to asymmetric reduction of aromatic ketones by transfer hydrogenation, *New. J. Chem.* **2011**, 35, 2622-2631.
17. For a general review on the addition of organozinc reagent to aldehyde, see: a) K. Soai, S. Niwa. Enantioselective addition of organozinc reagents to aldehydes, *Chem. Rev.* **1992**, 92, 833-856. (b) L. Pu, H.-B. Yu. Catalytic asymmetric vinylation of ketones, *Chem. Rev.*, **2001**, 101, 757-824.
18. Huynh, K. D., H. Ibrahim, L. Bouchardy, C. Bournaud, E. Kolodziej, M. Toffano, Vo-Thanh G. [Biosourced ligands from isosorbide for the ethylation of aldehydes or alkynylation of imines](#), *Asian J. Org. Chem.*, **2016**, 5, 1242-1246.
19. H. Ibrahim, C. Bournaud, R. Guillot, M. Toffano, Vo-Thanh G. [Synthesis of novel chiral monophosphine ligands derived from isomannide and isosorbide. Application to enantioselective hydrogenation of olefins](#), *Tetrahedron Lett.*, **2012**, 53, 4900-4902.

20. (a) K. Yuan, H.-L. Song, Y. Hu, X.-Y. Wu. [Chiral phosphinothiourea-catalyzed asymmetric Morita–Baylis–Hillman reactions of acrylates with aromatic aldehydes](#), *Tetrahedron*, **2009**, 65, 8185-8190. (b) Y.-Q. Fang, E. N. Jacobsen. Cooperative, highly enantioselective phosphinothiourea catalysis of imine–allene [3 + 2] cycloadditions, *J. Am. Chem. Soc.* **2008**, 130, 5660-5561. (c) K. Yuan, H.-L. Song, Y. Hu, J.-F. Fang, X.-Y. Wu. [Enantioselective intramolecular Morita–Baylis–Hillman reaction using chiral bifunctional phosphinothiourea as an organocatalyst](#), *Tetrahedron : Asymmetry*, **2010**, 21, 903-908. (d) H.-P. Deng, M. Shi. Preparation of chiral multifunctional thiourea–phosphanes and synthesis of chiral allylic phosphites and phosphane oxides through asymmetric allylic substitution reactions of Morita–Baylis–Hillman carbonates, *Eur. J. Org. Chem.* **2012**, 183-187. (e) H.-P. Deng, Y. Wei, M. Shi. [Enantioselective synthesis of highly functionalized trifluoromethyl-bearing cyclopentenes: Asymmetric \[3+2\] Annulation of Morita–Baylis–Hillman carbonates with trifluoroethylidene malonates catalyzed by multifunctional thiourea-phosphines](#), *Adv. Synth. Catal.*, **2012**, 354, 783-789. (f) X.-N. Zhang, M. Shi. Phosphine-catalyzed [3 + 2]cycloaddition of 4,4-dicyano-2-methylenebut-3-enoates with benzyl buta-2,3-dienoate and penta-3,4-dien-2-one, *ACS Catal.*, **2013**, 3, 507-512. (f) Y. Wei, M. Shi. Multifunctional chiral phosphine organocatalysts in catalytic asymmetric Morita–Baylis–Hillman and related reactions, *Acc. Chem. Res.*, **2010**, 43, 1005-1018.
21. (a) S.-X. Wang, X. Han, F. Zhong, Y. Wang, Y. Lu. Novel amino acid based bifunctional chiral phosphines, *Synlett.*, **2011**, 2766-2778. (b) X. Han, F. Zhong, Y. Wang, Y. Lu. Versatile enantioselective [3+2] cyclization between Imines and allenates catalyzed by dipeptide-based phosphines, *Angew. Chem. Int. Ed.*, **2012**, 51, 767-770. (c) F. Hu, Y. Wei, M. Shi. [Enantioselective synthesis of spirocyclic cyclopentenes: asymmetric \[3+2\] annulation of 2-arylideneindane-1,3-diones with MBH carbonates derivatives catalyzed by multifunctional thiourea–phosphines](#), *Tetrahedron*, **2012**, 68, 7911-7919.
22. (a) S. Mukherjee, J.-W. Yang, S. Hoffmann, B. List. Asymmetric enamine catalysis, *Chem. Rev.*, **2007**, 107, 5471-5569. (b) A. Berkessel, H. Gröger. Asymmetric organocatalysis; Wiley-VCH: Berlin, Germany, **2005**. (c) F. Tanaka, C.-F. Barbas. Enantioselective organocatalysis; Dalko, P.-I., Ed. Wiley-VCH: Berlin, Germany, **2007**. (d) S. Bertelsen, K.-A. Jørgensen. Organocatalysis—after the gold rush, *Chem. Soc. Rev.* **2009**, 38, 2178-2189. (e) P.-M. Pihko, I. Majander, A. Erkkilä. Asymmetric organocatalysis; List, B., Ed. Springer-Verlag: Berlin, Germany, **2010**.
23. Nguyen T.H., M. Toffano, C. Bournaud, Vo-Thanh G. Synthesis of chiral thiourea–phosphine organocatalysts derived from L-proline, *Tetrahedron Lett.*, **2014**, 55, 6377-6380.
24. (a) C. W. Cho, M. J. Krische. *Angew. Chem. Int. Ed.*, Regio- and stereoselective construction of gamma-butenolides through phosphine-catalyzed substitution of Morita–Baylis–Hillman acetates: an organocatalytic allylic alkylation, **2004**, 43, 6689-6691. (b) Y. Q. Jiang, Y. L. Shi, M. Shi. Chiral phosphine-catalyzed enantioselective construction of γ -butenolides through substitution of Morita–Baylis–Hillman acetates with 2-trimethylsilyloxy furan, *J. Am. Chem. Soc.* **2008**, 130, 7202-7203. (c) H. L. Cui, X. Feng, J. Peng, J. Lei, K. Jiang, Y. C. Chen. Chemoselective asymmetric N-allylic alkylation of indoles with Morita–Baylis–Hillman carbonates, *Angew. Chem. Int. Ed.*, **2009**, 48, 5737-5740. (d) S. J. Zhang, H. L. Cui, K. Jiang, R. Li, Z. Y. Ding, Y. C. Chen. [Enantioselective allylic amination of Morita–Baylis–Hillman carbonates catalysed by modified cinchona alkaloids](#), *Eur. J. Org. Chem.*, **2009**, 2009(33), 5804-5809. (e) H. L. Cui, J. R. Huang, J. Lei, Z. F. Wang, S. Chen, L. Wu, Y. C. Chen. [Direct asymmetric allylic alkylation of butenolides with Morita–Baylis–Hillman carbonates](#), *Org. Lett.*, **2010**, 12, 720-723. (f) L. Hong, W. S. Sun, C. X. Liu, D. P. Zhao, R. Wang. Enantioselective construction of allylic phosphine oxides through substitution of Morita–Baylis–Hillman carbonates with phosphine oxides, *Chem. Commun.*, **2010**, 46, 2856-2858. (g) W. S. Sun, L. Hong, C. X. Liu, R. Wang. [Base-accelerated enantioselective substitution of Morita–Baylis–Hillman carbonates with dialkyl phosphine oxides](#), *Org. Lett.*, **2010**, 12, 3914-3917. (h) J. Peng, X. Huang, H. L. Cui, Y. C. Chen. [Organocatalytic and electrophilic approach to oxindoles with C3-quaternary stereocenters](#), *Org. Lett.*, **2010**, 12, 4260-4263. (i) J. R. Huang, H. L. Cui, J. Lei, X. H. Sun, Y. C. Chen. Organocatalytic chemoselective asymmetric N-allylic alkylation of enamides, *Chem. Commun.*, **2011**, 47, 4784-4786. (j) H. P. Deng, Y. Wei, M. Shi. Chiral bifunctional thiourea–phosphane organocatalysts in asymmetric allylic amination of Morita–Baylis–Hillman acetates, *Eur. J. Org. Chem.*, **2011**, 2011, 1956-1960. (k) Y. L. Yang, C. K. Pei, M. Shi. Multifunctional chiral phosphines-catalyzed highly diastereoselective and enantioselective substitution of Morita–Baylis–Hillman adducts with oxazolones, *Org. Biomol. Chem.*, **2011**, 9, 3349-3358.
25. P. Chauhan, S. Mahajan, D. Enders. Organocatalytic carbon–sulfur bond-forming reactions, *Chem. Rev.* **2014**, 114, 8807-8864.
26. X.-W. Liu, W.-Y. Han, X.-L. Liu, Y. Zhou, X.-M. Zhang, W.-C. Yuan. Highly regioselective synthesis of 3-alkylthio-2-oxindoles via DABCO-catalyzed allylic α -substitution of Morita–Baylis–Hillman carbonates of isatins with various thiols, *Tetrahedron*, **2014**, 70, 9191-9197.
27. A. Lin, H. Mao, X. Zhu, H. Ge, R. Tan, C. Zhu, Y. Cheng. [Organocatalytic asymmetric C-S bond formation: Synthesis of \$\alpha\$ -methylene- \$\beta\$ -mercapto esters with simple alkyl thiols](#), *Adv. Synth. Catal.*, **2011**, 353, 3301-3306.
28. Ngo, T.-T.-D., Nguyen, T.-H., C. Bournaud, R. Guillot, M. Toffano, Vo-Thanh G. [Phosphine–thiourea-organocatalyzed asymmetric C–N and C–S bond formation reactions](#), *Asian J. Org. Chem.*, **2016**, 5, 895-899.
29. (a) Ma, S. *Chem. Rev.*, **2005**, 105, 2829. (b) B. J. Cowen, S. J. Miller. Enantioselective catalysis and complexity generation from allenates, *Chem. Soc. Rev.*, **2009**, 38, 3102-3116. (c) S. Yu, S. Ma. Allenes in catalytic asymmetric Synthesis and Natural Product Syntheses, *Angew. Chem. Int. Ed.*, **2012**, 51, 3074-3112. (d) F. López, J. L. Mascareñas. *Chem. Soc. Rev.*, **2014**, 43, 2904-2915. (e) Y. Xiao, Z. Sun, H. Guo, O. Kwon. Chiral phosphines in nucleophilic organocatalysis, *Beilstein J. Org. Chem.*, **2014**, 10, 2089-2121.
30. For recent reports in [2+2] annulation of allenates, see: (a) G.-L. Zhao, J.-W. Huang, M. Shi. [Abnormal Aza-Baylis–Hillman reaction of N-tosylated imines with ethyl 2,3-butadienoate and penta-3,4-dien-2-one](#), *Org. Lett.*, **2003**, 5, 4737-4739. (b) J. B. Denis, G. Masson, P. Retailleau, J. Zhu. Cinchona alkaloid amide catalyzed enantioselective formal [2+2] cycloadditions of allenates and imines: Synthesis of 2,4-disubstituted azetidines, *Angew. Chem. Int. Ed.*, **2011**, 50, 5356-5360. (c) L. B. Saunders, S. J. Miller. Divergent reactivity in amine- and phosphine-catalyzed C–C bond-forming reactions of allenates with 2,2,2-trifluoroacetophenones,

- ACS Catal., **2011**, 1, 1347-1350. (d) T. Wang, X.-Y. Chen, S. Ye. DABCO-catalyzed [2+2] cycloaddition reactions of allenates and trifluoromethylketones: synthesis of 2-alkyleneoxetanes, *Tetrahedron Lett.*, **2011**, 52, 5488-5490. (e) P. Selig, A. Turočkin, W. Raven. Synthesis of highly substituted oxetanes via [2+2] cycloaddition reactions of allenates catalyzed by a guanidine Lewis base, *Chem. Commun.*, **2013**, 49, 2930-2932. (f) H.-B. Yang, Y. C. Yuan, Y. Wei, M. Shi. Amine-catalyzed tunable reactions of allenates with dithioesters: formal [4+2] and [2+2] cycloadditions for the synthesis of 2,3-dihydro-1,4-oxathiines and enantioenriched thietanes, *Chem. Commun.*, **2015**, 51, 6430-6433.
31. For recent reports in [4+2] annulation of allenates, see: a) X.-Y. Chen, M.-Y. Wen, S. Ye, Z.-X. Wang. Unusual formal [4 + 2] cycloaddition of ethyl allenate with arylidenoxindoles: Synthesis of dihydropyran-fused indoles, *Org. Lett.*, **2011**, 13, 1138. (b) K. D. Ashtekar, R. J. Staples, B. Borhan. [Development of a formal catalytic asymmetric \[4 + 2\] addition of ethyl-2,3-butadienoate with acyclic enones](#), *Org. Lett.*, **2011**, 13, 5732-5735. (c) X. Wang, T. Fang, X. Tong. Enantioselective amine-catalyzed [4+2] annulations of allenates and oxo-dienes: An asymmetric synthesis of dihydropyrans, *Angew. Chem. Int. Ed.*, **2011**, 50, 5361-5634. (d) C. K. Pei, L. Wu, L.; Lian, Z.; Shi, M. *Org. Biomol. Chem.*, **2012**, 10, 171. (e) F. Wang, Z. Li, J. Wang, X. Li, J.-P. Cheng. *J. Org. Chem.*, **2015**, 80, 5279-5286. (f) F. Wang, C. Luo, Y.-Y. Shen, Z.-D. Wang, X. Li, J.-P. Cheng. [Highly enantioselective \[4 + 2\] cycloaddition of allenates and 2-olefinic benzofuran-3-ones](#), *Org. Lett.*, **2015**, 17, 338-341. (g) Y. Liu, Y. Du, A. Yu, D. Qin, X. Meng. [Diverse synthesis of pyrano\[2,3-b\]indol and dihydropyrano\[2,3-b\]indol via tunable Lewis bases catalyzed domino reactions](#), *Tetrahedron*, **2015**, 71, 7706-7716. (h) F. Wang, C. Yang, X.-S. Xue, X. Li, J.-P. Cheng. A highly efficient chirality switchable synthesis of dihydropyran-fused benzofurans by fine-tuning the phenolic proton of β -isocupreidine (β -ICD) catalyst with methyl, *Chem. Eur. J.*, **2015**, 21, 10443.
32. Ngo.T.-T.-Duong, K. Kishi, M. Sako, M. Shigenobu, C. Bournaud, M. Toffano, R. Guillot, J.-P. Baltaze, S. Takizawa, H. Sasai, G. Vo-Thanh. Organocatalyzed [4+2] annulation of all-carbon tetrasubstituted alkenes with allenates: Synthesis of highly functionalized 2H- and 4H-Pyran derivatives, *Chemistry Select*, **2016**, 1, 5414-5420.