



# South Adriatic Recipes: Estimating the Vertical Mixing in the Deep Pit

Vanessa Cardin, Achim Wirth, Maziar Khosravi, Miroslav Gačić

## ► To cite this version:

Vanessa Cardin, Achim Wirth, Maziar Khosravi, Miroslav Gačić. South Adriatic Recipes: Estimating the Vertical Mixing in the Deep Pit. *Frontiers in Marine Science*, 2020, 7, pp.565982. 10.3389/fmars.2020.565982 . hal-03132961

**HAL Id: hal-03132961**

**<https://cnrs.hal.science/hal-03132961>**

Submitted on 7 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# South Adriatic Recipes: estimating the vertical mixing in the deep pit

Vanessa Cardin<sup>1,\*</sup>, Achim Wirth<sup>2</sup>, Maziar Khosravi<sup>1,3</sup> and Miroslav Gačić<sup>1</sup>

<sup>1</sup> National Institute of Oceanography and Applied Geophysics - OGS, Italy

<sup>2</sup> Univ. Grenoble Alpes, CNRS, Grenoble INP, LEGI, 38000 Grenoble, France

<sup>3</sup> TRIL programme, Abdus Salam International Center for Theoretical Physics - ICTP, 34151 Trieste, Italy

Correspondence\*:

V. Cardin

vcardin@inogs.it

## 2 ABSTRACT

3 The available historical oxygen data show that the deepest part of the South Adriatic Pit remains  
 4 well-ventilated despite the winter convection reaching only the upper 700 m depth. Here, we  
 5 show that the evolution of the vertical temperature structure in the deep South Adriatic Pit (dSAP)  
 6 below the Otranto Strait sill depth (780 m) is described well by continuous diffusion, a continuous  
 7 forcing by heat fluxes at the upper boundary (Otranto Strait sill depth) and an intermittent forcing  
 8 by rare (several per decade) deep convective and gravity-current events. The analysis is based  
 9 on two types of data: (i) 13-year observational data time series (2006 – 2019) at 750 m, 900  
 10 m, 1000 m and 1200 m depths of the temperature from the E2M3A Observatory and (ii) 55  
 11 vertical profiles (1985 – 2019) in the dSAP. The analytical solution of the gravest mode of the heat  
 12 equation compares well to the temperature profiles, and the numerical integration of the resulting  
 13 forced heat equation compares favorably to the temporal evolution of the time-series data. The  
 14 vertical mixing coefficient is obtained with three independent methods. The first is based on  
 15 a best fit of the long-term evolution by the numerical diffusion-injection model to the 13-year  
 16 temperature time series in the dSAP. The second is obtained by short-time (daily) turbulent  
 17 fluctuations and a Prandtl mixing length approximation. The third is based on the zero and first  
 18 modes of an Empirical Orthogonal Function (EOF) analysis of the time series between 2014 and  
 19 2019. All three methods are compared, and a diffusivity of approximately  $\kappa = 5 \cdot 10^{-4} \text{m}^2 \text{s}^{-1}$  is  
 20 obtained. The eigenmodes of the homogeneous heat equation subject to the present boundary  
 21 conditions are sine functions. It is shown that the gravest mode typically explains 99.5% of the  
 22 vertical temperature variability (the first three modes typically explain 99.85%) of the vertical  
 23 temperature profiles at 1 m resolution. The longest time scale of the dissipative dynamics in  
 24 the dSAP, associated with the gravest mode, is found to be approximately 5 years. The first  
 25 mode of the EOF analysis (85%) represents constant heating over the entire depth, and the zero  
 26 mode is close to the parabolic profile predicted by the heat equation for such forcing. It is shown  
 27 that the temperature structure is governed by continuous warming at the sill depth and deep  
 28 convection and gravity current events play less important roles. The simple model presented here  
 29 allows evaluation of the response of the temperature in the dSAP to different forcings derived  
 30 from climate change scenarios, as well as feedback on the dynamics in the Adriatic and the  
 31 Mediterranean Sea.


**Keywords:** ocean observations, vertical mixing, deep ocean, South Adriatic, time-series analysis, parameterization, climate variability

## 1 INTRODUCTION

The paradigm of the vertical temperature structure of the deep ocean is given by the “Abyssal recipes” of Munk (1966). It is based on a balance between an upward advective cooling and a downward diffusion of heat, which leads to an exponential profile of the temperature in the vertical direction. It explains large parts of the world’s ocean temperature and salinity stratification below the thermocline. The vertical turbulent diffusion of heat needed to maintain the structure is at least two orders of magnitude larger than the molecular diffusion values of sea water, and today, it is not entirely clear where the energy that supports the vertical mixing originates from (Munk and Wunsch (1998)). Tides are supposed to play an important role in the mixing process due to the breaking of internal tidal waves, generated by the interaction of the barotropic tide with the topography (see, i.e., Vic et al. (2019) and de Lavergne et al. (2019)). Mixing in the boundary layers and generation by wind-driven horizontal turbulence are other possible sources of mixing (see Vallis (2017) for a pedagogical discussion). These considerations can be applied to the Adriatic Sea (see fig. 1), whose dynamics have been widely studied in a large number of publications (see, e.g., the review by Cushman-Roisin et al. (2001)). As far as the deep South Adriatic Pit (dSAP) is concerned, historical studies have shown that even though the winter convection very rarely goes deeper than approximately 700 m, the bottom layer is well oxygenated (see e.g., Zore-Armanda et al. (1991), Bensi (2012)). Ventilation events are associated with the outbreaks of the cold North Adriatic deep water but do not occur every year. Thus, the ventilation should also be associated with some other processes, such as vertical diffusion.

More specifically, we consider the temperature structure in the area of the dSAP based on a considerable number of observations, starting from 1985 and increasing in number over the last two decades. The dSAP has a maximal depth of approximately 1230 m and is delimited at the south by the Otranto Sill (780 m deep)(see fig. 1). We focus on the temperature structure below the sill depth. It is significantly different from an exponential structure, as is shown below, and the theory put forward by Munk (1966) has to be supplemented. The Adriatic is also interesting because the tidal motion is negligible (see, e.g., Cushman-Roisin et al. (2001) and references therein), which simplifies the internal dynamics. An advantage of considering the dSAP is that it is easily accessible (only approximately 60 nautical miles from the coast) and has been extensively observed with considerable data coverage in space and time over the last 40 years. The present work relies on two types of data: first, in-situ ship-borne data measurements, vertical temperature and salinity profiles at a 1 m resolution, collected in the dSAP from 1985 to 2019 (Cardin et al. (2011)) and second, time series of high-frequency point measurements from 2006 to 2019 at 750 m, 900 m, 1000 m and 1200 m nominal depths from the mooring at the E2M3A Observatory, (<http://nettuno.ogs.trieste.it/e2-m3a/>). The latter are positioned in the dSAP and are particularly dedicated to the study of the long-term thermohaline and biogeochemical properties. In the present work, the diffusivity processes in the deep layer are analyzed, and thus, only measurements at 750 m (considered as the reference level for the Otranto sill depth) and deeper are taken into consideration.

It was shown by Querin et al. (2016) that the dynamics in the dSAP are forced by intermittent gravity current and deep convection events that reach below the sill depth every few years and that only last from a week to a few months. The most recent events of this type occurred in March and April of 2012 and in March 2017. The former event is due to cold and very dense water from the northern Adriatic (Bensi et al. (2013)), whereas the latter event is due to salty and warm water and has, so far, not been discussed in a



**Figure 1.** Left: Region of the South Adriatic Pit (SAP). The red rectangle shows the area where all the profiles originated. After 2010, data were collected within the black rectangle. The yellow dot indicates the position of the E2M3A observatory of the thermohaline time series. Right: All 55 profiles of the temperature dependence with the depth. The two profiles during the deep convection/gravity current events are emphasized in blue and red.

74 scientific publication. The subject of the present work is to determine what happens between such events,  
 75 when the vertical temperature structure is governed mainly by continuous vertical turbulent diffusion as  
 76 shown by Querin et al. (2016). The different time scales associated with short-lasting convection/gravity  
 77 current events and continuous diffusion allow separately consideration of the two regimes. In the present  
 78 work, we investigate continuous mixing based on observational data using the dynamics of the heat equation.  
 79 The aim is to show that in between deep convection/gravity current events, the dynamics are governed by  
 80 turbulent diffusion and that a bulk mixing coefficient can be determined to allow efficient parameterization.

81 The theory is presented in the next section, followed by the description of the experimental data and  
 82 methodology in section 3. The results for the profiles are given in section 4.3, and those for the time series  
 83 are given in section 4.4. The results are discussed in section 5, and conclusions are given in section 6.

## 2 THEORY

84 We consider the vertical structure of the dSAP and ignore the horizontal variations. This is justified by  
 85 the dominance of horizontal (isopycnal) mixing over vertical (diapycnal) mixing (Ledwell et al. (1998)).  
 86 Conceptually, the Munk solution is not valid in the dSAP, as there is no continuous injection in the deep  
 87 waters, which justifies a continuous upward velocity. Furthermore, fig. 1 shows that the vertical gradient  
 88 of the temperature vanishes towards the bottom, which indicates intense mixing without restratification  
 89 effects. The data from a depth of 750 m down to the bottom have a clearly non-exponential functional form,  
 90 indicating that the Munk theory of a balance between the upward advection and downward diffusion of heat  
 91 does not apply. Without upward advection, only vertical turbulent diffusion remains, with no process to  
 92 balance it and the only stationary solution is a constant temperature in space and time. This is not observed.  
 93 Instead, the density structure is evolving in space and time with dynamics described by the heat equation,  
 94 as is demonstrated below.

## 2.1 Heat Equation

We are interested in the evolution of the temperature  $T(z, t)$  in the dSAP deeper than the sill depth ( $\approx 780$  m). The underlying assumption of the present work is, that the dynamics in the dSAP are governed by the forced heat equation:

$$\partial_t T(z, t) = \kappa \partial_{zz} T(z, t) + I(z, t), \quad (1)$$

where the only free parameter in the system is the vertical eddy diffusivity  $\kappa$ . The last term represents the intermittent intrusions in the form of deep convection or gravity-current events and is supposed to vanish over extended periods in time. The dynamics evolve between  $z_s = -750$  m, the measurement point closest to the sill depth, and the bottom,  $z_b = -1220$  m. The boundary conditions are  $T(z, t) = T_{\text{obs}}(z_s, t)$  at the sill depth and  $\partial_z T(z_b, t) = 0$  at the bottom. This shows that the model is forced continuously by a time-dependent heat flux at its upper boundary that deeply diffuses, whereas there is no heat flux at the bottom. This model is called the diffusion-intrusion model in the sequel. It is linear and therefore allows us to separately obtain analytical solutions for different types of forcing and then add them to obtain the overall solution.

The unforced heat equation, eq. (1), is given by  $I(z, t) = 0$  and  $T_{\text{obs}}(z_s, t) = \text{const}$ . In the long term, the temperature relaxes to the imposed temperature at the sill depth down to the bottom, the only time independent solution. The time-dependent evolution is given by  $T^h(z, t) = T_{\text{obs}}(z_s) + r(t)m(z)$ , where  $r(t)$  and  $m(z)$  are related by:

$$\frac{\partial_t r}{r} = \kappa \frac{\partial_{zz} m}{m} = \lambda \quad (2)$$

and  $\lambda$  does not depend on  $z$  or  $t$ , as  $m$  is not dependent on  $t$  and  $r$  is not dependent on  $z$ . (Please see Zachmanoglou and Thoe (1986) chapter 9.2 for more details on the calculations and the solution using the “separation of variables”). We further define the following:  $\Delta T^h(z, t) = T^h(z, t) - T_{\text{obs}}(z_s, t)$ . The boundary conditions of  $\Delta T^h(z, t) = 0$  at the upper boundary and a vanishing temperature gradient at the lower boundary lead to the solution of the unforced heat equation of the form ( $z$  is given in meters):

$$\Delta T^h(z, t) = \sum_k A_k \sin\left(\frac{\pi(2k-1)(z-z_s)}{2(z_s-z_b)}\right) \exp(-\mu(2k-1)^2 t). \quad (3)$$

Note that only the wavenumbers, with odd integers  $2k-1$ , satisfy the bottom boundary-condition and eq. (2) imposes

$$\mu = \kappa \left( \frac{\pi}{2(z_s - z_b)} \right)^2. \quad (4)$$

The characteristic e-folding decay time scale for a vertical eddy diffusivity of  $\kappa = 5 \cdot 10^{-4} \text{m}^2 \text{s}^{-1}$  (see section 4) and for the first mode,  $k = 1$ , is  $\mu^{-1} = \kappa^{-1}(\pi/2)^{-2}(470\text{m})^2 \approx 5.7$  years. For the following modes satisfying the boundary conditions,  $k = 2, 3$  the time scales are 9 and 25 times shorter, respectively; the modes are damped in less than a year. The higher order modes ( $k > 3$ ) are damped within less than a month. The interannual dynamics are therefore dominated by the mode  $k = 1$  (see section 4). This shows that the unforced dynamics consists of a sum of noninteracting sine modes that decay exponentially in time at a rate that is proportional to the square of their wavenumber.

When a time-dependent forcing is applied, the dynamics of each mode are still independent. The coefficients are no longer constant in time, and we define  $B_k(t) = A_k(t) \exp(-\mu(2k-1)^2 t)$ . The first coefficient  $B_1(t)$  is obtained by projection on the first mode:

$$B_1(t) = \frac{4}{\pi(z_s - z_b)} \int_{z_b}^{z_s} \Delta T(z, t) \sin\left(\frac{\pi(z - z_b)}{2(z_s - z_b)}\right) dz \quad (5)$$

The projection of the data for modes  $k = 2$  and  $3$  and for data records that do not go to the total depth of 1220 m is given in appendix 1. The temporal evolution of the coefficients  $B_k$  is governed by the initial conditions at  $t = 0$ , the heat flux at the sill depth and the intermittent forcing.

In the case of the intermittently forced heat equation, through deep convection or gravity currents, the temperature structure changes, and higher order modes may also have large amplitudes. This intermittent forcing is represented in eq. (1) through the term  $I(z, t)$ . After a short (several months) intermittent event, the dynamics evolve unforced, and the amplitude of the higher modes decreases rapidly, as explained above. Only a few months after the intermittent forcing, the dynamics are again dominated by the lower order modes.

When the continuously forced heat equation is driven by increasing temperature at a depth of 750 m at a constant rate  $\gamma = \partial_t T_{\text{obs}}(-750\text{m}, t)$ , the solution is:

$$T^{cf}(z, t) = \gamma t + \frac{\gamma}{2\kappa} (z - z_b)^2, \quad (6)$$

which can be determined by introducing the above equation in eq. (1) with a vanishing inhomogeneous term,  $I(z, t) = 0$ . This corresponds to the constant temperature increase of a parabolic shaped temperature profile. The temperature difference between the sill depth and the bottom of the dSAP is constant in time and is related to the temperature increase at the sill depth by:

$$T^{cf}(z_s, t) - T^{cf}(z_b, t) = \frac{\gamma}{2\kappa} (z_s - z_b)^2 \quad (7)$$

It is important to note that the quadratic shape is not very different from the first sine mode (see fig. 2), and when it is projected on the sine modes, the amplitudes decay quickly with the wavenumber as  $(2k-1)^{-3}$ . The first sine mode explains more than 92.7% of the total variance; for the first three modes, it is almost 97%. This means that from the shape of a single temperature profile at a given time  $t_0$ , we cannot significantly determine whether the dynamics are freely decaying or forced by a constant-in-time vertical temperature flux. In other words,  $T^{cf}(z, t_0)$  resembles  $T^h(z, t_0)$  at a time  $t_0$ ; however, the evolution in time for the unforced and continuously forced cases is different. In the former case, the amplitude decays exponentially, while in the latter, it stays constant.

## 2.2 Heat Content and Heat Flux

When a linear equation of state is supposed, the heat content, its evolution, the potential energy and its evolution as well as the increase in potential energy by mixing can be obtained independently for every mode  $k$  and are determined by the coefficients  $B_k$ . If the vertical temperature profile is close to the first sine mode, the heat flux per unit area at the sill depth is

$$H = c_p \kappa \rho \partial_z T(z_s) = c_p \kappa \rho \frac{\pi(T(z_s) - T(z_b))}{2(z_s - z_b)} \quad (8)$$

157 and the heat content per unit area is:

$$\mathcal{H} = c_p \rho \frac{2(z_s - z_b)}{\pi} (T(z_s) - T(z_b)). \quad (9)$$

158 The ratio of the heat content and the heat flux is the characteristic time scale of the first mode, which is  
159 given in eq. (4).

### 3 EXPERIMENTAL DATA AND METHODS

160 In order to validate Munk's theory using experimental data we have considered the area of the South  
161 Adriatic (see fig. 1) where mixing and winter convection play an important role in homogenizing the  
162 physical and chemical properties of seawater and spreading heat along the water column. To analyze  
163 the importance of these processes, two different data sets are used: The first one consists of temperature  
164 profiles, taken during ship cruises, covering a long period of time and a large spatial area, and the second  
165 is times-series data, with high frequency point measurements along the water column by the E2M3A  
166 Observatory. The latter is located in the dSAP and is particularly dedicated to the study of long-term  
167 thermohaline and biogeochemical properties. In this study, we analyze the diffusivity processes in the deep  
168 layer; therefore, we consider only measurements deeper than 750 m.

#### 169 3.1 Profiles

170 The long-term variability of the thermal characteristics of the deep layer in the South Adriatic area  
171 is determined based on temperature data collected during several oceanographic campaigns between  
172 1985 and 2019. The cruises were undertaken as part of Italian and European projects, and other datasets  
173 were kindly provided by the NATO UNDERSEA RESEARCH CENTER, PANGAEA Data Repository  
174 (<https://www.pangaea.de/>) and MEDATALAS Dataset (<https://nodc.inogs.it/nodc/>). All the profiles were  
175 processed following the methodology described by Cardin et al. (2011) and applied for the data between  
176 1990 and 2010, i. e. the oceanographic condition during a specific campaign is represented by a profile  
177 obtained by averaging all the CTD casts deeper than 1000 m measured in the area between 41.25° and  
178 42.25° N and 17.25 ° and 18.25 ° E in the SAP (fig. 1). For the studied period, we consider data from 55  
179 temperature profiles deeper than 750 m. The potential temperature (hereafter referred to as temperature)  
180 is calculated from the original profiles with the surface as the reference depth. Profiles after 2010 were  
181 collected in a narrower area of 0.5°x0.5° degrees Lat/Lon located in the dSAP and near the E2M3A  
182 Observatory (black square in fig. 1). Further information about these cruise operations, materials and  
183 methods can be found at <http://www2.ogs.trieste.it:8585/biblioteca/>.

#### 184 3.2 Time Series

185 The data collected at the E2M3A Observatory enables the monitoring of changes that can be linked to  
186 variations in the general circulation of the Eastern Mediterranean Sea or, on a larger time scale, to the  
187 climate variability in the area, demonstrating the importance of high-frequency measurements to resolving  
188 events and rapid processes.

189 The Observatory has been working almost continuously since the end of 2006, and therefore, the high-  
190 frequency time series of physical and biochemical parameters measured at the site can be considered the  
191 longest almost continuous offshore data set available in the region (<http://nettuno.ogs.trieste.it/e2-m3a/>).  
192 The temperature was measured using CT-CTD sensors of the SBE37 at a sampled rate of 1 hour and SBE16  
193 with a sampling interval of 3 hours. For homogeneity, all the time series were averaged on a daily basis,  
194 and the temperature values were converted to the potential temperature. A low-pass filter PL33 with a

cutoff period of 33 hours (Flagg et al. (1976)) was applied to remove inertial oscillations (inertial period of 17.593 hours at 42° N in the South Adriatic Pit) so that tidal and higher frequency fluctuations do not mask the low-frequency fluctuations in the time series driven by winds and the density field. The filter removes more than 99% of the amplitude at the semidiurnal tidal periods and more than 90% of the amplitude at the diurnal tidal periods. These subinertial non-tidal time series are considered as the starting point for the analysis.

For this study, the time series considered along the mooring line come from 4 instruments positioned at 750 m, 900 m, 1000 m and 1200 m (Cardin and Bensi (2014), Cardin et al. (2014), Cardin et al. (2015) and Cardin et al. (2018)). To simplify the analysis, the depths of the series are nominal. The available data used are from November 2006 to October 2019, with a gap between September 2010 and May 2011. The time series at 900 m started in November 2013. An exhaustive description of the site, data availability and data analysis can be found in Bensi (2012), Bensi et al. (2014) and at the Observatory webpage (<http://nettuno.ogs.trieste.it/e2-m3a/>).

All instruments were calibrated and controlled for quality before and after each recovery and deployment, following the procedures indicated in Coppola et al. (2016) and Pearlman et al. (2019). In addition, during recovery or deployment operations, a CTD cast was performed to assess the accuracy of the measured data and to identify possible instrument drift.

### 3.3 Empirical Orthogonal Function Analysis

To assess the temporal variability and the vertical structure of the water column in the dSAP, we applied empirical orthogonal function analysis (EOF) (Preisendorfer and Mobley (1988)) to the temperature time series. For this purpose, time series at 750 m, 900 m, 1000 m, and 1200 m were analyzed.


Unfortunately, the time series shows gaps (not simultaneous) at different levels due to instrument failures, especially in the deepest part of the mooring line in the period from 2006 to 2010 and from 2011 to 2014. Since the EOF analysis requires data at all levels for its application, the data collected before 2014 can not be used for this analysis. Thus, we considered only the time series from September 2014 to August 2019, as it was the longest period without important gaps (only short ones -no longer than 5 days -due to the Observatory service). To eliminate these gaps, the daily time series were subsequently averaged on a weekly time scale. The filtered time series allowed us to study the time scales of interest (e.g. seasonal and interannual).

## 4 RESULTS

### 4.1 Numerical Integration

We performed a numerical integration of the heat equation with an imposed temperature at approximately the sill depth  $T_{\text{obs}}(-750\text{m}, t)$  (Dirichlet boundary condition) and a vanishing gradient at the bottom (1220 m) (Neumann boundary condition). For illustration, an unforced integration that starts from a linear stratification is represented in fig. 2. All values are divided by the temperature difference between 750 m and 1000 m depths to emphasize the evolution of the shape and the convergence to the first mode. Snapshots are shown at one and six months and at one, two and ten years ( $\kappa = 5 \cdot 10^{-4} \text{m}^2 \text{s}^{-1}$ ). The convergence to the first sine-mode, shown in fig. 2, with a characteristic time scale of several months, is clearly visible. In the same figure the shape of the parabolic profile for the constant forcing case is also given for comparison.

In the numerical integration, the two types of forcing, with time-dependent temperature variation at the sill depth  $\gamma$  and time and space dependent intermittent forcing  $I(z, t)$ , can be easily combined. To



**Figure 2.** Temperature profiles from the numerical model given by the unforced heat equation (1) (all values are divided by the temperature difference between 750 m and 1000 m depths) for the initial condition (black line), after one month (red), six months (green), one year (blue), two years (yellow) and ten years (brown) of the integration time ( $\kappa = 5 \cdot 10^{-4} \text{m}^2 \text{s}^{-1}$ ). The black dashed line is the analytic solution of the first mode, to which the dynamics converge. The red dashed line is the parabolic profile of the continuous forcing.

reproduce the space-time dynamics of the observed data, both types of forcing are applied, and eq. (1) is integrated numerically.

## 4.2 Eddy Diffusivity

The only free parameter in the heat equation is the diffusivity. The vertical exchange of heat is at least two orders of magnitude larger than that predicted by molecular motion only, and most is performed by turbulent fluid motion. Its parameterization is often achieved by assuming a vertical eddy diffusivity. It cannot be calculated from first principles but has to be determined from observations. It can be obtained based on different data analysis methods. The first method is a best fit of the time evolution of a numerical integration described in section 4.1 to the time-series data discussed in section 3.2. More precisely, we numerically solve eq. (1), which is forced at the upper boundary by imposing the temperature observed at 750 m depth and supplemented by two intermittent injections in 2012 and 2017, modeling the deep convection/gravity current events that penetrate the dSAP. The value of the diffusion constant  $\kappa$  is then estimated so that the evolution of the model best fits the time-series data. The second method is based on Prandtl's mixing-length concept (see: Prandtl (1925), Taylor (1922) Taylor (1959) and Bradshaw (1974)). We first calculate a typical r.m.s. vertical scale  $\Delta z_{rms} = \sqrt{\langle T'^2 \rangle / \partial_z \bar{T}}$  based on the temperature fluctuations  $T'$  at 900 m depth and an average temperature gradient  $\partial_z \bar{T}$  at the same depth. We also determine the correlation time of the vertical motion by looking at the autocorrelation function of the temperature fluctuations  $C(\tau) = \langle T'(t + \tau)T'(t) \rangle_t / \langle (T'(t))^2 \rangle_t$ , which is independent of time  $t$  for statistically stationary dynamics. We then obtain the correlation time:  $\tau_{corr} = \int_0^{\tau_0} C(\tau) d\tau$ , where  $\tau_0$  is the first zero crossing of  $C(\tau)$ . The vertical eddy diffusivity can then be estimated by:

$$\kappa_{turb} = C_s \frac{(\Delta z_{rms})^2}{\tau_{corr}}, \quad (10)$$

where  $C_s$  is a non-dimensional constant. The third method is based on the continuously forced dynamics, which leads to a balance between the temperature increase and the temperature difference between the sill depth and the bottom, as expressed by eq. (7). The three approaches for estimating the vertical eddy diffusivity are performed in sections 4.4 and 4.5.

### 4.3 Profiles


When considering the 55 temperature profiles taken during the period from 1985 to 2019, which are presented in fig. 1, a similarity of most of them to the first sinusoidal mode depicted in fig. 2 is discernible. For a quantitative analysis, we calculated the coefficients  $B_k$  for  $k = 1, 2$  and 3, which are given in fig. 3. Only the amplitude of the first mode has a value which is significantly different from zero, varying around  $0.5^\circ \text{ K}$ , while the amplitudes of the second and third mode are typically more than an order of magnitude smaller and have no definite sign. We further calculated the remaining variance not explained by the first mode or the first three modes in fig. 3. For over 2/3 of the profiles, more than 99.5% of the variance can be explained by the first mode and more than 99.85% by the first three modes. Note that the 2017 event is clearly depicted by the atypical profile from 22/03/2017, given in red in fig. 1. The event also has a clear signature in the amplitude of the second and third modes (fig. 3) and in a strong increase in the variability neither explained by the first mode nor by the first three modes (fig. 3). This means that for this profile, higher order modes have an increased amplitude. In the subsequent profile only 4 months later (27/07/2017), the amplitude of the second and third mode as well as the unexplained variability has decreased to the values prior to the event. This is coherent with the fact that the decay times of modes two and three are only a few months, as explained in section 2.1. The decay of even higher modes happens at scales shorter than a month. The 2012 event, however, has no conspicuous signature in the amplitude or variability data. The profile from 31/03/2012 (given in blue in fig. 1)) shows that the dense gravity current water is just starting to arrive, and it is seen at the very bottom of the data but does not significantly impact the overall shape of the profile. No profile is available for the rest of the year, and the next profile is one year later (23/03/2013); at this time, the second, third and higher modes have faded away.

### 4.4 Time Series

We first numerically integrate the diffusion-intrusion model. An integration with  $\kappa_{\text{fit}} = 5 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-1}$  and two convection or gravity-current events in March 2012 centered at the bottom and in March 2017 centered at  $z = 1100 \text{ m}$  manages to reproduce, to a reasonable degree, the observed temperature and salinity evolution, as shown in fig. 4.

Furthermore, the data at 750 m and 1200 m allows determining the amplitude of the first mode. Using this information and supposing that the higher-order modes are of low amplitude, allows to interpolate for the depths in-between. The obtained values were compared to the observed data at 900 m and 1000 m. Again, we remark a good correspondence between the observational data and the interpolation in fig. 5. A slight deviation is observed between years 2014 and 2017.

The eddy diffusion can be estimated based on the correlation time and the vertical r.m.s displacement of a fluid parcel, as explained in section 4.2. The former is given by the integral of the correlation function from zero to the first zero crossing. The first zero crossing is at 4 days, and the integral gives  $\tau_{\text{corr}} = 1.6$  days (not shown). The latter is calculated based on the temperature fluctuations and the mean temperature gradient (box-averaged over 30 days); its average value lies around  $\Delta z_{\text{rms}} = 20 \text{ m}$ . When we impose  $\kappa_{\text{turb}} = \kappa_{\text{fit}}$  we find, using eq. (10), that  $C_s \approx 0.18$ . There is no surprise that it is smaller than one, as some of the vertical motion is due to internal waves, which do not lead to mixing if they do not break.




**Figure 3.** Upper row: evolution of the amplitudes of the sine modes  $B_1$  (black),  $B_2$  (red) and  $B_3$  (green). The vertical scale gives the temperature difference between the 750 m time series and the maximum profile depth (1219 m) of the corresponding mode. Lower row: part of variability not explained by  $B_1$  (black),  $B_1 + B_2$  (red) or  $B_1 + B_2 + B_3$  (green) (left) and zoom (right). Figures on the right show the same data magnified.


297 Interestingly, the former is obtained based on turbulent fluctuations on a time scale of a few days, while the  
 298 latter is obtained based on the evolution of the temperature profile over several years.

#### 299 4.5 EOF analysis

300 The modes of the EOF analysis of the time series and the temporal evolution of EOF mode 1 are presented  
 301 in fig. 6. Note that the time average is given by EOF mode 0 and that its profile resembles the parabolic  
 302 profile put forward in section 2.1 for the case with a constant forcing. The dominant evolution is given by  
 303 EOF mode 1 (describing 87% of the variance), which is close to a constant-in-depth temperature increase.  
 304 In this case, the temperature increase at the sill depth and the amplitude of mode 0 are related by eq. (7). The  
 305 amplitude for mode 0 is  $\Delta T = 0.5^\circ \text{ K}$  (see fig. 6a). The warming trend is obtained by taking the product of  
 306 the vertical average of mode 1 (which is close to constant see fig. 6b) and the trend of its amplitude (see fig.




**Figure 4.** The observational data (thin lines) and the corresponding solutions of the injection-diffusion model (thick lines) for different depths (black 750 m, red 900 m, green 1000 m, blue 1200 m) for temperature.



**Figure 5.** Observed data at 900 m (red) and 1000 m (light green) and their interpolations 900 m (orange) and 1000 m (dark green). The right figure is a magnification of the data starting in 2017.

307 6c), that is:  $\gamma = 0.5 \cdot 0.64^\circ \text{K}/(5\text{years}) = 6.4 \cdot 10^{-2}^\circ \text{K/year}$ . A diffusivity of  $\kappa_{EOF} = 4.5 \cdot 10^{-4} \text{m}^2 \text{s}^{-1}$  is  
 308 obtained from eq. (7), which is close to  $\kappa_{\text{turb}}$  and  $\kappa_{\text{fit}}$ . We remind that the Otranto Sill depth is 780 m.

309 The higher order EOF modes 1, 2 and 3 closely resemble discretized versions of the analytic sine modes  
 310  $k=1, 2$  and  $3$  (wavenumbers 1,3 and 5). If we suppose that they are subject to random forcing of equal  
 311 strength and diffusive damping, their relative variance should decrease as 1, 1/9 and 1/25. The observed  
 312 values are 1, 1/8 and 1/48, which is within discretization and statistical error (the variance of the last EOF  
 313 mode is only 0.24%).



**Figure 6.** Temporal average of data, EOF mode 0 (a) and the EOF modes 1 to 4 with their variances given in the legend (b). Temporal evolution of the amplitude of EOF mode 1 (c).

## 5 DISCUSSION

All the above results based on temperature measurements show that the action of turbulent mixing in the dSAP can be modeled well by a heat equation with a constant diffusion coefficient in space and time, a prescribed heat flux at the sill depth and a forcing term that represents rare gravity current and deep convection events. The similarity between the empirical EOF analysis results (applied only to a subset of the time series between 2014 and 2019) and a solution of the forced heat equation, with a forcing that imposes a warming trend through a heat flux at 750 m depth, confirms that the heat equation satisfactorily describes the processes in the dSAP.

When the functional form of the first mode for the temperature stratification is assumed to have an amplitude of  $0.5^{\circ}\text{K}$ , the heat flux of the dSAP is  $H = 3.3\text{Wm}^{-2}$  according to eq. (8), and the heat content is  $\mathcal{H} = 6 \cdot 10^8\text{Jm}^{-2}$  according to eq. (9). These values characterize the heat-budget of the dSAP and can help to validate numerical models and climate dynamics.

The power per square meter necessary to perform turbulent mixing of the mass (based on temperature only) is

$$P = g\alpha\rho\kappa_{\text{turb}}\Delta T. \quad (11)$$

For a  $\Delta T = 0.5^\circ \text{ K}$ , this parameter is  $P = 5 \cdot 10^{-4} \text{ W/m}^2$ . A mixing efficiency of 0.2 leads to a drain of five times more kinetic energy (see, i.e., Osborn (1980) and Peltier and Caulfield (2003)). If the turbulent Prandtl number is unity, we obtain a characteristic vertical shear of  $\approx 3 \cdot 10^{-3} \text{ s}^{-1}$ .

The presented theory and observations show that from 2006 to 2019, the temperature stratification in the dSAP is determined by the continuous warming in the upper part (above the Otranto Sill depth) and the vertical turbulent diffusion. Other processes such as deep convection and gravity currents that penetrate the dSAP are only visible during a few months on two occasions. We obtain an upper (diffusive) limit for the characteristic time scale for the density structure of the dSAP, which is important as it tells us that the long-term changes in the circulation in the area are due to external forcing. The upper 5-year limit is also important for numerical modeling, as it gives a maximum spin-up time for numerical integrations. When the first mode is correctly estimated in the initial conditions of the numerical integration, a spin up of less than a year becomes sufficient, as only modes higher than  $k = 1$  have to spin up.

The main result is that at a high accuracy, the spatio-temporal evolution of the temperature in the dSAP is described by a heat equation based on a constant (in space and time) vertical diffusivity of  $\kappa = 5 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-1}$ . This value is similar to published values of vertical diffusivities in other regions of the world oceans. Munk (1966), considering several stations in the interior Pacific Ocean over much larger depths, found that the dominant vertical dynamics acting on the temperature were advective diffusive, with  $\kappa = 1.3 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-1}$ . The eddy diffusivity in deep basins of the North Aegean was estimated to lie between  $\kappa = 2 \cdot 10^{-4}$  to  $10 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-1}$  below a depth of 500 m ((Zervakis et al. (2003)). In the deepest part of the Calypso Deep (below 4400 m), the vertical diffusivity was found to vary around  $\kappa = (4 \pm 3) \cdot 10^{-3} \text{ m}^2 \text{ s}^{-1}$  by Kontoyiannis et al. (2016).

## 6 CONCLUSIONS AND PERSPECTIVES

We showed that the theory proposed by the “Abyssal recipes” of Munk (1966) for the world ocean is valid for the dSAP, a small oceanic basin, when forcing is added. The hypothesis of the present work is that the dynamics in the dSAP are governed by turbulent diffusion and forced by a continuous heat flux at the upper boundary and sporadic deep convection/gravity current events. A hierarchy of models, all based on the heat equation, is used. The simplest are based on the gravest or the three gravest modes of the analytically solved unforced heat equation. It is shown that the heat equation forced by a constant heat flux at the upper boundary has a functional form close to the gravest mode. The most evolved model includes a time-dependent forcing at the upper boundary as well as intermittent convection events that are solved numerically.

The models successfully describe the profiles their spatio-temporal evolution. The functional form of the profile is described by the gravest mode of the analytical solution of the homogeneous heat equation, while the time series can be fitted well by the numerical solution of the forced heat equation. Two major deep convection/gravity current events are identified in the time series in 2012 and 2017 between 2007 and 2019 (during the winter/spring season). The longest dissipative time scale in the dSAP, associated with the gravest mode, is approximately 5 years. This does not only indicate that within this period, stratification in the dSAP disappears when forcing resides, but it represents an upper bound for the absence of deep convection events penetrating the dSAP. More precisely, when the temperature at the sill depth does not increase and no deep convection event reaches below the Otranto Sill depth within 5 years, the dSAP becomes essentially unstratified, and subsequent convection is likely to reach the bottom of the dSAP. In other words, a constant warming of the water mass above the sill depth prevents deep convection events

from reaching the dSAP, and only diffusion remains; however, when there is no warming at the sill depth, a deep convection event will occur at least every five years.

The above developed theory about the dynamics of the dSAP allows a simple injection diffusion model of the dSAP to be built. This can be used to study the evolution of the dSAP under climate change scenarios, as well as to parameterize the feedback of the dSAP on the surrounding ocean dynamics. More precisely, the decomposition of modes with different characteristic time scales can be used to parameterize the response of the deep South Adriatic Pit to a specific process. For interannual variability, only the dynamics of the first mode have to be considered, while for a seasonal forcing, only the first two modes have to be taken into account, considerably reducing the complexity of the problem. The next step is to attempt to extrapolate the above findings to the more involved dynamics above the sill depth, where water masses are subject to horizontal advection and air-sea fluxes.

The performance of the theory and data analysis developed in this publication are not limited to the dSAP but can be equally applied to other enclosed pits in the world ocean, away from the direct influence of atmospheric forcing and strong currents. The temperature variation in the dSAP is significantly different from the exponential profile put forward by Munk (1966). As in Munk and Wunsch (1998), we have to ask the question of which process provides the power to perform the mixing in the case of the dSAP, where tides are excluded. This is one of the key open questions in physical oceanography. To answer this question, the synergy of laboratory experiments on the Coriolis platform (Grenoble) of idealized numerical simulations and of fine resolution (in time-and-space) velocity measurements in the dSAP is currently being employed.

The proposed empirical model for the vertical, one-dimensional, structure of the temperature profile, its evolution in time, as well as the determination of the vertical mixing coefficients and the forcing, relies on the synergy between mooring data with dense temporal coverage and a substantial number of profiles, providing fine resolution in the vertical direction. This dataset has been collected during the last four decades, at a single oceanic site and shows the importance of a continuous effort of collecting data over several generations of funding programs.

## 1 PROJECTION

The discretized version of the projection to obtain the amplitude of the modes is given for the first three modes, it is defined iteratively by:

$$B_1(t) = \frac{\lambda_1}{2 \cdot 469} \sum_{i=750}^{1219} \Delta T(i, t) \sin\left(\frac{\pi(i - z_s)}{2 \cdot 469m}\right) \quad (12)$$

$$B_2(t) = \frac{\lambda_2}{2 \cdot 469} \sum_{i=750}^{1219} \left( \Delta T(i, t) - B_1(t) \sin\left(\frac{\pi(i - z_s)}{2 \cdot 469m}\right) \right) \sin\left(3 \frac{\pi(i - 750m)}{2 \cdot 469m}\right) \quad (13)$$

$$B_3(t) = \frac{\lambda_3}{2 \cdot 469} \sum_{i=750}^{1219} \left( \Delta T(i, t) - B_1(t) \sin\left(\frac{\pi(i - z_s)}{2 \cdot 469m}\right) - B_2(t) \sin\left(3 \frac{\pi(i - 750m)}{2 \cdot 469m}\right) \right) \sin\left(5 \frac{\pi(i - 750m)}{2 \cdot 469m}\right) \quad (14)$$

If the record is over the total length  $\lambda_k = 1$ . If the record stops at length  $z'_B < 1219m$ , so does the summation in eqs. (12) - (14) and:

$$\lambda_k = \left( \frac{z'_B - z_s}{469m} - \frac{\sin(2(k-1)\pi \frac{z'_B - z_s}{469m})}{2(k-1)\pi \cdot 469m} \right)^{-1} \quad (15)$$

## CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## AUTHOR CONTRIBUTIONS

All authors have contributed to the paper. The numerical projection and integration was performed by AW, the EOF analysis by MK.

## FUNDING

This work was funded by Labex OASUG@2020 (Investissement d'avenir - ANR10 LABX56) and OGS internal funding.

## ACKNOWLEDGMENTS

Part of this work was performed when AW visited OGS and VC visited LEGI. We thank the OGS ExO and TECDEV Group so as the Captain and crew of the RV OGS Explora and RV Laura Bassi for the valuable support and work during the maintenance of the E2M3A Observatory.

## DATA AVAILABILITY STATEMENT

Data sets are referenced in the references.

## REFERENCES

- Bensi, M. (2012). *Thermohaline variability and mesoscale dynamics observed at the E2M3A deep-site in the South Adriatic Sea*. Ph.D. thesis, Università degli studi di Trieste, <https://www.openstarts.units.it/handle/10077/7387>
- Bensi, M., Cardin, V., and Rubino, A. (2014). Thermohaline variability and mesoscale dynamics observed at the deep-ocean observatory e2m3a in the southern adriatic sea. *The Mediterranean Sea: Temporal Variability and Spatial Patterns, Geophysical Monograph Series. John Wiley & Sons, Inc., Oxford, UK*, 139–155
- Bensi, M., Cardin, V., Rubino, A., Notarstefano, G., and Poulain, P. (2013). Effects of winter convection on the deep layer of the southern adriatic sea in 2012. *Journal of Geophysical Research: Oceans* 118, 6064–6075
- Bradshaw, P. (1974). Possible origin of prandtl's mixing-length theory. *Nature* 249, 135–136
- [Dataset] Cardin, V. and Bensi, M. (2014). E2m3a-2006-2010-time-series-southadriatic. doi:10.6092/36728450-4296-4E6A-967D-D5B6DA55F306
- Cardin, V., Bensi, M., and Pacciaroni, M. (2011). Variability of water mass properties in the last two decades in the south adriatic sea with emphasis on the period 2006–2009. *Continental Shelf Research* 31, 951–965

- 424 [Dataset] Cardin, V., Bensi, M., Siena, G., and Ursella, L. (2014). E2m3a-2011-2013-time-series-  
425 southadriatic. doi:10.6092/84CB588D-97E5-4C64-91BB-BA6109DFA530
- 426 [Dataset] Cardin, V., Bensi, M., Ursella, L., and Siena, G. (2015). E2m3a-2013-2015-time-series-  
427 southadriatic. doi:10.6092/F8E6D18E-F877-4AA5-A983-A03B06CCB987
- 428 [Dataset] Cardin, V., Bensi, M., Ursella, L., and Siena, G. (2018). E2m3a-2015-2017-time-series-  
429 southadriatic. doi:10.6092/238B5903-A173-4FC3-AC5F-5FEDF1064A39
- 430 Coppola, L., Ntoumas, M., Bozzano, R., Bensi, M., Hartman, S., Charcos Llorens, M., et al. (2016).  
431 Handbook of best practices for open ocean fixed observatories.
- 432 Cushman-Roisin, B., Gacic, M., Poulain, P.-M., and Artegiani, A. (2001). *Physical oceanography of the*  
433 *Adriatic Sea: past, present and future* (Springer Science & Business Media)
- 434 de Lavergne, C., Falahat, S., Madec, G., Roquet, F., Nycander, J., and Vic, C. (2019). Toward global maps  
435 of internal tide energy sinks. *Ocean Modelling* 137, 52–75
- 436 Flagg, C., Vermersch, J., and Beardsley, R. (1976). 1974 mit new england shelf dynamics experiment  
437 (march, 1974). *Data Report, Part II: the Moored Array, MIT Report 76-1*
- 438 Kontoyiannis, H., Lykousis, V., Papadopoulos, V., Stavrakakis, S., Anassontzis, E., Belias, A., et al. (2016).  
439 Hydrography, circulation, and mixing at the calypso deep (the deepest mediterranean trough) during  
440 2006–09. *Journal of Physical Oceanography* 46, 1255–1276
- 441 Ledwell, J. R., Watson, A. J., and Law, C. S. (1998). Mixing of a tracer in the pycnocline. *Journal of*  
442 *Geophysical Research: Oceans* 103, 21499–21529
- 443 Munk, W. and Wunsch, C. (1998). Abyssal recipes ii: Energetics of tidal and wind mixing. *Deep-sea*  
444 *research. Part I, Oceanographic research papers* 45, 1977–2010
- 445 Munk, W. H. (1966). Abyssal recipes. In *Deep Sea Research and Oceanographic Abstracts* (Citeseer),  
446 vol. 13, 707–730
- 447 Osborn, T. (1980). Estimates of the local rate of vertical diffusion from dissipation measurements. *Journal*  
448 *of physical oceanography* 10, 83–89
- 449 Pearlman, J. S., Bushnell, M., Coppola, L., Buttigieg, P. L., Pearlman, F., Simpson, P., et al. (2019).  
450 Evolving and sustaining ocean best practices and standards for the next decade. *Frontiers in Marine*  
451 *Science* 6, 277
- 452 Peltier, W. and Caulfield, C. (2003). Mixing efficiency in stratified shear flows. *Annual review of fluid*  
453 *mechanics* 35, 135–167
- 454 Prandtl, L. (1925). 7. bericht über untersuchungen zur ausgebildeten turbulenz. *ZAMM-Journal of Applied*  
455 *Mathematics and Mechanics/Zeitschrift für Angewandte Mathematik und Mechanik* 5, 136–139
- 456 Preisendorfer, R. W. and Mobley, C. D. (1988). Principal component analysis in meteorology and  
457 oceanography. *Developments in atmospheric science* 17
- 458 Querin, S., Bensi, M., Cardin, V., Solidoro, C., Bacer, S., Mariotti, L., et al. (2016). Saw-tooth modulation  
459 of the deep-water thermohaline properties in the southern adriatic sea. *Journal of Geophysical Research:*  
460 *Oceans* 121, 4585–4600
- 461 Taylor, G. (1959). The present position in the theory of turbulent diffusion. In *Advances in Geophysics*  
462 (Elsevier), vol. 6. 101–112
- 463 Taylor, G. I. (1922). Diffusion by continuous movements. *Proceedings of the london mathematical society*  
464 2, 196–212
- 465 Vallis, G. K. (2017). *Atmospheric and oceanic fluid dynamics* (Cambridge University Press)
- 466 Vic, C., Garabato, A. C. N., Green, J. M., Waterhouse, A. F., Zhao, Z., Melet, A., et al. (2019). Deep-ocean  
467 mixing driven by small-scale internal tides. *Nature communications* 10, 1–9

- 468 Zachmanoglou, E. C. and Thoe, D. W. (1986). *Introduction to partial differential equations with*  
469 *applications* (Courier Corporation)
- 470 Zervakis, V., Krasakopoulou, E., Georgopoulos, D., and Souvermezoglou, E. (2003). Vertical diffusion  
471 and oxygen consumption during stagnation periods in the deep north aegean. *Deep Sea Research Part I:*  
472 *Oceanographic Research Papers* 50, 53–71
- 473 Zore-Armanda, M., Bone, M., Dadic, V., Morovic, M., Ratkovic, D., Stojanoski, L., et al. (1991).  
474 Hydrographic properties of the adriatic sea in the period from 1971 through 1983. *Acta adriatica*. 1991.