



General third order Chapman-Enskog expansion of lattice Boltzmann schemes

François Dubois

► To cite this version:

François Dubois. General third order Chapman-Enskog expansion of lattice Boltzmann schemes. 16th International Conference for Mesoscopic Methods in Engineering and Science, Jul 2019, Edinburgh, United Kingdom. hal-03434871

HAL Id: hal-03434871

<https://cnrs.hal.science/hal-03434871>

Submitted on 18 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

General third order Chapman-Enskog expansion of lattice Boltzmann schemes

François Dubois^{ab}

^a Laboratoire de Mathématiques d'Orsay, Université Paris-Sud,

CNRS, Université Paris-Saclay, Orsay, France.

^b Conservatoire National des Arts et Métiers, LMSSC laboratory, F-75003 Paris, France.

08 May 2019 *

The lattice Boltzmann scheme in his actual form has been developed with the contributions of Lallemand, Succi, d'Humières, Luo [HSB91, QHL92, DDH92, LL00] and many others. In order to derive the equivalent partial differential equations, a classical of the Chapman Enskog expansion is popular in the lattice Boltzmann community (see *e.g.* [LL00]) A main drawback of this approach is the fact that multiscale expansions are used without a clear mathematical signification of the various variables and functions. Independently of this framework, we have proposed in [FD07, FD08] the Taylor expansion method to obtain formally equivalent partial differential equations. The infinitesimal variable is simply the time step (proportional to the space step with the acoustic scaling). This approach has been experimentaly validated in various contributions [DL10, DL11]. A third order extension for fluid flow has been proposed in [FD09] and an efficient implementation up to fourth order accuracy is presented in [ADGL14].

In this contribution, we consider a regular lattice \mathcal{L} composed by vertices x separated by distances that are simple expressions of the space step Δx . A discrete time t is supposed to be an integer multiple of a time step $\Delta t > 0$.

* Edition 22 March 2021. This contribution has been presented to the 16th International Conference for Mesoscopic Methods in Engineering and Science, Edinburgh, 22–26 July 2019.

A very general lattice Boltzmann scheme with q discrete velocities of the form

$$f_j(x, t + \Delta t) = f_j^*(x - v_j \Delta t, t), \quad 0 \leq j < q.$$

The distribution f^* after relaxation is defined with moments m such that

$$m_k = \sum_j M_{k\ell} f_j.$$

The d'Humières matrix [DDH92] M is invertible and we decompose the moments in the following way:

$$m \equiv \begin{pmatrix} W \\ Y \end{pmatrix}.$$

The conserved variables W are not modified after relaxation: $W^* = W$. The microscopic variables Y are changed in a nonlinear way by the relaxation process:

$$Y^* = Y + S(\Phi(W) - Y).$$

The matrix S is invertible, and often chosen as diagonal. It is supposed to be fixed in the asymptotic process presented hereafter. The equilibrium values $Y^{\text{eq}} = \Phi(W)$ are given smooth functions of the conserved variables. When Y^* is evaluated, we have simply

$$f^* = M^{-1} m^*.$$

We introduce the momentum-velocity operator matrix Λ defined by the relation

$$\Lambda_{k\ell} = \sum_{j, \alpha} M_{kj} v_j^\alpha (M^{-1})_{j\ell} \partial_\alpha, \quad 0 \leq k, \ell < q.$$

It is nothing else than the advection operator seen in the space of moments. Then we have an exponential form of the discrete iteration of the lattice Boltzmann scheme:

$$m(x, t + \Delta t) = \exp(-\Delta t \Lambda) m^*(x, t).$$

With this general framework, we follow in this contribution the Chapman-Enskog formalism proposed by Chen-Doolen [CD98] and Qian-Zhou [QZ00]. We suppose that $\Delta t \equiv \varepsilon$ is an infinitesimal parameter and we expand the nonconserved moments as differential nonlinear function of the conserved variables:

$$Y = \Phi(W) + \varepsilon S^{-1} (\Psi_1(W) + \varepsilon^2 \Psi_2(W)) + O(\varepsilon^3).$$

Then we suppose that a multi-scale approach is present for the time dynamics:

$$\partial_t = \partial_{t_1} + \varepsilon \partial_{t_2} + \varepsilon^2 \partial_{t_3} + O(\varepsilon^3).$$

Then we prove that the conserved quantities W follow the following multi-time dynamics:

$$\partial_{t_1} W + \Gamma_1(W) = 0, \quad \partial_{t_2} W + \Gamma_2(W) = 0, \quad \partial_{t_3} W + \Gamma_3(W) = 0.$$

The differential operators $\Gamma_1(W)$, $\Psi_1(W)$, $\Gamma_2(W)$, $\Psi_2(W)$ and $\Gamma_3(W)$ of this expansion are recursively determined as a function of the data v_j , M , $\Phi(W)$ and S . We compare our result with the particular third order expansion proposed in [FD19] and the linear approach presented in [ADGL14]. The previous operators $\Gamma_j(W)$ and $\Psi_i(W)$ are of order j and are exactly the ones derived in our fourth order expansion of lattice Boltzmann schemes with the Taylor expansion method [FD19].

Acknowledgments

The author thanks Bruce Boghosian and Pierre Lallemand for enlightening discussions in Medford (MA, USA) in July 2018 and in Paris (France) in April 2019.

References

- [HSB91] F.J. Higuera, S. Succi, R. Benzi, “Lattice gas dynamics with enhanced collisions”, *Europhysics Letters*, volume 9, issue 4, p. 345-349, 1991.
- [QHL92] Y. H. Qian, D d’Humières, P. Lallemand, “Lattice BGK models for Navier-Stokes equation”, *Europhysics Letters*, volume 17, issue 6, p. 479-484, 1992.
- [DDH92] D. d’Humières, “Generalized Lattice-Boltzmann Equations”, in *Rarefied Gas Dynamics: Theory and Simulations*, volume 159 of *AIAA Progress in Astronautics and Astronautics*, p. 450-458, 1992.
- [LL00] P. Lallemand, L.-S. Luo, “Theory of the Lattice Boltzmann Method: Dispersion, Dissipation, Isotropy, Galilean Invariance, and Stability”, *Physical Review E*, volume 61, 6546, 2000.
- [FD07] F. Dubois, “Une introduction au schéma de Boltzmann sur réseau”, *ESAIM Proceedings*, volume 18, p. 181-215, 2007.
- [FD08] F. Dubois, “Equivalent partial differential equations of a lattice Boltzmann scheme”, *Computers and Mathematics with Applications*, volume 55, p. 1441-1449, 2008.
- [DL10] F. Dubois, P. Lallemand, “Towards higher order lattice Boltzmann schemes”, *Journal of Statistical Mechanics: Theory and Experiment*, P06006, 2009.

FRANÇOIS DUBOIS

- [DL11] F. Dubois, P. Lallemand, “Quartic Parameters for Acoustic Applications of Lattice Boltzmann Scheme”, *Computers and Mathematics with Applications*, volume 61, issue 12, p. 3404-3416, 2011.
- [FD09] F. Dubois, “Third order equivalent equation of lattice Boltzmann scheme”, *Discrete and Continuous Dynamical Systems-Series A*, volume 23, issue 1/2, p. 221-248, 2009.
- [ADGL14] A. Augier, F. Dubois, B. Graille et P. Lallemand, “On rotational invariance of Lattice Boltzmann schemes”, *Computers and Mathematics with Applications*, volume 67, issue 2, p. 239-255, 2014.
- [CD98] S. Chen, G. D. Doolen, “Lattice Boltzmann Method for Fluid Flows”, *Annual Review of Fluid Mechanics*, vol. **30**, p. 329-364, 1998.
- [QZ00] Y. H. Qian, Y. Zhou, “On Higher Order Dynamics in Lattice-based Models Using Chapman-Enskog Method”, *Physical Review E*, 61(2):2103-6, 2000.
- [FD19] F. Dubois, “Nonlinear fourth order Taylor expansion of lattice Boltzmann schemes”, *Hal archives ouvertes hal-02081116 and arXiv 1903.12417*, march 2019 ; submitted for publication.

General third order Chapman-Enskog expansion of lattice Boltzmann schemes

François Dubois^{*†‡}


ICMMES 2019 conference
Edinburgh University
23 July 2019

* Conservatoire National des Arts et Métiers, LMSSC, Paris


† Department of Mathematics, University Paris-Sud, Orsay, France

‡ Thanks to Bruce Boghosian and Pierre Lallemand for enlightening discussions!

Lattice Boltzmann scheme



advection



collision

advection

Outlook

- Multiple Relaxation Times Lattice Boltzmann scheme
- Compact expression
- A general expansion result at third order
- Proof for Chapman-Enskog
- Proof for Taylor expansion method
- Link with the previous expansion at third-order
(DCDS-A, 2009)
- Survey

Multiple Relaxation Times Lattice Boltzmann scheme

Two steps for one time iteration

(i) Nonlinear relaxation

the particle distribution f is modified locally
into a new distribution f^*

(ii) Linear advection

method of characteristics when it is exact !

Compact description of the lattice Boltzmann scheme:

$$f_j(x, t + \Delta t) = f_j^*(x - v_j \Delta t, t), \quad v_j \in \mathcal{V}, \quad x \in \mathcal{L}^0.$$

d'Humières matrix M for moments: $m_k \equiv M_{k\ell} f_\ell$, $m \equiv \begin{pmatrix} W \\ Y \end{pmatrix}$

Conserved variables W ; after relaxation, $W^* = W$

Non-Conserved moments Y

equilibrium value: $Y^{\text{eq}} = \Phi(W)$

after relaxation: $Y^* = Y + S(Y^{\text{eq}} - Y)$, $m^* = \begin{pmatrix} W \\ Y^* \end{pmatrix}$

Important hypotheses

the discrete function $f(x, t)$ for
 x vertex of the lattice
 t discrete time

is supposed to be the restriction of a very regular function
denoted in the same way $f(x, t, \Delta t, s_k, \dots)$


but defined for

- x a point of the continuous space \mathbb{R}^d
- t continuous time
- Δt the time step
- s_k the relaxation parameters

The numerical velocity $\lambda \equiv \frac{\Delta x}{\Delta t}$ is fixed

The relaxation parameters s_k are fixed

Example of D2Q9



$$M = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & \lambda & 0 & -\lambda & 0 & \lambda & -\lambda & -\lambda & \lambda \\ 0 & 0 & \lambda & 0 & -\lambda & \lambda & \lambda & -\lambda & -\lambda \\ -4\lambda^2 & -\lambda^2 & -\lambda^2 & -\lambda^2 & -\lambda^2 & 2\lambda^2 & 2\lambda^2 & 2\lambda^2 & 2\lambda^2 \\ 0 & \lambda^2 & -\lambda^2 & \lambda^2 & -\lambda^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \lambda^2 & -\lambda^2 & \lambda^2 & -\lambda^2 \\ 0 & -2\lambda^3 & 0 & 2\lambda^3 & 0 & \lambda^3 & -\lambda^3 & -\lambda^3 & \lambda^3 \\ 0 & 0 & -2\lambda^3 & 0 & 2\lambda^3 & \lambda^3 & \lambda^3 & -\lambda^3 & -\lambda^3 \\ 4\lambda^4 & -2\lambda^4 & -2\lambda^4 & -2\lambda^4 & -2\lambda^4 & \lambda^4 & \lambda^4 & \lambda^4 & \lambda^4 \end{bmatrix} \begin{array}{l} \rho \\ J_x \\ J_y \\ \varepsilon \\ XX \\ XY \\ q_x \\ q_y \\ h \end{array}$$

Advection operator in the basis of moments

Momentum-velocity operator matrix $\Lambda \equiv M \text{diag} \left(\sum_{\alpha} v^{\alpha} \partial_{\alpha} \right) M^{-1}$
 $1 \leq \alpha \leq d = \text{space dimension}$

Block decomposition $\Lambda \equiv \begin{pmatrix} A & B \\ C & D \end{pmatrix}$

Example of Λ operator matrix for the thermal D2Q9 scheme

0	∂_x	∂_y	0	0	0	0	0
$\frac{2\lambda^2}{3} \partial_x$	0	0	$\frac{1}{6} \partial_x$	$\frac{1}{2} \partial_x$	∂_y	0	0
$\frac{2\lambda^2}{3} \partial_y$	0	0	$\frac{1}{6} \partial_y$	$-\frac{1}{2} \partial_y$	∂_x	0	0
0	$\lambda^2 \partial_x$	$\lambda^2 \partial_y$	0	0	0	∂_x	∂_y
0	$\frac{\lambda^2}{3} \partial_x$	$-\frac{\lambda^2}{3} \partial_y$	0	0	0	$-\frac{1}{3} \partial_x$	$\frac{1}{3} \partial_y$
0	$\frac{2}{3} \lambda^2 \partial_y$	$\frac{2}{3} \lambda^2 \partial_x$	0	0	0	$\frac{1}{3} \partial_y$	$\frac{1}{3} \partial_x$
0	0	0	$\frac{\lambda^2}{3} \partial_x$	$-\lambda^2 \partial_x$	$\lambda^2 \partial_y$	0	0
0	0	0	$\frac{\lambda^2}{3} \partial_y$	$\lambda^2 \partial_y$	$\lambda^2 \partial_x$	0	0
0	0	0	0	0	0	$\lambda^2 \partial_x$	$\lambda^2 \partial_y$

Compact expression of the lattice Boltzmann scheme

Proposition

$$m(x, t + \varepsilon) = \exp(-\varepsilon \Lambda) m^*(x, t)$$

$$\begin{aligned}
 m_k(x, t + \varepsilon) &= \sum_j M_{kj} f_j^*(x - v_j \varepsilon, t) \\
 &= \sum_{j \ell} M_{kj} (M^{-1})_{j\ell} m_{\ell}^*(x - v_j \varepsilon, t) \\
 &= \sum_{j \ell} M_{kj} (M^{-1})_{j\ell} \sum_{n=0}^{\infty} \frac{1}{n!} \left(-\varepsilon \sum_{\alpha} v_j^{\alpha} \partial_{\alpha} \right)^n m_{\ell}^*(x, t) \\
 &= \sum_{\ell} \sum_{n=0}^{\infty} \frac{1}{n!} \sum_j M_{kj} \left(-\varepsilon \sum_{\alpha} v_j^{\alpha} \partial_{\alpha} \right)^n (M^{-1})_{j\ell} m_{\ell}^*(x, t) \\
 &= \sum_{\ell} \left[\sum_{n=0}^{\infty} \frac{1}{n!} (-\varepsilon \Lambda)_{k\ell}^n \right] m_{\ell}^*(x, t) \\
 &= \sum_{\ell} \exp(-\varepsilon \Lambda)_{k\ell} m_{\ell}^*(x, t) \\
 &= \left(\exp(-\varepsilon \Lambda) m^*(x, t) \right)_k
 \end{aligned}$$

Chapman-Enskog framework

Introduce a small parameter ε . For numerical schemes, $\varepsilon = \Delta t$

$$f = f^{\text{eq}} + \varepsilon f^1 + \varepsilon^2 f^2 + O(\varepsilon^3)$$

Chapman-Enskog hypothesis : the perturbation terms f^ℓ
are only function of the equilibrium $f = f^{\text{eq}}$
apply the d'Humières matrix:

$$Mf = Mf^{\text{eq}} + \varepsilon Mf^1 + \varepsilon^2 Mf^2 + O(\varepsilon^3)$$

take the first conserved component : $W = W + 0$

take the second nonconserved component :

$$Y = Y^{\text{eq}} + \varepsilon (Mf^1)_Y + \varepsilon^2 (Mf^2)_Y + O(\varepsilon^3)$$

The perturbation terms $\varepsilon^\ell (Mf^\ell)_Y$ depend only
on the conserved moments W .

Then $Y = \Phi(W) + S^{-1} (\varepsilon \Psi_1(W) + \varepsilon^2 \Psi_2(W)) + O(\varepsilon^3)$

Suppose also a multi-scale approach for the time dynamics:

$$\partial_t = \partial_{t_1} + \varepsilon \partial_{t_2} + \varepsilon^2 \partial_{t_3} + O(\varepsilon^3)$$

Chapman-Enskog expansion

The conserved quantities W follow a multi-time dynamics :

$$\begin{aligned}\partial_{t_1} W + \Gamma_1(W) &= 0 \\ \partial_{t_2} W + \Gamma_2(W) &= 0 \\ \partial_{t_3} W + \Gamma_3(W) &= 0\end{aligned}$$

The differential operators

$\Gamma_1(W)$, $\Psi_1(W)$, $\Gamma_2(W)$, $\Psi_2(W)$ and $\Gamma_3(W)$
of this expansion are recursively determined

coefficients at first order

$$\Gamma_1 = A W + B \Phi(W)$$

coefficients at second order

$$\Psi_1 = d\Phi(W). \Gamma_1 - (C W + D \Phi(W))$$

$$\Gamma_2 = B \Sigma \Psi_1 \quad \text{Hénon matrix } \Sigma \equiv S^{-1} - \frac{1}{2} I$$

coefficients at third order

$$\Psi_2 = \Sigma d\Psi_1. \Gamma_1 + d\Phi(W). \Gamma_2 - D \Sigma \Psi_1$$

$$\Gamma_3 = B \Sigma \Psi_2 + \frac{1}{12} B_2 \Psi_1 - \frac{1}{6} B d\Psi_1. \Gamma_1$$

Taylor vs Chapman-Enskog expansion methods

Asymptotic hypothesis: emerging partial differential equations

$$\partial_t W + \Gamma_1 + \Delta t \Gamma_2 + \Delta t^2 \Gamma_3 = O(\Delta t^3)$$

Γ_j : vector obtained after j space derivations
of the conserved moments W
and the equilibrium vector $\Phi(W)$.

Non-Conserved moments:

$$Y = \Phi(W) + S^{-1} (\Delta t \Psi_1 + \Delta t^2 \Psi_2) O(\Delta t^2)$$

Φ_j analogous to Γ_j but not with the same dimension!

General nonlinear Taylor expansion method

Coefficients at first order

$$\Gamma_1 = A W + B \Phi(W)$$

Coefficients at second order

$$\Psi_1 = d\Phi(W) \cdot \Gamma_1 - (C W + D \Phi(W))$$

$$\Gamma_2 = B \Sigma \Psi_1 \quad \text{with } \Sigma \text{ the Hénon matrix: } \Sigma \equiv S^{-1} - \frac{1}{2} I$$

Coefficients at third order

$$\Psi_2 = d\Phi(W) \cdot \Gamma_2 + \Sigma d\Psi_1 \cdot \Gamma_1 - D \Sigma \Psi_1$$

$$\Gamma_3 = B \Sigma \Psi_2 + \frac{1}{6} B d\Psi_1 \cdot \Gamma_1 - \frac{1}{12} B_2 \Psi_1$$

We find the **same formulas** than with Chapman-Enskog:

the two expansions are equivalent!

Chapman-Enskog expansion : order zero

$$m(t + \varepsilon) = \exp(-\varepsilon \Lambda) m^*$$

$$m + O(\varepsilon) = m^* + O(\varepsilon)$$

first component : $W + O(\varepsilon) = W^* + O(\varepsilon)$

no more information because $W^* = W$

second component : $Y + O(\varepsilon) = Y^* + O(\varepsilon)$

relaxation: $Y^* = Y + S(\Phi(W) - Y)$

The matrix S is supposed fixed

then $Y = \Phi(W) + O(\varepsilon)$

and $Y^* = \Phi(W) + O(\varepsilon)$

Chapman-Enskog expansion : order one

$$m(t + \varepsilon) = \exp(-\varepsilon \Lambda) m^*$$

$$m + \varepsilon \partial_t m + O(\varepsilon^2) = m^* - \varepsilon \Lambda m^* + O(\varepsilon^2)$$

with $\partial_t = \partial_{t_1} + \varepsilon \partial_{t_2} + \varepsilon^2 \partial_{t_3} + O(\varepsilon^3)$

then $m + \varepsilon \partial_{t_1} m + O(\varepsilon^2) = m^* - \varepsilon \Lambda m^* + O(\varepsilon^2)$

first component

$$W + \varepsilon \partial_{t_1} W + O(\varepsilon^2) = W^* - \varepsilon (A W + B Y^*) + O(\varepsilon^2)$$

with $W^* = W$ and $Y^* = \Phi(W) + O(\varepsilon)$

then $\partial_{t_1} W = -(A W + B \Phi(W))$

$\partial_{t_1} W + \Gamma_1(W) = 0$ with $\Gamma_1 = A W + B \Phi(W)$

Chapman-Enskog expansion : order one bis

second component of $m + \varepsilon \partial_{t_1} m + O(\varepsilon^2) = m^* - \varepsilon \Lambda m^* + O(\varepsilon^2)$

$$Y + \varepsilon \partial_{t_1} Y + O(\varepsilon^2) = Y^* - \varepsilon (C W + D Y^*) + O(\varepsilon^2)$$

known: $Y^* = Y + S(\Phi(W) - Y)$
 $Y = \Phi(W) + O(\varepsilon)$ and $Y^* = \Phi(W) + O(\varepsilon)$

$$S(Y - \Phi(W)) = Y - Y^*$$

$$\begin{aligned} &= -\varepsilon \partial_{t_1} (\Phi(W) + O(\varepsilon)) - \varepsilon (C W + D (\Phi(W) + O(\varepsilon))) + O(\varepsilon^2) \\ &= \varepsilon [- d\Phi(W) \cdot \partial_{t_1} W - (C W + D \Phi(W))] + O(\varepsilon^2) \\ &= \varepsilon [d\Phi(W) \cdot \Gamma_1 - (C W + D \Phi(W))] + O(\varepsilon^2) \end{aligned}$$

Then $Y = \Phi(W) + \varepsilon S^{-1} \Psi_1(W) + O(\varepsilon^2)$

$$\Psi_1 = d\Phi(W) \cdot \Gamma_1 - (C W + D \Phi(W))$$

$$Y^* = \Phi(W) + \varepsilon (S^{-1} - I) \Psi_1(W) + O(\varepsilon^2)$$

Chapman-Enskog expansion : Hénon's matrix

$$\Sigma = S^{-1} - \frac{1}{2} I$$

$$Y = \Phi(W) + \varepsilon S^{-1} \Psi_1(W) + O(\varepsilon^2)$$

Then $Y = \Phi(W) + \varepsilon (\Sigma + \frac{1}{2} I) \Psi_1(W) + O(\varepsilon^2)$

$$Y^* = \Phi(W) + \varepsilon (\Sigma - \frac{1}{2} I) \Psi_1(W) + O(\varepsilon^2)$$

$$\Psi_1 = d\Phi(W) \cdot \Gamma_1 - (C W + D \Phi(W))$$

Chapman-Enskog expansion : order two

$$m + \varepsilon \partial_t m + \frac{1}{2} \varepsilon^2 \partial_t^2 m + O(\varepsilon^3) = m^* - \varepsilon \Lambda m^* + \frac{1}{2} \varepsilon^2 \Lambda^2 m^* + O(\varepsilon^3)$$

then

$$\begin{aligned} m + \varepsilon (\partial_{t_1} + \varepsilon \partial_{t_2}) m + \frac{1}{2} \varepsilon^2 (\partial_{t_1} + O(\varepsilon))^2 m + O(\varepsilon^3) \\ = m^* - \varepsilon \Lambda m^* + \frac{1}{2} \varepsilon^2 \Lambda^2 m^* + O(\varepsilon^3) \end{aligned}$$

first component

$$\begin{aligned} W + \varepsilon (\partial_{t_1} + \varepsilon \partial_{t_2}) W + \frac{1}{2} \varepsilon^2 \partial_{t_1}^2 W + O(\varepsilon^3) \\ = W - \varepsilon (A W + B Y^*) + \frac{1}{2} \varepsilon^2 (A_2 W + B_2 Y^*) + O(\varepsilon^3) \\ \text{with } Y^* = \Phi(W) + \varepsilon (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) + O(\varepsilon^2) \end{aligned}$$

second order term

$$\partial_{t_2} W + \frac{1}{2} \partial_{t_1}^2 W = -B (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) + \frac{1}{2} (A_2 W + B_2 \Phi)$$

Chapman-Enskog expansion : order two (ii)

$$\partial_{t_2} W + \frac{1}{2} \partial_{t_1}^2 W = -B \Sigma \Psi_1 + \frac{1}{2} B \Psi_1 + \frac{1}{2} (A_2 W + B_2 \Phi)$$

$$\begin{aligned} \text{with } \partial_{t_1}^2 W &= -\partial_{t_1} (\Gamma_1(W)) = -\partial_{t_1} (A W + B \Phi(W)) \\ &= A \Gamma_1 + B d\Phi(W). \Gamma_1 = A (A W + B \Phi) + B d\Phi(W). \Gamma_1 \end{aligned}$$

$$\begin{aligned} \partial_{t_2} W + \frac{1}{2} (A^2 W + AB \Phi) + \frac{1}{2} B d\Phi(W). \Gamma_1 \\ = -B \Sigma \Psi_1 + \frac{1}{2} B \Psi_1 + \frac{1}{2} (A_2 W + B_2 \Phi) \\ \text{with } A_2 = A^2 + BC, \quad B_2 = AB + BD \end{aligned}$$

$$\begin{aligned} \partial_{t_2} W + B \Sigma \Psi_1 &= -\frac{1}{2} (A^2 W + AB \Phi) - \frac{1}{2} B d\Phi(W). \Gamma_1 \\ &\quad + \frac{1}{2} B \Psi_1 + \frac{1}{2} (A^2 + BC) W + \frac{1}{2} (AB + BD) \Phi \\ &= -\frac{1}{2} B d\Phi(W). \Gamma_1 + \frac{1}{2} B (d\Phi(W). \Gamma_1 - CW - D\Phi) \\ &\quad + \frac{1}{2} BCW + \frac{1}{2} BD\Phi \\ &= 0 \\ \partial_{t_2} W + \Gamma_2(W) &= 0 \quad \text{with } \Gamma_2(W) = B \Sigma \Psi_1(W) \end{aligned}$$

Chapman-Enskog expansion : order two bis

$$\begin{aligned} m + \varepsilon (\partial_{t_1} + \varepsilon \partial_{t_2})m + \frac{1}{2} \varepsilon^2 \partial_{t_1}^2 m + O(\varepsilon^3) \\ = m^* - \varepsilon \Lambda m^* + \frac{1}{2} \varepsilon^2 \Lambda^2 m^* + O(\varepsilon^3) \end{aligned}$$

second component

$$\begin{aligned} Y + \varepsilon \partial_{t_1} Y + \varepsilon^2 \partial_{t_2} Y + \frac{1}{2} \varepsilon^2 \partial_{t_1}^2 Y + O(\varepsilon^3) \\ = Y^* - \varepsilon (C W + D Y^*) + \frac{1}{2} \varepsilon^2 (C_2 W + D_2 Y^*) + O(\varepsilon^3) \end{aligned}$$

$$S(Y - \Phi(W)) = Y - Y^*$$

$$\begin{aligned} &= -\varepsilon \partial_{t_1} Y - \varepsilon^2 (\partial_{t_2} Y + \frac{1}{2} \partial_{t_1}^2 Y) - \varepsilon (C W + D Y^*) \\ &\quad + \frac{1}{2} \varepsilon^2 (C_2 W + D_2 Y^*) + O(\varepsilon^3) \\ \text{with } &Y = \Phi(W) + \varepsilon (\Sigma \Psi_1 + \frac{1}{2} \Psi_1) + O(\varepsilon^2) \end{aligned}$$

$$\begin{aligned} &= -\varepsilon \partial_{t_1} \Phi(W) - \varepsilon^2 (\partial_{t_1} (\Sigma \Psi_1 + \frac{1}{2} \Psi_1) + \partial_{t_2} \Phi(W) + \frac{1}{2} \partial_{t_1}^2 \Phi(W)) \\ &\quad - \varepsilon (C W + D Y^*) + \frac{1}{2} \varepsilon^2 (C_2 W + D_2 Y^*) + O(\varepsilon^3) \end{aligned}$$

Chapman-Enskog expansion : order two bis (ii)

$$\begin{aligned} S(Y - \Phi(W)) &= -\varepsilon \partial_{t_1} \Phi(W) \\ &\quad - \varepsilon^2 (\partial_{t_1} (\Sigma \Psi_1 + \frac{1}{2} \Psi_1) + \partial_{t_2} \Phi(W) + \frac{1}{2} \partial_{t_1}^2 \Phi(W)) \\ &\quad - \varepsilon (C W + D Y^*) + \frac{1}{2} \varepsilon^2 (C_2 W + D_2 Y^*) + O(\varepsilon^3) \\ \text{with } &Y^* = \Phi(W) + \varepsilon (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) + O(\varepsilon^2) \end{aligned}$$

we have by definition $S(Y - \Phi(W)) = \varepsilon \Psi_1 + \varepsilon^2 \Psi_2 + O(\varepsilon^3)$

second order term

$$\begin{aligned} \Psi_2 &= -\Sigma \partial_{t_1} \Psi_1 - \frac{1}{2} \partial_{t_1} \Psi_1 - \partial_{t_2} \Phi(W) - \frac{1}{2} \partial_{t_1}^2 \Phi(W) \\ &\quad - D (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) + \frac{1}{2} C_2 W + \frac{1}{2} D_2 \Phi(W) \end{aligned}$$

$$\text{with } \partial_{t_1} \Psi_1 = d\Psi_1 \cdot \partial_{t_1} W = -d\Psi_1 \cdot \Gamma_1$$

$$\begin{aligned} \text{and } \partial_{t_1} \Psi_1 &= \partial_{t_1} (d\Phi \cdot \Gamma_1 - C W - D \Phi(W)) \\ &= \partial_{t_1} (d\Phi \cdot \Gamma_1) - C \partial_{t_1} W - D d\Phi \cdot \partial_{t_1} W \\ &= \partial_{t_1} (d\Phi \cdot \Gamma_1) + C \Gamma_1 + D d\Phi \cdot \Gamma_1 \end{aligned}$$

$$\begin{aligned} \Psi_2 &= \Sigma d\Psi_1 \cdot \Gamma_1 - \frac{1}{2} (\partial_{t_1} (d\Phi \cdot \Gamma_1) + C \Gamma_1 + D d\Phi \cdot \Gamma_1) - \partial_{t_2} \Phi(W) \\ &\quad - \frac{1}{2} \partial_{t_1}^2 \Phi(W) - D (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) + \frac{1}{2} C_2 W + \frac{1}{2} D_2 \Phi(W) \end{aligned}$$

Chapman-Enskog expansion : order two bis (iii)

$$\begin{aligned}\Psi_2 = \sum d\Psi_1 \cdot \Gamma_1 - \frac{1}{2} (\partial_{t_1}(d\Phi \cdot \Gamma_1) + C \Gamma_1 + D d\Phi \cdot \Gamma_1) - \partial_{t_2} \Phi(W) \\ - \frac{1}{2} \partial_{t_1}^2 \Phi(W) - D (\sum \Psi_1 - \frac{1}{2} \Psi_1) + \frac{1}{2} C_2 W + \frac{1}{2} D_2 \Phi(W) \\ \text{with } \partial_{t_2} \Phi(W) = d\Phi(W) \cdot \partial_{t_2} W = -\Phi(W) \cdot \Gamma_2\end{aligned}$$

$$\partial_{t_1}^2 \Phi(W) = \partial_{t_1} (\partial_{t_1} \Phi(W)) = \partial_{t_1} (d\Phi \cdot \partial_{t_1} W) = -\partial_{t_1} (d\Phi \cdot \Gamma_1)$$

$$\begin{aligned}\Psi_2 = \sum d\Psi_1 \cdot \Gamma_1 - \frac{1}{2} (\partial_{t_1}(d\Phi \cdot \Gamma_1) + C \Gamma_1 + D d\Phi \cdot \Gamma_1) + \Phi(W) \cdot \Gamma_2 \\ + \frac{1}{2} \partial_{t_1}(d\Phi \cdot \Gamma_1) - D \sum \Psi_1 + \frac{1}{2} D \Psi_1 + \frac{1}{2} C_2 W + \frac{1}{2} D_2 \Phi(W) \\ \text{with } C_2 = CA + BD, \quad D_2 = CB + D^2\end{aligned}$$

$$\begin{aligned}\Psi_2 = \sum d\Psi_1 \cdot \Gamma_1 - \frac{1}{2} (C \Gamma_1 + D d\Phi \cdot \Gamma_1) + \Phi(W) \cdot \Gamma_2 - D \sum \Psi_1 + \frac{1}{2} D \Psi_1 \\ + \frac{1}{2} C(AW + B\Phi) + \frac{1}{2} D(CW + B\Phi) \\ \text{with } CW + B\Phi = d\Phi \cdot \Gamma_1 - \Psi_1\end{aligned}$$

$$\begin{aligned}\Psi_2 = \sum d\Psi_1 \cdot \Gamma_1 - \frac{1}{2} D d\Phi \cdot \Gamma_1 + \Phi(W) \cdot \Gamma_2 - D \sum \Psi_1 + \frac{1}{2} D \Psi_1 \\ + \frac{1}{2} D(d\Phi \cdot \Gamma_1 - \Psi_1)\end{aligned}$$

$$\Psi_2 = \sum d\Psi_1 \cdot \Gamma_1 + d\Phi(W) \cdot \Gamma_2 - D \sum \Psi_1$$

Chapman-Enskog expansion : order three

$$\begin{aligned}m + \varepsilon \partial_t m + \frac{1}{2} \varepsilon^2 \partial_t^2 m + \frac{1}{6} \varepsilon^3 \partial_t^3 m + O(\varepsilon^4) \\ = m^* - \varepsilon \Lambda m^* + \frac{1}{2} \varepsilon^2 \Lambda^2 m^* - \frac{1}{6} \varepsilon^3 \Lambda^3 m^* + O(\varepsilon^4)\end{aligned}$$

then

$$\begin{aligned}m + \varepsilon (\partial_{t_1} + \varepsilon \partial_{t_2} + \varepsilon^2 \partial_{t_3}) m + \frac{1}{2} \varepsilon^2 (\partial_{t_1} + \varepsilon \partial_{t_2} + O(\varepsilon^2))^2 m \\ + \frac{1}{6} \varepsilon^3 (\partial_{t_1} + O(\varepsilon))^3 m + O(\varepsilon^4) = m^* - \varepsilon \Lambda m^* \\ + \frac{1}{2} \varepsilon^2 \Lambda^2 m^* - \frac{1}{6} \varepsilon^3 \Lambda^3 m^* + O(\varepsilon^4)\end{aligned}$$

first component

$$\begin{aligned}W + \varepsilon (\partial_{t_1} + \varepsilon \partial_{t_2} + \varepsilon^2 \partial_{t_3}) W + \frac{1}{2} \varepsilon^2 (\partial_{t_1}^2 + \varepsilon \partial_{t_1} \partial_{t_2} + \varepsilon \partial_{t_2} \partial_{t_1}) W \\ + \frac{1}{6} \varepsilon^3 \partial_{t_1}^3 W + O(\varepsilon^4) = W - \varepsilon (AW + B\textcolor{blue}{Y}^*) \\ + \frac{1}{2} \varepsilon^2 (A_2 W + B_2 \textcolor{blue}{Y}^*) - \frac{1}{6} \varepsilon^3 (A_3 W + B_3 \textcolor{blue}{Y}^*) + O(\varepsilon^4)\end{aligned}$$

$$\text{with } \textcolor{blue}{Y}^* = \Phi(W) + \varepsilon (\sum \Psi_1 - \frac{1}{2} \Psi_1) + \varepsilon^2 (\sum \Psi_2 - \frac{1}{2} \Psi_2) + O(\varepsilon^3)$$

Chapman-Enskog expansion : order three (ii)

$$\begin{aligned}
& W + \varepsilon (\partial_{t_1} + \varepsilon \partial_{t_2} + \varepsilon^2 \partial_{t_3}) W + \frac{1}{2} \varepsilon^2 (\partial_{t_1}^2 + \varepsilon \partial_{t_1} \partial_{t_2} + \varepsilon \partial_{t_2} \partial_{t_1}) W \\
& + \frac{1}{6} \varepsilon^3 \partial_{t_1}^3 W + O(\varepsilon^4) = W - \varepsilon A W \\
& - \varepsilon B (\Phi(W) + \varepsilon (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) + \varepsilon^2 (\Sigma \Psi_2 - \frac{1}{2} \Psi_2)) + \frac{1}{2} \varepsilon^2 A_2 W \\
& + \frac{1}{2} \varepsilon^2 B_2 (\Phi(W) + \varepsilon (\Sigma \Psi_1 - \frac{1}{2} \Psi_1)) - \frac{1}{6} \varepsilon^3 (A_3 W + B_3 \Phi) + O(\varepsilon^4)
\end{aligned}$$

third order term

$$\begin{aligned}
& \partial_{t_3} W + \frac{1}{2} (\partial_{t_1} \partial_{t_2} W + \partial_{t_2} \partial_{t_1} W) + \frac{1}{6} \partial_{t_1}^3 W \\
& = -B (\Sigma \Psi_2 - \frac{1}{2} \Psi_2) + \frac{1}{2} B_2 (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) - \frac{1}{6} (A_3 W + B_3 \Phi)
\end{aligned}$$

with

$$\begin{aligned}
\partial_{t_1} \partial_{t_2} W &= \partial_{t_1} (-B \Sigma \Psi_1) = -B \Sigma d\Psi_1 \cdot \partial_{t_1} W = B \Sigma d\Psi_1 \cdot \Gamma_1 \\
\partial_{t_2} \partial_{t_1} W &= \partial_{t_2} (-A W - B \Phi) = A \Gamma_2 - B d\Phi \cdot \partial_{t_2} W \\
&= AB \Sigma \Gamma_1 + B d\Phi \cdot \Gamma_2
\end{aligned}$$

$$\begin{aligned}
& \partial_{t_3} W + \frac{1}{2} B \Sigma d\Psi_1 \cdot \Gamma_1 + \frac{1}{2} (AB \Sigma \Gamma_1 + B d\Phi \cdot \Gamma_2) + \frac{1}{6} \partial_{t_1}^3 W \\
& = -B (\Sigma \Psi_2 - \frac{1}{2} \Psi_2) + \frac{1}{2} B_2 (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) - \frac{1}{6} (A_3 W + B_3 \Phi)
\end{aligned}$$

Chapman-Enskog expansion : order three (iii)

$$\begin{aligned}
& \partial_{t_3} W + \frac{1}{2} (B \Sigma d\Psi_1 \cdot \Gamma_1 + AB \Sigma \Gamma_1 + B d\Phi \cdot \Gamma_2) + \frac{1}{6} \partial_{t_1}^3 W \\
& = -B (\Sigma \Psi_2 - \frac{1}{2} \Psi_2) + \frac{1}{2} B_2 (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) - \frac{1}{6} (A_3 W + B_3 \Phi)
\end{aligned}$$

with

$$\begin{aligned}
\partial_{t_1}^3 W &= \partial_{t_1} (A \Gamma_1 + B d\Phi \cdot \Gamma_1) = \partial_{t_1} (A (A W + B \Phi) + B d\Phi \cdot \Gamma_1) \\
&= -A (A \Gamma_1 - B d\Phi \cdot \Gamma_1) - B d(d\Phi \cdot \Gamma_1) \cdot \Gamma_1 \\
&= -A^2 \Gamma_1 - AB d\Phi \cdot \Gamma_1 - B d(d\Phi \cdot \Gamma_1) \cdot \Gamma_1
\end{aligned}$$

$$\text{and } A_3 = A_2 A + B_2 C, \quad B_3 = A_2 B + B_2 D$$

$$\begin{aligned}
A_3 W + B_3 \Phi &= A_2 (A W + B \Phi) + B_2 (C W + D \Phi) \\
&= (A^2 + B C) \Gamma_1 + (AB + BD) (d\Phi \cdot \Gamma_1 - \Psi_1) \\
&= A (A \Gamma_1 + B d\Phi \cdot \Gamma_1) + B (C \Gamma_1 + D d\Phi \cdot \Gamma_1) - B_2 \Psi_1 \\
&= A (A \Gamma_1 + B d\Phi \cdot \Gamma_1) + B (d(d\Phi \cdot \Gamma_1) \cdot \Gamma_1 - d\Psi_1 \cdot \Gamma_1) - B_2 \Psi_1 \\
&= -\partial_{t_1}^3 W - B d\Psi_1 \cdot \Gamma_1 - B_2 \Psi_1
\end{aligned}$$

Chapman-Enskog expansion : order three (iv)

$$\begin{aligned}\partial_{t_3} W + \frac{1}{2} (B \Sigma d\Psi_1 \cdot \Gamma_1 + AB \Sigma \Gamma_1 + B d\Phi \cdot \Gamma_2) &= -B \Sigma \Psi_2 \\ &+ \frac{1}{2} B \Psi_2 + \frac{1}{2} B_2 (\Sigma \Psi_1 - \frac{1}{2} \Psi_1) + \frac{1}{6} (B d\Psi_1 \cdot \Gamma_1 + B_2 \Psi_1)\end{aligned}$$

with

$$\Psi_2 = \Sigma d\Psi_1 \cdot \Gamma_1 + d\Phi \cdot \Gamma_2 - D \Sigma \Psi_1$$

$$\begin{aligned}\partial_{t_3} W + \frac{1}{2} (B \Sigma d\Psi_1 \cdot \Gamma_1 + AB \Sigma \Gamma_1 + B d\Phi \cdot \Gamma_2) &= -B \Sigma \Psi_2 \\ &+ \frac{1}{2} B (\Sigma d\Psi_1 \cdot \Gamma_1 + d\Phi \cdot \Gamma_2 - D \Sigma \Psi_1) + \frac{1}{2} B_2 \Sigma \Psi_1 \\ &- \left(\frac{1}{4} - \frac{1}{6}\right) B_2 \Psi_1 + \frac{1}{6} B d\Psi_1 \cdot \Gamma_1\end{aligned}$$

then $\partial_{t_3} W + B \Sigma \Psi_2 + \frac{1}{12} B_2 \Psi_1 - \frac{1}{6} B d\Psi_1 \cdot \Gamma_1 = 0$

$$\Gamma_3 = B \Sigma \Psi_2 + \frac{1}{12} B_2 \Psi_1 - \frac{1}{6} B d\Psi_1 \cdot \Gamma_1$$

Taylor second-order expansion

$$\begin{aligned}m + \Delta t \partial_t m + \frac{1}{2} \Delta t^2 \partial_t^2 m + O(\Delta t^3) &= \\ &= m^* - \Delta t \Lambda m^* + \frac{1}{2} \Delta t^2 \Lambda^2 m^* + O(\Delta t^3)\end{aligned}$$

We replace the vector m by its two components W and Y

$$\begin{aligned}W + \Delta t \partial_t W + \frac{1}{2} \Delta t^2 \partial_t^2 W + O(\Delta t^3) &= \\ &= W - \Delta t (A W + B Y^*) + \frac{1}{2} \Delta t^2 (A_2 W + B_2 Y^*) + O(\Delta t^3)\end{aligned}$$

$$\begin{aligned}Y + \Delta t \partial_t Y + \frac{1}{2} \Delta t^2 \partial_t^2 Y + O(\Delta t^3) &= \\ &= Y^* - \Delta t (C W + D Y^*) + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^*) + O(\Delta t^3)\end{aligned}$$

At zero-order: $Y - Y^* = O(\Delta t)$ and $Y^* \equiv Y + S(\Phi(W) - Y)$

The matrix S is supposed fixed

then $Y = \Phi(W) + O(\Delta t)$ and $Y^* = \Phi(W) + O(\Delta t)$

Taylor second-order expansion (ii)

$$Y^* = \Phi(W) + O(\Delta t)$$

Second-order partial differential equations:

$$\begin{aligned}\partial_t W + \frac{1}{2} \Delta t \partial_t^2 W + O(\Delta t^2) &= \\ &= -(A W + B Y^*) + \frac{1}{2} \Delta t (A_2 W + B_2 Y^*) + O(\Delta t^2)\end{aligned}$$

at first-order:

$$\partial_t W + O(\Delta t) = -A W - B \Phi(W) + O(\Delta t)$$

$$\text{then } \partial_t W = -\Gamma_1 \quad \text{with} \quad \Gamma_1 = A W + B \Phi(W)$$

$$\text{then } \partial_t Y = d\Phi(W). \partial_t W + O(\Delta t) = -d\Phi(W). \Gamma_1 + O(\Delta t)$$

$$Y - Y^* = -\Delta t \partial_t Y - \Delta t (C W + D Y^*) + O(\Delta t^2)$$

$$\text{then } S(Y - \Phi) = \Delta t (d\Phi(W). \Gamma_1 - C W - D \Phi) + O(\Delta t^2)$$

$$\begin{aligned}Y &= \Phi + \Delta t S^{-1} (d\Phi(W). \Gamma_1 - C W - D \Phi) + O(\Delta t^2) \\ \text{and } \Psi_1 &= d\Phi(W). \Gamma_1 - (C W + D \Phi)\end{aligned}$$

Taylor second-order expansion (iii)

$$\partial_t W = -\Gamma_1 + O(\Delta t) = -(A W + B \Phi(W)) + O(\Delta t)$$

$$\begin{aligned}\text{Then } \partial_t^2 W &= \partial_t (-\Gamma_1 + O(\Delta t)) = -d\Gamma_1. \partial_t W + O(\Delta t) \\ &= A \Gamma_1 + B d\Phi. \partial_t W + O(\Delta t) \\ &= A \Gamma_1 + B d\Phi. \Gamma_1 + O(\Delta t)\end{aligned}$$

Second-order partial differential equations:

$$\begin{aligned}\partial_t W + \frac{1}{2} \Delta t \partial_t^2 W + O(\Delta t^2) &= \\ &= -(A W + B Y^*) + \frac{1}{2} \Delta t (A_2 W + B_2 Y^*) + O(\Delta t^2)\end{aligned}$$

$$\begin{aligned}\partial_t W &= -A W - B (\Phi + (\Sigma - \frac{1}{2} I) \Delta t \Psi_1) - \frac{1}{2} \Delta t (A \Gamma_1 + B d\Phi. \Gamma_1) \\ &\quad + \frac{1}{2} \Delta t ((A^2 + B C) W + (AB + BD) \Phi) + O(\Delta t^2) \\ &= -A W - B \Phi + \Delta t (-B \Sigma \Psi_1 + \frac{1}{2} B (d\Phi. \Gamma_1 - C W - D \Phi)) \\ &\quad - \frac{1}{2} (A (A W + B \Phi)) - \frac{1}{2} B d\Phi. \Gamma_1 + \frac{1}{2} (A^2 + B C) W \\ &\quad + \frac{1}{2} (AB + BD) \Phi + O(\Delta t^2)\end{aligned}$$

$$\partial_t W = -\Gamma_1 - \Delta t B \Sigma \Psi_1 + O(\Delta t^2) \quad \text{and} \quad \Gamma_2 = B \Sigma \Psi_1$$

Link with the original Taylor expansion method (2007) 29

First-order $\partial_t W = -\Gamma_1 + O(\Delta t) = -A W - B \Phi(W) + O(\Delta t)$

id est $\partial_t W_i + \Lambda_{ij}^\beta \partial_\beta W_j + \Lambda_{i\ell}^\beta \partial_\beta \Phi(W)_\ell = O(\Delta t)$

or $\partial_t W_i + \Lambda_{i\ell}^\beta \partial_\beta m_\ell^{\text{eq}} = O(\Delta t)$

with the notation $m_k^{\text{eq}} \equiv \Phi(W)_k$

Defect of conservation $\theta_k \equiv \partial_t m_k^{\text{eq}} + \sum_{\ell\beta} \Lambda_{k\ell}^\beta \partial_\beta m_\ell^{\text{eq}}$

vector of conservation defect

$$\begin{aligned} \theta &= \partial_t \Phi(W) + \Lambda \cdot m^{\text{eq}} \\ &= d\Phi.(-\Gamma_1 - \Delta t \Gamma_2 + \dots) + (C W + D \Phi(W)) \\ &= (-d\Phi. \Gamma_1 + O(\Delta t)) + (C W + D \Phi(W)) \\ &= -\Psi_1 + O(\Delta t) \end{aligned} \quad \theta = -\Psi_1 + O(\Delta t)$$

Second-order term

$$(B \Sigma \Psi_1)_i = \sum_k B_{ik} \sigma_k (-\theta_k + O(\Delta t)) = \sum_k \Lambda_{ik}^\beta \sigma_k \partial_\beta \theta_k + O(\Delta t)$$

and $\partial_t W_i + \Lambda_{i\ell}^\beta \partial_\beta m_\ell^{\text{eq}} = \Delta t \sum_k \Lambda_{ik}^\beta \sigma_k \partial_\beta \theta_k + O(\Delta t^2)$

Taylor third-order expansion

$$\begin{aligned} m + \Delta t \partial_t m + \frac{1}{2} \Delta t^2 \partial_t^2 m + \frac{1}{6} \Delta t^3 \partial_t^3 m + O(\Delta t^4) &= \\ &= m^* - \Delta t \Lambda m^* + \frac{1}{2} \Delta t^2 \Lambda^2 m^* + \frac{1}{6} \Delta t^3 \Lambda^3 m^* + O(\Delta t^4) \end{aligned}$$

We replace the vector m by its two components W and Y

$$\begin{aligned} W + \Delta t \partial_t W + \frac{1}{2} \Delta t^2 \partial_t^2 W + \frac{1}{6} \Delta t^3 \partial_t^3 W + O(\Delta t^4) &= \\ &= W - \Delta t (A W + B Y^*) + \frac{1}{2} \Delta t^2 (A_2 W + B_2 Y^*) \\ &\quad - \frac{1}{6} \Delta t^3 (A_3 W + B_3 Y^*) + O(\Delta t^4) \end{aligned}$$

then $\partial_t W = -A W - B Y^* + \frac{1}{2} \Delta t (A_2 W + B_2 Y^* - \partial_t^2 W)$
 $\quad - \frac{1}{6} \Delta t^2 (A_3 W + B_3 Y^* - \partial_t^3 W) + O(\Delta t^3)$

$$\begin{aligned} Y + \Delta t \partial_t Y + \frac{1}{2} \Delta t^2 \partial_t^2 Y + O(\Delta t^3) &= \\ &= Y^* - \Delta t (C W + D Y^*) + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^*) + O(\Delta t^3) \end{aligned}$$

and

$$\begin{aligned} Y - Y^* &\equiv S(Y - \Phi) \\ &= -\Delta t (C W + D Y^* + \partial_t Y) + \frac{1}{2} \Delta t^2 (C_2 W + D_2 Y^* - \partial_t^2 Y) + O(\Delta t^3) \end{aligned}$$

Computation of the coefficient Ψ_2

$$\begin{aligned} S(Y - \Phi) = & \\ & -\Delta t(CW + DY^* + \partial_t Y) + \frac{1}{2}\Delta t^2(C_2 W + D_2 Y^* - \partial_t^2 Y) \\ & + O(\Delta t^3) \end{aligned}$$

with [see Annex -1- page (iii)]

$$\begin{aligned} CW + DY^* &= -\Psi_1 + d\Phi.\Gamma_1 + \Delta t D (\Sigma - \frac{1}{2} I) \Psi_1 + O(\Delta t^2) \\ C_2 W + D_2 Y^* &= d(d\Phi.\Gamma_1 - \Psi_1).\Gamma_1 - D \Psi_1 + O(\Delta t) \end{aligned}$$

and [see Annex -2- pages (iii) and (iv)]

$$\begin{aligned} \partial_t Y &= -d\Phi.\Gamma_1 - \Delta t \left(d\Phi.\Gamma_2 + (\Sigma + \frac{1}{2} I) d\Psi_1.\Gamma_1 \right) + O(\Delta t^2) \\ \partial_t^2 Y &= d(d\Phi.\Gamma_1).\Gamma_1 + O(\Delta t) \end{aligned}$$

Then

$$\begin{aligned} S(Y - \Phi) &= \Delta t \Psi_1 + \Delta t^2 \left(-D (\Sigma - \frac{1}{2} I) \Psi_1 + d\Phi.\Gamma_2 \right. \\ &\quad \left. + (\Sigma + \frac{1}{2} I) d\Psi_1.\Gamma_1 - \frac{1}{2} (d\Psi_1.\Gamma_1 + D\Psi_1) \right) + O(\Delta t^3) \\ &= \Delta t \Psi_1 + \Delta t^2 (\Sigma d\Psi_1.\Gamma_1 + d\Phi.\Gamma_2 - D\Sigma\Psi_1) + O(\Delta t^3) \\ \Psi_2 &= \Sigma d\Psi_1.\Gamma_1 + d\Phi.\Gamma_2 - D\Sigma\Psi_1 \end{aligned}$$

Computation of the coefficient Γ_3

$$\begin{aligned} \partial_t W &= -AW - BY^* + \frac{1}{2}\Delta t (A_2 W + B_2 Y^* - \partial_t^2 W) \\ &\quad - \frac{1}{6}\Delta t^2 (A_3 W + B_3 Y^* + \partial_t^3 W) + O(\Delta t^3) \end{aligned}$$

with [see Annex -1- pages (i) and (ii)]

$$\begin{aligned} AW + BY^* &= AW + B\Phi + \Delta t B (\Sigma - \frac{1}{2} I) \Psi_1 \\ &\quad + \Delta t^2 B (\Sigma - \frac{1}{2} I) \Psi_2 + O(\Delta t^3) \end{aligned}$$

$$\begin{aligned} A_2 W + B_2 Y^* &= A\Gamma_1 + B(d\Phi.\Gamma_1 - \Psi_1) + \Delta t B_2 (\Sigma - \frac{1}{2} I) \Psi_1 \\ &\quad + O(\Delta t^2) \end{aligned}$$

$$A_3 W + B_3 Y^* = \partial^2\Gamma_1.\Gamma_1 - B d\Psi_1.\Gamma_1 - B_2 \Psi_1 + O(\Delta t)$$

and [see Annex -2- pages (i) and (ii)]

$$\partial_t^2 W = d\Gamma_1.\Gamma_1 + \Delta t (d\Gamma_1.\Gamma_2 + d\Gamma_2.\Gamma_1) + O(\Delta t^2)$$

$$\partial_t^3 W = -\partial^2\Gamma_1.\Gamma_1 + O(\Delta t)$$

$$\partial_t W = -\Gamma_1 - \Delta t B \Sigma \Psi_1$$

$$\begin{aligned} &+ \Delta t^2 \left(-B (\Sigma - \frac{1}{2} I) \Psi_2 + \frac{1}{2} (B_2 (\Sigma - \frac{1}{2} I) \Psi_1 - d\Gamma_1.\Gamma_2 - d\Gamma_2.\Gamma_1) \right. \\ &\quad \left. + \frac{1}{6} (B d\Psi_1.\Gamma_1 + B_2 \Psi_1) \right) + O(\Delta t^3) \end{aligned}$$

Computation of the coefficient Γ_3 (ii)

$$\partial_t W + \Gamma_1 + \Delta t \Gamma_2 = \Delta t^2 \left(-B \left(\Sigma - \frac{1}{2} I \right) \Psi_2 + \frac{1}{2} \left(B_2 \left(\Sigma - \frac{1}{2} I \right) \Psi_1 - d\Gamma_1 \cdot \Gamma_2 - d\Gamma_2 \cdot \Gamma_1 \right) + \frac{1}{6} (B d\Psi_1 \cdot \Gamma_1 + B_2 \Psi_1) \right) + O(\Delta t^3)$$

and the coefficient of Δt^2 is given by

$$\begin{aligned} \Gamma_3 &= B \Sigma \Psi_2 - \frac{1}{2} B \Psi_2 + \frac{1}{2} (\textcolor{red}{AB} + BD) \Sigma \Psi_1 + \left(\frac{1}{4} - \frac{1}{6} \right) B_2 \Psi_1 \\ &\quad - \frac{1}{2} (\textcolor{red}{A}\Gamma_2 + B d\Phi \cdot \Gamma_2) - \frac{1}{2} B \Sigma d\Psi_1 \cdot \Gamma_1 - \frac{1}{6} B d\Psi_1 \cdot \Gamma_1 \\ &\qquad\qquad\qquad \text{because } \Gamma_2 = B \Sigma \Psi_1 \\ &= B \Sigma \Psi_2 - \frac{1}{2} B (D \Sigma \Psi_1 - d\Phi \cdot \Gamma_2 - \Sigma d\Psi_1 \cdot \Gamma_1) + \frac{1}{2} \textcolor{red}{BD} \Sigma \Psi_1 \\ &\quad \frac{1}{12} B_2 \Psi_1 - \frac{1}{2} \textcolor{green}{B} d\Phi \cdot \Gamma_2 - \frac{1}{2} \textcolor{blue}{B} \Sigma d\Psi_1 \cdot \Gamma_1 - \frac{1}{6} B d\Psi_1 \cdot \Gamma_1 \end{aligned}$$

Finally,

$$\Gamma_3 = B \Sigma \Psi_2 + \frac{1}{12} B_2 \Psi_1 - \frac{1}{6} B d\Psi_1 \cdot \Gamma_1$$

Link with the previous formal expansion at third-order 34

$$\text{Defect of conservation } \theta_k \equiv \partial_t m_k^{\text{eq}} + \sum_{\ell\beta} \Lambda_{k\ell}^\beta \partial_\beta m_\ell^{\text{eq}}$$

$$\text{vector of conservation defect } \theta = \partial_t \Phi(W) + \Lambda \cdot m^{\text{eq}}$$

$$\text{then } \theta = -\Psi_1 - \Delta t d\Phi \cdot \Gamma_2 + O(\Delta t)$$

$$\text{and } \partial_t \theta = -\partial_t \Psi_1 + O(\Delta t) = d\Psi_1 \cdot \Gamma_1 + O(\Delta t)$$

We have the following calculus:

$$\begin{aligned} \Gamma_2 + \Delta t \Gamma_3 &= B \Sigma \textcolor{blue}{\Psi_1} + \Delta t (B \Sigma \textcolor{red}{\Psi_2} + \frac{1}{12} B_2 \Psi_1 - \frac{1}{6} B d\Psi_1 \cdot \Gamma_1) \\ &= B \Sigma (-\textcolor{blue}{\theta} - \Delta t d\Phi \cdot \Gamma_2 + O(\Delta t^2)) + \Delta t B \Sigma (\Sigma d\Psi_1 \cdot \Gamma_1 + d\Phi \cdot \Gamma_2 \\ &\quad - D \Sigma \textcolor{red}{\Psi_1}) + \frac{1}{12} \Delta t B_2 \Psi_1 - \frac{1}{6} \Delta t B d\Psi_1 \cdot \Gamma_1 \\ &= -B \Sigma \theta + \Delta t \left(-(B \Sigma D \Sigma - \frac{1}{12} B_2) \Psi_1 \right. \\ &\quad \left. + B (\Sigma^2 - \frac{1}{6}) d\Psi_1 \cdot \Gamma_1 \right) + O(\Delta t^2) \\ &= -B \Sigma \theta + \Delta t \left((B \Sigma D \Sigma - \frac{1}{12} B_2) \theta \right. \\ &\quad \left. + B (\Sigma^2 - \frac{1}{6}) \partial_t \theta \right) + O(\Delta t^2) \end{aligned}$$

Link with the previous formal expansion at third-order (ii)

$$\Gamma_2 + \Delta t \Gamma_3 = -B \Sigma \theta + \Delta t \left((B \Sigma D \Sigma - \frac{1}{12} B_2) \theta + B (\Sigma^2 - \frac{1}{6}) \partial_t \theta \right) + O(\Delta t^2)$$

Third-order partial equivalent equations:

$$\partial_t W + \Lambda m^{\text{eq}} - \Delta t B \Sigma \theta + \Delta t^2 \left((B \Sigma D \Sigma - \frac{1}{12} B_2) \theta + B (\Sigma^2 - \frac{1}{6}) \partial_t \theta \right) = O(\Delta t^3)$$

Explicitation with the cartesian components:

$$(B \Sigma D \Sigma \theta)_i = \Lambda_{ik}^\beta \sigma_k (\Lambda_{kl}^\gamma) \sigma_l \partial_\beta \partial_\gamma \theta_\ell = \Lambda_{ik}^\beta \sigma_k \Lambda_{kl}^\gamma \sigma_l \partial_\beta \partial_\gamma \theta_\ell$$

$$(B_2 \theta)_i = (\Lambda_{ik}^\beta \partial_\beta) (\Lambda_{kl}^\gamma \partial_\gamma) \theta_\ell = \Lambda_{ik}^\beta \Lambda_{kl}^\gamma \partial_\beta \partial_\gamma \theta_\ell$$

$$((B \Sigma D \Sigma - \frac{1}{12} B_2) \theta)_i = \Lambda_{ik}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell$$

$$(B (\Sigma^2 - \frac{1}{6}) \partial_t \theta)_i = \Lambda_{ik}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k$$

Other form of the third-order partial equivalent equations:

$$\partial_t W_i + \Lambda_{ik}^\alpha \partial_\alpha m_k^{\text{eq}} - \Delta t \Lambda_{ik}^\beta \sigma_k \partial_\beta \theta_k + \Delta t^2 (\Lambda_{ik}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell + \Lambda_{ik}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k) = O(\Delta t^3)$$

Link with the previous formal expansion at third-order (iii)

$$\partial_t W_i + \Lambda_{ik}^\beta \partial_\beta m_k^{\text{eq}} - \Delta t \Lambda_{ik}^\beta \sigma_k \partial_\beta \theta_k + \Delta t^2 (\Lambda_{ik}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell + \Lambda_{ik}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k) = O(\Delta t^3)$$

Classical lattice Boltzmann scheme : $M_{0j} = 1$ and $M_{\alpha j} = v_j^\alpha$

$$\text{then } \Lambda_{0\ell}^\alpha = M_{0j} v_j^\alpha (M^{-1})_{j\ell} = M_{\alpha j} (M^{-1})_{j\ell} = \delta_{\alpha\ell}$$

$$\text{and } \Lambda_{0k}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell = \delta_{\beta k} \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell = \Lambda_{\beta\ell}^\gamma (\sigma_\beta \sigma_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell$$

Thermal case: $\sigma_\beta \neq 0$ and

$$\Lambda_{0k}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell = \Lambda_{\beta\ell}^\gamma (\sigma_\beta \sigma_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell$$

Fluid case; the momentum is conserved: $\sigma_\beta = 0$. Then

$$\Lambda_{0k}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell = -\frac{1}{12} \Lambda_{\beta\ell}^\gamma \partial_\beta \partial_\gamma \theta_\ell$$

$$\Lambda_{0k}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k = \delta_{\beta k} (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k = (\sigma_\beta^2 - \frac{1}{6}) \partial_t (\partial_\beta \theta_\beta)$$

Link with the previous formal expansion at third-order (iv)

$$\partial_t W_i + \Lambda_{ik}^\beta \partial_\beta m_k^{\text{eq}} - \Delta t \Lambda_{ik}^\beta \sigma_k \partial_\beta \theta_k + \Delta t^2 (\Lambda_{ik}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_l + \Lambda_{ik}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k) = O(\Delta t^3)$$

$$\Lambda_{0k}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k = \begin{cases} \Lambda_{\beta\ell}^\gamma (\sigma_\beta \sigma_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell & \text{thermics} \\ -\frac{1}{12} \Lambda_{\beta\ell}^\gamma \partial_\beta \partial_\gamma \theta_\ell & \text{fluid} \end{cases}$$

$$\Lambda_{0k}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k = (\sigma_\beta^2 - \frac{1}{6}) \partial_t (\partial_\beta \theta_\beta)$$

Thermal case: $\sigma_\beta \neq 0$ and

$$\Lambda_{0k}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k = (\sigma_\beta^2 - \frac{1}{6}) \partial_t (\partial_\beta \theta_\beta)$$

Fluid case; the momentum is conserved: $\theta_\beta = O(\Delta t)$. Then

$$\Lambda_{0k}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k = O(\Delta t)$$

Link with the previous formal expansion at third-order (v)

$$\partial_t W_i + \Lambda_{ik}^\beta \partial_\beta m_k^{\text{eq}} - \Delta t \Lambda_{ik}^\beta \sigma_k \partial_\beta \theta_k + \Delta t^2 (\Lambda_{ik}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_l + \Lambda_{ik}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k) = O(\Delta t^3)$$

Classical notations: $W_0 = \rho \equiv \sum_j f_j$, $W_\alpha = J_\alpha = \sum_j v_j^\alpha f_j$

$$\Lambda_{0k}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k = \begin{cases} \Lambda_{\beta\ell}^\gamma (\sigma_\beta \sigma_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell & \text{thermics} \\ -\frac{1}{12} \Lambda_{\beta\ell}^\gamma \partial_\beta \partial_\gamma \theta_\ell & \text{fluid} \end{cases}$$

$$\Lambda_{0k}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k = \begin{cases} (\sigma_\beta^2 - \frac{1}{6}) \partial_t (\partial_\beta \theta_\beta) & \text{thermics} \\ 0 & \text{fluid} \end{cases}$$

Mass conservation in the thermal case :

$$\partial_t \rho + \partial_\alpha J_\alpha^{\text{eq}} - \Delta t \sigma_\alpha \partial_\alpha \theta_\alpha + \Delta t^2 (\Lambda_{\beta\ell}^\gamma (\sigma_\beta \sigma_\ell - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_\ell + (\sigma_\beta^2 - \frac{1}{6}) \partial_t (\partial_\beta \theta_\beta)) = O(\Delta t^3)$$

[relation (35) of DCDS-A, 2009].

Mass conservation in the fluid case :

$$\partial_t \rho + \partial_\alpha J_\alpha - \frac{1}{12} \Delta t^2 \Lambda_{\beta\ell}^\gamma \partial_\beta \partial_\gamma \theta_\ell = O(\Delta t^3)$$

[relation (40) of DCDS-A, 2009].

Link with the previous formal expansion at third-order (vi)

$$\partial_t W_i + \Lambda_{ik}^\beta \partial_\beta m_k^{\text{eq}} - \Delta t \Lambda_{ik}^\beta \sigma_k \partial_\beta \theta_k \\ + \Delta t^2 (\Lambda_{ik}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_l + \Lambda_{ik}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k) = O(\Delta t^3)$$

Momentum conservation for the fluid case: $i = \alpha$

$$\partial_t J_\alpha + \Lambda_{\alpha k}^\beta \partial_\beta m_k^{\text{eq}} - \Delta t \Lambda_{\alpha k}^\beta \sigma_k \partial_\beta \theta_k \\ + \Delta t^2 (\Lambda_{\alpha k}^\beta \Lambda_{kl}^\gamma (\sigma_k \sigma_l - \frac{1}{12}) \partial_\beta \partial_\gamma \theta_l \\ + \Lambda_{\alpha k}^\beta (\sigma_k^2 - \frac{1}{6}) \partial_\beta \partial_t \theta_k) = O(\Delta t^3)$$

[relation (41) of DCDS-A, 2009].

- Compact iteration of lattice Boltzmann schemes
- Block decomposition of the moment-velocity operator matrix
- Asymptotic expansion of the equivalent partial differential equations and of the non conserved moments
- Explication of the coefficients up to order 3 with recursive formulas containing less than 3 terms
- Intensive use of differential calculus
- Validation of the nonlinear expansion for fluid flow and thermal problems up the the order 3
- Validation of the non linear expansion at second order for Navier Stokes flows with two en three space dimensions
[Icmmes 2018, ICIAM 2019, ...]
- New applications soon !