



**HAL**  
open science

## Effect of water restriction stress on social behavior in mice: Impact on serotonergic system plasticity

Houari Boukersi, Nemcha Lebaili, Anne Nosjean, Nathalie Samson, Alexis Faure, Sylvie Granon

### ► To cite this version:

Houari Boukersi, Nemcha Lebaili, Anne Nosjean, Nathalie Samson, Alexis Faure, et al.. Effect of water restriction stress on social behavior in mice: Impact on serotonergic system plasticity. Behavioural Brain Research, 2021, 399:, pp.113022. 10.1016/j.bbr.2020.113022 . hal-03452677

**HAL Id: hal-03452677**

**<https://cnrs.hal.science/hal-03452677v1>**

Submitted on 27 Nov 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

## **Effect of water restriction stress on social behavior in mice:**

### **Impact on serotonergic system plasticity**

Houari Boukersi<sup>1,2,3</sup>, Nemcha Lebaili<sup>1</sup>, Anne Nosjean<sup>3</sup>, Nathalie Samson<sup>3</sup>, Alexis Faure<sup>3</sup>, Sylvie Granon<sup>3</sup>.

1-Laboratoire d'écophysiologie animale, école normale supérieure de Kouba, Alger, Algérie

2-Département de biologie, faculté des sciences de la nature et de la vie, université Hassiba ben Bouali, CHLEF, Algérie

3-Institut des Neurosciences Paris-Saclay, Université Paris-Saclay, CNRS 9197, Orsay, France

#### **Abstract**

We explored here the hypothesis that temporary chronic water restriction in mice affects social behavior, via its action on the density of 5-HT neurons in dorsal and median raphe nuclei (DRN and MRN). For that, we submitted adult C57BL/6J mice to mild and controlled temporary dehydration, i.e., 6h of water access every 48h for 15 days. We investigated their social behavior in a social interaction task known to allow free and reciprocal social contact and to rely on prefrontal cortex activity and monoaminergic prefrontal input. Results showed that temporary dehydration increases significantly time spent in social contact and social dominance. It also expands 5-HT neuron density within both DRN and MRN and the behavioral and neuronal plasticity were positively correlated. Our findings suggest that disturbance in 5-HT innervation caused by temporary dehydration stress unbalances choice processes of animals in social context.

**Keywords:** Raphe nucleus; Serotonin; Dehydration; Social behavior; Immunohistochemistry.

#### **Disclosure of interest**

The authors declare that they have no conflicts of interest concerning this article.

#### **Acknowledgments**

This study was supported by The Algerian Directorate-General of Scientific Research and Technological Development (DGRSDT) and the Paris-Saclay Institute of Neuroscience (Neuro-PSI).

## **Introduction**

In the wild, rodents are able to survive on relatively dry diets and may go into temporary dehydration states for various periods (Tirado et al., 2008). Temporary dehydration alters serotonergic innervation in multiple mammal species such as rats (Chatoui et al., 2012), merions (Elgot et al., 2009; 2012<sub>a, b</sub>), and gerbils (Boukersi et al., 2018). A temporary controlled dehydration in the laboratory is thus an ecological modulator of serotonin activity that mimics what happens in real life context. Serotonin (5-hydroxytryptamin or 5-HT) is the most widely distributed transmitter in the brain (Dahlstrom and Fuxe, 1964) and is well known for its role in mood regulation (David and Gardier, 2016) since abnormalities in its neurotransmission are associated with neuropathologies like depression and anxiety (Cowen, 2008; Quesseveur et al., 2013) and other neurological or psychiatric disorders (Jayamohanam et al., 2019). However, it is still unclear how various environmental factors such as temporary dehydration can physiologically control brainstem neuromodulatory systems, including the dorsal and medial raphe (DRN and MRN) serotonergic nuclei, the main sources of 5-HT (Michelsen et al., 2008; Steinbusch, 1981). It is well established that 5-HT containing neurons of DRN and MRN project to forebrain areas (Kosofsky and Molliver, 1987; O'Hearn and Molliver, 1984; Steinbusch and de Vente, 1997) and receive descending projections from forebrain regions (Celada et al., 2001; Peyron et al., 1998). The MRN innervates several subcortical areas such as the hippocampus and the nucleus accumbens, and sends axons to dopaminergic (DA) neurons of the ventral tegmental area which in turn innervates the prefrontal cortex (PFC). The DRN innervates the PFC, the amygdala, the nucleus accumbens and the hippocampus and sends axons to DA neurons of the substantia nigra that innervate the amygdala and the dorsal striatum. Both DRN and MRN innervate several hypothalamic nuclei, therefore triggering endocrine release (Lechin et al., 2006). The 5-HT system is thus ideally placed to support integration of emotional, motivational and cognitive behaviours (Muzerelle et al., 2014; Suri et al., 2015). For example, in animal models, 5-HT in limbic forebrain areas was altered by multiple environmental stressful conditions, such as temperature variation (Linthorst et al., 2008), swim stress (Kelly et al., 2011), or social defeat (Paul et al., 2011). In humans, patients with anxiety and affective disorders show an increase of brain 5-HT turnover compared to healthy subjects (Esler et al. 2007; Barton et al. 2008). However, there are little studies reporting alteration of serotonergic innervation in 5-HT main source of production, namely DRN and MRN.

Social interactions are important components of healthy behaviours (Monica et al., 2017) since they enable animals to communicate with and learn from one another, and can increase the likelihood of survival and reproduction (Ferri et al., 2016). We previously showed that social interaction relies on PFC integrity and triggers its activation (Avale et al., 2011; Nosjean et al., 2018), and that both cholinergic integrity (de Chaumont et al., 2012; Nosjean et al., 2015) and the PFC monoaminergic input are necessary for displaying adapted and non aggressive reciprocal social interaction (Coura et al., 2013). A recent study provides proteomic evidences that serotonergic and dopaminergic activities were increased by social play in rats (Alugubelly et al., 2019). In addition, dorsal raphé activation was specifically associated with social dominance (Kim et al., 2015) while optogenetic stimulation of dorsal raphé nuclei elevated prefrontal 5-HT levels along with social behavior frequency (Balazsfi et al., 2018). Despite these recent data, the precise role of serotonergic activity on social behavior remains largely unknown.

In this study, we tested the behavioral and neurobiological consequences of dehydration on social behavior and on DRN and MRN neuronal reactivity.

## **Materials and methods**

### **Animals**

20 adult male C57BL/6J mice obtained from Charles Rivers Laboratories (L'Arbresle Cedex, France) were used in our study. We randomly divided them into three groups: the first group (n=5) served as a control with water access *ad libitum*, the second group (n=5) was submitted to a controlled dehydration protocol by allowing them water access only for 6 hours per 48 hours for 15 consecutive days, the third group (n=10) was used as social partner (see below) and had access to water and food *ad libitum*. All animals had food access *ad libitum*.

Animals of groups 1 & 2 were socially isolated for 4 weeks prior to social tests (transparent plexiglas cage, 50x20 cm, covered with clean sawdust) while animals of group 3 were maintained 3 to 5 per cage under constant room temperature ( $23\pm 2^{\circ}\text{C}$ ), humidity and 12/12 h light-dark cycle.

Behavioural procedures were carried out in accordance with European Commission guidelines. All efforts were made to minimize animal suffering and reduce the number of animals used.

### **Social interaction task (SIT)**

At the end of the dehydration procedure, we followed the experimental protocol previously described for social interactions (Avale et al., 2011; Nosjean et al., 2015). Briefly, each

isolated mouse (thereafter called host mouse) was put in a novel transparent open field (50 x20 cm) containing a handful of clean sawdust, where it stayed alone exploring this novel environment for 30 minutes. This procedure promotes the development of limited dominance from the host mouse (De Chaumont et al., 2012; Faure et al., 2017; Nosjean et al., 2015). After that, a visitor male mouse of the same strain and age and not isolated before (maintained in social cages of 3 to 5 animals), was introduced into the open field for a 8-min social interaction test that was recorded by a camera connected to a computer for offline analysis.

We scored the time spent in contact, the number of follow behaviors (i.e., bouts of behaviors during which the host mouse follows the visitor mouse while making nose-genital contact), the number of escape behavior (i.e. behavioral sequences initiated by the host mouse consisting of interrupting the social contact and going away from the visitor mouse) and the number of rearing (lifting the two forepaws) performed by host mice.

### **Tissue preparation and immunostaining for 5-HT**

At the end of the experiment, as previously described (Nosjean et al., 2015; Boukersi et al., 2018), all mice were deeply anesthetized by intraperitoneal injection of sodium pentobarbital (40 mg/kg and transcardially perfused with ice-cold phosphate buffered saline (PBS, 0.1 M, pH 7.4) followed by 4% paraformaldehyde in (PBS, 0.1 M, pH 7.4). Brains were removed and post-fixed for one week in the same fixative at 4 °C. For each brain serial coronal sections (40µm thickness), including the raphe nuclei in the brainstem were cut using Leica vibratome. Free floating sections were collected in 6-well tissue culture plates in 1.5 ml of cryoprotection solution and stored at -20°C until the day of primary antibody incubation. Sections processed in midbrain through the DRN and MRN were chosen for 5-HT immunolabelling. Free-floating sections from both hydrated and dehydrated groups were immunostained simultaneously using the same reagents and same incubation time. Free-floating sections were rinsed 3 times in (PBS, 0.1 M) before being incubated in a solution containing methanol and hydrogen peroxide during 30 minutes at room temperature, rinsed in 0.1 M PBS followed by one wash in (PBS, 0.1 M) containing 0.2% Triton X-100 (PBST), and preincubated in 5% normal goat serum with PBST for 1 h. Sections were then incubated overnight at 4°C with rabbit primary antibody anti 5-HT (Euromedex) diluted 1:2000 in PBST and goat serum 5%. The next day tissues were washed 3 times in PBST the incubated for 2 h in goat secondary biotinylated anti-rabbit IgG (Eurobio) diluted 1:200 in PBST and goat serum 5%. Tissue was then washed 3 times in PBST followed by 2 h incubation with an avidin-biotin-peroxidase complex (ABC, Vectastain ABC kit, Vector laboratories). After washing sections in BPST

then twice in Tris HCL 0.1 M, the peroxidase activity was revealed by incubating them in 0.03% DAB (3,3' diaminobenzidine, Sigma) + Nickel (NiCl<sub>2</sub>) 0.06% and 0.02% H<sub>2</sub>O<sub>2</sub> in 0.05 M Tris HCl, pH 7.5. All immunochemical reactions and washings described above were possessed simultaneously in four 12-well tissue culture plates, 10 sections in each well, to insure identical condition staining for all sections. Immediately after the chromogen reaction, sections were washed in (PB, 0.1M) and then in 0.15% gelatine in H<sub>2</sub>O before being mounted on glass microscope slides (Thermo Scientific Super Frost Plus). Lastly, slides were air dried for 24 hours, dehydrated through 2 min serial ethanol baths (70%, 95% and 100% ethanol), and cleared with xylene. Coverslips were placed on the sections using Eukit for optic microscopy observation.

### **5-HT immunostaining assessment**

The selected coronal brain sections permitted evaluation of raphe regions known to be heavily populated with serotonergic neurons. Anatomical landmarks were used to ensure that comparable brain sections were analyzed for each region in each animal. Using atlas brain of mouse (Paxinos and Franklin, 2001), three sections corresponding to the same three levels of bregma (-4.72, -4.60 and -4.48) were chosen for each brain in which the whole DRN and MRN could be seen clearly and most entirely. Examination of the slices was performed by Olympus light microscope coupled to an image-analysis workstation (Mercator; Explora Nova software). Photomicrographic digital images were taken for the three sections of each animal's brain, and the average density of 5-HT immunopositive neurons (number of somas per 1mm<sup>2</sup>) was calculated in DRN and in MRN by counting automatically all neurons having colour density more than threshold defined at start of counting.

### **Statistical analysis**


Because of the small number of animals in each group (n=5), Mann-Whitney test was chosen to compare results of hydrated and dehydrated groups. A p value <0.05 was considered to indicate statistical significance between groups. Finally, to examine potential relationships between SIT and 5-HT immunostaining we applied the non-parametric test of correlation of Spearman, considering a significant correlation at p<0.05.

## **Results**

### **Effect of water restriction on social interaction task**

Our data showed an increase of time spent in social contact between animals in dehydration condition as compared to normal hydration (p= 0.015) (Fig 1). The number of follow


behavior, index of social dominance, was significantly higher in the dehydrated group as compared to normally hydrated one (control) ( $p= 0.007$ ) (Figure 1). The number of rearings and of escape behavior were not significantly altered in dehydrated animals (group effect respectively  $p= 0.221$  and  $p= 0,149$ ).




**Figure 1.** Social interaction task in dehydrated and control mice. Data is presented as means  $\pm$  SEM. Significant group effect (i.e. effect of dehydration) is indicated by \* ( $p < 0.05$ , Mann-Whitney test).

### **Serotonin immunolabeling in DRN and in MRN**

We examined the 5-HT immunolabeling in the DRN and MRN. Positive 5-HT immunolabeled neurons density in both nuclei was significantly increased in dehydrated mice as compared to controls ( $p= 0.007$  and  $p= 0.015$ , Figures 2 and 3)



**Figure 2.** Representative microphotographs of 5-HT immunolabeled neurons in the dorsal raphe nucleus (DRN, A & C) and median raphe nucleus (MRN, B & D, bregma -4.72 mm) in dehydrated (top) and control mice (bottom). DRD: dorsal part of dorsal raphe nucleus; DRVL: ventro lateral part of dorsal raphe nucleus; DRV: ventral part of dorsal raphe nucleus; DRI: interfascicular part of dorsal raphe nucleus; mlf: medial longitudinal fasciculus.




**Figure 3.** 5-HT immunolabeled neuron density in the DRN and MRN in dehydrated and control mice. Data is presented as means  $\pm$  SEM. Significant group effect (i.e. effect of dehydration) was indicated by \* ( $p < 0.05$ , Mann-Whitney test).




### Correlation between serotonin immunolabeling in DRN and MRN and social behavior in SIT


Across both DRN, density of 5-HT immunolabeled neurons correlates positively and significantly with time spent in social contact ( $R^2 = 65\%$ ;  $p < 0.01$ , Figure 4) and number of follow behaviors ( $R^2 = 65\%$ ;  $p < 0.01$ , fig 5). However, in MRN, density of 5-HT immunolabeled neurons correlates positively and significantly only with time spent in social contact ( $R^2 = 52\%$ ;  $p < 0.05$ , Figure 4) but not with the number of follow behaviors ( $R^2 = 41.5\%$ ;  $p > 0.05$ , fig.5). The density of 5-HT neurons in DRN and MRN was not correlated with the number of escape behavior ( $R^2 = 5.9\%$ ;  $p > 0.05$  and  $R^2 = 56.8\%$ ;  $p > 0.05$  respectively, Figure 6) nor with the number of rearings ( $R^2 = 3.1\%$ ;  $p > 0.05$  and  $R^2 = 42.1\%$ ;  $p > 0.05$  respectively, Figure 7).




**Figure 4.** Positive correlation between 5-HT neuron density in the DRN and MRN and contact time in the social interaction task. Black dots represent dehydrated mice, grey dots represent control mice. Significant correlation was indicated by \* ( $p < 0.05$ ,  $R^2$  Spearman correlation coefficient).



**Figure 5.** Positive correlation between 5-HT neuron density in the DRN and MRN and the number of follow behaviour displayed by the host mice. Black dots represent dehydrated mice, grey dots represent control mice. Significant correlation was indicated by\* ( $p < 0.05$ ,  $R^2$  Spearman correlation coefficient).



**Figure 6.** No correlation between 5-HT neuron density in the DRN and MRN and the number of escapes displayed by the host mice. Black dots represent dehydrated mice, grey dots represent control mice ( $p > 0.05$ ,  $R^2$  Spearman correlation coefficient).



**Figure 7.** No correlation between 5-HT neuron density in the DRN and MRN and the number of rearings displayed by the host mice. Black dots represent dehydrated mice, grey dots represent control ( $p > 0.05$ ,  $R^2$  Spearman correlation coefficient).

## Discussion

In the present study, we aim to explore the hypothesis that temporary chronic water restriction could affect social behavior, via its action on the density of 5-HT neurons in DRN and in MRN. The C57Bl/6j strain that we used is known for its sociability and is thus perfectly suitable for studying social behavior (Faure et al. 2017; Lawrence et al. 2017).

Results reveal significant increases of the time spent in social contact and of dominance behavior, and of 5-HT neurons density within both DRN and MRN in dehydrated animals as compared to control animals.

Our findings suggest that disturbance in 5-HT innervation caused by temporary dehydration stress could unbalance choice processes of animals in the social task. Indeed, we previously showed that this social task relies on the ability to make choices between two concurrent motivations, i.e., getting a social contact and exploring a novel environment (Granon et al., 2003; Avale et al., 2011; De Chaumont et al., 2012;), processes supported by prefrontal activity (Avale et al., 2011; Nosjean et al., 2018) and prefrontal monoaminergic modulation (Cambon et al., 2010; Coura et al., 2013). Results of correlation between 5-HT immunolabelling in DRN and MRN and social contact parameters support this hypothesis, showing that higher number of 5-HT neurons in DRN and MRN is associated with longer

social contact and more dominance behavior. We previously evidenced that follow behavior, an index of social dominance, largely contributes to social contact (Coura et al., 2013; Faure et al., 2017) and is promoted by acute stress (Nosjean et al., 2018). Our current data therefore supports and reinforces previous works showing that social dominance is positively related to 5-HT neuronal activity (Kiser et al., 2012; Kim et al., 2015), and that while decreased levels of 5-HT promote social isolation (Higley et al., 1996), its increase promote social cooperation and contact (Paula et al. 2015; Anstey et al., 2009) and affiliative behaviors (Aan het Rot et al., 2006; Tse and Bond, 2002).

It is noticeable that aggressive behavior was not observed in any mouse that was socially isolated before the social task, independently of their hydration condition (except one brief aggression from one dehydrated animal). This supports our previous work (Nosjean et al., 2015) showing that acute stress, but not temporary social isolation during adulthood, can trigger aggressive behavior in some mice. Social dominance is a normal behavior dissociated from aggressive behavior that we previously showed to be modulated by the cholinergic system activity (Nosjean et al., 2015) while aggressive behavior was modulated by the noradrenergic system activity (Cambon et al., 2010; Coura et al., 2013). However, other studies suggested causal links between increased aggression and low 5-HT activity (Audero, et al, 2013; Bjork et al., 1999; Caramaschi et al., 2007). One hypothesis to explain this discrepancy is that these authors consider extracellular 5-HT and its metabolite 5-HIAA (Dekeyne et al., 2000; Mir and Taylor, 1997; Sambunaris et al., 1997), whereas we look here at 5-HT neurons. We have recently shown that acute sleep restriction produced a decrease of 5-HT level in the prefrontal cortex, together with altered decision-making processes (Pittaras et al., 2018). It could therefore be of interest in future works to measure not only the cellular plasticity of 5-HT after dehydration but its consequences on 5-HT level and its metabolites in various brain areas. Besides, further study should clarify the signalling consequences of intracellular 5-HT increase after dehydration and its temporality, i.e., whether synthesis, degradation, reuptake and release of 5-HT are altered, by which mechanisms and which duration of dehydration is needed for them to take place. Conversely, it would be valuable to investigate other cognitive consequences of temporary dehydration. Nevertheless, to our knowledge, this is the first work showing neuronal plasticity of 5-HT in DRN and MRN after temporary dehydration stress and its detrimental effect on social interaction.

5-HT neurons had been found in higher density under dehydration in *Gerbillus tarabuli* (Boukersi et al., 2018), a rodent adapted to dry environment. The fact that we showed a

similar plasticity of the serotonergic system in mice, rodents not particularly adapted to lack of water, suggests that this monoaminergic system may play a pivotal role in the rapid brain response to water homeostasis. This is in accordance with several reports providing evidence for involvement of this neurotransmitter in controlling vasopressin secretion (Popova et al., 2001), drinking behaviour (Reis et al., 1990), salt appetite (Cooper and Ciccocioppo, 1993), water intake and urine production (Olivares et al., 2003; Reis et al., 1994), supporting the view that serotonergic circuits originating from the mesencephalic raphe are implicated in hydromineral homeostasis (Fitzsimons, 1998; Franchini et al. 2002; McCann et al., 2003; Phillips et al., 1982; Reis et al. 1994).

In addition, it is well known that midbrain 5-HT neurons establish reciprocal interactions with thirst and water balance control areas, such as the subfornical organ (Tanaka et al., 2003), organum vasculosum laminae terminalis (Thrasher and Keil, 1987), and hypothalamo-pituitary axis (Maslova et al., 1990; Popova et al., 2001). The latter is known to be markedly affected when animals are dehydrated (Langle et al., 2002) since it secretes vasopressin which controls water homeostasis in mammals (Elgot et al. 2012<sub>a, b</sub>; Jones and Pickering, 1969). Furthermore, some areas involved in stress management and social behavior such as the prefrontal cortex (Abrams et al., 2004; Avale et al., 2011; Loiseau et al., 2008), the amygdala (Imai et al., 1986; Nosjean et al., 2018; Olsson and Phelps, 2007 ) and the hippocampus (McEwen et al., 2016; Kohler and Steinbusch, 1982) make important reciprocal anatomical connections with raphe nuclei (Groenewegen and Uylings, 2000; Hajós et al., 1998; Sesack et al., 1989), hence playing an important role in regulating executive function (Puig and Gullledge, 2011). Consequently, the wide distribution of 5-HT in the brain may make any change in its source (DRN and MRN) a potential cause of change in all functions controlled by areas innervated by 5-HT.

## References

1. Aan het Rot, M., Moskowitz, D.S., Pinard, G., Young, S.N., 2006. Social behaviour and mood in everyday life: the effects of tryptophan in quarrelsome individuals. *J.Psychiatry Neurosci.* 31, 253–262.
2. Abrams, J.K., Johnson, P.L., Hollis, J.H., Lowry, C.A., 2004. Anatomic and functional topography of the dorsal raphe nucleus. *Ann N Y Acad Sci.* 1018, 46–57.
3. Alugubelly, N., Mohammad, A.N., Edelman, M.J., Nanduri, B., Sayed, M., Won Park, J., Carr, R.L., 2019. Proteomic and transcriptional profiling of rat amygdala following social play. *Behavioural Brain Research.* 376, 112210.
4. Anstey, M.L., Rogers, S.M., Ott, S.R., Burrows, M., Simpson, S.J., 2009. Serotonin mediates behavioral gregarization underlying swarm formation in desert locusts. *Science.* 323, 627–630.
5. Audero, E., Mlinar, B., Baccini, G., Skachokova, Z. K., Corradetti, R., and Gross, C., 2013. Suppression of serotonin neuron firing increases aggression in mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience.* 33(20), 8678–8688.
6. Avale, M.E., Chabout, J., Pons, S., Serreau, P., De Chaumont, F., Olivo-Marin, J.C., Bourgeois, J.P., Maskos, U., Changeux, J.P., Granon, S. 2011. Prefrontal nicotinic receptors control novel social interaction between mice. *FASEB J.* 25(7), 2145–55.
7. Barton, D.A., Esler, M.D., Dawood, T., Lambert, E.A., Haikerwal, D., Brenchley, C., Socratous, F., Hastings, J., Guo, L., Wiesner, G., Kaye, D.M., Bayles, R., Schlaich, M.P., Lambert, G.W., 2008. Elevated brain serotonin turnover in patients with depression: effect of genotype and therapy. *Arch Gen Psychiatry.* 65, 38–46.
8. Bjork, J.M., Dougherty, D.M., Moeller, F.G., Cherek, D.R., Swann, A.C., 1999. The effects of tryptophan depletion and loading on laboratory aggression in men: time course and a food-restricted control. *Psychopharmacology.* 142, 24 –30.
9. Boukersi, H., Lebaili, N., Samson, N., Granon, S., 2018. Implication of regional brain serotonergic neurons in dorsal and median raphe nuclei in adaptation to water lacking in *Gerbillus tarabuli*. *Journal of Chemical Neuroanatomy.* 93, 57–61.
10. Cambon, K., Dos-Santos C.R., Groc, L., Carbon, A., Weissmann, D., Changeux, J.P., Pujol, J.F., Granon, S., 2010. Aggressive behavior during social interaction in mice is controlled by the modulation of tyrosine hydroxylase expression in the prefrontal cortex. *Neuroscience.* 171(3), 840–51.
11. Caramaschi, D., de Boer, S.F., Koolhaas, J.M., 2007. Differential role of the 5-HT1A receptor in aggressive and nonaggressive mice: an across-strain comparison. *Physiol Behav.* 90, 590–601.

12. Celada, P., Puig, M.V., Casanovas, J.M., Guillazo, G., Artigas, F., 2001. Control of dorsal raphe serotonergic neurons by the medial prefrontal cortex: involvement of serotonin-1A, GABA(A), and glutamate receptors. *J. Neurosci.* 21, 9917–9929.
13. Chatoui, H., El Hiba, O., Elgot, A., Gamrani, H., 2012. The rat SCO responsiveness to prolonged water deprivation Implication of Reissner's fiber and serotonin system. *C. R. Biol.* 335, 253–260.
14. Cooper, S. J. and Ciccocioppo, R., 1993. Effects of selective 5-HT1 receptor agonists in water-deprived rats on salt intake in two-choice tests, *Pharmacol. Biochem. Behav.* 45, 513–518
15. Coura, R.S., Cressant, A., Xia, J., de Chaumont, F., Olivo-Marin, J.C., Pelloux, Y., Dalley, J.W., Granon, S., 2013. Nonaggressive and adapted social cognition is controlled by the interplay between noradrenergic and nicotinic receptor mechanisms in the prefrontal cortex, *FASEB J.* 27, 4343–4354.
16. Cowen, P.J., 2008. Serotonin and depression: pathophysiological mechanism or marketing myth? *Trends Pharmacol. Sci.* 29, 433–436.
17. Dahlstrom, A., Fuxe, K., 1964. Localization of monoamines in the lower brain stem. *Experientia.* 20, 398–399.
18. David, D.J., Gardier, A.M., 2016. The pharmacological basis of the serotonin system: Application to antidepressant response. *Encephale.* 42(3), 255–63.
19. De Chaumont, F., Coura, R.D., Serreau, P., Cressant, A., Chabout, J., Granon, S., Olivo-Marin, J.C., 2012. Computerized video analysis of social interactions in mice. *Nature Methods.* 9 (4), 410–417.
20. Dekeyne, A., Denorme, B., Monneyron, S., Millan, M.J., 2000. Citalopram reduces social interaction in rats by activation of serotonin (5-HT)<sub>2C</sub> receptors. *Neuropharmacology.* 39, 1114–1117.
21. Elgot, A., Ahboucha, S., Bouyatas, M.M., Fèvre-Montange, M., Gamrani, H., 2009. Water deprivation affects serotonergic system and glycoprotein secretion in the sub-commissural organ of a desert rodent *Meriones shawi*, *Neuroscience Letters.* 466, 6–10.
22. Elgot, A., El Hiba, O., Gamrani, H., 2012<sub>a</sub>. nuclei of hypothalamus of a desert rodent *Meriones Shawi* after a severe dehydration versus opposite treatment by rehydration: GFAP and vasopressin immunohistochemical study. *Neuroscience Letters.* 515, 55–60.
23. Elgot, A., El hiba, O., Gamrani, H., 2012<sub>b</sub>. The anxiogenic-like effects of dehydration in a semi-desert rodent *Meriones shawi* indicating the possible involvement of the serotonergic system. *Acta Histochem.* 114, 603–607.

24. Esler, M., Lambert, E., Alvarenga, M., Socratous, F., Richards, J., Barton, D., Pier, C., Brenchley, C., Dawood, T., Hastings, J., Guo, L., Haikerwal, D., Kaye, D., Jennings, G., Kalff, V., Kelly, M., Wiesner, G., Lambert, G., 2007. Increased brain serotonin turnover in panic disorder patients in the absence of a panic attack: reduction by a selective serotonin reuptake inhibitor. *Stress*. 10, 295–304
25. Faure, A., Pittaras, E., Nosjean, A., Chabout, J., Cressant, A., Granon, S., 2017. Social behaviors and acoustic vocalizations in different strains of mice. *Behavioural Brain Research*. 320,383–390
26. Ferri, S.L., Kreibich, A.S., Torre, M.A., Piccoli, C.T., Dow, H., Pallathra, A.A., Li, H., Bilker, W.B., Gur, R.C., Abel, T., and Brodtkin, E.S. 2016. Activation of basolateral amygdala in juvenile c57bl/6j mice during social approach behavior. *Neuroscience*. 335, 184–194
27. Fitzsimons, J.T., 1998. Angiotensin, thirst, and sodium appetite. *Physiological Reviews*. 78, 583-686.
28. Franchini, L.F., Johnson, A.K., and Vivas, L., 2002. Sodium appetite and Fos activation in serotonergic neurons. *Am J Physiol*. 282, R235-R243.
29. Granon, S., Faure, P., Changeux, J. P. 2003. Executive and social behaviors under nicotinic receptor regulation. *Proc. Natl. Acad. Sci. U.S.A.* 100, 9596–9601.
30. Groenewegen, H.J., Uylings, H.B., 2000. The prefrontal cortex and the integration of sensory, limbic and autonomic information. *Prog Brain Res*. 126, 3–2.
31. Hajós, M., Richards, C.D. Székely, A.D., Sharp, T., 1998. An electrophysiological and neuroanatomical study of the medial prefrontal cortical projection to the midbrain raphe nuclei in the rat. *Neuroscience*. 87, 95–10.
32. Higley, J.D., Mehlman, P.T., Poland, R.E., Taub, D.M., Vickers, J., Suomi, S.J., Linnoila, M. 1996. CSF testosterone and 5-HIAA correlate with different types of aggressive behaviors. *Biol Psychiatry*. 40, 1067–1082.
33. Imai, H., Steindler, D.A., Kitai, S.T., 1986. The organization of divergent axonal projections from the midbrain raphe nuclei in the rat. *J Comp Neurol*. 243,363–380.
34. Jayamohan, H., Kalappurackal, M., Kumar, M., Aneesh, T.P., 2019. 5-HIAA as a Potential Biological Marker for Neurological and Psychiatric Disorders. *Adv Pharm Bull*. 9(3), 374–381.
35. Jones, C.W., Pickering, B.T., 1969. Comparison of the effects of water deprivation and sodium chloride imbibition of the hormone content of the neurohypophysis on the rat, *J. Physiol*. 203, 444–458.
36. Kelly, K.J., Donner, N.C., Hale, M.W., Lowry, C.A., 2011. Swim stress activates serotonergic and nonserotonergic neurons in specific subdivisions of the rat dorsal



raphe nucleus in a temperature-dependent manner. *Neuroscience*. 197, 251–268.

Kim, Y., Venkataraju, K. U., Pradhan, K., Mende, C., Taranda, J., Turaga, S. C., 2015. Mapping social behavior-induced brain activation at cellular resolution in the mouse. *Cell Rep.* 10, 292–305.

37. Kiser, D., Steemers, B., Branchi, I., Homberg, J.R., 2012. The reciprocal interaction between serotonin and social behaviour. *Neuroscience and Biobehavioral Reviews*. 36,786–798.
38. Kohler, C., Steinbusch, H., 1982. Identification of serotonin and nonserotonin-containing neurons of the mid-brain raphe projecting to the entorhinal area and the hippocampal formation. A combined immunohistochemical and fluorescent retrograde tracing study in the rat brain. *Neuroscience*. 7,951–975.
39. Kosofsky, B.E., Molliver, M.E., 1987. The serotonergic innervation of cerebral cortex: different classes of axon terminals arise from dorsal and median raphe nuclei. *Synapse*. 1, 153–168.
40. Langle, S.L. Poulain, D.A Theodosis, D.T. 2002. Neuronal-glia remodeling: a structural basis for neuronal-glia interactions in the adult hypothalamus. *J Physiol (Paris)*. 96, 169–75.
41. Lawrence S.H., John H. We, Laura M., Paul J.L., and Angélique, B., 2017, Outbred CD1 mice are as suitable as inbred C57BL/6J mice in performing social tasks, *Neuroscience Letters*. 637, 142–147.
42. Lechin, F., van der Dijs, B., Hernández-Adrián, G., 2006. Dorsal raphe vs. median raphe serotonergic antagonism. Anatomical, physiological, behavioral, neuroendocrinological, neuropharmacological and clinical evidences: Relevance for neuropharmacological therapy. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. 30, 565–585
43. Linthorst, A.C., Flachskamm, C., Reul, J.M., 2008, Water temperature determines neurochemical and behavioural responses to forced swim stress: an in vivo microdialysis and biotelemetry study in rats. *Stress* 11, 88–100.
44. Loiseau, F., Dekeyne, A., Millan, M.J., 2008. Pro-cognitive effects of 5-HT<sub>6</sub> receptor antagonists in the social recognition procedure in rats: implication of the frontal cortex. *Psychopharmacology (Berl)*.196, 93–104.
45. Maslova, L.N., Chernigovskaya, E.V., Belen'kiĭ, M.A., Danilova, O.A., Naumenko, E.V., Polenov, A.L., 1990. The effects of serotonin on hypothalamic structures involved in the regulation of the hypophyseal-adrenocortical system. *Fiziol. Zh. SSSR*. (76)3, 331–337.

46. McCann, S.M., Gutkowska, J., and Antunes-Rodrigues, J., 2003. Neuroendocrine control of body fluid homeostasis. *Braz J Med Biol Res.* 36, 165–181.
47. McEwen, B.S., Nasca, C., Gray, J.D., 2016, Stress Effects on neuronal structure: hippocampus, amygdala, and prefrontal cortex. *Neuropsychopharmacology.* 41, 3–23.
48. Michelsen, K.A., Prickaerts, J., Steinbusch, H.W.M., 2008. The dorsal raphe nucleus and serotonin: implications for neuroplasticity linked to major depression and Alzheimer's disease. *Prog. Brain Res.* 172, 233–264.
49. Mir, S., Taylor, D., 1997. The adverse effects of antidepressants. *Current Opinion in Psychiatry.* 10, 88–94.
50. Monica, M. G., Ning, Z., Adam, H.H., Surbhi., Min, Z., Courtney, D.A., 2017. Social interaction modulates the neuroinflammatory response to global cerebral ischemia in male mice. *Brain Research.* 1673 ,86–94
51. Muzerelle, A., Scotto-Lomassese, S., Bernard, J.F., Soiza-Reilly, M., and Gaspar, P. 2014. Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5-B9) to the forebrain and brainstem. *Brain Struct Funct.* 221, 535–561
52. Nosjean, A., Cressant, A., de Chaumont, F., Olivo-Marin, J.C., Chauveau, F., Granon, S., 2015. Acute stress in adulthood impoverishes social choices and triggers aggressiveness in preclinical models. *Front Behav Neurosci.* 8,447
53. Nosjean, A., de Chaumont, F., Olivo-Marin, J.C. Granon., S., 2018. Stress-induced brain activation: buffering role of social behavior and neuronal nicotinic receptors. *Brain Struct Funct.* 223, 4259–4274.
54. O'Hearn, E. and Molliver, M.E., 1984. Organization of raphe-cortical projections in rat: a quantitative retrograde study. *Brain Res Bull.* 13(6), 709–726.
55. Olivares EL, Costa-e-Sousa RH, Cavalcante-Lima HR, Lima HRC, Cedraz-Mercez PL, Reis LC. 2003. Effect of electrolytic lesion of the dorsal raphe nucleus on water intake and sodium appetite. *Braz J Med Biol Res.* 36, 1709–16.
56. Olsson, A., Phelps, E.A., 2007. Social learning of fear. *Nat. Neurosci.* 10, 1095–1102.
57. Paul, E.D., Hale, M.W., Lukkes, J.L., Valentine, M.J., Sarchet, D.M., Lowry, C.A., 2011. Repeated social defeat increases reactive emotional coping behavior and alters functional responses in serotonergic neurons in the rat dorsal raphe nucleus. *Physiol Behav.* 104, 272–282.
58. Paula, J.R., Messias, J.P., Grutter, A.S., Bshary, R., Soares, M.C., 2015. The role of serotonin in the modulation of cooperative behavior. *Behavioral Ecology.* 26(4),

1005–1012.

59. Paxinos, G., Franklin, K.B.J., 2001. The mouse brain in stereotaxic coordinates, 2nd ed. Academic Press, London.
60. Peyron, C., Petit, J.M., Rampon, C., Jouvet, M., Luppi, P.H., 1998. Forebrain afferents to the rat dorsal raphe nucleus demonstrated by retrograde and anterograde tracing methods. *Neuroscience* 82, 443–468.
61. Phillips, M.I, Hoffman, W.E., Bealer, S.L., 1982. Dehydration and fluid balance: central effects of angiotensin. *Federation Proceedings*. 41, 2520–2527.
62. Pittaras, E., Callebert, J., Dorey, R., Chennaoui, M., Granon, S., Rabat, A., 2018. Mouse Gambling Task reveals differential effects of acute sleep debt on decision-making and associated neurochemical changes. 41(11).
63. Popova, N.K., Ivanova, L.N., Amstislavskaya, T.G., Melidi, N.N., Naumenko, K.S., Maslova, L.N., Bulygina, V.V., 2001, Brain Serotonin Metabolism during Water Deprivation and Hydration in Rats. *Neuroscience and Behavioral Physiology*. 31, 327–332.
64. Puig, M.V., and Gullledge, A.T., 2011. Serotonin and Prefrontal Cortex Function: Neurons, Networks, and Circuits, *Mol Neurobiol*, 44, 449–464.
65. Quesseveur, G., David, D.J., Gaillard, M.C., Pla, P., Wu M.V., Nguyen, H.T., Nicolas, V., Auregan, G., David, I., Dranovsky, A., Hantraye, P., Hen, R., Gardier, A.M., Déglon, N., Guiard, B.P., BDNF overexpression in mouse hippocampal astrocytes promotes local neurogenesis and elicits anxiolytic-like activities. *Transl Psychiatry*. 3(4), e253
66. Reis, L.C., Ramalho, M.J., Antunes-Rodrigues, J., 1990. Central serotonergic modulation of drinking behavior induced by water deprivation: effect of serotonergic agonist (MK-212) administered intracerebroventricularly. *Braz. Med. Biol. Res.* 23, 1335–1338.
67. Reis, L.C., Ramalho, M.J., Favaretto, A.L., Gutkowska, J., McCann, S.M., and Antunes-Rodrigues, J., 1994. Participation of the ascending serotonergic system in the stimulation of atrial natriuretic peptide release. *Proc Natl Acad Sci USA* 91, 12022–12026.
68. Sambunaris, A., Hesselink, J.K., Pinder, R., Panagides, J., Stahl, S.M., 1997. Development of new antidepressants. *Journal of Clinical Psychiatry*. 58, 40–53.
69. Sesack, S.R., Deutch, A.Y., Roth, R.H., Bunney, B.S., 1989. Topographical organization of the efferent projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study with *Phaseolus vulgaris* leucoagglutinin. *J Comp Neurol*. 290, 213–242.

70. Steinbusch, H.W., 1981. Distribution of serotonin-immunoreactivity in the central nervous system of the rat-cell bodies and terminals. *Neuroscience*. 6, 557–618.
71. Steinbusch, H.W.M., de Vente, J., 1997. New vistas on the neurobiology of depression: Colocalization of serotonin, dopamine and nitric oxide synthase containing neurons in the dorsal raphe nucleus. In: Honig, A., van Praag, H.M., editors. *Depression: Neurobiological, Psychopathological and Therapeutic Advances*. New York: John Wiley. 179–196.
72. Suri, D., Teixeira, C.M., Cagliostro, M.K.C., Mahadevia, D., and Ansorge, M.S., 2015. Monoamine-sensitive developmental periods impacting adult emotional and cognitive behaviors. *Neuropsychopharmacology*. 40, 88–112.
73. Tanaka, J., Kariya, K. Nomura, M., 2003. Angiotensin II reduces serotonin release in the rat subfornical organ area. *Peptides*. 24, 881–887.
74. Thrasher, T.N., Keil, L.C., 1987. Regulation of drinking and vasopressin secretion: role of organum vasculosum laminae terminalis. *American Journal of Physiology*, 253, 108–120.
75. Tirado, C., Cortes, A., Bozinovic, F., 2008. Water balance in two South American Phyllotis desert rodents, *P. xanthopygus rupestris* and *P. darwini darwini*, *J. Arid Environ.* 72, 664–670.
76. Tse, W.S., Bond, A.J., 2002. Serotonergic intervention affects both social dominance and affiliative behaviour. *Psychopharmacology (Berl)*. 161, 324–330.