



**HAL**  
open science

## **Earliest humans in Europe: the age of TD6 Gran Dolina, Atapuerca, Spain**

Christophe Falguères, Jean-Jacques Bahain, Yuji Yokoyama, Juan Luis Arsuaga, Jose Maria Bermudez de Castro, Eudald Carbonell, James L Bischoff, Jean-Michel Dolo

► **To cite this version:**

Christophe Falguères, Jean-Jacques Bahain, Yuji Yokoyama, Juan Luis Arsuaga, Jose Maria Bermudez de Castro, et al.. Earliest humans in Europe: the age of TD6 Gran Dolina, Atapuerca, Spain. *Journal of Human Evolution*, 1999, 37, pp.343-352. 10.1006/jhev.1999.0326 . hal-03739352

**HAL Id: hal-03739352**

**<https://cnrs.hal.science/hal-03739352>**

Submitted on 13 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Earliest Humans in Europe : the age of TD6 Gran Dolina, Atapuerca, Spain**

**Christophe Falguères, Jean-Jacques Bahain & Yuji Yokoyama**

Laboratoire de Préhistoire du Muséum National d'Histoire Naturelle, UMR 6569 du CNRS,

Institut de Paléontologie Humaine, 1, Rue R. Panhard, 75013, Paris, France.

e-mail : falguere@mnhn.fr

**Juan Luis Arsuaga**

Departamento de Paleontologia, Universidad Complutense, 28040, Madrid, Spain.

e-mail : azara@eucmax.sim.ucm.es

**Jose Maria Bermudez de Castro**

Museo Nacional de Ciencias Naturales, CSIC, Calle Jose Abascal, 2, 28006, Madrid, Spain.

e-mail : mcnc54@mncn.csic.es

**Eudald Carbonell**

Universitat Rovira i Virgili, Facultats de Química i de Lletres, Imperial Tarraco 1,

43005, Tarragona, Spain. e-mail : ecr@astor.urv.es

**James L. Bischoff**

US Geological Survey, 345, Middlefield Road, Menlo Park, CA 94025, United States.

e-mail : jimb@octopus.wr.usgs.gov

**Jean-Michel Dolo**

Laboratoire de Mesure des Rayonnements Ionisants, BP 52, CEA, 91193, Gif-sur-Yvette, France. e-mail : dolo@babaorum.cea.fr

**Abstract**

Hominid remains found in 1994 from the stratified Gran Dolina karst-filling at the Atapuerca site in NE Spain were dated to somewhat greater than 780 ka based on palaeomagnetic measurements, making these the oldest known hominids in Europe (*sensu stricto*). We report new ESR and U-series results on teeth from four levels of the Gran Dolina deposit which confirm

the palaeomagnetic evidence, and indicate that TD6 (from which the human remains have been recovered) dates to the end of the Early Pleistocene. The results for the other levels are consistent with estimates based mainly on microfaunal evidence, and suggest that TD8, TD10 and TD11 date to the Middle Pleistocene.

**Key-words** : Early Pleistocene, Gran Dolina, Atapuerca, Combined U-series / ESR dating method.

## Introduction

The Sierra de Atapuerca, near the city of Burgos in north-central Spain, contains several archaeological and fossiliferous deposits of major importance for European prehistory. Those exposed in the abandoned railroad cut, Trinchera, are fillings of karst collapses that have been excavated since 1978 (Aguirre *et al.*, 1987 ; Bermudez de Castro *et al.*, 1995). The most important of these is the 18 m thick section at Gran Dolina (TD), which has yielded human bones associated with primitive lithic tools and a rich fauna characteristic of an Early Pleistocene age (*Mimomys savini* zone, Laplana & Cuenca-Bescos, 1997). The Gran Dolina is divided into eleven stratigraphic levels (TD1-TD11, see Parés & Perez-Gonzales, 1995), increasing in number upward (Figure 1). The fossil hominids excavated from TD6 (Aurora Stratum) have been proposed as a new species *Homo antecessor*, a possible common ancestor of modern humans and Neanderthals (Bermudez de Castro *et al.*, 1997). TD6 lies 1m below TD7 in which the Matuyama-Brunhes boundary was observed (Parés & Perez-Gonzales, 1995), dated to 780 ka (Shackleton *et al.*, 1990). Ideal dating materials, such as volcanic rocks or speleothems, are lacking. We have, therefore, applied combined electron spin resonance (ESR) and U-series dating methods to fossil tooth enamel of ungulates from levels TD6, TD8, TD10 and TD11.

## Methods

We utilized U-series (see details in Ivanovich & Harmon, 1992) and ESR methods (Grün, 1989; Ikeya, 1993; Rink, 1997), both independently and "combined". The ESR method is based on the detection of accumulated natural radiation damage resulting from environmental radioactivity. Both U-series and ESR methods are dependent on the mode of uranium uptake in the fossil teeth. Several mathematical models have been proposed to describe the effects of uranium accumulation, for example early uptake (EU) and linear uptake (LU). The EU model assumes that uranium is incorporated shortly after burial (Bischoff & Rosenbauer, 1981). The LU model supposes that uranium was incorporated at a constant rate since the time of burial (Ikeya, 1982). The recent uptake (RU) model corresponds to a late sublinear uranium uptake, continuously accelerating (Blackwell *et al.*, 1992). However, in most cases the uranium accumulation is somewhere between the first two cases. By combining U-series and ESR, it is possible to determine the history of uranium uptake using an uptake parameter  $p$  (Grün *et al.*, 1988). This describes more accurately the geochemical history of the tooth (Grün & Mc Dermott, 1994; Falguères *et al.*, 1997). This correction procedure does not apply to situations in which uranium has been lost. To use this procedure, the EU-ESR age must be older than the EU-U-series age.

## Samples and methodology

We analyzed ungulate teeth from TD6, TD8, TD10 and TD11, taking three coeval samples from each layer. Those from TD6 came from the bottom of the Aurora Stratum which is a muddy layer including some clasts and calcarenite fragments. This level contains the human remains which are contemporaneous with the analyzed teeth. Samples from TD8 were taken stratigraphically from below a hiatus marked by the disappearance of several micromammal species, most notably *Mimomys savini* (Laplana & Cuenca-Bescos, 1997). TD10 and TD11 levels mark a change in lithic assemblage (Carbonell *et al.*, 1995) and the appearance of characteristic Middle Pleistocene rodents. TD7 and TD9 are archaeologically and palaeontologically sterile. Most of the samples are equid molars that contain enamel, dentine and cementum.

The samples were in a good state of preservation. The different tissues were separated mechanically, then the outer surface of the enamel was removed using an electric drill to eliminate the effect of external alpha radiation. Nine enamel aliquots were irradiated using a calibrated  $^{60}\text{Co}$  gamma-ray source with dose of 50, 100, 200, 300, 400, 800, 1500, 3000, and 6000 Gy. The ESR measurements were performed on an EMX Bruker ESR spectrum (X-band, 9.82 GHz) with the following parameters : 10mW microwave power, 1G modulation amplitude, room temperature.

The equivalent doses (ED) were determined from the  $g=2.0018$  enamel signal with the additive method using an exponential fitting (see details in Yokoyama *et al.*, 1985). Each ESR measurement was repeated three times for each dose.

U-series analyses using alpha-ray spectrometry were made at the Institut de Paléontologie Humaine (IPH), Paris, according to the standard methods described by Bischoff *et al.* (1988).

The calculation for ESR ages was made using the following parameters :

- 1) A k-value of  $0.15 \pm 0.05$  (De Cannière *et al.*, 1986).
- 2) The age calculation was made following the cases : dentine-enamel-cementum and dentine-enamel-sediment for teeth without cementum.
- 3) The water content was estimated to be 3 wt%, 10 wt% and 25 wt% in the enamel, dentine, cementum, and sediment respectively. We give further details in "Results and discussion".
- 4) The initial thickness of the enamel layers varies by between 0.1 and 0.2 cm. The amount removed on each side was taken into account in the age calculation following Grün (1986).
- 5) Gamma-ray spectrometry with a high purity germanium detector was used to evaluate the external doses of the surrounding sediments which were collected at the exact position from where the teeth were excavated (see Table 2 caption). The dose rate is calculated from the U, Th and K contents of the environmental soil. These contents were measured by TL dosimetry, field gammametry, gamma-ray spectrometry in the laboratory, and other methods such as plasma atomic emission photometry, flame atomic emission spectrometry in order to evaluate the

uncertainties in annual dose rate determination. The method was tested for homogeneous rocks (granite, sandstone, lavas) (Faïn *et al.*, 1997), and on an archaeological site (Bechtel *et al.*, 1997).

The precision of the dose rate was in a range of 2 to 5% and showed a good agreement of the gamma-ray spectrometry results and those of the other methods.

6) The effect of Ra and Rn losses in enamel, dentine and cementum was determined by combining alpha-ray and gamma-ray results (Bahain *et al.*, 1992).

7) The cosmic dose was evaluated using the formula given by Yokoyama *et al.* (1982).

## Results and discussion

Table 1 summarizes the radiometric data of the TD samples. The U content in enamel varies between 0.1 and 2.9 ppm. For TD6 and TD8 samples, the U content in dentine and cementum, ranges from 7 to 27 ppm except for AT9602 (rhinoceros tooth) in which uranium content rises 57 ppm. TD10 and TD11 show higher U contents, ranging from 33 to 107 ppm, perhaps caused by closer proximity to the modern soil zone. The Ra and Rn losses in enamel are slight for most samples, but vary between 25 to 75% in the other two tissues, and are systematically higher in cementum than in dentine. All the samples show  $^{230}\text{Th} / ^{232}\text{Th}$  ratios of greater than 100 indicating that detrital thorium contamination is negligible. Table 2 details ESR dose rates. The external dose rate, measured by gamma-ray spectrometry at IPH includes the cosmic dose rate for which an average depth was determined for each layer. The contribution of external dose rate to the total dose rate, for US model ranges between 71 and 90% for TD6 and TD8 layers and between 38 to 62% for TD10 and TD11.

Nominal (EU) ESR and U-Th ages are shown in Table 3. The ESR ages were determined assuming a 25% water content in the sediment for all the samples. This value seems reasonable in light of measurements averaging around 30% for present-day water content of sediments sampled in the karst (see Bischoff *et al.*, 1997). However, because the Gran Dolina cave was open to the air after TD8 deposits and before TD10 and TD11, we used lower values for the age determination of TD10 and TD11 (Table 4). The results do not change beyond the error range for what we consider as a minimum water content versus the values used in this analysis (Table 4). The ESR results are considered as minimum ages, in as much as we used Grün's data (DATA) for beta attenuation (Grün, 1986). New measurements of beta attenuation, ROSY (Brennan *et al.*, 1997 ; Yang *et al.*, 1998), have shown that when there is little uranium uptake by teeth, which is the case for TD6 and TD8, ROSY ages (EU and LU) are about 5-10% older than DATA ages.

All samples (Table 3) have EU-ESR dates greater than EU uranium-series dates. This implies no U loss or U cycling, so it was possible to apply the coupled U/Th - ESR method based on the formula given by Grün *et al.* (1988). LU-ESR dates are generally 30-40% older than EU dates, but within-group variation among coeval samples is rather small and between group dates show a

coherent stratigraphic progression (Table 3). For example, EU-ESR ages from the three coeval samples of TD6 range between 626-685 ka while LU -ESR dates range between 765-807 ka. EU U-series dates are systematically younger than the ESR dates and appear to have little stratigraphic coherence. For instance, two samples from TD8 are at full isotopic equilibrium (>400 ka) whereas those from TD6 range from only 176-279 ka. Combined U-series / ESR age estimates (Table 3) are intermediate between the EU and LU ESR dates, show reasonably small within-group variation, and increase coherently with stratigraphic depth (Figure 1).

Dates of  $337 \pm 51$  ka,  $379 \pm 57$  ka and  $418 \pm 63$  ka (weighted mean age of  $372 \pm 33$  ka) for TD10 and  $308 \pm 46$  ka,  $332 \pm 50$  ka and  $390 \pm 59$  ka (weighted mean age of  $337 \pm 29$  ka) for TD11 have overlapping ranges suggesting possible rapid sedimentation for these units, and a correlation with oxygen isotope stages 9 to 11. All p-values range between 0 (LU) and -1 (EU), except for one which  $p = +0.365$ . The emergence of Middle Palaeolithic production technology in TD10 and TD11 (Carbonell *et al.*, 1995), dated between 300 and 400 ka, is consistent with that observed at the French sites of la Micoque and Orgnac 3 (Falguères *et al.*, 1997, 1988). Dates on these upper levels of Gran Dolina can also be correlated with radiometric dates on the hominid bone assemblage at Sima de los Huesos (Bischoff *et al.*, 1997).

Dates for TD8 are  $586 \pm 88$  ka,  $606 \pm 91$  ka and  $615 \pm 92$  ka (weighted mean age of  $602 \pm 52$  ka). The mode of U-uptake for TD8 samples is distinctly different from those above. For example, p enamel and p dentine, for AT9604, indicate a closed system, while p cementum suggests a sublinear late uptake. Pollen, such as *Castanea* and *Quercus* associated with *Betula*, *Acer*, and *Alnus* (Garcia Anton, 1998), and the presence of a stalagmitic crust, suggest a warm to temperate climate for this level. These results and the coupled-dates suggest correlation of TD8 to oxygen isotope stage 15.

For TD6, two of the three coupled US-ESR dates are  $770 \pm 116$  ka and  $762 \pm 114$  ka, and the third is  $676 \pm 101$  ka. Sample AT9601 exhibits a younger age than the two others because of cementum and dentine p-values indicating an early u-uptake. The US-ESR ages of AT9602 and AT9603 are close to those of the LU-ESR, with p-values showing a later uptake. The presence of an inverse negative polarity indicates an age of >780 ka for TD6. The US-ESR mean age of  $731 \pm 63$  ka is within error of these earlier observations. The palynological record, with *Pinus*, *Quercus* and Cupressaceae, associated with Mediterranean taxa, such as *Olea*, *Ceratonia*, *Celtis* and *Pistacia*, suggest temperate and wet climatic conditions for the hominid-bearing level (Garcia Anton, 1998). In addition, the rodent assemblage (with *Mimomys savini*, *Pliomys episcopalpis*, *Iberomys huescarensis*) is considered characteristic of the late Early Pleistocene (Cuenca-Bescos *et al.*, 1999).

The combination of palaeomagnetic data and US-ESR ages provides confirmation of an age range of between 780 and 857 ka for TD6 (2-sigma error on the weighted mean age). Because the palaeoenvironmental record suggests that the upper TD6 level was deposited under wet, temperate conditions our results indicate that the Aurora Stratum correlates to oxygen isotope stages 21 or 19. These dates obtained directly on the level which yielded the human remains corroborate the presence of *Homo antecessor* at the end of the Early Pleistocene and furthermore confirm the applicability of ESR for dating ancient sites. The agreement between the ESR dates and independent palaeomagnetic measurements confirms the reliability of ESR dating in the 800 ka time range at this site.

In their entirety, the combined US-ESR age estimates at Gran Dolina indicate human occupation from about 800 to 300 ka. These results strengthen the claim of an early settlement in southwest Europe, and support claims for other Pre-Acheulian archaeological sites, such as Vallonnet (Lumley *et al.*, 1988) and Monte Poggiolo (Gagnepain *et al.*, 1996), but at which no human fossils have yet been found.

#### Acknowledgements

This research is supported by a grant from the Centre National de la Recherche Scientifique, France. We are grateful to N. Mehidi and B. Morin for their help concerning the U-series and ESR analyses respectively. We thank Terry Harrison for his valuable editorial comments and three anonymous referees for critical comments on the manuscript.



**Figures caption**




Figure 1 : Combined ESR-U-series ages with errors ( $\pm 15\%$ ) of fossil teeth from TD6 to TD11 in relation to oxygen isotope stages taken from Bassinot *et al.* (1994). Shaded regions correspond to archaeologically sterile layers (TD7 and TD9). The dotted line in TD8 represents a hiatus marked by the disappearance of microfauna.

## References

- Aguirre, E., Carbonell, E. & Bermúdez de Castro, J.M. Eds. (1987). *El Hombre fósil de Ibeas y el Pleistoceno de la Sierra de Atapuerca*. Junta de Castilla y León, Consejería de Cultura y Bienestar Social, Valladolid, Spain.
- Bahain, J.J., Yokoyama, Y., Falguères, C. & Sarcia, M.N. (1992). ESR dating of tooth enamel: a comparison with K-Ar dating. *Quat. Sci. Rev.* **11**, 245-250.
- Bassinot, F.C., Labeyrie, L., Vincent, E., Quidelleur, X., Shackleton, N.J. & Lancelot, Y. (1994). The astronomical theory of climate and the age of the Brunhes-Matuyama magnetic reversal. *Earth Planet. Sci. Lett.* **126**, 91-108.
- Bechtel, F., Guibert, P., Schvoerer, M., Vartanian, E., Faïn, J., Miallier, D., Montret, M., Pilleyre, T., Sanzelle, S., Bahain, J.J., Falguères, C., Tripiet, J., Poupeau, G., Mercier, N. & Valladas, H. (1997). Evaluation de l'incertitude de mesure de la dose annuelle en datation par luminescence (TL et OSL) et par RPE. *Revue d'Archéométrie* **21**, 21-27.
- Bermúdez de Castro, J.M., Arsuaga, J.L., Carbonell, E., Rosas, A., Martínez, I. & Mosquera, M. (1997). A hominid from the Lower Pleistocene of Atapuerca, Spain : Possible ancestor to Neandertals and modern humans. *Science* **276**, 1392-1395.
- Bermúdez de Castro, J.M., Carbonell, E. & Arsuaga, J.L. Eds. (1995). *Human Evolution in Europe and the Atapuerca Evidence*. Consejería de Cultura y Turismo, Junta de Castilla y León, Valladolid, Spain.
- Bischoff, J.L. & Rosenbauer, R.J. (1981). Uranium-series dating of human skeletal remains from the Del Mar and Sunnyvale sites, California. *Science* **213**, 1003-1005.

- Bischoff, J.L., Rosenbauer, R.J., Tavano, A. & Lumley, H. de (1988). A test of uranium-series dating of fossil tooth enamel : results from Tournal cave, France. *Applied Geochem.* **3**, 135-141.
- Bischoff, J.L., Fitzpatrick, J.A., Léon, L., Arsuaga, J.L., Falguères, C., Bahain, J.J. & Bullen, T. (1997). Geology and preliminary dating of the hominid-bearing sedimentary fill of the Sima de los Huesos chamber, Cueva Mayor of the Sierra de Atapuerca, Burgos, Spain. *J. hum. Evol.* **33**, 129-154.
- Blackwell, B., Porat, N., Schwarcz, H.P. & Debénath, A. (1992). ESR dating of tooth enamel : comparison with  $^{230}\text{Th}$  /  $^{234}\text{U}$  speleothem dates at La Chaise-de-Vouthon (Charente), France. *Quat. Sci. Rev.* **11**, 231-244.
- Brennan, B.J., Rink, W.J., McGuirl, E.L., Schwarcz, H.P. & Prestwich, W.V. (1997). Beta doses in tooth enamel by "One-Group" theory and the Rosy ESR dating software. *Radiation Measurements* **27**, 307-314.
- Carbonell, E., Bermúdez de Castro, J.M., Arsuaga, J.L., Diez, J.C., Rosas, A., Cuenca-Bescos, G., Sala, R., Mosquera, M. & Rodríguez, X.P. (1995). Lower Pleistocene hominids and artifacts from Atapuerca-TD6 (Spain). *Science* **269**, 826-830.
- Cuenca-Bescos, G., Laplana, C & Canudo, J.I. (1999). Biochronological implications of the Arvicolidae (Rodentia, Mammalia) from the Lower Pleistocene hominid-bearing level Trincheras Dolina 6 (TD6, Atapuerca, Spain). *J. Hum. Evol.* , this volume.
- De Cannière, P., Debuyst, R., Dejehet, F., Apers, D. & Grün, R. (1986). ESR dating : a study of  $^{210}\text{Po}$ -coated geological and synthetic sample. *Nuclear Tracks and Radiation Measurements* **11**, 211-220.

Faïn, J., Miallier, D., Montret, M., Pilleyre, T., Sanzelle, S., Soumana, S., Bechtel, F., Guibert, P., Schvoerer, M., Vartanian, E., Mercier, N., Valladas, H., Bahain, J.J., Falguères, C., Tripier, J. & Joron, J.L. (1997). Intercomparaisons dosimétriques dans le cadre de la datation par thermoluminescence et par résonance paramagnétique électronique. Validation de milieux naturel-témoins de la région de Clermont-Ferrand (Puy-de-Dôme, France). *Revue d'Archéométrie* **21**, 29-34.

Falguères, C., Shen, G. & Yokoyama, Y. (1988). Datation de l'aven d'Orgnac III : comparaison par les méthodes de la résonance de spin électronique (ESR) et du déséquilibre des familles de l'uranium. *L'Anthropologie* **92**, 727-730.

Falguères, C., Bahain, J.J. & Saleki, H. (1997). U-series and ESR dating of teeth from Acheulian and Mousterian levels at la Micoque (Dordogne, France). *J. Archaeol. Sci.* **24**, 537-545.

Gagnepain, J., Hedley, I., Bahain, J.J., Falguères, C., Laurent, M., Peretto, C., Wagner, J.J. & Yokoyama, Y. (1996). Synthèse des données radiochronologiques et paléomagnétiques du site de Ca'Belvedere di Monte Poggiolo (Romagna, Italie) et de son environnement géologique. *XIIIe Congrès Union Internationale des Sciences Préhistoriques et Protohistoriques*, abstract, **2**, 129-130.

Garcia Anton, M. (1998). Reconstrucciones de paleovegetacion en Atapuerca según analisis polinico. *In : Atapuerca y la evolucion humana* (E. Aguirre, Ed.), Fundacion Ramon Areces, 63-71.

Grün, R. (1986). Beta-dose attenuation in thin layers. *Ancient TL* **4**, 1-8.

Grün R. (1989). Electron spin resonance (ESR) dating. *Quat. Internat.* **1**, 65-109.

- Grün, R. & Mc Dermott, F. (1994). Open system modelling for U-series and ESR dating of teeth. *Quat. Geochronol. (Quat. Sci. Rev.)* **13**, 121-125.
- Grün, R., Schwarcz, H.P. & Chadam, J. (1988). ESR dating of tooth enamel : coupled correction for U-uptake and U-series disequilibrium. *Nuclear tracks* **14**, 237-241.
- Ikeya, M. (1982). A model of linear accumulation of uranium for ESR age of Heidelberg (Mauer) and Tautavel bones. *Japan. J. Applied Physics* **22**, 763-765.
- Ikeya, M. (1993). *New Applications of Electron Spin Resonance-Dating, Dosimetry and Microscopy*. Singapore: World Scientific.
- Ivanovich, M. & Harmon, R.S. (1992). *Uranium-series Disequilibrium : Applications to Earth, Marine and Environmental Sciences*. 2nd Edition. Oxford: Clarendon Press.
- Laplana, C. & Cuenca-Bescos, G. (1997). Los arvicolidos (Arvicolidae, Rodentia) del limite Pleistoceno Inferior-Medio en el relleno carstico Trinchera Dolina (Sierra de Atapuerca, Burgos, Espana). *XIII Jornadas de Paleontologia* , 192-195.
- Lumley, H. de, Fournier, A., Krzepakowska, J. & Echassoux, A. (1988). L'industrie du Pléistocène inférieur de la grotte du Vallonnet, Roquebrune-Cap-Martin, Alpes-Maritimes. *L'Anthropologie* **92**, 501-614.
- Parés, J.M. & Pérez-Gonzales, A. (1995). Palaeomagnetic age for hominid fossils at Atapuerca archaeological site, Spain. *Science* **269**, 830-832.
- Rink, W.J. (1997). Electron spin resonance (ESR) dating and ESR applications in Quaternary science and archaeometry. *Radiation Measurements* **27**, 975-1025.

Shackleton, N.J., Berger, A. & Peltier, W.R. (1990). An alternative astronomical calibration of the lower Pleistocene timescale based on ODP Site 677. *Trans. Royal Soc. Edinburgh : Earth Sciences* **81**, 251-261.

Yang, Q., Rink, W.J., & Brennan, B.J. (1998). Experimental determinations of beta attenuation in planar dose geometry and application to ESR dating of tooth enamel. *Radiation Measurements* **29**, 663-671.

Yokoyama, Y., Nguyen, H.V., Quaegebeur, J.P. & Poupeau, G. (1982). Some problems encountered in the evaluation of annual dose rate in the Electron Spin Resonance dating of fossil bones. *PACT* **6**, 103-115.

Yokoyama, Y., Falguères, C. & Quaegebeur, J.P. (1985). ESR dating of quartz from Quaternary sediments : first attempt. *Nuclear Tracks* **10**, 921-928.



Sample	U content (ppm)	$^{234}\text{U}/^{238}\text{U}$	$^{230}\text{Th}/^{234}\text{U}$	$^{226}\text{Ra}/^{230}\text{Th}$	$^{222}\text{Rn}/^{230}\text{Th}$	T enamel ( $\mu\text{m}$ )	Removed enamel ( $\mu\text{m}$ )	ED (Gray)
<b>TD11</b>								
AT9610E	1.2	$1.38 \pm 0.05$	$0.87 \pm 0.03$	1	1	1400	400	$549 \pm 22$
D	56.9	$1.36 \pm 0.03$	$0.80 \pm 0.02$	0.60	0.35			
C	33.6	$1.42 \pm 0.03$	$0.86 \pm 0.02$	0.44	0.28			
AT9705E	1.4	$1.33 \pm 0.04$	$0.77 \pm 0.03$	0.82	0.82	1200	200	$436 \pm 13$
D	41.5	$1.29 \pm 0.03$	$0.73 \pm 0.02$	0.45	0.24			
AT9707E	1.2	$1.30 \pm 0.04$	$0.96 \pm 0.03$	0.77	0.77	1700	100	$512 \pm 13$
D	39.9	$1.33 \pm 0.03$	$0.94 \pm 0.03$	0.41	0.24			
<b>TD10</b>								
AT9606E	2.9	$1.41 \pm 0.02$	$0.85 \pm 0.03$	1	1	1200	200	$1\ 083 \pm 44$
D	95.4	$1.33 \pm 0.02$	$0.89 \pm 0.03$	0.53	0.31			
C	66.8	$1.40 \pm 0.02$	$0.92 \pm 0.02$	0.37	0.24			
AT9608E	2.6	$1.23 \pm 0.03$	$0.70 \pm 0.02$	1	1	1800	400	$836 \pm 40$
D	101.6	$1.32 \pm 0.02$	$0.86 \pm 0.02$	0.41	0.24			
C	72.8	$1.23 \pm 0.02$	$0.89 \pm 0.02$	0.40	0.24			
AT9703E	2.2	$1.10 \pm 0.03$	$0.73 \pm 0.03$	0.98	0.98	1200	200	$726 \pm 14$
D	107.4	$1.13 \pm 0.01$	$0.78 \pm 0.01$	0.4	0.26			
<b>TD8</b>								
AT9604E	0.2	$1.20 \pm 0.08$	$1.15 \pm 0.07$	1	1	1000	100	$476 \pm 13$
D	9.1	$1.06 \pm 0.04$	$0.99 \pm 0.04$	0.73	0.46			
C	10.6	$1.01 \pm 0.03$	$0.74 \pm 0.03$	0.45	0.30			
AT9605E	0.5	$0.95 \pm 0.05$	$1.03 \pm 0.05$	1	0.76	1500	500	$650 \pm 52$
D	19.3	$1.04 \pm 0.03$	$0.74 \pm 0.03$	0.60	0.46			
C	27.1	$1.11 \pm 0.03$	$0.90 \pm 0.03$	0.82	0.50			
AT9702E	0.6	$1.20 \pm 0.06$	$0.83 \pm 0.04$	1	0.59	2000	500	$607 \pm 17$
D	27.2	$1.08 \pm 0.01$	$0.73 \pm 0.01$	0.44	0.27			
<b>TD6</b>								
AT9601E	0.4	$1.16 \pm 0.06$	$0.96 \pm 0.05$	1	1	1100	100	$844 \pm 30$
D	13.9	$1.20 \pm 0.03$	$1.02 \pm 0.04$	0.66	0.36			
C	10.7	$1.22 \pm 0.05$	$1.07 \pm 0.05$	0.30	0.22			
AT9602E	0.3	$1.16 \pm 0.06$	$0.83 \pm 0.05$	1	1	2000	500	$840 \pm 42$
D	56.9	$1.10 \pm 0.02$	$0.76 \pm 0.02$	0.65	0.39			
AT9603E	0.4	$1.18 \pm 0.05$	$0.97 \pm 0.04$	0.88	0.60	1200	200	$846 \pm 56$
D	7.0	$1.36 \pm 0.06$	$0.84 \pm 0.04$	0.71	0.36			

---

C	7.2	$1.15 \pm 0.04$	$0.89 \pm 0.03$	0.44	0.26
---	-----	-----------------	-----------------	------	------

---

Table 1 : ESR and U-series data on fossil ungulate teeth from Atapuerca Gran Dolina. E = enamel, D = dentine, C = cementum. Uncertainties for isotopic activity ratios are given with  $\pm 1$  sigma. The initial (T) and removed enamel thicknesses are used for the age calculation. Equivalent doses (ED) have been determined using an exponential fitting with the additive dose method.

Sample	$(\beta + \gamma)$ sediment + cosmic ( $\mu\text{Gy} / \text{a}$ )	Early U-uptake (EU model)				Linear U-uptake (LU model)				Uranium-series U-uptake		
		Internal dose rate ( $\alpha + \beta$ ) enamel ( $\mu\text{Gy} / \text{a}$ )	$\beta$ dentine ( $\mu\text{Gy} / \text{a}$ )	$\beta$ cementum ( $\mu\text{Gy} / \text{a}$ )	Total dose rate ( $\mu\text{Gy} / \text{a}$ )	Internal dose rate ( $\alpha + \beta$ ) enamel ( $\mu\text{Gy} / \text{a}$ )	$\beta$ dentine ( $\mu\text{Gy} / \text{a}$ )	$\beta$ cementum ( $\mu\text{Gy} / \text{a}$ )	Total dose rate ( $\mu\text{Gy} / \text{a}$ )	Internal dose rate ( $\alpha + \beta$ ) enamel ( $\mu\text{Gy} / \text{a}$ )	$\beta$ dentine ( $\mu\text{Gy} / \text{a}$ )	$\beta$ cementum ( $\mu\text{Gy} / \text{a}$ )
<b>TD11</b>												
AT9610	1110.0	617.9	219.0	125.2	2072.1	269.0	106.6	61.2	1546.8	440.5	140.0	
AT9705	817.2	708.6	210.5	----	1736.3	316.8	104.1	----	1238.1	384.3	111.1	
AT9707	656.9	652.4	179.4	----	1488.7	287.7	87.9	----	1032.5	507.5	138.7	
<b>TD10</b>												
AT9606	1076.1	1882.8	517.2	349.3	3825.4	859.2	255.6	172.9	2363.8	1141.9	373.8	
AT9608	1120.2	1457.8	290.8	205.8	3074.6	672.4	144.5	102.7	2039.8	543.9	189.1	
AT9703	1289.5	1101.4	537.3	----	2928.2	501.9	265.8	----	2057.2	536.3	329.1	
<b>TD8</b>												
AT9604	577.5	144.3	66.0	69.0	856.8	57.2	31.8	34.1	700.6	146.3	62.5	
AT9605	795.4	211.8	64.3	95.6	1167.1	99.1	31.4	45.8	971.7	195.4	23.0	
AT9702	847.2	296.2	63.4	----	1206.8	123.1	31.3	----	1001.6	117.7	22.1	
<b>TD6</b>												
AT9601	922.4	248.4	106.1	74.0	1350.9	98.5	48.5	34.3	1103.7	159.4	93.2	
AT9602	987.7	210.0	143.4	----	1341.1	83.4	69.5	----	1140.6	67.8	46.2	
AT9603	922.5	225.6	46.7	39.6	1234.4	87.4	20.4	18.8	1049.1	139.3	17.3	

Table 2 : Components of dose rates for three models of teeth and sediment from TD6 to TD11. External dose rates correspond to both sediment ( $\beta + q$ ) and cosmic dose. About 100g of sediment including rock fragments when present were measured at least one month after packing in a box. Sediments have been sampled in the same square and at the same height as the teeth except for TD6 for which we sampled in the nearest square because this level has been completely excavated and does not exist anymore. The same was done for AT9610 which was extracted in 1984. AT9602 is a rhinoceros molar. This species does not develop cementum layer. AT9702 is a bovid molar in which the amount of cementum was not sufficient to be analyzed. A similar observation was made for equid molars AT9703, AT9705 and AT9707. For these samples, the enamel was directly in contact with the sediment. When the case dentine-enamel-cementum was used, the cementum layer was thick enough to shield the enamel from sediment beta particles.

	ESR Age (ka)		U-Th Age (ka)			ESR and U-series combined data			Age (ka)
	EU	LU	Enamel	Dentine	Cementum	p enamel	p dentine	p cementum	
<b>TD11</b>									
AT9610	265 ± 28	355 ± 38	185+19/-16	154+9/-8	181+13/-12	- 0.675	- 0.469	- 0.656	308 ± 46
AT9705	251 ± 26	352 ± 36	145+12/-10	132±8	---	- 0.305	- 0.132	---	332 ± 50
AT9707	344 ± 35	496 ± 51	246+41/-29	231+24/-20	---	- 0.785	- 0.736	---	390 ± 59
<b>TD10</b>									
AT9606	283 ± 30	458 ± 49	175+15/-13	201+19/-16	212+16/-14	- 0.462	- 0.616	- 0.676	379 ± 57
AT9608	272 ± 30	410 ± 45	123+7/-6	185+12/-11	208+16/-14	+ 0.365	- 0.449	- 0.584	418 ± 63
AT9703	248 ± 26	353 ± 36	133+12/-11	156±5	---	- 0.121	- 0.370	---	337 ± 51
<b>TD8</b>									
AT9604	556 ± 67	680 ± 81	>400	398+∞/-102	147+15/-13	- 1.000	- 0.911	+ 0.564	586 ± 88
AT9605	557 ± 72	669 ± 86	>420	144+12/-11	229+35/-26	- 0.918	+ 0.683	- 0.432	606 ± 91
AT9702	503 ± 52	606 ± 63	175+28/-21	139±5	---	+ 0.067	+ 0.804	---	615 ± 92
<b>TD6</b>									
AT9601	626 ± 66	765 ± 78	279+148/-58	374+∞/-81	>350	- 0.663	- 0.896	- 0.995	676 ± 101
AT9602	626 ± 70	736 ± 82	176+37/-26	151+10/-9	---	+ 0.353	+ 0.940	---	762 ± 114
AT9603	685 ± 82	807 ± 97	279+96/-49	173+22/-18	215+33/-24	- 0.632	+ 0.324	- 0.159	770 ± 116

Table 3 : ESR and U-series age estimates for fossil ungulate teeth from Atapuerca Gran Dolina. A 25% water content in the sediment was used.

	15% sed 5% dent+cem	15% sed 10% dent+cem	20% sed 10% dent+cem	25% sed 10% dent+cem
<b>TD11</b>				
AT9610	280 ± 42	282 ± 42	294 ± 44	308 ± 46
AT9705	302 ± 45	303 ± 45	317 ± 48	332 ± 50
AT9707	363 ± 54	366 ± 55	378 ± 57	390 ± 59
<b>TD10</b>				
AT9606	352 ± 53	357 ± 54	368 ± 55	379 ± 57
AT9608	379 ± 57	383 ± 57	400 ± 60	418 ± 63
AT9703	304 ± 46	307 ± 46	321 ± 48	337 ± 51

Table 4 : US-ESR ages for TD10 and TD11 samples using different water contents in the sediment (sed), dentine (dent) and cementum (cem).