



HAL
open science

Mitochondrial oxidative phosphorylation response overrides glucocorticoid-induced stress in a reptile

Yann Voituron, Damien Roussel, Jean-François Le Galliard, Andréaz Dupoué, Caroline Romestaing, Sandrine Meylan

► **To cite this version:**

Yann Voituron, Damien Roussel, Jean-François Le Galliard, Andréaz Dupoué, Caroline Romestaing, et al.. Mitochondrial oxidative phosphorylation response overrides glucocorticoid-induced stress in a reptile. *Journal of Comparative Physiology B*, 2022, 192 (6), pp.765-774. 10.1007/s00360-022-01454-5 . hal-03764878

HAL Id: hal-03764878

<https://cnrs.hal.science/hal-03764878>

Submitted on 30 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Mitochondrial oxidative phosphorylation resilience overrides glucocorticoid-induced**
2 **stress in a reptile**

3
4
5
6 4 Yann Voituron^{1*}, Damien Roussel¹, Jean-François Le Galliard^{2,3}, Andréaz Dupoué², Caroline
7
8 Romestaing¹ and Sandrine Meylan²
9

10
11 6
12
13 7 1. Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés (U.M.R. CNRS 5023),
14
15 8 Université Claude Bernard Lyon1, Université de Lyon; Bd du 11 novembre 1918, Bât. Darwin
16
17 C; 69622 VILLEURBANNE Cedex, France
18
19

20
21 10 2. Sorbonne Université, CNRS, IRD, INRA, Institut d'écologie et des sciences de
22
23 11 l'environnement (IEES), Paris, France
24

25
26 12 3. Centre de recherche en écologie expérimentale et prédictive (CEREEP-Ecotron
27
28 13 IleDeFrance), Département de biologie, Ecole normale supérieure, CNRS, PSL University,
29
30 14 77140 Saint-Pierre-lès-Nemours, France
31
32

33 15
34
35
36 16
37
38 17 * To whom correspondence should be addressed: Tel: +33 4 72 43 15 20; Fax: +33 4 72 43 11
39
40 18 71; E-mail: yann.voituron@univ-lyon1.fr
41
42

43 19
44
45 20 **Running title:** Mitochondrial resilience under chronic stress in a reptile
46
47
48 21
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

22 **Abstract**

1
2
3 23 Stress hormones and their impacts on whole-organism metabolic rates are usually considered
4
5 24 as appropriate proxies for animal energy budget that is the basement of numerous concepts and
6
7 25 models aiming at predicting individual and population responses to environmental stress.
8
9
10 26 However, the kinetics of energy re-allocation under stress make the link between metabolism
11
12 27 and corticosterone complex and still unclear. Using ectopic application of corticosterone for 3
13
14 28 days, 11 days and 21 days, we simulated acute, mild-time and chronic stress in a lizard (*Zootoca*
15
16 29 *vivipara*). We then investigated whole-organism metabolism, muscle cellular O₂ consumption
17
18 30 and liver mitochondrial oxidative phosphorylation processes (O₂ consumption and ATP
19
20 31 production) and ROS production. The data showed that (i) while muscle is only slightly
21
22 32 impacted, stress regulates the mitochondrial functionality in a time-dependent manner with
23
24 33 contrasted pictures between acute, mild-time and chronic stress that is poorly described by the
25
26 34 whole-organism metabolism and (ii) an unexpected “pathway” to obtain mitochondrial
27
28 35 resilience. While acute stress is characterized by a decreased ATP synthesis rate and high H₂O₂
29
30 36 release with no change in the rate of oxygen consumption, the mild-time stress reduced all three
31
32 37 fluxes. Application of corticosterone for 21 days induced a return to initial oxidative
33
34 38 phosphorylation capacities with lower H₂O₂ production allowing a better capacity to respond to
35
36 39 the energetic demands imposed by the environment. So, the global mitochondrial functioning
37
38 40 has to be considered in order to better understand the proximal causes of resource allocation
39
40 41 under stress periods.
41
42
43
44
45
46
47
48
49
50
51

52 43 **Keywords:** Corticosterone, Lizard, Metabolism, Mitochondrial efficiency, ROS and ATP
53
54 44 production, Oxygen consumption
55
56
57
58
59
60
61
62
63
64
65

47 **Introduction**

48 Glucocorticoids (GCs) secreted by the hypothalamus–pituitary–adrenal axis are crucial
49 hormones orchestrating the tradeoff between survival and different physiological processes
50 when vertebrates are facing stressors. However, physiological consequences are known to be
51 different in accordance to short-term versus long-term elevations in GCs production. While
52 many of the acute effects of GCs consist of mobilizing energy, chronically elevated circulating
53 GCs enhance energy storage (Harris 2015, Sapolsky et al. 2000). Similarly, acute GC elevations
54 can activate the adaptive immune system, whereas chronic GC elevations are linked to
55 suppressed cell-mediated leucocyte trafficking (Sorrells et al. 2009; Dhabhar and Mc Ewen
56 1997). Such biphasic responses in which acute and chronic stress or corticosteroid exposure
57 induce contrasting and sometimes opposite effects have not yet been fully elucidated. While
58 some studies stipulate potential differential actions of different nuclear receptors of GCs (e.g.,
59 Joels and de Kloet 1992), others link it to larger magnitude changes in gene expression under
60 acute stress rather than chronic stress (e.g., Wang et al. 2004).

61 Other studies demonstrated rapid, non-genomic effects of GCs (de Kloet, 2013) explained at
62 least partly by the GCs translocation to the mitochondria that provide ATP through the Krebs
63 tricarboxylic-acid cycle and oxidative phosphorylation (Du, 2009, Lee et al. 2013). Indeed, *in*
64 *vitro* (with cortical neuronal cultures) and *in vivo* (with brains from male rats) experiments have
65 demonstrated that mitochondrial oxidation, membrane potential, mitochondrial complex I
66 activity and mitochondrial encoded genes (i.e. NADH dehydrogenase 1, 3, 6 and ATP synthase
67 6) are modulated by corticosterone in a dose and time-dependent manner (Du et al. 2009; Hunter
68 et al. 2016). In parallel, a reduction of the activity of specific mitochondrial electron transport
69 chain complexes together with an increased mitochondrial Reactive Oxygen Species (ROS)
70 production has been shown under chronic glucocorticoid treatment (Manoli et al. 2007), which
71 could impact the oxidative balance (Costantini 2011). Altogether, these physiological studies

1
2
3
4
5 72 thus demonstrated (i) a time adjustments of energetic resources in response to stress and (ii) a
6
7 73 direct linkage between GC secretion in response to stressor and mitochondrial physiology.
8
9 74 These observations led to the emergent concept of “mitochondrial allostatic load” that is the
10
11 75 mitochondrial analog of allostatic load in the broader context of stress biology (McEwen and
12
13 76 Wingfield 2003). The mitochondrial allostatic load can be defined as the “deleterious structural
14
15 77 and functional changes that mitochondria undergo” in response to elevated glucocorticoid
16
17 78 secretion under a chronic stress situation (Picard et al. 2014, 2018). The mitochondria are the
18
19 79 major ATP cell generators (Lehninger et al. 1993) sustaining physiological processes
20
21 80 underlying both maintenance and physiological responses to environmental variations.
22
23 81 Mitochondrial energy transduction system also continually produces ROS which can be
24
25 82 involved in cell signaling and/or physiopathology (Starkov, 2008). Unfortunately, at that time,
26
27 83 no direct assessments of mitochondrial functionality parameters like ATP and ROS under
28
29 84 chronic GCs production were available. Yet such data are crucial to understand functional
30
31 85 consequences of previously described responses to GC secretion, such as alteration of gene
32
33 86 expression and/or epigenetic modifications (Hunter 2016), increased oxidized mtDNA and
34
35 87 mitochondrial fission (Picard et al. 2014). A better knowledge of the temporal dynamics of
36
37 88 mitochondrial responses to GC secretion will undoubtedly have biomedical applications but
38
39 89 will also provide new visions in evolutionary biology (Eisner et al. 2018).

40
41
42
43
44
45 90 The objective of this study was to explore the time-dependent metabolic responses at
46
47 91 the whole-body and mitochondrial levels in lizards (*Zootoca vivipara*) exposed to 3 days, 11
48
49 92 days and 21 days of corticosterone treatment. This protocol provides the temporal dynamic of
50
51 93 the mitochondrial responses to GC secretion and answers the following three questions. First
52
53 94 question: do short-term *versus* long-term elevations in GCs production induce similar
54
55 95 physiological responses in ectotherms than in endotherms? Indeed, the commonly accepted
56
57 96 conclusions on time effects on GCs impacts are strongly biased toward endotherms (mammals
58
59
60
61
62
63
64
65

1 97 and birds), neglecting the vast majority of ectothermic metazoans. Recent experimental
2 98 approaches suggest different mitochondrial responses under stress in some ectotherms (e.g.,
3
4 99 Cote et al. 2010; Voituron et al. 2017). Comparative analyses demonstrate that evolution shapes
5
6
7 100 GCs variations which leads to a different impact between ectotherms and endotherms (Jessop
8
9 101 et al. 2013; Vitousek et al. 2019). Second question: are ATP and ROS productions equivalently
10
11 102 impacted by GCs? Third question: given that the liver and muscle contribute about 50% of body
12
13
14 103 O₂ consumption at rest and more during activity (Rolfe and Brown, 1997), does whole-
15
16 104 organism metabolism correctly reflect energetic adjustments at a cellular level of organization?
17
18
19
20 105

21 22 106 **Materials and methods**

23 24 25 107 *Capture and rearing condition*

26
27 108 The common lizard (*Zootoca vivipara*) is a small lacertid species (adult's snout-vent length
28
29 109 SVL ranging from 50–70 mm) widely distributed across Eurasia. In spring 2016, 49 sub-adult
30
31 110 males (1-year-old) were captured by hand in outdoor enclosures (10 × 10 m) at CEREEP
32
33 111 (Centre de Recherche en Ecologie Expérimentale et Prédictive; Saint-Pierre-lès-Nemours,
34
35 112 France, 48°17'N, 2°41'E) field station from May 16th to May 25th. Animals were measured
36
37 113 for body size (SVL, ± 0.5 mm) and body mass (± 1 mg). All animals were maintained in
38
39 114 individual terraria (25 × 15 × 16 cm) with a shelter, peat soil as substrate and opportunities for
40
41 115 optimal thermoregulation. We used incandescent light bulbs (25 W) for 8 hours per day from
42
43 116 09:00 to 17:00 local time to ensure a thermal gradient ranging from 17-23°C to 35-38°C. We
44
45 117 provided lizards with water *ad libitum* and, every other day, with 300 ± 20 mg of food (*Acheta*
46
47 118 *domestica*).
48
49
50
51

52 119 *Experimental design and whole-organism assays*

53
54
55 120 The study was performed between June 12th and July 12th 2016. At the start of the experiment,
56
57 121 animals were distributed in six groups corresponding to two experimental treatments
58
59
60
61
62
63
64
65

122 (corticosterone enhancement [CORT] and placebos [CONT]) and three treatment exposure
123 groups (3 days [3D], 11 days [11D] and 21 days [21D] of exposure to experimental treatments
124 until the functional analyses of mitochondria). Similar sample sizes (N=8) were used in each
125 group except in the CORT-21D group (N=9). We repeatedly measured the body mass and
126 whole-organism metabolic expenditure 3 days before the start of the experiments, and the day
127 before the cellular and mitochondrial measurements meaning days 2, 10 and 20 (in 3D, 11D
128 and 21D groups respectively) after the start of the experiment. Before each whole-organism
129 assay, the animals were left without food for 3 days to reach the post-absorptive state. After the
130 final assay, the animals were fed and treated with corticosterone according to their treatment
131 group. Then, the animals were euthanized to perform functional analyses of mitochondria (see
132 below and Figure 1).

133 Whole-organism metabolic expenditure at night was quantified with closed respirometry
134 techniques as previously described (Foucart et al., 2014). We measured oxygen consumption
135 and carbon production overnight (approximately 20:00–08:00h) in a dark climatic chamber
136 (AQUALYTIC® TC 135S, Dortmund, Germany). Trials were carried out at $25\pm 1^\circ\text{C}$, which
137 correspond to body temperature at which functional analyses of mitochondria were performed
138 *in vitro* (see below). The lizards were placed individually into glass jars (ca. 1,000 ml) within
139 the chamber and allowed to acclimatize for 1h. A baseline air sample (two 140 ml syringes)
140 was collected at the onset of the trial, and the glass vial was then carefully sealed. The trial
141 duration was set to achieve adequate oxygen suppression based on previous studies and
142 preliminary trials (mean=11.8 h, range=11.2-12.1h). A final sample of air was collected with
143 two 140 ml syringes connected to a stopcock. Oxygen and CO₂ concentrations (% of total
144 volume) of the air samples were determined using fuel-cell O₂ and infrared CO₂ sensors
145 (FOXBOX, Sable Systems, Las Vegas, NV, USA) at room temperature with a constant flow (60
146 ml.min⁻¹) after water absorption through a column of Drierite. Oxygen consumption (VO₂, in

147 ml min⁻¹) and CO₂ production (VCO₂) were calculated as: (final % – initial %) × exact chamber
148 volume (ml)/trial duration (min). Body mass was recorded to the nearest mg before each trial.
149 Respiratory quotient (RQ) was calculated like VCO₂/VO₂ and ranged from 0.60 to 0.82 [mean
150 = 0.72, 95% CI= 0.71-0.73]. We found no treatment effect on RQ (ANOVA, F_{1,47}=0.65, P =
151 0.42), but parallel, temporal changes in RQ for both control and treated lizards (F_{3,96}=36.59, P
152 < 0.0001). Since the theoretical RQ value for lipid substrates is 0.71, we therefore assumed that
153 most lizards used stored lipids as the main energy source and used energy equivalents for lipids
154 (19.8 J.mL⁻¹) to convert VO₂ into standard metabolic rates (SMR) values (in J.min⁻¹). These
155 metabolic measurements proved to be extremely reliable and repeatable compared to our earlier
156 studies (Foucart et al., 2014).

157 ***Hormonal manipulation and plasma corticosterone assays***

158 Circulating levels of corticosterone were enhanced using a non-invasive method designed by
159 Meylan et al. (2003). We diluted corticosterone (Sigma-Aldrich, France, C2505-500mg 92%,
160 C₂₁H₃₀O₄) in commercial sesame oil according to 3μg of corticosterone per 1μL of sesame oil.
161 Each evening between 20:00 h and 21:00 local time (when lizards are mostly inactive), 4.5 μL
162 of corticosterone mixture (CORT) or pure sesame oil (CONT, a placebo) were topically applied
163 on the backs of the lizards. To check effects on plasma levels of corticosterone, blood was
164 sampled from the infraorbital sinus of the lizards 7 days before the start of the experiment and
165 at the end of the treatment exposure using 2–3 20 μL microhematocrit tubes. In order to
166 standardize the measurements, all samples were collected between 15:00 h and 16:00 local time
167 within 3 min of removal of an animal from its home cage to avoid the handling-induced increase
168 in plasma corticosterone levels (Dauphin-Villemant, 1987). Plasma was obtained by
169 centrifugation at c.a. 5,000 g for 5 minutes of the blood samples and was stored at -40°C for
170 subsequent measurements of plasma levels of corticosterone. Corticosterone level was
171 measured with a competitive enzyme-immunoassay method using corticosterone EIA (IDS

172 Corticosterone EIA kit, ref AC-14F1, IDS EURL Paris, France) after 1:10 dilution of all
173 samples following previously published guidelines (Mugabo, 2017).

174 ***Respiratory capacities of muscular permeabilized fibres***

175 At the end of the experiment, animals were euthanized by decapitation. Skeletal muscles were
176 rapidly and entirely dissected, weighed and then placed in an ice-cold isolation buffer (BIOPS
177 containing 2.77 mM Ca-EGTA, 7.23 mM, 20 mM imidazole, 20 mM taurine, 50 mM K-MES,
178 0.5 mM DTT, 6.56 mM MgCl₂, 5.77 mM ATP, 15 mM phosphocreatine, pH 7.2). Skeletal
179 muscle fibres were permeabilized in BIOPS solution supplemented with saponin (50 µg ml⁻¹)
180 according to a standard protocol (Pesta and Gnaiger, 2012). Permeabilized fibres were weighed
181 and their respiration were monitored with a high-resolution respirometer (Oxygraph-2k,
182 Oroboros Instruments; Austria) in a hyper-oxygenated respiratory buffer maintained at 25 °C
183 (110 mM sucrose, 0.5 mM EGTA, 3 mM MgCl₂, 60 mM K-lactobionate, 20 mM taurine, 10
184 mM KH₂PO₄, 1 g l⁻¹ fatty acid-free bovine serum albumin, 20 mM Hepes, pH 7.1). A mixture
185 of respiratory substrates (5 mM pyruvate, 2.5 mM malate and 5 mM succinate) was added to
186 obtain the basal respiration (state 2). Cellular ATP synthesis was initiated by the addition of 1
187 mM ADP. The maximal fully uncoupled respiration (state 3_{unc}) was initiated by addition of 2
188 µM carbonyl cyanide p-tri-fluoro-methoxy-phenyl-hydrazone (FCCP), in presence of
189 oligomycin (2 µg ml⁻¹), an inhibitor of ATP synthase.

190 ***Extraction of liver and functional analyses of mitochondria***

191 After weighing the liver, the mitochondria were isolated in an ice-cold isolation buffer (250
192 mM sucrose, 1 mM EGTA, 20 mM Tris-HCl, pH 7.4 at 4°C) as previously described (Voituron
193 et al., 2017). Briefly, the isolation procedure involved Potter-Elvehjem homogenization (three
194 passages) and differential centrifugations (all steps at 4°C), with the liver mitochondria being
195 pelleted at 9,000×g (10 min). The protein concentration of mitochondrial suspension was
196 spectrophotometrically determined at 540 nm by a Biuret method with bovine serum albumin

197 as a standard. Because lizard mitochondrial preparations contained a dark pigment that
198 absorbed at 540 nm, the absorbance of the same volume of mitochondria in a solution
199 containing 0.06% Na deoxycholate, 0.6% K-Na-tartrate and 3% NaOH was subtracted.

200 Rates of oxygen consumption and ATP synthesis were measured in a closed glass cell fitted
201 with a Clark oxygen electrode (Rank Brothers Ltd, UK) at 25°C in a respiratory buffer
202 containing 120 mM KCl, 5 mM KH₂PO₄, 1 mM EGTA, 2 mM MgCl₂, 0.3 % fatty acid-free
203 bovine serum albumin, 1.6 U/ml hexokinase, 20 mM glucose, 3 mM Hepes (pH 7.4). Liver
204 mitochondria (0.5-1.5 mg/mL) were energized with a mixture of substrates (5 mM pyruvate,
205 2.5 mM malate and 5 mM succinate). Mitochondrial ATP synthesis was initiated by the addition
206 of 500 µM ADP. After recording the phosphorylating respiration rate for 3 min, four 100 µL
207 samples were withdrawn from the suspension every 1 min and were quenched in 100 µL ice-
208 cold perchloric acid solution consisting of 10% HClO₄ and 25 mM EDTA. After centrifugation
209 (15,000×g, 5 min) and neutralization of the resulting supernatant with a KOH solution (2 M
210 KOH, 0.3 M MOPS), ATP production was determined from the slope of the linear accumulation
211 of glucose-6-phosphate content of samples (Voituron et al., 2017). To make sure that the rates
212 we measured were specific of the mitochondrial ATP synthase activity and not associated with
213 contaminated ATPase activity such as adenylate kinase, we determined oxygen consumption
214 and ATP synthesis rates in the presence of oligomycin (2 µg/mL). These values were taken into
215 account to calculate the rate of mitochondrial ATP synthesis that is associated with
216 mitochondrial ATP synthase activity and oxygen consumption.

217 **Mitochondrial reactive oxygen species production**

218 The rate of H₂O₂ released by isolated liver mitochondria was measured at 25°C in a respiratory
219 buffer supplemented with 5 U/mL horseradish peroxidase and 1 µM Amplex red fluorescent
220 dye using a fluorescence spectrophotometer (SFM-25, Kontron Instrument) at excitation and
221 emission wavelengths of 560 nm and 584 nm, respectively. The fluorescent signal was

222 calibrated using a standard curve obtained after successive addition of H₂O₂ (up to 35 pmoles).
1
2 223 Amounts of H₂O₂ release were corrected from background rate of product formation in the
3
4 224 absence of exogenous substrate as described previously (Voituron et al., 2017). Free radical
5
6
7 225 electron leak (ROS/O ratio) was calculated as the fraction (%) of basal non-phosphorylating
8
9
10 226 oxygen consumption that is reduced into H₂O₂ at the respiratory chain instead of H₂O at the
11
12 227 cytochrome-c oxidase. Similarly, the oxidative cost of mitochondrial ATP synthesis was
13
14 228 calculated from the ratio of H₂O₂ generation under phosphorylating state divided by the
15
16
17 229 corresponding rate of ATP synthesis (ROS/ATP ratio).
18

19 230 **Statistical analyses**

21
22 231 All statistical analyses were performed with linear models in R 3.3.2 (R Development Core
23
24 232 Team, 2015). We first checked the effects of treatment groups on change in plasma
25
26 233 corticosterone levels from before the end of the experiment with a linear model fitted with the
27
28
29 234 *lm* package (Venables and Ripley, 2002). We included initial plasma corticosterone level and
30
31 235 body mass as covariates in this model. Next, repeated measurements (body mass and SMR)
32
33
34 236 were analyzed with mixed-effects linear models in the *lme* package. Dependence among
35
36 237 repeated measurements were accounted using random intercept model where the intercept of
37
38
39 238 the model is allowed to vary randomly among individuals. A compound symmetry covariance
40
41 239 structure that includes within-subject correlated errors was used and the random term was
42
43
44 240 always kept in the models. This model assumes that individuals react in the same way to
45
46 241 treatments and provides information about intra-class correlations and therefore consistency of
47
48
49 242 inter-individual differences. In addition, all models included fixed effects of experimental
50
51 243 treatments, treatment exposure groups (categorical variable) and their interactions. Since SMR
52
53 244 increases exponentially with body mass, we log transformed SMR prior to the analyses and
54
55
56 245 included log-transformed body mass as a covariate in the analyses. The normality and
57
58 246 homogeneity of variance of residuals was systematically checked in the full models and was
59
60
61
62
63
64
65

247 found to be satisfactory. In a third set of analyses, we compared final measurements of liver
248 mass and mitochondria functioning across treatment groups with a linear model including fixed
249 effects of experimental treatments, treatment exposure groups and their interactions. In all
250 models, the significance of fixed effects was tested with type III F statistics using the *Anova*
251 procedure in the *car* package. We removed non-significant variables ($\alpha=0.05$) one by one using
252 a backward elimination procedure. Whenever significant differences were found among
253 treatment groups, we used Tukey's procedure to conduct post hoc tests (pairwise comparisons
254 between the experimental groups) with the *lsmeans* package (Lenth, 2016). Results are
255 presented as mean \pm standard error unless otherwise stated.

257 **Results**

258 **Corticosterone supplementation increases body and liver masses**

259 Before the experiment, animals had similar SVL, body mass and body condition, SMR and
260 initial plasma corticosterone levels among treatment groups (ANOVA, all $P > 0.66$). Analysis
261 of intra-individual change in plasma corticosterone levels confirmed that treated groups
262 exhibited higher concentration over the entire experiment (treatment: $F_{2,44}=71.6$, $P < 0.0001$;
263 exposure group: $F_{2,44}=6.49$, $P = 0.003$). On average, plasma corticosterone levels were 154.8
264 ng.mL^{-1} (± 18.3) higher in treated than in control lizards, similar to previous effects reported
265 with the same protocol (Voituron et al. 2017). In addition, treated lizards exhibited an increase
266 of body mass over time (+2% at 3D, +5% at 11D and +17% at 21D), whereas the body mass of
267 control lizards did not change significantly over time (treatment \times exposure group: $F_{3,94}=18.13$,
268 $P < 0.0001$) (see Table 1). Total liver mass (treatment \times exposure group: $F_{2,43}=8.24$, $P = 0.0009$)
269 and relative liver mass (g of liver per 100 g body mass; liver mass controlled for variation in
270 body mass; treatment \times exposure group: $F_{2,42}=8.33$, $P = 0.0009$) differed also between

271 treatments after 11 days of exposure only. There was a significant increase of liver mass at D11
272 and D21 in treated lizards relative to controls (Table 1).

273 **Corticosterone supplementation and whole-organism oxygen consumption**

274 Overall, SMR remained relatively similar between treatments apart for a slight, close to
275 significant decrease in SMR from treated lizards relative to controls at D21 (Table 1; treatment
276 \times exposure group: $F_{3,92}=2.69$, $P = 0.05$; Student's t test at D21: $P = 0.07$). However, when
277 controlling for the positive relationship between individual body mass and individual SMR
278 (log-log slope= 0.67 ± 0.13 ; $F_{1,90}=25.78$, $P < 0.0001$), the best fit model indicated that mass-
279 corrected SMR differences over time between treatments ranged from a small positive
280 difference for CORT at D3, followed by a return to normal at D11, and a significant decrease
281 in mass-corrected SMR from treated lizards at D21 (treatment \times exposure group: $F_{3,90}=5.01$,
282 $P=0.003$). These results can be interpreted as evidence that corticosterone treated lizards
283 maintained relatively similar whole-organism SMR despite their increase in body mass during
284 the 21-day long exposure period, thus had lower mass-corrected SMR through time.

285 **Corticosterone supplementation and liver mitochondrial functions**

286 The treatment affected mitochondrial activities with contrasted effects between the rates of ATP
287 synthesis and oxygen consumption over the course of the study (Table 1 and Figure 2, see Table
288 S1 for statistical details). The rate of ATP synthesis was significantly lower in the corticosterone
289 treatment irrespective of treatment exposure. In contrast, both basal and phosphorylating rates
290 of oxygen consumption did not change after 3 days of treatment, then were significantly
291 decreased in 11-day treated lizards compared with control animals, and eventually returned to
292 same levels than in controls at day 21 (Table 1). Consequently, mean values of the mitochondrial
293 coupling efficiency (ATP/O ratio) were significantly decreased after 3 days of corticosterone
294 treatment, but were not significantly different from control values after 11 days and 21 days of
295 treatment (treatment \times exposure group: $F_{2,41}=4.14$, $P = 0.02$, see Figure 2).

296 The rates of mitochondrial reactive oxygen species production under both basal non-
297 phosphorylating and active phosphorylating states were not significantly altered after 3 days of
298 glucocorticoid treatment but exhibited between 40% to 50% decrease in treated lizards relative
299 to controls after 11 and 21 days (Table 1, see Table S1 for statistical details). Overall, the
300 electron leak defined by the ROS/O ratio was not significantly changed by the glucocorticoid
301 treatment neither in basal non-phosphorylating state (treatment \times exposure group: $F_{2,41}=2.09$, P
302 = 0.14; treatment: $F_{1,43}=0.13$, $P = 0.82$; exposure group: $F_{2,44}=10.21$, $P = 0.0002$) nor in active
303 phosphorylating state (treatment \times exposure group: $F_{2,41}=1.81$, $P = 0.18$; treatment: $F_{1,43}=0.89$,
304 $P = 0.35$; exposure group: $F_{2,44}=2.26$, $P = 0.12$). In contrast, the oxidative cost of ATP synthesis
305 (ROS/ATP ratio) was marginally affected by hormonal treatment, being higher in the 3-day
306 treated group than in the control group, but returning to the level of control groups after 11 days
307 and 21 days of treatment (treatment \times exposure group: $F_{2,41}=3.02$, $P = 0.06$; see Figure 3).

308 **Corticosterone supplementation and muscle mitochondria functioning**

309 Whatever the state of activation (basal non-phosphorylating, phosphorylating and maximal),
310 cellular oxygen consumption of lizard muscle was not different between control and treated
311 individuals neither at 3 days, 11 days nor at 21 days of treatment (Table 1 and Table S1).

313 **Discussion**

314 This study aimed at unraveling the time-dependent mechanisms linking corticosterone and
315 metabolism at different levels of organization (whole-organism, cellular and mitochondrial) in
316 an ectothermic organism. The data presented here provide first clear evidence of GC-dependent
317 regulation of the liver mitochondrial functionality in a time-dependent manner leading to a
318 “functional resilience” after 21 days of chronic corticosterone increase. Indeed, short-term GC
319 supplementation induced a significant decrease of mitochondrial ATP synthesis without change
320 of oxygen consumption and H_2O_2 release, thus causing a lower efficiency (ATP/O) and a higher

1 321 oxidative cost of an ATP molecule production (ROS/ATP). In contrast, longer-term GC
2 322 supplementation induced a significant decrease of mitochondrial H₂O₂ release with
3
4 323 mitochondrial oxidative phosphorylation characteristics to returning to the same level than
5
6
7 324 controls. In an important synthesis, Mc Ewen and Wingfiels (2003) defined the allostatic state
8
9 325 as the “*altered and sustained activity levels of the primary mediators, e.g., glucocorticosteroids,*
10
11 326 *that integrate physiology and associated behaviors in response to changing environments and*
12
13
14 327 *challenges such as social interactions, weather, disease, predators, pollution, etc*”. The present
15
16 328 data show that liver mitochondria decrease first their efficiency and then their fluxes (both in
17
18
19 329 terms of O₂ and ATP) but restore their functional capacity in response to the energetic
20
21
22 330 dysfunction imposed by the endocrine variations. In addition, the data also suggest a tissue
23
24 331 dependent response, with corticosterone treatment affecting functioning of mitochondria in the
25
26 332 liver but not that of skeletal muscles. Interestingly, changes in metabolism of mitochondria *in*
27
28
29 333 *vitro* were not reflected by variation in the whole-organism metabolism *in vivo*.

30
31 334 The present data expand results of a previous study on this species (Voituron et al. 2017)
32
33
34 335 providing the time dynamics of functional regulation of the mitochondrial oxygen consumption,
35
36 336 ATP and H₂O₂ productions under GC secretion. In response to the glucocorticoid
37
38
39 337 supplementation, the liver mitochondria first lowered its ATP synthesis rate without changing
40
41 338 its oxygen consumption (see Figure 2). Even if mechanisms underpinning this lower efficiency,
42
43
44 339 the cellular ATP/AMP ratio being modified, a cascade of genes upregulation required for
45
46 340 mitochondrial respiratory chain expression and function is triggered (Teperino et al. 2010;
47
48
49 341 Mouchiroud et al. 2014). At 11 days of treatment, the mitochondrial pattern is characterized by
50
51 342 low values of all fluxes reflecting a diminished activity of the respiratory chain. Interestingly,
52
53 343 this mitochondrial hypometabolism restored both the efficiency (ATP/O) and oxidative cost of
54
55
56 344 an ATP molecule production (ROS/ATP). Altogether, these elements are in line with scenarios
57
58 345 of downregulation of the respiratory chain subunits (Pandya et al., 2004) and of the activity of
59
60
61
62
63
64
65

346 substrate oxidation reported after glucocorticoids treatment in rodents (Roussel et al., 2004;
1
2 347 Arvier et al., 2007). The nearly total recovering of both oxygen and ATP after 21 days of
3
4 348 treatment associated with lower ROS release might be ascribed to *de novo* mitogenesis
5
6
7 349 (Jornayvaz and Shulman 2010). Indeed, glucocorticoids can induce higher expression of key
8
9
10 350 nuclear genes that are required to produce new mitochondria (Psarra and Sekeris 2011). Such a
11
12 351 functional resilience in terms of ATP production associated with lower H₂O₂ production
13
14 352 corroborates previous results on male common lizards that showed lower superoxide dismutase
15
16
17 353 activity in corticosterone-treated individuals after 21 days (Cote et al. 2010). These results are
18
19 354 however not congruent with data on endotherms in which chronically elevated GC levels
20
21
22 355 accelerate aging and reduce lifespan (Schoenle et al. 2018), thus imposing fitness costs
23
24 356 potentially through oxidative balance (Costantini 2011).

26 357 All these mitochondrial adjustments occurred without modification of whole-organism
27
28
29 358 metabolism (present study; Voituron et al. 2017). Even if only two tissues were tested in the
30
31
32 359 study, the data thus strongly suggest that whole-organism oxygen consumption cannot be used
33
34 360 as an accurate proxy for neither ATP production nor ROS release by tissue-specific
35
36 361 mitochondria (Salin et al. 2015). However, the GC effect on whole-organism metabolism, an
37
38
39 362 important parameter of the total energy budget, still remains an open question since it has been
40
41 363 demonstrated that chronic increase of corticosterone slightly increases metabolic rates at rest in
42
43
44 364 pregnant female common lizards (Meylan et al. 2010) when other lizard species reduce their
45
46 365 total oxygen consumption rate when exposed to an increase in corticosterone (Miles et al. 2007;
47
48
49 366 Durant et al. 2008). Nevertheless, the functional mitochondrial resilience observed *in vivo*
50
51 367 might constitute a proximal explanation of the increase in survival of male common lizards with
52
53 368 corticosterone enhancement (Meylan and Clobert 2005; Cote et al. 2006).

55
56 369 Even if oxygen consumption between tissues has been assessed at different levels of
57
58 370 organization, the data suggests that muscles and liver showed differential response patterns.
59
60
61
62
63
64
65

1 371 This statement would be wrong only if mitochondrial content in muscles strongly increase
2 372 during acute and mild-time stress that has never been reported and not congruent with
3
4 373 mitogenesis dynamic (Jornayvaz and Shulman 2010). The data available on mammals are
5
6
7 374 puzzling with no muscular effect in rodents under dexamethasone (Dumas et al. 2003, Roussel
8
9 375 et al. 2004) or muscle wasting mediated, in part, by GR-dependent transactivation of genes that
10
11 376 drive myocyte atrophy (Patel et al. 2014). In addition, the muscular impact of GC is dose-
12
13
14 377 dependent with moderate or transient exposure to GCs enhancing muscle performance (Caruso
15
16 378 et al. 2014). Even if muscles of common lizards do not exhibit a mitochondrial response,
17
18
19 379 muscles could be involved as amino-acids source for liver neoglucogenesis under
20
21
22 380 corticosterone.

23
24 381

25 26 382 **Conclusions and perspectives**

27
28
29 383 This study demonstrates that liver mitochondrial energetics of ectotherms is directly influenced
30
31 384 in a time-dependent manner by exposure to higher plasma concentrations of circulating GCs.
32
33
34 385 After chronic exposure to GCs, the liver mitochondria reached a functional resilience with a
35
36 386 complex interplay between O₂ consumption, ATP production and H₂O₂ release. The coupling
37
38
39 387 between these three mitochondrial processes is thus plastic and differentially regulated leading
40
41 388 to a critical period when stress is acute (low ATP production with high H₂O₂ release), an
42
43
44 389 intermediate period when efficiency returns to the initial levels but with low respiratory chain
45
46 390 activity, and finally a complete “recovery” to the initial fluxes and ratios. This time-dependent
47
48
49 391 relationship between GC and mitochondria in ectotherms needs to be taken into account to
50
51 392 better understand mechanisms that ensure and drive the flow of energy during
52
53 393 physiopathological responses (e.g., Rohleder 2012) but also towards adaptive processes.
54
55
56
57
58
59
60
61
62
63
64
65

394

1
2
3 **Acknowledgements**

4
5 396 We are thankful to field assistants and technical staff for their support, especially Beatriz
6
7
8 397 Decencière, Simon Agostini, Victorien, Virginie and Amandine. We thank Olivier Lourdais for
9
10 398 renting the respirometry equipment.

11
12
13 399

14
15 **Declarations**

16
17
18 **Funding**

19
20 402 This study was funded by the Centre National de la Recherche Scientifique (CNRS), the Agence
21
22 403 Nationale de la Recherche (ANR-13-JSV7-0011-01 to S.M.) and the Région Île-de-France
23
24 404 R2DS program (grant 2013-08 to S.M., J.F.L.G. and R.J.). This work has also benefited from
25
26
27 405 technical and human resources provided by CEREEP-Ecotron IleDeFrance (CNRS/ENS UMS
28
29 406 3194), which is supported by Regional Council of Ile-de-France under the DIM Program R2DS
30
31 407 bearing the reference I-05-098/R and the program " Investissements d'Avenir" launched by the
32
33
34 408 French government and implemented by ANR with the reference ANR-11-INBS-0001 AnaEE
35
36
37 409 France.

38
39
40 **Conflict of interest/Competing interests**

41
42 411 There is no conflict of interest to declare

43
44 **Availability of data and material**

45
46
47 413 The datasets generated during the current study are available from the corresponding author on
48
49 414 reasonable request.

50
51
52 **Author contributions**

53
54 416 Conceptualization: Y.V., S.M., D.R., J.F.L.G., C.R.; Methodology: D.R., Y.V., S.M., C.R.,
55
56 417 J.F.L.G.; Validation: D.R., Y.V.; Formal analysis: D.R., J.F.L.G., A.D., C.R., Y.V.; Investigation:

1
2 418 D.R., J.F.L.G., S.M., A.D., C.R., Y.V.; Writing - original draft: Y.V., D.R.; Writing - review &
3 editing: Y.V., S.M., D.R., J.F.L.G., A.D.

4
5 420 **Ethics approval**

6
7 421 The present investigation was carried out according to the ethical principles of the Préfecture
8
9 422 de Seine-et-Marne under agreement A77-341-1.

10
11
12 423

424 **Bibliography**

- 1
2
3 425 Arvier M, Lagoutte L, Johnson G, Dumas JF, Sion B, Grizard G, Malthiery Y, Simard G, Ritz
4
5 426 P (2007) Adenine nucleotide translocator promotes oxidative phosphorylation and mild
6
7 427 uncoupling in mitochondria after dexamethasone treatment. *Am J Physiol Endocrinol*
8
9
10 428 *Metab* 293: E1320-1324.
- 11
12 429 Casuso RA, Melskens L, Bruhn T, Secher NH, Nordsborg NB (2014) Glucocorticoids
13
14
15 430 improve high-intensity exercise performance in humans. *Eur J Appl Physiol* 114: 419–
16
17 431 424.
- 18
19 432 Costantini D, Marasco V, Møller AP (2011) A meta-analysis of glucocorticoids as modulators
20
21
22 433 of oxidative stress in vertebrates. *J Comp Physiol B* 181:447-56. doi: 10.1007/s00360-
23
24 434 011-0566-2.
- 25
26
27 435 Cote J, Clobert J, Meylan, S, Fitze PS (2006) Experimental enhancement of corticosterone
28
29 436 levels positively affects subsequent male survival. *Horm Behav* 49: 320-327.
30
31
32 437 doi:10.1016/j.yhbeh.2005.08.004.
- 33
34 438 Cote J, Meylan S, Clobert, J, Voituron Y (2010) Carotenoid-based coloration, oxidative stress
35
36
37 439 and corticosterone in common lizards. *J Exp Biol* 213, 2116-2124.
38
39 440 doi:10.1242/jeb.040220.
- 40
41
42 441 de Kloet ER (2013) Functional profile of the binary brain corticosteroid receptor system:
43
44 442 mediating, multitasking, coordinating, integrating. *Eur J Pharmacol* 719:53–62.
- 45
46 443 Dhabhar FS, McEwen BS (1997) Acute stress enhances while chronic stress suppresses cell-
47
48
49 444 mediated immunity in vivo: A potential role for leukocyte trafficking. *Brain Behav*
50
51 445 *Immun* 65:360-368.
- 52
53
54 446 Du J, Wang Y, Hunter R, Wei Y, Blumenthal R, Falke C, Khairova R, Zhou R, Yuan P,
55
56 447 Machado-Vieira R, McEwen BS, Manji HK (2009) Dynamic regulation of
57
58
59 448 mitochondrial function by glucocorticoids. *Proc Natl Acad Sci USA* 106: 3543-3548.

- 449 Dumas JF, Simard G, Roussel D, Douay O, Foussard F, Malthiery Y, Ritz P (2003)
1
2 450 Mitochondrial energy metabolism in a model of undernutrition induced by
3
4 451 dexamethasone. *Brit J Nutri* 90: 969-977.
5
6
7 452 Durant SE, Romero LM, Talent LG, Hopkins, WA (2008) Effect of exogenous corticosterone
8
9 453 on respiration in a reptile. *Gen Comp Endo* 156:126-133.
10
11 454 doi:10.1016/j.ygcen.2007.12.004
12
13
14 455 Eisner V, Picard M, Hajnóczky G (2018) Mitochondrial dynamics in adaptive and
15
16 456 maladaptive cellular stress responses. *Nat Cell Biol* 20:755–765.
17
18 457 doi.org/10.1038/s41556-018-0133-0
19
20
21 458 Foucart T, Lourdais O, DeNardo DF, Heulin B (2014) Influence of reproductive mode on
22
23 459 metabolic costs of reproduction: insight from the bimodal lizard *Zootoca vivipara*. *J Exp*
24
25 460 *Biol* 217:4049-56. 2014. doi: 10.1242/jeb.104315
26
27
28
29 461 Harris RBS (2015) Chronic and acute effects of stress on energy balance: are there
30
31 462 appropriate animal models? *Am J Physiol – Regul Integ Comp Physiol* 308: R250-
32
33 463 R265.
34
35
36 464 Hunter RG, Seligsohn M, Rubin TG, Griffiths BB, Ozdemir Y, Pfaff DW, Datson NA, McEwen
37
38 465 BS (2016) Stress and corticosteroids regulate rat hippocampal mitochondrial DNA gene
39
40 466 expression via the glucocorticoid receptor. *Proc Natl Acad Sci USA*. 113: 9099-9104. doi:
41
42 467 10.1073/pnas.1602185113.
43
44
45
46 468 Jani MS, Telang SD, Katyare SS (1991) Effect of corticosterone treatment on energy
47
48 469 metabolism in rat liver mitochondria. *J Steroid Biochem Mol Biol* 38:587-591.
49
50
51 470 Jessop TS, Woodford R, Symonds MRE (2013) Macrostress: do large-scale ecological
52
53 471 patterns exist in the glucocorticoid stress response of vertebrates? *Funct Ecol* 27:120–
54
55 472 130. 2013doi: 10.1111/j.1365-2435.2012.02057.x
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 473 Joels M, De Kloet ER (1992) Coordinative mineralocorticoid and glucocorticoid receptor-
474 mediated control of responses to serotonin in rat hippocampus. *Neuroendocrinology*
475 55:344–350.
- 476 Jornayvaz FR, Shulman GI (2010) Regulation of mitochondrial biogenesis. *Essays Biochem*
477 47:69-84. doi: 10.1042/bse0470069
- 478 Lee SR, Kim HK, Song IS, Youm J, Dizon LA, Jeong SH, Ko TH, Heo HJ, Ko KS, Rhee BD,
479 Kim N, Han J (2013) Glucocorticoids and their receptors: insights into specific roles in
480 mitochondria. *Prog Biophys Mol Biol* 112:44-54.
- 481 Lehninger AL, Nelson DL, Cox MM (1993) *Principles of Biochemistry*, 2nd edn. New York,
482 USA: Worth Publishers.
- 483 Lenth RV (2016) Least-Squares Means: The R Package lsmeans. *J Stat Soft* 69:1-33. doi:
484 10.18637/jss.v069.i01
- 485 McEwen BS, Wingfield JC (2003) The concept of allostasis in biology and biomedicine.
486 *Horm Behav* 43: 2–15.
- 487 Manoli I, Alesci S, Blackman MR, Su YA, Rennert OM, Chrousos GP (2007) Mitochondria as
488 key components of the stress response. *Trends Endo Metab* 18:190-198.
489 doi:10.1016/j.tem.2007.04.004
- 490 Meylan S, Clobert J (2005) Is corticosterone-mediated phenotype development adaptive?
491 Maternal corticosterone treatment enhances survival in male lizards. *Horm Behav*
492 48:44-52. doi:10.1016/j.yhbeh.2004.11.022.
- 493 Meylan S, Dufty AM, Clobert J (2003) The effect of transdermal corticosterone application on
494 plasma corticosterone levels in pregnant *Lacerta vivipara*. *Comp Biochem Physiol A*
495 13:497-503.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 496 Meylan S, Haussy C, Voituron Y (2010) Physiological actions of corticosterone and its
497 modulation by an immune challenge in reptiles. *Gen Comp Endo* 169:158-166.
498 doi:10.1016/j.ygcen.2010.08.002.
- 499 Miles DB, Calsbeek R, Sinervo B (2007) Corticosterone, locomotor performance, and
500 metabolism in side-blotched lizards (*Uta stansburiana*). *Horm Behav* 51: 548-554.
501 doi:10.1016/j.yhbeh.2007.02.005.
- 502 Mouchiroud L, Eichner LJ, Shaw RJ, Auwerx J (2014) Transcriptional Coregulators: Fine-
503 Tuning Metabolism. *Cell Metab* 20:26-40.
504 doi:<https://doi.org/10.1016/j.cmet.2014.03.027>
- 505 Mugabo M, Le Galliard JF, Perret S, Decencière B, Haussy C, Meylan S (2017) Sex-
506 specific density- dependent secretion of glucocorticoids in lizards: insights from
507 laboratory and field experiments. *Oikos* 126:1051-1061.
508 <https://doi.org/10.1111/oik.03701>
- 509 Pandya JD, Agarwal NA, Katyare SS (2004) Effect of dexamethasone treatment on oxidative
510 energy metabolism in rat liver mitochondria during postnatal developmental periods.
511 *Drug Chem Toxicol* 27:389-403.
- 512 Patel, R., Williams-Dautovich, J. and Cummins, C. L. (2014). New molecular mediators of
513 glucocorticoid receptor activity in metabolic tissues. *Mol Endocrinol* 28, 999–1011.
- 514 Pesta D, Gnaiger E (2012) High-resolution respirometry: OXPHOS protocols for human cells
515 and permeabilized fibers from small biopsies of human muscle. *Methods Mol Biol*
516 810:25-58. doi: 10.1007/978-1-61779-382-0_3.
- 517 Picard M, Juster RP, McEwen BS (2014) Mitochondrial allostatic load puts the 'gluc' back in
518 glucocorticoids. *Nat Rev Endocrinol* 10: 03-310.
- 519 Picard M, McEwen BS, Epel ES, Sandi C (2018) An energetic view of stress: Focus on
520 mitochondria. *Front Neuroendocrinol* 49:72–85.

521 Picard M, McEwen BS (2018) Psychological Stress and Mitochondria: A Systematic Rev
1
2 522 Psychosom Med 80:141-153. doi: 10.1097/PSY.0000000000000545.
3
4 523 Psarra AM, Sekeris CE (2011) Glucocorticoids induce mitochondrial gene transcription in
5
6
7 524 HepG2 cells: role of the mitochondrial glucocorticoid receptor. Bioch Biophys Acta
8
9
10 525 1813:1814-1821. doi:10.1016/j.bbamcr.2011.05.014.
11
12 526 Rohleder N (2012) Acute and chronic stress induced changes in sensitivity of peripheral
13
14 527 inflammatory pathways to the signals of multiple stress systems.
15
16
17 528 Psychoneuroendocrinol 37:307-316.
18
19 529 Rolfe DFS, Brown GC (1997) Cellular energy utilization and molecular origin of standard
20
21
22 530 metabolic rate in mammals. Physiol Rev 77:731-58. doi:10.1152/physrev.1997.77.3.731
23
24 531 Roussel, D., Dumas, J. F., Simard, G., Malthiery, Y. and Ritz, P. (2004). Kinetics and control
25
26
27 532 of oxidative phosphorylation in rat liver mitochondria after dexamethasone treatment.
28
29 533 Biochem. J. 382, 491-499.
30
31 534 Salin K, Luquet E, Rey B, Roussel D, Voituron Y (2012) Alteration of mitochondrial
32
33
34 535 efficiency affects oxidative balance, development and growth in frog (*Rana temporaria*)
35
36 536 tadpoles. J Exp Biol 215:863-869.
37
38
39 537 Salin K, Auer SK, Rey B, Selman C, Metcalfe NB (2015) Variation in the link between
40
41 538 oxygen consumption and ATP production, and its relevance for animal performance.
42
43
44 539 Proc Biol Sci 282:20151028. doi:10.1098/rspb.2015.1028.
45
46 540 Sapolsky RM, Romero LM, Munck AU (2000) How do glucocorticoids influence stress
47
48
49 541 responses? Integrating permissive, suppressive, stimulatory, and preparative actions.
50
51 542 Endo Rev 1:55–89. doi:10.1210/er.21.1.55
52
53 543 Schoenle LA, Zimmer C, Vitousek MN (2018) Understanding context dependence in
54
55
56 544 glucocorticoid-fitness relationships: the role of the nature of the challenge, the intensity
57
58 545 and frequency of stressors, and life history. Int Comp Biol 58:777–789.
59
60
61
62
63
64
65

- 546 Sorrells SF, Caso JR, Munhoz CD, Sapolsky RM (2009) The stressed CNS: when
1
2 547 glucocorticoids aggravate inflammation. *Neuron* 64:33-39. doi:
3
4 548 10.1016/j.neuron.2009.09.032.
5
6
7 549 Starkov AA (2008) The role of mitochondria in reactive oxygen species metabolism and
8
9 550 signaling. *Ann N Y Acad Sci* 1147:37-52. doi: 10.1196/annals.1427.015.
10
11
12 551 Strack AM, Bradbury MJ, Dallman MF (1995) Corticosterone decreases nonshivering
13
14 552 thermogenesis and increases lipid storage in brown adipose tissue. *Am J Physiol*
15
16 553 268:R183-191.
17
18
19 554 Team RDC (2008) R: A language and environment for statistical computing. R Foundation for
20
21 555 Statistical Computing, Vienna, Austria.
22
23
24 556 Teperino R, Schoonjans K, Auwerx J (2010) Histone methyl transferases and demethylases;
25
26 557 can they link metabolism and transcription? *Cell Metab* 12:321-327. doi:
27
28 558 10.1016/j.cmet.2010.09.004.
29
30
31
32 559 Venables WN, Ripley BD (2002) *Modern Applied Statistics with S*, Fourth edition. Springer,
33
34 560 New York. ISBN 0-387-95457-0
35
36
37 561 Vitousek MN, Johnson MA, Downs CJ, Miller ET, Martin LB, Francis CD, Donald JW,
38
39 562 Fuxjager MJ, Goymann W, Hau M, Husak JF, Kircher BK, Knapp R, Schoenle LA,
40
41 563 Williams TD (2019) Macroevolutionary Patterning in Glucocorticoids Suggests
42
43 564 Different Selective Pressures Shape Baseline and Stress-Induced Levels. *Am Nat*
44
45 565 193:866-880. doi: 10.1086/703112.
46
47
48
49 566 Voituron Y, Josserand R, Le Galliard JF, Haussy C, Roussel D, Romestaing C, Meylan S
50
51 567 (2017) Chronic stress, energy transduction and free radical production in a reptile.
52
53 568 *Oecologia* 185:195-203.
54
55
56
57 569 Wang XL, Herzog B, Waltner-Law M, Hall RK, Shiota M, Granner, DK (2004) The
58
59 570 synergistic effect of dexamethasone and all-trans-retinoic acid on hepatic
60
61
62
63
64
65

571 phosphoenolpyruvate carboxykinase gene expression involves the coactivator p300. J

1

2 572

3

Biol Chem 279:34191–34200.

4

5 573

6

7 574

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

575 **Figure 1:** Experimental design

576

577 **Figure 2:** Effect of 3 days, 11 days and 21 days of exogenous treatment with corticosterone

578 (CORT: 3µg of corticosterone per µL of sesame oil) on liver metabolism mitochondria from

579 *Zootoca vivipara*. (A) Mitochondrial oxygen consumption and ATP production; the values are

580 represented as % of control values (\pm SEM represented by the horizontal gray-shaded

581 rectangle). (B) Mitochondrial efficiency (ATP/O) in function of treatment time. Circles

582 represent individuals' data and squares represent means for each group. * and † Significantly

583 different from the corresponding control.

584

585 **Figure 3:** Effect of exogenous treatment of corticosterone (CORT: 3µg of corticosterone per

586 µL of sesame oil; control: only sesame oil (white circles and square)) during 3 days (black

587 circles and square), 11 days (black circles and square) and 21 days (black circles and square)

588 in the oxidative cost of an ATP production. Circles represent individuals' data and squares

589 represent means for each group. Values are means \pm s.e.m. for 8-9 animals.

590

591 **Table 1.** Effects of exogenous treatment of corticosterone (CORT: 3µg of corticosterone per µL

592 of sesame oil; CONT: only sesame oil) during 3 days, 11 days and 21 days in *Zootoca vivipara*.

593 The metabolic traits studied include whole-organism standard metabolic rates (SMR), muscle

594 fibers and mitochondrial liver oxygen consumptions, liver mitochondrial ATP and ROS

595 production. Data are reported as mean \pm SE ($n = 8-9$ per group). Significant difference among

596 groups from Tukey post-hoc tests of the best model (see main text) are specified with different

597 letters.

598

Figure 1




Figure 2





Figure 3



1 **Table 1.** Effects of exogenous treatment of corticosterone (CORT: 4.5 μ g of corticosterone per μ L of sesame oil; CONT: only sesame oil) during 3 days, 11 days
 2 and 21 days in *Zootoca vivipara*. The metabolic traits studied include whole-organism standard metabolic rates (SMR), muscle fibers and mitochondrial liver
 3 oxygen consumptions, liver mitochondrial ATP and ROS production. Data are reported as mean \pm SE ($n = 8-9$ per group). Significant difference among groups
 4 from Tukey post-hoc tests of the best model (see main text) are specified with different letters.

5

Morphological and Metabolic traits	Treatment	Day 0	Day 3	Day 11	Day 21
Body and liver mass					
Body mass (g)	CONT	2.00 \pm 0.03 ^a	2.01 \pm 0.04 ^a	1.95 \pm 0.04 ^a	2.03 \pm 0.04 ^a
	CORT	2.00 \pm 0.04 ^a	2.02 \pm 0.04 ^a	2.06 \pm 0.04 ^a	2.26 \pm 0.06 ^b
Liver mass (mg)	CONT		66.13 \pm 3.04 ^a	56.12 \pm 2.18 ^b	57.12 \pm 1.68 ^b
	CORT		60.25 \pm 2.40 ^a	69.12 \pm 4.34 ^b	71.78 \pm 2.44 ^b
Relative liver mass (% of BW)	CONT		3.07 \pm 0.15 ^a	2.64 \pm 0.10 ^b	2.71 \pm 0.09 ^b
	CORT		2.88 \pm 0.12 ^a	3.30 \pm 0.19 ^b	3.46 \pm 0.09 ^b
SMR (J.min⁻¹)					
Whole-organism SMR	CONT	6.70 \pm 0.19 ^a	7.00 \pm 0.16 ^a	7.30 \pm 0.22 ^b	7.26 \pm 0.32 ^{ab}
	CORT	6.69 \pm 0.17 ^a	7.29 \pm 0.18 ^{ab}	7.58 \pm 0.22 ^b	6.96 \pm 0.20 ^a
Muscle fibers oxygen consumption (pmol O₂.s⁻¹.mg⁻¹ fresh mass)					
Maximal respiration	CONT		19.92 \pm 1.59 ^a	19.01 \pm 0.70 ^a	20.53 \pm 0.85 ^a
	CORT		19.26 \pm 1.75 ^a	20.97 \pm 0.93 ^a	20.54 \pm 1.54 ^a

Phosphorylating respiration	CONT		15.24 ± 1.07 ^a	14.18 ± 0.71 ^a	15.55 ± 0.57 ^a
	CORT		15.18 ± 1.37 ^a	15.0 ± 0.85 ^a	16.29 ± 1.41 ^a
Basal respiration	CONT		4.80 ± 0.25 ^a	4.96 ± 0.20 ^a	5.23 ± 0.09 ^b
	CORT		5.05 ± 0.34 ^a	4.99 ± 0.32 ^a	6.11 ± 0.34 ^b
Liver mitochondrial oxygen consumption and ATP fluxes (nmol ATP/min.mg protein or nmol O.min⁻¹.mg protein⁻¹)					
ATP synthesis	CONT		37.21 ± 4.39 ^a	42.06 ± 6.08 ^a	44.43 ± 7.52 ^a
	CORT		27.0 ± 2.72 ^b	21.87 ± 1.94 ^b	35.09 ± 4.43 ^b
Phosphorylating respiration	CONT		21.35 ± 2.35 ^{ab}	24.94 ± 3.39 ^a	22.34 ± 2.90 ^{ab}
	CORT		21.10 ± 2.44 ^{ab}	12.40 ± 1.18 ^b	20.32 ± 2.40 ^{ab}
Basal respiration	CONT		3.85 ± 0.44 ^a	3.93 ± 0.39 ^a	2.33 ± 0.41 ^b
	CORT		3.92 ± 0.45 ^a	2.12 ± 0.19 ^b	1.99 ± 0.22 ^b
Liver mitochondrial ROS production (pmol H₂O₂/min.mg protein)					
Basal state	CONT		167.69 ± 22.4 ^a	129.86 ± 13.05 ^a	164.86 ± 27.92 ^{ab}
	CORT		181.80 ± 30.6 ^a	63.92 ± 8.85 ^b	97.57 ± 7.35 ^b
Phosphorylating state	CONT		50.84 ± 6.83 ^{ab}	42.87 ± 4.58 ^{ab}	57.24 ± 10.72 ^a
	CORT		60.26 ± 11.5 ^a	20.64 ± 3.13 ^b	29.97 ± 4.14 ^b