



Rare earth elements as new biogeochemical proxies in deep-sea mussels

Jean-Alix Barrat, Germain Bayon, Robert Carney, Laurent Chauvaud

► To cite this version:

Jean-Alix Barrat, Germain Bayon, Robert Carney, Laurent Chauvaud. Rare earth elements as new biogeochemical proxies in deep-sea mussels. *Chemical Geology*, 2022, 610, pp.121102. 10.1016/j.chemgeo.2022.121102 . hal-03840641

HAL Id: hal-03840641

<https://cnrs.hal.science/hal-03840641>

Submitted on 16 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rare Earth Elements as new biogeochemical proxies in deep-sea mussels

By

Jean-Alix Barrat^{1,2}, Germain Bayon³, Robert S. Carney⁴,
Laurent Chauvaud¹

¹Univ Brest, CNRS, LEMAR, Institut Universitaire Européen de la Mer (IUEM), Place
Nicolas Copernic, 29280 Plouzané, France. E-Mail: barrat@univ-brest.fr. (corresponding
author).

²Institut Universitaire de France.

³Univ Brest, CNRS, Ifremer, Geo-Ocean, F-29280 Plouzané, France

⁴Department of Oceanography and Coastal Sciences, Louisiana State University, Baton
Rouge, LA 70803, USA

Submitted to Chemical Geology, 12/04/2022

Revised version, 25/08/2022

Abstract

We report on the abundances of REE in a comprehensive suite of shells of deep-sea chemosynthetic mussels from hydrothermal vents and cold seeps. Except for mussel shells from oceanic hydrothermal sites that often show extremely pronounced Eu anomalies ($(\text{Eu}/\text{Sm})_{\text{sn}}=2-200$), and abundances for this element that can occasionally exceed 200 ng/g, REE concentrations are usually low and typically between 10^{-4} and 10^{-3} times the shale reference.

In addition to exhibiting commonly high Eu anomalies, mussel shells from hydrothermal vents are depleted in light-REE and heavy-REE compared to reference shales (e.g., $(\text{Pr}/\text{Sm})_{\text{sn}} < 1$, $(\text{Tb}/\text{Yb})_{\text{sn}} = 1-6$). These features are inherited from hydrothermal fluids. Mussel shells from cold seeps have very different REE concentrations, which also reflect the compositions of the waters they filter: their $(\text{Eu}/\text{Sm})_{\text{sn}}$ ratios are much lower (< 2) and are much less heavy-REE fractionated ($(\text{Tb}/\text{Yb})_{\text{sn}} = 0.5-2.7$). Furthermore, the REE distributions show a clear dichotomy between thiotrophic and methanotrophic mussels. The latter show marked enrichments in light-REE and even sometimes La enrichments much larger than those classically observed in deep-sea waters, leading to huge positive La anomalies ($\text{La}/\text{La}^* > 10$). These light-REE enrichments are likely related to REE-dependent methanol dehydrogenase enzymes used by the symbionts that these mussels host in their gills. These data show that REE chemistry is a promising tool to study chemosynthetic faunas living near hydrothermal vents or cold seeps. Furthermore, REE chemistry, coupled with stable isotopes, should reveal the footprint of aerobic methanotrophy in carbonates formed in cold seeps, but also potentially in ancient sediments.

Key words:

rare earth elements, bivalve, shell chemistry, hydrothermal system, cold seep, methanotrophy

55 **1/ Introduction**

56 Over the last fifty years, rare earth elements (REEs) have become one of the most
57 studied groups of elements by geochemists, finding applications in all fields, from the
58 petrogenesis of rocks, terrestrial or not, to the origin of the first condensed solids in the solar
59 system, through rock dating and isotope tracing (e.g., Henderson, 1984). In marine
60 geochemistry, REEs and their isotopes have been also extensively studied since they allow,
61 among other things, the characterization of the different water masses composing the oceans
62 and their circulation (e.g., Elderfield, 1988). For decades, it has been assumed that these
63 elements do not participate in any biochemical cycle and therefore that their distributions at
64 the Earth's surface are not impacted by life. Nature is certainly more complex. Notable
65 biological effects of the REEs have been observed in plants, and REEs are even used as
66 fertilizers (Tommasi et al., 2021). Furthermore, some species can accumulate large amounts
67 of these elements, as exemplified by the fern *Dicranopteris linearis* (Ozaki et al., 1997). More
68 recently, it was established that light REEs are vital for aerobic methanotrophic and
69 methylotrophic bacteria that play an essential role in global carbon cycling (e.g., Pol et al.,
70 2014; Semrau et al., 2018; Cotruvo, 2019). Methanotrophic bacteria first convert methane to
71 methanol, which is subsequently degraded into formaldehyde using Ca-dependent (MxaF
72 type) or REE-dependent (XoxF type) methanol dehydrogenase enzymes (Skovran *et al.*,
73 2011). Ramachandran and Walsh (2015) and Taubert et al. (2015) have shown that XoxF type
74 enzymes are more frequently used by marine bacteria. These recent findings clearly indicate
75 the ability of micro-organisms to fractionate and redistribute REE, emphasizing the
76 importance of life in the REE cycling in Earth's surface environments. For instance, there is
77 growing evidence that bacterial activity and chemosynthetic ecosystems can influence the
78 REEs distribution in seawater, at least locally. During the blowout of the Deepwater Horizon
79 well (Gulf of Mexico), the depletion of light REE in a submerged plume of methane-rich
80 water was explained by an uptake of these elements by methanotrophic bacteria (Shiller et al.,
81 2017). More recently, light REE depletions were observed in seawater from the Sargasso Sea,
82 at depths between approximately 200 and 500 m, ascribed to the activity of methanotrophic or
83 methylotrophic microorganisms (Meyer et al., 2021). Therefore, methanotrophic bacteria, at
84 least those using enzymes containing REEs, display distinctive REE distributions, as do
85 animals living in symbiosis with them.

Within this scope, deep-sea fauna associated with black smokers and cold seeps are ideal targets to evaluate the distinctive chemical signatures developed by organisms. In these environments, reduced sulfur compounds, methane and/or hydrogen fuel primary productivity sustained by the symbiotic bacteria that form here the basis of the trophic web (Cavanaugh et al., 1981, 1987; Felbeck, 1981; Childress et al., 1986; Petersen et al., 2011), which are vital for many thiotrophic or methanotrophic animals. Recently, two investigations of chemosynthetic fauna have provided direct support for an active consumption of light-REE associated with aerobic microbial oxidation of methane at cold seeps. Bayon et al. (2020a) determined trace element abundances along a one-meter-long chitin tube of a siboglinid worm (*Escarpia southwardae*) collected at the Regab pockmark (Congo margin), identifying distinctive light REE enrichments that were attributed to methanotrophic activity in the branchial plume region of the worm. Additionally, similar light-REE enrichments were found in the gills of methanotrophic mussels collected at seeps from the South China Sea, but not in thiotrophic clams or mussels from the same sites (Wang et al., 2020). Because gills are the organs hosting the methanotrophic symbionts in mussels, Wang et al. (2020) proposed that these light-REE enrichments were related to aerobic methanotrophy. In that latter study, other soft tissues and the shell of studied methanotrophic mussels, while displaying lower light-REE abundances, also show higher positive La anomalies relative to seawater or pore water.

Rare earth element data for deep-sea shells are still sparse. To date, mussels from only three hydrothermal vent fields (Bau et al., 2010) and two cold seeps (Wang et al., 2020) have been investigated. Here we present a comprehensive review for the distribution of REE in mussel shells from deep-sea chemosynthetic environments, which combine previously published literature data (Bau et al., 2010; Wang et al., 2020) and new results obtained on a suite of deep-sea mussels from both submarine methane seeps and hydrothermal vents (Fig. 1). The aim of this study was to further investigate the factors influencing the REE distribution in marine mollusks and assess the degree to which REE abundances in shells can be used to monitor the activity of methanotrophic symbiotic bacteria.

2/ Sampling

Forty-five mussel shells belonging to ten different species were analyzed here, of which nine reside in the genus *Bathymodiulus*. Application of DNA analyses to newly discovered species has led to the assignment of the intensively studied *Bathymodiulus*

childressi to the genus *Gigantidas* (Xu et al., 2019). Twenty-five shell samples, belonging to nine species of *Bathymodiolus*, were sourced from the Scripps Institution of Oceanography Benthic Invertebrate Collection (SIO-BIC) in La Jolla (California, USA). Our sampling includes both thiotrophic and methanotrophic species. Descriptive data for each sample (site, depth, species, museum catalog number, etc.) are given in Supplementary Table 1.

2.1/ Hydrothermal vents

We analyzed three shells from three different sites in the Pacific Ocean for this study, which complement the dataset obtained by Bau et al. (2010) on thiotrophic mussels from the Mid-Atlantic Ridge. Two mussels (*Bathymodiolus thermophilus*) originate from the East Pacific Rise (SIO-BIC M16462) and the Rose Garden vent field (SIO-BIC M8223) located east of the Galápagos Islands, which was among the first submarine hydrothermal sites discovered (e.g., Corliss and Ballard, 1977). The third shell (*Bathymodiolus septemdierum*; SIO-BIC M15353) comes from the Nifonea vent field (Schmidt et al., 2017), located within the caldera of a large shield-type volcano of the Vate Trough, a young extensional rift in the New Hebrides back-arc (Vanuatu, SW Pacific).

2.2/ Cold seeps

We analyzed samples from seven seeps located in three different regions: the Gulf of Mexico, the western margin of Costa Rica, and Melanesia.

The Gulf of Mexico seeps (Brine Pool, Bush Hill, and West Florida Escarpment) have been extensively studied both geologically and biologically (Paull et al., 1984, 1985; Martens et al. 1991; Chanton et al., 1993), and were the basis for seminal observations on chemoautotrophic life, including the demonstration of methane-based symbiosis between animals and intracellular bacteria (Childress et al., 1986). The mussels of these sites are particularly well characterized, and the reader is referred to the numerous papers concerning in particular their symbionts or their isotopic compositions allowing analysis of their diet (e.g., Duperron et al., 2007; Macavoy et al., 2008; Riekenberg et al., 2018; Feng et al., 2018). Twenty *Gigantidas childressi* shells from Brine Pool and Bush Hill, and one *Bathymodiolus heckerae* shell from West Florida Escarpment (SIO-BIC M8018), were analyzed. *Gigantidas childressi* is associated with only one type of methanotrophic symbionts, and consumes primarily methane. The metabolism of *Bathylomodius heckerae*, predominantly

methanotrophic, is however potentially much more complex as its gills harbor a greater diversity of symbionts, including one methanotrophic, one methylotrophic, and two thiotrophic polytypes (Duperron et al., 2007). However, $\delta^{13}\text{C}$ values of these mussels confirm that seep hydrocarbons represent the dominant carbon source in their soft tissues (Cavanaugh et al., 1987; Paull et al., 1989; Duperron et al., 2007).

Cold seeps at the continental margin offshore Costa Rica harbor a varied fauna (Sahling et al., 2008; Levin et al., 2012, 2015), including mytilid species whose distribution shows some stratification with water depth (McCowin et al., 2020): *Bathymodiolus billschneideri* (~1400–1900 m), *B. earlougheri* (~1000–1900 m), and *B. nancyschneiderae* (1000–1100 m). Furthermore, *Bathymodiolus thermophilus*, a thiotrophic mussel typically encountered at hydrothermal vents along the Galápagos Rift Zone and the East Pacific Rise (e.g., Kenk and Wilson, 1985), has been also identified in this area (McCowin et al., 2020). These four species are all thiotrophic (Levin et al., 2012; Brzechffa and Goffredi, 2021). A total of twenty shells from three seeps (Jaco Scar, Mound 12, and Parrita Seep) were analyzed (SIO-BIC specimens listed in Table S-1).

Two mussel shells (*Bathymodiolus edisonensis*, SIO-BIC M18361; and *B. anteumbonatus*, SIO-BIC M15360) from a cold seep in the vicinity of Edison Seamount (New Ireland Basin, Melanesia, Herzig et al., 1998) complement our sampling of methane seep mollusks. Mussels from this location have been described by von Cosei and Janssen (2008), but to the best of our knowledge, the nature of their symbionts and their isotopic compositions are unknown. The abundance of methane at these sites (Herzig et al., 1998), as well as the similarities of these shells to *Gigantidas childressi* from Gulf of Mexico, strongly suggest that they are methanotrophic (von Cosei and Janssen, 2008).

3/ Analytical procedures

Periostracum was removed from the shells by scraping using a clean steel blade. The determination of the REE abundances follows the procedure designed by Barrat et al. (2020). About 150–200 mg of shell were precisely weighted, spiked with Tm, and dissolved with HNO_3 . REE and Y were separated using ion-exchange chromatographic columns loaded with about 1 ml DGA resin. The measurements were performed on a Thermo Scientific ELEMENT XRTM spectrometer located at the “Pôle Spectrométrie Ocean”, Institut

Universitaire Européen de la Mer (IUEM), Plouzané (France). The reader is referred to previous papers where our analytical methodology for carbonate samples (Barrat et al., 2020), calibration, isobaric interference corrections and calculation of concentrations based on Tm addition are extensively described, and in which reference values for many international rock standards are reported (e.g., Barrat et al., 1996, 2012, 2016; Charles et al., 2021).

Measured REE concentrations were normalized to the Post Archean Australian Shale (PAAS; Pourmand et al., 2012), adjusted to standard results obtained in our laboratory (Barrat et al., 2020). The La, Ce, and Gd anomalies were calculated using the La/La^* , Ce/Ce^* , Gd/Gd^* ratios, where X^* is the extrapolated concentration for a smooth PAAS-normalized REE pattern and X_{sn} is the concentration of element X normalized to PAAS: $La_{sn}^* = Pr_{sn}^3/Nd_{sn}^2$, $Ce_{sn}^* = Pr_{sn}^2/Nd_{sn}$, $Gd_{sn}^* = Tb_{sn}^2/Dy_{sn}$. Replicate measurements obtained for two carbonate standards (CAL-S and BEAN) during the analytical sessions are given in Supplementary Table 1, together with reference values from the literature. The precision (RSD) on measured abundances and elemental ratios was generally better than 3 %. Note that there is no analytical bias between the REE concentrations presented here and those obtained by Wang et al. (2020), which were obtained using the same procedure in our laboratory; or those from Bau et al. (2010) obtained at Jacobs University (Bremen), which are also of very high quality. However, Y concentrations obtained in Bremen on different standards (e.g., Bau et al., 2010, 2014, 2018) were systematically lower than those obtained in Brest (e.g., Barrat et al., 2012, 2020, Charles et al., 2021). This calibration bias was corrected here by multiplying all Y concentrations obtained by Bau et al. (2010) by 1.10.

4/ Results

Sixty-six shell analyses were obtained here (supplementary Table 1), greatly increasing the number of available data for deep-sea mussels. These shells are very poor in REEs. Only rarely did concentrations of an element exceed 100 or 200 ng/g (e.g. Eu for some samples from hydrothermal vents). PAAS-normalized REY patterns from the available data set (Bau et al., 2010; Wang et al., 2020 and this work) are given in Figures 2-5 and provide an overview of the diversity of compositions of these shells.

4.1/ Hydrothermal vent mussels

As shown by Bau et al. (2010), the shells of hydrothermal vent mussels have REY patterns showing large positive Eu anomalies, inherited from hydrothermal fluids (Fig. 2). In this regard, the shells analyzed here are no exception. The East Pacific shells show identical patterns (Fig. 2b), except for Eu, with the EPR sample having an Eu-anomaly similar to those from Logatchev (Fig. 2a) and the Rose Garden sample similar to those from Golden Valley (Fig. 2c) obtained by Bau et al. (2010). Both samples are also marked by positive Y and negative Ce anomalies that are unquestionably the fingerprint of seawater. Two samples of the same shell from Nifonea were analyzed and the results are almost identical, suggesting a homogeneity of the latter (Fig. 2e). The patterns obtained show positive Eu anomalies lower than for the two other samples from Pacific, as well as negative anomalies in Ce and positive in Y quite pronounced. These patterns are similar to those of Lilliput (Fig. 2b). The less pronounced Eu anomalies here could be explained by the composition of hydrothermal fluids, whose Eu anomalies are much lower than those of typical oceanic hydrothermal fluids (Schmidt et al., 2017), but also by a more important contribution of ambient seawater than for the other sites.

4.2/ Cold seep mussels

4.2.1/ Thiotrophic mussels

Only three thiotrophic mussel shells from the China Sea (Site F, Fig. 1) had been analyzed before this study (Wang et al., 2020). Here we studied a much larger series of such mussels collected from three sites located in western Costa Rica. Four species of mussels were analyzed. For these three sites, whatever the species considered, the REY-patterns are quite similar (Fig. 3), and comparable in abundance and shape to those of *Bathymodiolus aduloides* collected in the China Sea (Fig. 4 b). Unlike shells from hydrothermal sites, these shells display low $(\text{Eu}/\text{Sm})_{\text{sn}}$ ratios ($=1.21\text{-}1.92$). The shape of their REY patterns resembles those of marine carbonates (e.g., Northdurft et al., 2004). The patterns are characterized by marked negative Ce anomalies ($\text{Ce}/\text{Ce}^* = 0.34\text{-}0.77$ for Costa Rica samples, and $0.77\text{-}0.81$ for China Sea samples) and positive Y anomalies ($\text{Y}/\text{Ho} = 51\text{-}80$ for Costa Rica samples, and $50\text{-}64$ for China Sea samples). They also display positive anomalies in La ($\text{La}/\text{La}^* = 1.9\text{-}3.0$) in the range of seawater values, and apparent positive anomalies in Gd ($\text{Gd}/\text{Gd}^* = 1.1\text{-}1.37$).

4.2.1/ Methanotrophic mussels

The two *Bathymodiolus* shells from the Edison Seamount display lower REE than the *Gigantidas* from the China Sea (Wang et al., 2020), but their patterns show quite similar characteristics (Fig. 4). Two fragments per shell were analyzed, and the results are very consistent, suggesting that both shells do not show marked heterogeneities for REE and Y. As expected, they show positive anomalies in Y ($Y/Ho=59.8-69.5$) and negative in Ce ($Ce/Ce^*=0.39-0.46$), but also striking La anomalies ($La/La^*=7.9-11.9$). Wang et al. (2020) previously showed that the shells of methanotrophic mussels from the China Sea display exceptionally high positive anomalies in La ($La/La^*=2.5-5.1$). Those measured in Edison Seamount shells are even higher, and to the best of our knowledge, represent the most pronounced La anomalies ever measured to date in natural environments not affected by emerging micropollutants. In addition, these shells also show apparent anomalies in Gd ($Gd/Gd^*=1.26-1.77$).

Different REY patterns were obtained for mussels from the Gulf of Mexico (Fig. 5):

-The patterns of *Gigantidas* shells from Brine Pool and Bush Hill show distinctive light REE enrichments, moderate negative Ce anomalies, variable Gd anomalies ($Gd/Gd^*=0.88-1.53$), as well as usual seawater-like positive Y anomalies ($Y/Ho=43.8-72.5$). Their La anomalies ($La/La^*=1.73-2.70$) are not as high as those discussed above, being comparable to typical marine values. Mussels from these two sites display similar REE concentrations, showing that the brines at Brine Pool do not affect the REE composition of shells. Note that the mussels at this site do not live in direct contact with the brines but within at ~10-cm-thick layer of seawater immediately overlying the brine pool or the surficial sediment at the pool edge. However, a sample from Brine Pool (BP7a) stands out from other shells by displaying higher heavy REE contents. We analyzed a second fragment from the same shell (BP7b), and obtained results that are closer to the rest of studied samples, suggesting that the first sub-sample may have been contaminated by impurities. Alternatively, the range of shell composition could be larger than suggested by the other samples.

-Two fragments of the same *Bathymodiolus heckeræ* shell from the West Florida Escarpment were analyzed, with identical results (Fig. 5c). The corresponding patterns are different from those of other methanotrophic mussels. They do not show notable enrichments in light REE like those obtained from the Brine Pool or Bush Hill *Gigantidas*, nor large

positive La anomalies like the methanotrophic mussels from China Sea or Melanesia. Their La/La^* ($=2.35\text{--}2.40$) are in the range of seawater values. They are similar to those of thiotrophic mussels (e.g., those from Costa Rican seeps, Fig. 3), with similar negative Ce and positive Y anomalies ($\text{Ce/Ce}^*=0.39\text{--}0.41$) and positive Y ($\text{Y/Ho}=55.2\text{--}56.3$).

5/ Discussion

5.1/ REE characteristics of hydrothermal *versus* cold seep mussels

Notable differences are apparent between the REE concentrations of mussel shells from hydrothermal vents and those from cold seeps. Bau et al. (2010) showed that mussel shells living in the vicinity of Atlantic hydrothermal vents exhibit characteristic REY patterns, with large positive Eu anomalies [$(\text{Eu/Sm})_{\text{sn}}$ up to 5, and >2 in most cases], which are inherited from hydrothermal fluids. The analyses obtained here confirm this observation, showing that all shells from submarine hydrothermal vents display positive Eu anomalies. In contrast, significant positive Eu anomalies are totally absent in shells from cold seeps (Fig. 6), which exhibit $(\text{Eu/Sm})_{\text{sn}}$ ratios with little variability (between 1.1 to 1.9). Furthermore, ‘hydrothermal’ shells show much more variable and generally much higher REE concentrations than those from cold seeps, as illustrated by the histograms of Sm concentrations (Fig. 7). Their patterns also sometimes show enrichments in middle REE compared to heavy REE, which can be quantified with $(\text{Tb/Yb})_{\text{sn}}$ ratios frequently exceeding 2 (Fig. 8). These mid-REE enrichments, combined with presence of significant positive Eu anomalies, reflect the contribution of hydrothermal fluids with REY distributions very different from seawater abundances (Bau and Dulski, 1999; Douville et al., 1999, Schmidt et al., 2007, 2017; Craddock et al. (2010) ; Cole et al., 2014), with no evidence for any presumed biological effect.

5.2/ Differences between thiotrophic and methanotrophic mussels at cold seeps

At cold seeps, the REE distributions in mussel shells appear, at least in part, to reflect some characteristics of the surrounding water that they filter (Fig. 8, Fig. 9). Thiotrophic mussels have REE ratios falling in the range of seawater and pore water values. Some samples, however, have slightly higher $(\text{Tb/Sm})_{\text{sn}}$ ratios than pore waters, but we do not consider this discrepancy to be very significant since the fluid composition at the sites where our shells originate is not known. As for the mussels from hydrothermal sites, there is no evidence for measurable imprint of any biochemical process involving REE for these animals.

The situation is quite different for methanotrophic mussels. While their heavy-REE ratios are similar to those for seawater or pore waters (Fig. 8), light-REE abundances depart substantially from both seawater and pore water values (Fig. 9). They are, in most cases, much more enriched in light-REE than waters, but also more than oil that can be present in some seeps like those of the Gulf of Mexico (Smrzka et al., 2019). As mentioned above, Wang et al (2020) showed that the shells of methanotrophic mussels at methane seeps from the South China Sea display high La anomalies relative to nearby thiotrophic mussel species, clams and seawater. The decoupling of La from other trivalent REE at cold seeps has been interpreted as reflecting the enzymatic activity of associated methanotrophic symbionts, hence indicating aerobic methanotrophy. The new data obtained in this study provide further support to this hypothesis. First, the thiotrophic mussel shells are not particularly enriched in light-REE, and do not show positive La anomalies exceeding usual seawater values, confirming that the enzymatic activity of their symbionts has no notable effect on these elements. On the other hand, the REY patterns obtained for methanotrophic mussel shells are much more variable than those associated with thiotrophic species. Light REE provide a clear-cut separation between most thiotrophic and methanotrophic mussels (Figs. 8 and 9). Except for one sample that will be discussed later, methanotrophic mussels show light-REE enrichments (illustrated by $(\text{Pr}/\text{Sm})_{\text{sn}}$ or $(\text{Pr}/\text{Nd})_{\text{sn}}$ ratios, Figs. 8 and 9) commonly accompanied by even more pronounced La enrichments (illustrated by the ratios $(\text{La}/\text{Nd})_{\text{sn}}$ and La/La^* , Fig. 9). The range of measured light-REE enrichments or La anomalies cannot be explained by local or environmental factors because it far exceeds the range of values known for sediments and marine waters (e.g., Fig. 8 and 9). In agreement with Wang et al. (2020), these enrichments are most likely related to the enzymatic activity of methanotrophic symbionts hosted by the gills of these mussels. It is logical to link them to the degradation of methane to methanol and then to formaldehyde, since REE-dependent (XoxF type) methanol dehydrogenase enzymes are used by these organisms (e.g., Skovran et al., 2011; Semrau et al., 2018; Cotruvo et al., 2019). The large variability of REY patterns obtained for methanotrophic shells suggests that the use of light-REE by these organisms differs from one species to another. For example, some animals generate very large La anomalies in their shells (e.g., *Gigantidas platifrons* from China Sea or *Bathymodiolus* from Edison Seamount), whereas others display only light REE enrichments, with La anomalies similar to those in seawater (e.g., *Gigantidas childressi* from Gulf of Mexico). These differences, which are extremely marked, necessarily reflect different needs in light-REE for the enzymatic activity of these animals. It should be noted, however, that the transfer of these elements from the gills (which are the seat of symbiont

activity) to the mantle (which synthesizes the shell) has never been considered until now, and that REE fractionation during this transfer cannot be ignored.

One sample, a *Bathymodiolus heckeræ* shell from the West Florida Escarpment (Gulf of Mexico), differs from all other methanotrophic mussel shells analyzed to date (Wang et al., 2020 and this study) in that it lacks significant light REE enrichment, and has an anomaly in La within the range of conventional marine values. This sample even shows REE distributions very similar to those of thiotrophic mussel shells (Fig. 5c). Note that these particular REE signatures associated with *Bathymodiolus heckeræ* cannot be related to analytical artefacts because two separate shell fragments were analyzed, both returning similar results (Supplementary Table 1). Clearly, the symbionts of this mussel did not require light-REE to degrade methane. It is logical to assume that they do not use REE-dependent (XoxF type) methanol dehydrogenase enzymes, but rather Ca-dependent (MxaF type) methanol dehydrogenase enzymes. However, only one shell of this species was available for our study. It would be appropriate at this stage to analyze other samples from this site, and obviously to identify the enzymes used by the mussel symbionts of this species.

5.3/ Gd anomalies in mussels from cold seeps

An unexpected result of our study concerns the presence of marked positive Gd anomalies in many cold seep shells (Fig. 10), whereas those from hydrothermal vents show a much narrower range of Gd/Gd* values.

Gd anomalies are a common feature of REE sorption in the presence of strong solution complexation, for example with carbonate (dominant in the ocean) or with organic ligands (e.g., Quinn et al., 2006; Christenson and Schijf, 2011). Gd/Gd* ratios on the order of 1.2 are typically encountered in the marine environment. The cold seep mussels investigated here typically display Gd/Gd* ratios exceeding 1.2. We are confident that these anomalies do not correspond to analytical artifacts resulting from biased correction of interferences produced by Ce and Pr oxides. Our procedure effectively corrects for these interferences (e.g., Charles et al., 2021), and the results obtained on standards during the course of the study demonstrate the quality of our analyses (Supplementary Table 1). Moreover, these anomalies were confirmed through multiple measurements of different fragments of the same shells, and cannot possibly relate to inappropriate normalization values, because similar Gd/Gd* values are calculated by normalizing abundances to other shale references or to chondrites (e.g.,

Barrat et al., 2012; Bau et al., 2018, not shown). We are also confident that measured Gd excesses in this study were not caused by an artefact related to the calculation of Gd anomalies. Here, Gd anomalies have been calculated by extrapolating Gd* using Tb and Dy concentrations, but similar results are obtained when Gd* is interpolated between Sm and Tb (not shown).

Positive Gd anomalies have been measured in mollusk shells living in waters polluted by Gd-based contrast agents (GBCA) used for medical imaging by magnetic resonance (e.g., Merschel and Bau, 2015; Ponnurangam et al., 2016; Akagi and Edanami, 2017; Le Goff et al., 2019). Even if this pollution is now common in coastal waters worldwide, it is highly unlikely that anthropogenic Gd pollution affects the deep sea at which our samples were collected. We tentatively propose that these Gd anomalies are the result of either biological fractionation or that they reflect the composition of the water filtered by the mussels. Since Gd anomalies are present in the shells of both thiotrophic and methanotrophic animals, they are probably not the result of any specific enzymatic activity related to nutritional symbiosis. It is possible that cold seep mussels fractionate Gd for an undiscovered use in other biological functions. One may wonder why these anomalies are not ubiquitous features of bivalve shells, including in thiotrophic species encountered at hydrothermal vents, where mussel shells lack the marked Gd anomalies. As an alternative, Gd anomalies could be inherited from water filtered by mussels. Recently, Möller et al. (2021) reported high Gd/Gd* ratios in natural, uncontaminated, spring waters and hypersaline waters (e.g., Dead Sea). Gd/Gd* ratios close to 1.7 or higher are unknown in seawaters and oceanic hydrothermal fluids. What about methane-rich fluids from cold seeps? There are limited REE data for methane-rich fluids at cold seeps (e.g., Himmler et al., 2013; Lemaitre et al., 2014). However, some of their characteristics can be inferred from the composition of the carbonates they deposit (e.g., Smrzrka et al., 2020; Bayon et al., 2020a). To date, it has not been convincingly shown that cold seeps carbonates can exhibit such anomalies, making it more than hypothetical that they are present in cold seeps fluids. The origin and significance of the Gd anomalies observed in cold seeps mussel shells is not yet understood.

6/ Conclusions and perspectives

In this study, we compared REE abundances in deep-water mussel shells from various hydrothermal and cold seep systems worldwide, confirming that animals living at

hydrothermal sites display compositional shell characteristics of surrounding fluids such as positive Eu anomalies (Bau et al., 2010). The REE distribution in bivalve shells at submarine methane seeps provides a window into the enzymatic activity of associated symbionts. Unlike thiotrophic mussels, methanotrophic species whose symbionts use light-REE to degrade methane, show characteristic light-REE enrichments in their shells. Our results confirm that only methanotrophic mussels display distinctive positive La anomalies, but selective La enrichments are not systematic to all species analyzed. The degree of light REE enrichment in shells probably depends on the distribution pattern of hosted symbionts and associated enzymatic activities, suggesting regional variability. For instance, only shells from western Pacific sites displayed strong positive La anomalies in this study. At this stage, it is unclear whether this characteristic relates to the presence of specific symbiont lineages. Future investigations of other mussel species from additional sites combined with symbiont characterization will be required to address this issue. In future studies, REE concentrations in abyssal mussel shells could be used to distinguish methanotrophic from thiotrophic species, hence serving as tools for identifying their general enzymatic pathway. This novel approach could be also applied to the sedimentary record and the investigation of fossil shells in ancient seep deposits, providing of course that diagenesis or any other post-depositional processes have not modified their REE abundances. Overall, these findings open up important multidisciplinary perspectives in biology, Earth and environmental sciences, requiring further improvements in our understanding of the REE biogeochemistry and inventory in organisms, and their role in enzymatic activities and metabolic pathways. Additional work will be also needed to assess the impact of life on REE cycling in the ocean, something that has been largely overlooked until now. Our new data reinforce the idea that methanotrophic organisms, at least those using REE-based enzymes, can leave a distinctive chemical fingerprint in rocks, in particular in methane-derived carbonates at cold seeps. Some of these authigenic carbonates contain organic-rich aragonitic cements exhibiting high light-REE enrichments (Zwicker et al., 2018). So far, these cements have been interpreted as fossil biofilms (Hagemann et al., 2013), but it is very tempting to see here the imprint of aerobic methanotrophic activity. Finally, we draw the attention of the large geochemical community studying ancient marine sedimentary formations of Precambrian age, to the interest of carrying out detailed investigations coupling REE to other conventional biogeochemical proxies such as stable isotopes in order to detect the emergence of aerobic methanotrophy on Earth.

Acknowledgements.

This project was founded by “Laboratoire d’Excellence” LabexMER (ANR-10-LABX-19) and co-funded by grants from the French Government. Shell fragments from museum specimens at SIO-BIC were provided by Charlotte Seid and Greg Rouse. Specimens from Costa Rica were collected under permits SINAC-CUS-PI-R-035-2017 and SINAC-CUSBSE-PI-R-032-2018, issued by the Ministerio de Ambiente y Energía (MINAE), Sistema Nacional de Areas de Conservación (SINAC). Collection of the Costa Rica specimens was funded by the United States National Science Foundation, grant # OCE-1634172 to Greg Rouse and Lisa Levin. We thank the captain and crew of the R/V *Atlantis*, the pilots and technicians of the HOV *Alvin*, chief scientist Erik Cordes, and the scientists of cruises AT37-13 and AT42-03 for assistance with specimen collection in Costa Rica. *Gigantidas childressi* samples were collected by R.S. Carney and made available through grant GM-09-01-08 from the US Bureau of Ocean Energy Management to R.S. Carney and B. Fry of Louisiana State University. We thank Balz Kamber for the editorial handling, Johan Schijf and Dong Feng for their constructive reviews.

References

- Akagi T., Edanami K. (2017) Sources of rare earth elements in shells and soft tissues of bivalves from Tokyo Bay. *Marine Chemistry* **194**, 55–62.
- Barrat J.A., Keller F., Amossé J., Taylor R.N., Nesbitt R.W., Hirata T. (1996) Determination of rare earth elements in sixteen silicate reference samples by ICP-MS after Tm addition and ion exchange separation. *Geostandards Newsletter* **20**, 1, 133-140.
- Barrat J.A., Zanda B., Moynier F., Bollinger C., Liorzou C., and Bayon G. (2012) Geochemistry of CI chondrites: Major and trace elements, and Cu and Zn isotopes. *Geochim. Cosmochim. Acta* **83**, 79-92.
- Barrat J.A., Dauphas N., Gillet P., Bollinger C., Etoubleau J., Bischoff A., Yamaguchi A. (2016) Evidence from Tm anomalies for non-CI refractory lithophile element proportions in terrestrial planets and achondrites. *Geochim. Cosmochim. Acta*, **176**, 1-17.
- Barrat J.A., Bayon G., Wang X., Le Goff S., Rouget M.L., Gueguen B., Ben Salem B. (2020) A new chemical separation procedure for the determination of rare earth elements and yttrium abundances in carbonates by ICP-MS. *Talanta* **219**, 121244.
- Bau M. and Dulski P. (1999) Comparing yttrium and rare earths in hydrothermal fluids from the Mid-Atlantic Ridge: implications for Y and REE behaviour during near-vent mixing and for the Y/Ho ratio of Proterozoic seawater. *Chem. Geol.* **155**, 77-90.
- Bau M., Balan S., Schmidt K., Koschinsky, A. (2010) Rare earth elements in mussel shells of the *Mytilidae* family as tracers for hidden and fossil high-temperature hydrothermal systems. *Earth Planet Sci. Lett.* **299**, 310-316.
- Bau M., Schmidt K., Koschinsky A., Hein J.R., Kuhn T., Usui A. (2014) Discriminating between different genetic types of marine ferro-manganese crusts and nodules based on rare earth elements and yttrium, *Chem. Geol.* **381**, 1-9.

- Bau M., Schmidt K., Pack A., Bendel B., Kraemer D. (2018) The European Shale: an improved data set for normalization of rare earth elements and yttrium concentrations in environmental and biological samples from Europe. *Applied Geochem.* **90**, 142-149.
- Bayon G., Lemaitre N., Barrat J.A., Wang X., Feng D., Duperron S. (2020a) Microbial utilization of rare earth elements at cold seeps related to aerobic methane oxidation. *Chem. Geol.* **555**, 119832.
- Bayon G., Lambert T., Vigier N., De Dekker P., Freslon N., Jang K., Larkin C.S., Piotrovski A.M., Tachikawa K., Thollon M., Tipper E.D. (2020b) Rare earth element and neodymium isotope tracing of sedimentary rock weathering. *Chem. Geol.* **553**, 119794.
- Brzechffa C., Goffredi S.K. (2021) Contrasting influences on bacterial symbiont specificity by co-occurring deep-sea mussels and tubeworms. *Environmental Microbiology Reports* **13**, 104-111.
- Cavanaugh, C. M., Gardiner, S. L., Jones, M. L., Jannasch, H. W., Waterbury, J. B. (1981) Prokaryotic cells in the hydrothermal vent tube worm *Riftia pachyptila* Jones: possible chemoautotrophic symbionts. *Science* **213**, 340-342.
- Cavanaugh, C. M., Levering, P. R., Maki, J. S., Mitchell, R., Lidstrom, M. E. (1987) Symbiosis of methylotrophic bacteria and deep-sea mussels. *Nature* **325**, 346-348.
- Chanton JP, Martens CS, Paull CK, Coston JA (1993) Sulfur isotope and porewater geochemistry of Florida Escarpment seep sediments. *Geochim. Cosmochim. Acta* **57**, 1253-1266
- Charles C., Barrat J.A., Pelleter E. (2021) Trace element determinations in Fe-Mn oxides by high resolution ICP-MS after Tm addition. *Talanta* **233**, 122446.
- Childress, J. J. et al. (1986) A methanotrophic marine molluscan (*Bivalvia*, *Mytilidae*) symbiosis: mussels fueled by gas. *Science* **233**, 1306-1308.
- Christenson E.A. and Schijf J. (2011) Stability of YREE complexes with the trihydroxamate siderophore desferrioxamine B at seawater ionic strength. *Geochim. Cosmochim. Acta* **75**, 7047-7062.
- Cole C.S., James R.H., Connelly D.P. and Hathorne E.C. (2014) Rare earth elements as indicators of hydrothermal processes within the East Scotia subduction zone system. *Geochim. Cosmochim. Acta* **140**, 20-38.
- Corliss J.B. and Ballard R.D. (1977) Oases of life in the cold abyss. *Natl. Geograph.* **152**, 441-453.
- Cotruvo, jr., J.A. (2019) The chemistry of lanthanides in biology: recent discoveries, emerging principles, and technological applications. *ACS Central Science* **5**, 1496-1506.
- Craddock P.R., Bach W., Seewald J.S., Rouxel O.J., Reeves E. and Tivey M.K. (2010) Rare earth element abundances in hydrothermal fluids from the Manus Basin, Papua New Guinea: indicators of sub-seafloor hydrothermal processes in back-arc basins. *Geochim. Cosmochim. Acta* **74**, 5494-5513.
- Douville, E., Bienvenu, P., Charlou, J.L., Donval, J.P., Fouquet, Y., Appriou, P., Gamo, T. (1999) Yttrium and the rare earth elements in fluids from various deep-sea hydrothermal systems. *Geochim. Cosmochim. Acta* **63**, 627-643.
- Duperron, S., Sibuet, M., MacGregor, B.J., Kuypers, M.M.M., Fisher, C.R., Dubilier, N. (2007) Diversity, relative abundance and metabolic potential of bacterial endosymbionts in three *Bathymodiolus* mussel species from cold seeps in the Gulf of Mexico. *Environmental Microbiology* **9**, 1423-1438.
- Elderfield H. (1988) The oceanic chemistry of the rare-earth elements. *Philosophical Transaction of the Royal Society of London. Series A, Mathematical and Physical Sciences* **325**, 105-126.
- Felbeck, H. (1981) Chemoautotrophic potential of the hydrothermal vent tube worm, *Riftia pachyptila* Jones (*Vestimentifera*). *Science* **213**, 336-338.

- Feng D., Peckmann J., Li N., Kiel S., Qiu J.W., Liang Q., Carney R.S., Peng Y, Tao J., Chen D. (2018) The stable isotope fingerprint of chemosymbiosis in the shell organic matrix of seep-dwelling bivalves. *Chem. Geol.* **479**, 241-250
- Hagemann A., Leefman T., Peckmann J., Hoffmann V.E., Thiel, V. (2013) Biomarkers from individual carbonate phases of an Oligocene cold-seep deposit, Washington State, USA. *Lethaia* **46**, 7-18.
- Henderson, P. (1984) General geochemical properties and abundances of the rare earth elements, in: P. Henderson (Ed.), *Rare Earth Element Geochemistry*, Elsevier, 1984, pp. 1-32.
- Herzig P.M., Hanningen M.D., Stoffers P. and the shipboard scientific party (1998) Petrology, gold mineralisation and biological communities at shallow submarine volcanoes of the New Ireland fore-arc (Papua-New Guinea): preliminary results of R/V Sonne cruise SO-133. *Inter-Ridge News* **7**, 34-38.
- Himmler, T., Haley B., Torres M.E., Klinkhammer G.P., Bohrmann G., Peckmann J. (2013) Rare earth element geochemistry in cold-seep pore waters of Hydrate Ridge, northeast Pacific Ocean. *Geo-Mar. Lett.* **33**, 369-379.
- Kenk V.C. & Wilson B.R. (1985). A new mussel (*Bivalvia*, *Mytilidae*) from hydrothermal vents, in the Galapagos Rift zone. *Malacologia* **26**, 253-271.
- Le Goff S., Barrat J.A., Chauvaud L., Paulet Y.M., Gueguen B., Ben Salem D. (2019) Compound-specific recording of gadolinium pollution in coastal waters by great scallops. *Scientific Reports* **9**, 8015.
- Lemaitre N., Bayon G., Ondréas H., Caprais J.C., Freslon N., Bollinger C., Rouget M.L., de Prunelé A., Ruffine L., Olu-Le Roy K., Sarthou G. (2014) Trace element behaviour at cold seeps and the potential export of dissolved iron to ocean. *Earth Planet. Sci. Lett.* **404**, 376-388.
- Levin, L.A., Orphan, V.J., Rouse, G.W., Rathburn, A.E., Ussler, W., Cook, G.S., Goffredi, S. K., Perez, E.M., Warren, A., Grupe, B.M., Chadwick, G., Strickrott, B. (2012) A hydrothermal seep on the Costa Rica margin: middle ground in a continuum of reducing ecosystems. *Proc. R. Soc. B Biol. Sci.* **279**, 2580-2588.
- Levin, L.A., Mendoza, G.F., Grupe, B.M., Gonzalez, J.P., Jellison, B., Rouse, G., Thurber, A.R., Warren, A. (2015) Biodiversity on the rocks: macrofauna inhabiting authigenic carbonate at Costa Rica methane seeps. *PloS One* **10**, e0131080.
- Macavoy S.E., Carney R.S., Morgan E., Macko S.A. (2008) Stable isotope variation among the mussel *Bathymodiolus childressi* and associated heterotrophic fauna at four cold-seep communities in the Gulf of Mexico. *J. Shellfish Res.* **27**, 147-151.
- Martens C.S., Chanton J.P., Paull C.K. (1991) Biogenic methane in the Florida Escarpment brine seeps. *Geology* **19**, 851-857.
- McCowin M.F., Feehery C., Rouse G.W. (2020) Spanning the depths or depth-restricted: three new species of *Bathymodiolus* (*Bivalvia*, *Mytilidae*) and a new record for the hydrothermal vent *Bathymodiolus* thermophiles at methane seeps along the Costa Rica margin. *Deep-Sea Res.* **164**, 103322.
- Merschel G., Bau M. (2015) Rare earth elements in the aragonitic shell of freshwater mussel *Corbicula fluminea* and the bioavailability of anthropogenic lanthanum, samarium and gadolinium in river water. *Sci. Total Environment* **533**, 91-101.
- Meyer A.C.S., Grundle D., Cullen J.T. (2021) Selective uptake of rare earth elements in marine systems as an indicator of and control on aerobic bacterial methanotrophy. *Earth Planet. Sci. Lett.* **558**, 116756.
- Möller P, Dulski P., De Lucia M. (2021) REY patterns and their natural anomalies in waters and brines: the correlation of Gd and Y anomalies. *Hydrology* **8**, 116, <https://doi.org/10.3390/hydrology8030116>.
- Northdurft L.D., Webb G.E., Kamber B.S. (2004) Rare earth element geochemistry of Late Devonian reefal carbonates, Canning Basin, Western Australia: confirmation of a seawater REE proxy in ancient limestones. *Geochim. Cosmochim. Acta* **68**, 263-283.

- Ozaki T., Enomoto S., Minai Y., Ambe S., Ambe F., Tominaga T. (1997) Determination of Lanthanides and other trace elements in ferns by instrumental neutron activation analysis. *J. Radioanalytical Nuclear Chem.* **217**, 117-124.
- Paull C.K., Hecker B., Commeau R., Freeman-Lynde R.P., Neumann C., Corso W.P., Golubic S., Hook J.E., Sikes E., Curray J. (1984) Biological communities at the Florida Escarpment resemble hydrothermal vent taxa. *Science* **226**, 965-967.
- Paull C.K., Jull A.J.T., Toolin L.J., Linick T. (1985) Stable isotope evidence for chemosynthesis in an abyssal seep community. *Nature* **317**, 709-711.
- Paull, C.K., Martena, C.S., Chanton, J.P., Neumann, A.C., Coston, J., Jull, A.J.T., Toolin, L.J. (1989) Old carbon in living organisms and young CaCO₃ cements from abyssal brine seeps. *Nature* **342**, 166-168.
- Petersen J.M., Zielinski F.U., Pape T., Seifert R., Moraru C., Amann R., Hourdez S., Girguis P.R., Wankel S.D., Barbe V., Pelletier E., Fink D., Borowski C., Bach W., Dubilier N. (2011) Hydrogen is an energy source for hydrothermal vent symbioses. *Nature* **476**, 176-180.
- Pol, A., Barends T.R.M., Dietl A., Khadem A.F., Eygensteyn J., Jetten M.S.M., Op Den Camp H.J.M. (2014) Rare earth metals are essential for methanotrophic life in volcanic mud pots. *Environmental Microbiology* **16**, 255-264.
- Ponnurangam, A., Bau, M., Brenner, M., Kochinsky, A. (2016) Mussel shells of *Mytilus edulis* as bioarchives of the distribution of rare earth elements and yttrium in seawater and the potential impact of pH and temperature on their partitioning behavior. *Biogeosciences* **13**, 751-760.
- Pourmand, A., Dauphas, N., Ireland, T.J. (2012) A novel extraction chromatography and MC-ICP-MS technique for rapid analysis of REE, Sc and Y: Revising CI-chondrite and Post-Archean Australian Shale (PAAS) abundances. *Chem. Geol.* **291**, 38-54.
- Quinn K.A., Byrne R.H., Schijf J. (2006) Sorption of yttrium and rare earth elements by amorphous ferric hydroxide: Influence of solution complexation with carbonate. *Geochim. Cosmochim. Acta* **70**, 4151-4165.
- Ramachandran, A. & Walsh, D. A. (2015) Investigation of XoxF methanol dehydrogenases reveals new methylotrophic bacteria in pelagic marine and freshwater ecosystems. *FEMS Microbiology Ecology* 91(10), fiv105.
- Riekenberg P.M., Carney R.S., Fry, B. (2018) Shell carbon isotope indicators of metabolic activity in the deep-sea mussel *Bathymodiolus childressi*. *Deep Sea res. Part. I* **134**, 48-54.
- Sahling, H., Masson, D.G., Ranero, C.R., Hühnerbach, V., Weinrebe, W., Klauke, I., Bürk, D., Brückmann, W., Suess, E. (2008) Fluid seepage at the continental margin offshore Costa Rica and southern Nicaragua. *Geochem. Geophys. Geosyst.* **9**, Q05S05.
- Schmidt K., Koschinsky A., Garbe-Schönberg D., Decarvallo L. and Seifert R. (2007) Geochemistry of hydrothermal fluids from the ultramafic-hosted Logatchev hydrothermal field, 15°N on the Mid-Atlantic Ridge: temporal and spatial investigation. *Chem. Geol.* **242**, 1-21.
- Schmidt K., Garbe-Schönberg D., Hannington M.D., Anderson M.O., Bühring B., Haase K., Haruel C., Lupton J., Koschinsky A. (2017) Boiling vapour-type fluids from the Nifonea vent field (New Hebrides Back-Arc, Vanuatu, SW Pacific): geochemistry of an early-stage, post-eruptive hydrothermal system. *Geochim. Cosmochim. Acta* **207**, 185-209.
- Semrau, J.D., Dispirito, A.A., GU, W., Yoon, S. (2018) Metals and Methanotrophy. *Applied and Environmental Microbiology* **84**, e02289-17.
- Shiller, A.M., Chan, E.W., Joung D.J., Redmond, M.C., Kessler, J.D. (2017) Light rare earth element depletion during *Deepwater Horizon* blowout methanotrophy. *Scientific Reports* **7**, 10389, DOI:10.1038/s41598-017-11060-z

- Skovran, E., Palmer, A. D., Rountree, A. M., Good, N. M., Lidstrom, M. E. (2011) XoxF is required for expression of methanol dehydrogenase in *Methylobacterium extorquens* AM1. *J. Bacteriol.* 193, 6032-6038.
- Smrzka D., Zwicker J., Misch D., Walkner C., Gier S., Monien P., Bohrmann G., Peckmann J. (2019) Oil seepage and carbonate formation: A case study from the southern Gulf of Mexico. *Sedimentology* **66**, 2318-2353.
- Smrzka D., Feng D., Himmler T., Zwicker J., Hu Y., Monien P., Tribovillard N., Chen D., Peckmann J. (2020) Trace elements in methane-seep carbonates: potentials, limitations, and perspectives. *Earth Sci Rev.* **208**, 103263.
- Taubert M. *et al.* (2015) XoxF encoding an alternative methanol dehydrogenase is widespread in coastal marine environments. *Environ. Microbiol.* 17, 3937–3948.
- Tommasi F., Thomas P.J., Pagano G., Perono G.A., Oral R., Lyons D.M., Toscanesi M., Trifuoggi M. (2021) Review of rare earth elements as fertilizers and feed additives: a knowledge gap analysis. *Archives of Environmental Contamination and Toxicology* **81**, 531-540.
- von Cosei R., Janssen R. (2008) Bathymodioline mussels of the *Bathymodiolus* (s.l.) *childressi* clade from methane seeps near Edison Seamount, New Ireland, Papua New Guinea. *Arch. Molluskenkunde* **137**, 195-224.
- Wang X., Barrat J.A., Bayon G., Chauvaud L., Feng D. (2020) Lanthanum anomalies as fingerprints of methanotrophy. *Geochem. Persp. Let.* **14**, 26–30.
- Xu T, Feng D., Tao J., Qiu J-W. (2019) A new species of deep-sea mussel (Bivalvia: Mytilidae; Gigantidas) from the South China Sea: Morphology, phylogenetic position, and gill-associated microbes. *Deep Sea Res., Pt. 1* **146**, 79-90.
- Zwicker J., Smrzka D., Himmler T., Monien P., Gier S., Goedert J.L., Peckmann J. (2018) Rare earth elements as tracers for microbial activity and early diagenesis: A new perspective from carbonate cements of ancient methane-seep deposits. *Chem. Geol.* **501**, 77-85.




Figure 1. World map and location of the hydrothermal vents and cold seeps where mussels have been analyzed for REE and Y abundances (Bau et al., 2010; Wang et al., 2020; this study).




Figure 2. PAAS-normalized REY patterns for shells of thiotrophic (T) mussels from hydrothermal vents.




Figure 3. PAAS-normalized REY patterns for mussel shells from vents located west of Costa Rica. Here, all the mussels are thiotrophic (T).




Figure 4. PAAS-normalized REY patterns for mussel shells from cold seeps located in the China Sea (site F and Haima, Wang et al., 2020), and from Edison Seamount (this work). *Bathymodiolus aduloides* mussels are thiotrophic (T). The others are methanotrophic (M).




Figure 5. PAAS-normalized REY patterns for mussel shells from the Gulf of Mexico (M: methanotrophic; T: thiotrophic).




Figure 6. Eu vs. Sm plot for mussel shells from hydrothermal vents and cold seeps (data from Bau et al., 2010, Wang et al., 2020, and this study).




Figure 7. Histograms of Sm abundances in mussel shells from hydrothermal vents and cold seeps (data from Bau et al., 2010, Wang et al., 2020, and this study). When two fragments of the same shell were analyzed, we used their average here.




Figure 8. $(Tb/Yb)_{sn}$ vs. $(Pr/Sm)_{sn}$ plot for mussels shells (data from Bau et al., 2010, Wang et al., 2020, and this study). Fields for seawater (SW), pore waters (PW) and vent fluids (VF) are drawn from a database of results of the literature (see Bayon et al. (2020b) for the references for seawaters and pore waters, and Bau and Dulski (1999, Douville et al. (1999), Schmidt et al. (2007, 2017), Craddock et al. (2010), and Cole et al. (2014) for vent fluids).




Figure 9. $(\text{La}/\text{Nd})_{\text{sn}}$ vs. $(\text{Pr}/\text{Nd})_{\text{sn}}$ plot for mussels (data from Bau et al., 2010, Wang et al., 2020, and this study). Fields for seawater (SW), pore waters (PW) and vent fluids (VF) are drawn from a database of results of the literature (see figure 8 for the references). Oils from seeps from the southern Gulf of Mexico are shown for comparison (Smrzka et al., 2019).




Figure 10. Histograms of Gd/Gd* ratios in mussels shells from hydrothermal vents and cold seeps (data from Bau et al., 2010, Wang et al., 2020, and this study). When two fragments of the same shell were analyzed, we used their average here.