



**HAL**  
open science

# AtEXT3 is not essential for early embryogenesis or plant viability in Arabidopsis

Nicolas Max Doll, Eduardo Berengue, Jekaterina Truskina, Gwyneth Ingram

## ► To cite this version:

Nicolas Max Doll, Eduardo Berengue, Jekaterina Truskina, Gwyneth Ingram. AtEXT3 is not essential for early embryogenesis or plant viability in Arabidopsis. *New Phytologist*, 2022, 236 (5), pp.1629-1633. 10.1111/nph.18452 . hal-03841277

**HAL Id: hal-03841277**

**<https://cnrs.hal.science/hal-03841277v1>**

Submitted on 7 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

**AtEXT3 is not essential for early embryogenesis or plant viability in Arabidopsis.**

Nicolas Max Doll<sup>\*†</sup>, Eduardo Berenguer<sup>†</sup>, Jekaterina Truskina and Gwyneth Ingram<sup>1\*</sup>

<sup>1</sup> Laboratoire Reproduction et Développement des Plantes, ENS de Lyon, CNRS, INRAE, UCBL, F-69342, Lyon, France.

<sup>†</sup> Authors contributed equally to the study.

\* Authors for correspondence : [Nicolas.Doll@ugent.vib.be](mailto:Nicolas.Doll@ugent.vib.be), [Gwyneth.Ingram@ens-lyon.fr](mailto:Gwyneth.Ingram@ens-lyon.fr)

*ORCID IDs*

N.M. Doll: 0000-0001-5429-8920

E. Berenguer: 0000-0002-0902-1057

G. Ingram: 0000-0002-1425-9545

Running title: *AtEXT3* is a non-essential gene

32 EXTENSINs (EXTs) are a subclass of the very diverse family of cell wall proteins known as  
33 hydroxyproline-rich glycoproteins. The EXT subclass is defined by the presence of at least  
34 two repeats of the Ser-Pro<sub>3-5</sub> motif in which the proline residues can be hydroxylated and,  
35 subsequently, O-glycosylated with arabinose containing oligosaccharides leading to the  
36 formation of hydrophilic domains (Cannon *et al.*, 2008; Velasquez *et al.*, 2011; Ogawa-  
37 Ohnishi *et al.*, 2013). In addition, many EXTs from land-plants also contain distinct and  
38 characteristic hydrophobic Tyr-containing motifs (Schnabelrauch *et al.*, 1996; Held *et al.*,  
39 2004; Cannon *et al.*, 2008). These have recently been termed cross-linking (CL)-EXTs,  
40 because of their ability to undergo peroxidase-mediated covalent cross linking *in vitro*  
41 through their Tyr residues, a phenomenon that has been proposed to underlie important  
42 structural functions in plant cell walls (Marzol *et al.*, 2018). EXTs have a remarkable ability to  
43 self-assemble into dendritic networks *in vitro* (Cannon *et al.*, 2008). This capacity, proposed  
44 to be based on self-ordering due to the alternation of hydrophobic and hydrophilic  
45 domains (Lampert *et al.*, 2011), may mediate observed interactions with other cell wall  
46 components, including pectins (Valentin *et al.*, 2010).

47 Despite an abundance of information regarding the biochemical behavior of EXTs *in*  
48 *vitro*, information on their biological functions *in vivo* remains remarkably scarce, possibly  
49 due to extensive redundancy between the multiple genes encoding EXTs and EXT-like  
50 proteins within plant genomes. Work in Arabidopsis has identified potential roles for AtEXTs  
51 in promoting root-hair elongation (Velasquez *et al.*, 2011), pollen development (Choudhary  
52 *et al.*, 2015) and early embryogenesis (Hall & Cannon, 2002; Saha *et al.*, 2013; Chen *et al.*,  
53 2015). This last function is our focus in this comment.

54 Following an enhancer-trap (transposon)-based mutagenesis screen, Hall and Cannon  
55 (Hall & Cannon, 2002) isolated a seedling-lethal mutant called *root-shoot-hypocotyl-*  
56 *defective (rsh)* which, when homozygous, produces disorganized embryos that arrest either  
57 before (Cannon *et al.*, 2008) or just after (Hall & Cannon, 2002) germination. Mutants show  
58 clear defects in cell-plate formation. A transposon insertion in the promoter of the *AtEXT3*  
59 gene was stated to be the causal lesion in *rsh* mutants. Consistent with this,  
60 complementation of the *rsh* phenotype using an *AtEXT3* genomic DNA fragment was  
61 reported. However, in two subsequent studies, “self-rescue” of the *rsh* phenotype to a fully  
62 wild-type phenotype was reported in up to 20% of genetically *atext3* homozygous plants  
63 (Saha *et al.*, 2013; Chen *et al.*, 2015). No intermediate phenotypes were reported, and  
64 “rescue” was transmitted stably to all the progeny in subsequent generations. This calls into  
65 question the validity of genetic complementation experiments, since the reported “self-  
66 rescue” phenomenon would interfere with this approach. A mechanism involving  
67 compensatory regulation of other AtEXT-encoding genes was evoked to explain this  
68 phenomenon. Nonetheless it appears difficult, based on the published data, to exclude the  
69 possibility that the reported *rsh* phenotype is caused by a lesion linked to the original  
70 transposon insertion in *AtEXT3*, but affecting a separate locus, and lost by segregation in  
71 “rescued” lines.

72 We became interested in the *AtEXT3* gene as a result of our discovery that the  
73 developing embryo of Arabidopsis is covered with an endosperm-derived “sheath” rich in  
74 epitopes detected by anti-EXT antibodies (Moussu *et al.*, 2017). *In silico* data (Le *et al.*, 2010)  
75 concur with published *in situ* hybridization data (Francoz *et al.*, 2016) in showing that *AtEXT3*  
76 is expressed primarily in the embryo-surrounding endosperm during Arabidopsis seed  
77 development, rather than in the embryo itself. We were therefore keen to further  
78 investigate the function of *AtEXT3* and we generated a series of new alleles in the Col-0

79 background using the CRISPR-Cas9 based technique. We selected 4 alleles containing  
80 deletions or insertions within the 5' coding sequence of the *AtEXT3* gene (Figure 1A). All are  
81 predicted to produce either strongly truncated or frame-shifted proteins (Figure 1B). Being  
82 aware of recent work suggesting that 5' mutations generated by CRISPR can lead to the  
83 production of functional proteins due to the use of downstream in-frame ATGs (Smits *et al.*,  
84 2019), we verified the *AtEXT3* coding sequence (which lacks introns) for such sequences.  
85 None are present. We therefore conclude that all four alleles are null. Homozygous plants  
86 were all fully phenotypically wild-type in terms of root growth, plant growth and  
87 reproductive development (Figure C-E). No defective seeds or seedlings were detected for  
88 any line, in any generation.

89 Our results suggest that loss of AtEXT3 do not lead to detectable defects in development  
90 in the Col-0 background. However, the *rsh* mutant originally described by Hall and Cannon  
91 was identified in the Landsberg *erecta* (L-*er*) background. To eliminate the possibility that a  
92 suppressor exists in Col-0 that is not present in the L-*er* background, we carried out CRISPR  
93 experiments in the L-*er* background, this time using guides predicted to interrupt the  
94 EXTENSIN 2 domains of the AtEXT3 protein. Of multiple null alleles brought to homozygosity  
95 we selected three deletions predicted to produce either strongly truncated or frame-shifted  
96 proteins (Figure 2). We were again unable to detect any defects in root growth, plant growth  
97 or reproductive development in any of these lines. In addition, we confirmed the absence of  
98 phenotypes during seed development, and germination (Figure S1).

99 Taken together, our results indicate that AtEXT3 is not essential for embryogenesis and  
100 plant development in Arabidopsis, suggesting that its function is likely redundant with other  
101 related proteins. We feel that it is important that our result be communicated to the  
102 scientific community, since the role of AtEXT3 has been very widely cited as a proof of the  
103 important function of EXTs in plant development and particularly in cytokinesis. Although  
104 fully convinced that EXTs are of primary importance in plants, we propose that the roles of  
105 AtEXT3 may be almost fully redundant with those of other related proteins. Supporting this  
106 hypothesis, phenotypeless *atext3* mutants were shown to have extensive changes in the  
107 expression of other AtEXT-encoding genes, or the accumulation of AtEXT proteins (Saha *et al.*,  
108 2013; Chen *et al.*, 2015). It is possible that a similar phenomenon occurs in the CRISPR  
109 generated *atext3* mutants.

110 Finally, our study does not call into question the presence of AtEXT3 in cell walls post  
111 germination, or the beautiful biochemical studies of AtEXT3 behavior *in vitro* that have been  
112 published (Cannon *et al.*, 2008).

113

## 114 **Material and methods**

115

### 116 **CRISPR/Cas9 allele generation**

117 The CRISPR/Cas9 alleles of *AtEXT3* (*AT1G21310*) were generated through the method  
118 described by Wang and co-workers for double guides assembly (Wang *et al.*, 2015). The 2  
119 guides were designed on the CRISPR-P v2.0 software and chosen to target the 5' end of the  
120 gene (first guide: 5'-GAAGAATAGAAATAGTTAG-3', second guide: 5'-TATTCTTCTCCTCCACCAC-  
121 3'). For the cloning, the primers used were 5'-  
122 ATATATGGTCTCGATTGGAAGAATAGAAATAGTTAGGTT-3' and 5'-  
123 TGGAAGAATAGAAATAGTTAGGTTTTAGAGCTAGAAATAGC -3' for the first guide and 5'-  
124 AACGTGGTGGAGGAGAAGAATACAATCTCTTAGTTCGACTCTAC -3' and 5'-

125 ATTATTGGTCTCGAAACGTGGTGGAGGAGAAGAATACAA -3' for the second guide. The  
126 pHEE401 plasmid was used as the destination vector.

127

128 For the generation of the CRISPR/Cas9 alleles of *EXT3* in the *L-er* background the method  
129 described by Stuttmann and co-workers was used (Stuttmann *et al.*, 2021). Two guides (first  
130 guide: 5'-GGTGGGGAGTGGTATACCGG-3', second guide: 5'-ATACAAATCTCCACCTCCAC-3')  
131 were inserted into the pDGE332 and pDGE334 shuttle vectors and recombined into the  
132 pDGE347 recipient vector.

133

#### 134 **Stable transformation of *Arabidopsis thaliana* (L.) Heynh.**

135 The plasmids generated were first transformed into the C58PMP90 *Agrobacterium* strain by  
136 electroporation at 2,2kV in a 1mL cuvette (Eurogenetec). *Agrobacteria* were then grown at  
137 28°C for 2h in LB liquid media without antibiotics before being spread on Petri dishes  
138 containing YEB solid media with rifampicin (50 mg.L<sup>-1</sup>), gentamicin (20 mg.L<sup>-1</sup>) and kanamycin  
139 (50 mg.L<sup>-1</sup>) for the pHEE401 based plasmid or rifampicin (50 mg.L<sup>-1</sup>), gentamicin (20 mg.L<sup>-1</sup>)  
140 and spectinomycin (100 mg.L<sup>-1</sup>) for the pDGE347 based plasmid. After 2 days, the  
141 *Agrobacteria* were used to transform Col-0 plants by the floral-dip method (Logemann *et*  
142 *al.*, 2006). Selection of transgenic plants was performed on MS plates with 30 mg.L<sup>-1</sup> of  
143 hygromycin.

144

#### 145 **Identification of the mutant alleles and generation of homozygous plants.**

146 Amplification of the region around the cutting sites was performed by PCR on DNA extracted  
147 from transgenic Col-0 plants using the 5'-GGGTGTGAAGGGAAGGCACTAAATC-3' and 5'-  
148 GTAAACGTAGTGCTTCTTTGGTGG-3' primers. For the transgenic *L-er* plants the primers used  
149 were 5'-GGGTGTGAAGGGAAGGCACTAAATC-3' and 5'-AGGTGGGGGTGGGGAATGGTA-3'.  
150 Sequencing was carried out with the 5'-GTAAACGTAGTGCTTCTTTGGTGG-3' (Col-0) or and 5'-  
151 GGGTGTGAAGGGAAGGCACTAAATC-3' (*L-er*) primers. Homozygous mutants for *atext3* were  
152 selected from the T2 plants. In the Col-0 background the CRISPR/Cas9 cassette was removed  
153 by selection of hygromycin sensitive plants and validated by an absence of cassette  
154 amplification by PCR with the 5'-TGTCCCAGGATTAGAAATGATTAGGC-3' and 5'-  
155 AGCCCTCTTCTTCGATCCATCAAC-3' primers. Homozygous mutants were confirmed by  
156 sequencing. In *L-er* transgenic plants non-fluorescent seeds were selected in the T2 and  
157 homozygous mutants were confirmed by sequencing.

158

#### 159 **Plant growth**

160 For the selection processes and *in vitro* growth (Figure 1C and 2C), seeds were gas sterilized  
161 with Chlorine gas (3mL of HCl (33%) in 100mL of bleach) for at least 3 hours in a hermetic  
162 box. Seeds were then sown on MS medium with 0.5% of sucrose and, for the selection of  
163 Col-0 mutants, with 30 mg.L<sup>-1</sup> of hygromycin. Stratification was carried out 2 days at 4°C in  
164 darkness. Plates were then transferred into growth chambers in long day condition (16h  
165 light, 21°C). For the *in vitro* growth experiments, plants were grown for 14 days (Col 0) or 10  
166 days (*L-er*) and pictures were taken with an EPSON perfection V300 PHOTO scanner. For  
167 germination tests, plants were grown for 12 days (100 seeds per plate). In all the other  
168 cases, seedlings were transferred after 7 days onto soil (Argile 10 (favorit)) and grown in long  
169 day conditions (16h light, 21°C). Images of plants at 29 DAS and 46 DAS (Figure 1D and E) or  
170 28 DAS and 46 DAS (Figure 2D and 2E) were taken with a Cannon EOS 450D camera with a

171 Sigma 50 mm f/2,8 DG Macro objective. For analysis of silique filling, individual siliques were  
172 opened at the mature green stage, and seeds were counted under a binocular microscope.

173

#### 174 **Seed clearing**

175 To visualize developing seeds, the siliques at 7 DAP were dissected with a needle and forceps  
176 on adhesive tape on a microscope slide and mounted in the clearing solution (1 vol glycerol /  
177 7 vol chloral hydrate liquid solution, VWR Chemicals). After 48h incubation at 4°C samples  
178 were imaged under Zeiss Axio Imager M2 microscope.

179

180 **Acknowledgements:** We thank Alexis Lacroix, Patrice Bolland, Justin Berger, Isabelle  
181 Desbouchages and Hervé Leyral for technical assistance, Cindy Vial, Stéphanie Maurin,  
182 Laureen Grangier and Nelly Camilleri for administrative assistance.

183

184 **Funding:** The study was financed by the French Agence National de Recherche (ANR- 17-  
185 CE20-0027) (supporting J.T. and N.M.D.) and ANR- 20-CE20-CLEANSE (supporting E.B.).  
186 N.M.D. was also funded by a PhD fellowship from the Ministère de l'Enseignement Supérieur  
187 et de la Recherche.

188

189 **Author contributions:** G.I. and N.M.D. led the study. G.I. obtained funding for the study. G.I.,  
190 supervised the work. N.M.D., E.B and J.T carried out the experiments. G.I. and N.M.D. wrote  
191 the paper with input from all authors. N.M.D. and E.B. contributed equally.

192

193 **Data availability:** The data that support the findings of this study are available from the  
194 corresponding author upon reasonable request.

195

196 **Competing interests:** None declared.

197

#### 198 **References**

199 **Cannon MC, Terneus K, Hall Q, Tan L, Wang Y, Wegenhart BL, Chen L, Lampert DTA, Chen**  
200 **Y, Kieliszewski MJ. 2008.** Self-assembly of the plant cell wall requires an extensin scaffold.  
201 *Proceedings of the National Academy of Sciences of the United States of America* **105**: 2226–  
202 2231.

203 **Chen Y, Ye D, Held MA, Cannon MC, Ray T, Saha P, Frye AN, Mort AJ, Kieliszewski MJ. 2015.**  
204 Identification of the Abundant Hydroxyproline-Rich Glycoproteins in the Root Walls of Wild-  
205 Type Arabidopsis, an ext3 Mutant Line, and Its Phenotypic Revertant. *Plants (Basel,*  
206 *Switzerland)* **4**: 85–111.

207 **Choudhary P, Saha P, Ray T, Tang Y, Yang D, Cannon MC. 2015.** EXTENSIN18 is required for  
208 full male fertility as well as normal vegetative growth in Arabidopsis. *Frontiers in Plant*  
209 *Science* **6**: 553.

210 **Francoz E, Ranocha P, Pernot C, Le Ru A, Pacquit V, Dunand C, Burlat V. 2016.**  
211 Complementarity of medium-throughput in situ RNA hybridization and tissue-specific  
212 transcriptomics: case study of Arabidopsis seed development kinetics. *Scientific Reports* **6**:  
213 24644.

- 214 **Hall Q, Cannon MC. 2002.** The Cell Wall Hydroxyproline-Rich Glycoprotein RSH Is Essential  
215 for Normal Embryo Development in Arabidopsis. *The Plant Cell* **14**: 1161–1172.
- 216 **Held MA, Tan L, Kamyab A, Hare M, Shpak E, Kieliszewski MJ. 2004.** Di-isodityrosine Is the  
217 Intermolecular Cross-link of Isodityrosine-rich Extensin Analogs Cross-linked in Vitro\*.  
218 *Journal of Biological Chemistry* **279**: 55474–55482.
- 219 **Lampont DTA, Kieliszewski MJ, Chen Y, Cannon MC. 2011.** Role of the extensin superfamily  
220 in primary cell wall architecture. *Plant Physiology* **156**: 11–19.
- 221 **Le BH, Cheng C, Bui AQ, Wagmaister JA, Henry KF, Pelletier J, Kwong L, Belmonte M,  
222 Kirkbride R, Horvath S, et al. 2010.** Global analysis of gene activity during Arabidopsis seed  
223 development and identification of seed-specific transcription factors. *Proceedings of the  
224 National Academy of Sciences of the United States of America* **107**: 8063–8070.
- 225 **Logemann E, Birkenbihl RP, Ülker B, Somssich IE. 2006.** An improved method for preparing  
226 Agrobacterium cells that simplifies the Arabidopsis transformation protocol. *Plant Methods*  
227 **2**: 16.
- 228 **Marzol E, Borassi C, Bringas M, Sede A, Rodríguez Garcia DR, Capece L, Estevez JM. 2018.**  
229 Filling the Gaps to Solve the Extensin Puzzle. *Molecular Plant* **11**: 645–658.
- 230 **Moussu S, Doll NM, Chamot S, Brocard L, Creff A, Fourquin C, Widiez T, Nimchuk ZL,  
231 Ingram G. 2017.** ZHOUP1 and KERBEROS Mediate Embryo/Endosperm Separation by  
232 Promoting the Formation of an Extracuticular Sheath at the Embryo Surface. *The Plant Cell*  
233 **29**: 1642–1656.
- 234 **Ogawa-Ohnishi M, Matsushita W, Matsubayashi Y. 2013.** Identification of three  
235 hydroxyproline O-arabinosyltransferases in *Arabidopsis thaliana*. *Nature Chemical Biology* **9**:  
236 726–730.
- 237 **Saha P, Ray T, Tang Y, Dutta I, Evangelous NR, Kieliszewski MJ, Chen Y, Cannon MC. 2013.**  
238 Self-rescue of an EXTENSIN mutant reveals alternative gene expression programs and  
239 candidate proteins for new cell wall assembly in Arabidopsis. *The Plant Journal: For Cell and  
240 Molecular Biology* **75**: 104–116.
- 241 **Schnabelrauch LS, Kieliszewski M, Upham BL, Alizedeh H, Lampont DT. 1996.** Isolation of pl  
242 4.6 extensin peroxidase from tomato cell suspension cultures and identification of Val-Tyr-  
243 Lys as putative intermolecular cross-link site. *The Plant Journal: For Cell and Molecular  
244 Biology* **9**: 477–489.
- 245 **Smits AH, Ziebell F, Joberty G, Zinn N, Mueller WF, Clauder-Münster S, Eberhard D, Fälth  
246 Savitski M, Grandi P, Jakob P, et al. 2019.** Biological plasticity rescues target activity in  
247 CRISPR knock outs. *Nature Methods* **16**: 1087–1093.
- 248 **Stuttman J, Barthel K, Martin P, Ordon J, Erickson JL, Herr R, Ferik F, Kretschmer C, Berner  
249 T, Keilwagen J, et al. 2021.** Highly efficient multiplex editing: one-shot generation of 8×  
250 *Nicotiana benthamiana* and 12× Arabidopsis mutants. *The Plant Journal: For Cell and  
251 Molecular Biology* **106**: 8–22.

252 **Valentin R, Cerclier C, Geneix N, Aguié-Béghin V, Gaillard C, Ralet M-C, Cathala B. 2010.**  
253 Elaboration of extensin-pectin thin film model of primary plant cell wall. *Langmuir: the ACS*  
254 *journal of surfaces and colloids* **26**: 9891–9898.

255 **Velasquez SM, Ricardi MM, Dorosz JG, Fernandez PV, Nadra AD, Pol-Fachin L, Egelund J,**  
256 **Gille S, Harholt J, Ciancia M, et al. 2011.** O-glycosylated cell wall proteins are essential in  
257 root hair growth. *Science (New York, N.Y.)* **332**: 1401–1403.

258 **Wang Z-P, Xing H-L, Dong L, Zhang H-Y, Han C-Y, Wang X-C, Chen Q-J. 2015.** Egg cell-specific  
259 promoter-controlled CRISPR/Cas9 efficiently generates homozygous mutants for multiple  
260 target genes in Arabidopsis in a single generation. *Genome Biology* **16**: 144.

261

262 **Keywords:** cell wall, cytokinesis, development, embryogenesis, EXTENSIN

263

## 264 **Figure Legends:**

265 **Figure 1: Loss of AtEXT3 function gives no obvious phenotypes in the Arabidopsis Col-0**  
266 **background.**

267 A) DNA sequence of the AtEXTENSIN3 (AT1G21310) gene between position +60 and +118  
268 from the predicted start of translation. The CRISPR/Cas9 alleles generated are represented  
269 below and named *atext3-1 to -4*. The protospacer adjacent motif (PAM) sequences of the 2  
270 guides used are underlined. The mutations in the mutant alleles sequence are indicated in  
271 red. B) Predicted consequences of the mutations on the AtEXT3 protein. The 3 EXTENSIN-2  
272 domains predicted by the Protein Families database (Pfam) are represented as well as the  
273 position of the 2 guides used. For the predicted mutant proteins, sequences that differ from  
274 those of Col-0 due to a frameshift are represented in grey. The amino acid positions from the  
275 predicted first methionine are indicated. C) 14 days after stratification (DAS) seedlings,  
276 grown *in vitro* for Col-0 and for the four AtEXT3 null alleles. D) 29 DAS plants grown on soil  
277 for Col-0 and for the four AtEXT3 null alleles. E) 46 DAS plants grown on soil for Col-0 and for  
278 the four AtEXT3 null alleles. Scale bars in C and D = 2cm and in E = 3cm.

279

280 **Figure 2: Loss of AtEXT3 function gives no obvious phenotypes in the Arabidopsis L-er**  
281 **background.**




282 A) DNA sequence of the AtEXTENSIN3 (AT1G21310) gene between position +145 and +245  
283 from the predicted start of translation. The CRISPR/Cas9 alleles generated are represented  
284 below and named *atext3-5 to -7*. The protospacer adjacent motif (PAM) sequences of the 2  
285 guides used are underlined. The mutations in the mutant alleles are indicated in red. B)  
286 Predicted consequence of the mutations on the AtEXT3 protein. The 3 EXTENSIN-2 domains  
287 predicted by the Protein Families database (Pfam) are represented as well as the position of  
288 the 2 guides used. For the mutant proteins, sequences that differ from L-er due to a shift in  
289 the ORF are represented in grey. The amino acid positions from the predicted first  
290 methionine are indicated. C) 10 days after stratification (DAS) seedlings, grown *in vitro* for L-  
291 er and for the three AtEXT3 null alleles. D) 28 DAS plants grown on soil for L-er and for the  
292 three AtEXT3 null alleles. E) 46 DAS plants grown on soil for L-er and for the three AtEXT3

293 null alleles. Scale bars in C and D = 2cm and in E = 3cm.

**A**

```

+60                                     +118
Col-0  TTTGTATCTCAATCAACCGCTA-ACTATTTCTATTCTTCTCCTCCAC-CACCGGTTAAAC
atext3_1 TTTGTATCTCAATCAACCGCTA-A-----CACCGGTTAAAC
atext3_2 TTTG-----TATTCTTCTCCTCCACACACCGGTTAAAC
atext3_3 TTTGTATCTCAATCAACCGCTAAACTATTTCTATTCTTCTCCTCCACACACCGGTTAAAC
atext3_4 TTTGTATCTCAATCAA-----TCTATTTCTATTCTTCTCCTCCACTCACCGGTTAAAC
  
```



294

**A**

```

+145                                     +200
L-er   CTCCTCCACC-GGTATACCACTCCCCACCACCACCTAAGAAGCACTACGAATACAA
atext3-5 CTCCTCCACC-GGTTTACCATTCC-----CCACCTAAGAAGCACTACGAATACAA
atext3-6 CTCCTCCACC-GG-----CAA
atext3-7 CTCCTCCACCGGGTATACCACTCCCCACCACCACCTAAGAAGCACTACGAATACAA

+200                                     +245
L-er   ATCTCCACCTCC-ACCGGTTAAGCACTACTCACCTCCTCCGGTTTA
atext3-5 ATCTCCACCTCCCACCGGTTAAGCACTACTCACCTCCTCCGGTTTA
atext3-6 ATCTCCACCTCC-ACCGGTTAAGCACTACTCACCTCCTCCGGTTTA
atext3-7 ATCTCCACCTC-----GTTTA

```

