



Absolute Versus Relative Changes in Cardiac Troponin T: Corresponding Cutoffs Based on Quantile Generalized Additive Models (QGAM)

Denis Monneret, Dominique Bonnefont-Rousselot, Matteo Fasiolo

► To cite this version:

Denis Monneret, Dominique Bonnefont-Rousselot, Matteo Fasiolo. Absolute Versus Relative Changes in Cardiac Troponin T: Corresponding Cutoffs Based on Quantile Generalized Additive Models (QGAM). Clinical Laboratory, 2022, 68 (02/2022), 10.7754/Clin.Lab.2021.210607 . hal-03864166

HAL Id: hal-03864166

<https://cnrs.hal.science/hal-03864166>

Submitted on 21 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Absolute *versus* relative changes in cardiac troponin T: corresponding cut-offs based on quantile generalized additive models (qgam)

Running title: Absolute vs. relative cardiac troponin T changes

Denis Monneret^{1*}, PharmD, PhD; Dominique Bonnefont-Rousselot, PharmD, PhD^{2,3}; Matteo Fasiolo⁴, PhD.

¹ Service de Biochimie et Biologie Moléculaire, Laboratoire de Biologie Médicale Multisite (LBMMS), Hospices Civils de Lyon (HCL), Lyon, F-69000, France.

² Université de Paris, CNRS, INSERM, UTCBS, F-75006 Paris, France.

³ Service de Biochimie Métabolique, AP-HP, Hôpital Pitié-Salpêtrière, F-75013, Paris, France.

⁴ School of Mathematics, University of Bristol, Bristol BS8 1UG, UK.

*** Corresponding author:** Dr Denis Monneret

Current address: Service de Biochimie et Biologie Moléculaire, Laboratoire de Biologie Médicale Multisite (LBMMS), Hospices Civils de Lyon, F-69000, Lyon, France.

Former address: Service de Biochimie Métabolique, AP-HP, Hôpital Pitié-Salpêtrière, F-75013, Paris, France. E-mail: dmonneret2@gmail.com. Phone: (+33) 6 66 10 77 06

ORCID and email address:

Denis Monneret: <https://orcid.org/0000-0002-6491-0607>; dmonneret2@gmail.com

Dominique Bonnefont-Rousselot: <https://orcid.org/0000-0003-4689-9202>;
dominique.rousselot@aphp.fr

Matteo Fasiolo: <https://orcid.org/0000-0003-2335-5536>; matteo.fasiolo@gmail.com

Type of manuscript: Short communication

Description: ±1460 words, 2 figures, 1 table, 10 references.

Conflict of interest: None declared.

Research funding: None declared

Abstract

Background: The diagnosis of myocardial injury/infarction (MI) mainly relies on relative changes in cardiac troponin. However, absolute change cut-offs provide greater diagnostic sensitivity. We determined the absolute changes in high-sensitive cardiac troponin T concentrations ($\text{abs}\Delta\text{hs-cTnT}$) corresponding to the main relative cut-offs ($\text{rel}\Delta\text{hs-cTnT}$), using a quantile generalized additive model (qgam).

Methods: Plasma $\Delta\text{hs-cTnT}$ from patients selected with a time variation of 1 to 6 hours were collected over a 6-year period. The $\text{abs}\Delta\text{hs-cTnT}$ -to- $\text{rel}\Delta\text{hs-cTnT}$ relationship was fitted using qgam, after ordered quantile-based normalization (OQN) to reduce the influence of extreme values.

Results: The qgam regression curve was nonlinear. Classifying patients ($n=9753$) above the recommended $\text{rel}\Delta\text{hs-cTnT}$ and predicted $\text{abs}\Delta\text{hs-cTnT}$ cut-offs as positive, the MI diagnosis rates were similar, and more reliable using the OQN-transformed data-based qgam, as compared to the untransformed data-based one.

Conclusions: Through an optimized qgam-based approach accounting for heavy-tailed distributions, absolute $\Delta\text{hs-cTnT}$ are provided for the corresponding relative $\Delta\text{hs-cTnT}$ cut-offs.

Keywords: high-sensitivity cardiac troponin, variations, myocardial infarction.

Introduction

Cardiac troponin change (ΔcTn) is the cornerstone of myocardial infarction/injury (MI) diagnoses. When cTn values are $>99^{\text{th}}$ percentile URL (99PURL), a $>20\%$ rise and/or fall directs the diagnosis towards acute MI, whereas a $\leq 20\%$ change suggests chronic myocardial injury [1]. When the initial baseline cTn value is $\leq 99^{\text{th}}$ PURL, a relative change $>50\%$ is in favor of MI, this relative cut-off taking into account both analytical and biological variations. The 4th Universal Definition of MI (4UDMI) mainly considers relative ΔcTn , but it also presents the advantage of absolute ΔcTn . The extent to which absolute or relative change is preferable for MI diagnosis is an old debate. Ten years ago, Reichlin et al showed a 2-hour absolute change in high-sensitive cardiac troponin T (abs $\Delta\text{hs-cTnT}$) of higher diagnostic accuracy for acute MI than the relative change (rel $\Delta\text{hs-cTnT}$), regardless of whether the baseline cTn levels were low or high [2]. Similarly, Mueller et al demonstrated that abs $\Delta\text{hs-cTnT}$ performs better than rel $\Delta\text{hs-cTnT}$ for the diagnosis of acute coronary syndrome (ACS) and acute non-ACS-related troponin increases [3]. Indeed, given that the rise and/or fall of cTn often exceeds 20% in both these contexts, and does so systematically in acute cardiac diseases, the “diagnostic overlap” explains why rel $\Delta\text{hs-cTnT}$ fails to rule-in non-ST-segment elevation MI (NSTEMI). In accordance, and based on these two main publications, the 4UDMI underlines that “*absolute cTn changes appear superior to relative percent changes with hs-cTn assays (...) especially when the initial value is increased*”, and claims that “*the use of a fixed absolute value change criteria (...) provides greater sensitivity*” [1]. For greater changes, recommendations rely on abs $\Delta\text{hs-cTnT}$ -based diagnostic criteria, notably 5 times the 99PURL with normal baseline value for MI associated with percutaneous coronary intervention (PCI-related type 4a MI), or 10 times for MI associated with coronary artery bypass grafting (CABG-related type 5 MI) [1]. Furthermore, the recent 2020 ESC Guidelines for the management of ACS reminds that cTn elevations beyond 5-fold the URL have high ($>90\%$) positive predictive values for acute type 1 MI [4]. Many MI-related studies provided relevant abs $\Delta\text{hs-cTnT}$ results, for various specific contexts; therefore, the question that arises is to what relative variations these results correspond. To answer this, we modeled the abs $\Delta\text{hs-cTnT}$ -to-rel $\Delta\text{hs-cTnT}$ relationship using a quantile nonparametric additive regression model (qgam) [5], in order to provide different corresponding $\Delta\text{hs-cTnT}$ cut-offs.

Materials and Methods

A total of 218,063 hs-cTnT tests were assayed at our laboratory (Pitié Salpêtrière-Charles Foix University Hospital, AP-HP, Paris, France) over a 5.8-year period (April 2012 to January 2018), and extracted from the laboratory information system (GLIMS® software, MIPS-CliniSys, Chertsey-Surrey, UK). Of these, 9,753 were serial tests from patients aged 18-100 years and selected with serial time variations of 1 to 6 hours (Figure 1). Plasma hs-cTnT was assayed using

an electrochemiluminescent immunoassay on two Modular®E170 (Roche, Mannheim, Germany) [6]. Outliers were not removed, but ties were excluded (n=94 repeated pairs of $\text{abs}\Delta\text{hs-cTnT}$ and $\text{rel}\Delta\text{hs-cTnT}$). Nevertheless, given the extremely tail-heavy distributions of $\text{abs}\Delta\text{hs-cTnT}$ and $\text{rel}\Delta\text{hs-cTnT}$, which would make the qgam regression less reliable, both these datasets were transformed using an ordered quantile normalization (OQN), using the ‘orderNorm’ function from the ‘bestNormalize’ R package. This transformation is a rank-based procedure by which the values are mapped to their percentile, which is then mapped to the same percentile of the normal distribution. After the qgam regression, the transformed predicted values were inverted via the ‘predict’ function from the same R package.

The median quantile of the $\text{abs}\Delta\text{hs-cTnT}$ -to- $\text{rel}\Delta\text{hs-cTnT}$ regression was fitted using the ‘qgamV’ function of the ‘mgcViz’ R-package, which allows a graphical visualization of qgam models (‘qgam’ R-package) [7, 8]. The ‘qgamV’ function was programmed with cubic regression spline smooths (‘cr’), and base dimensions chosen high enough ($k=30$) to allow sufficient degrees of freedom. The ‘cqcheck’ function from the ‘qgam’ R package was used to visually check what proportion of $\text{abs}\Delta\text{hs-cTnT}$ falls below the fitted median quantile, named $\hat{P}(\text{abs}\Delta\text{hs-cTnT} < \hat{\mu})$.

The predicted $\text{abs}\Delta\text{hs-cTnT}$ were first determined for different $\text{rel}\Delta\text{hs-cTnT}$ cut-offs, using the ‘predict.gam’ function of the ‘mgcv’ R-package. Secondly, following recommendations [1], the $\text{rel}\Delta\text{hs-cTnT}$ values were categorized as suggestive of MI if greater than $\pm 50\%$ (rise and/or fall) with one hs-cTnT value above the 99PURL, or $\pm 20\%$ with two hs-cTnT values above the 99PURL, considering the well-known Roche 99PURL cut-off of 14 ng/L. Then, in order to test whether $\text{abs}\Delta\text{hs-cTnT}$ may lead to MI proportions similar to $\text{rel}\Delta\text{hs-cTnT}$, we reassessed the categorization by substituting the $\pm 50\%$ and $\pm 20\%$ cut-offs by the corresponding fitted $\text{abs}\Delta\text{hs-cTnT}$ values, for untransformed and OQN-transformed data-based models. A McNemar’s Chi-squared test was used for comparisons of paired proportions, considering a P -value < 0.05 as significant. All the statistics and qgam plots were computed in R (version 4.0.3, R Foundation, Vienna, Austria).

Results

Of the 9,753 $\Delta\text{hs-cTnT}$ (men 70.1%; women 29.9%), the proportions of negative (“*fall*”) and positive (“*rise*”) changes were 42.9 and 57.1%, respectively. A focus of qgam regression curves is depicted on Figure 2, for untransformed (2A) and OQN-transformed data (2B), both displaying a nonlinear profile. Table 1 provides the main corresponding values between the two types of variation with, notably, $\text{abs}\Delta\text{hs-cTnT}$ values of -90.0 , -43.8 , $+19.9$, $+52.0$ ng/L for $\text{rel}\Delta\text{hs-cTnT}$ cut-offs of -50 , -20 , $+20$, $+50\%$, respectively. The qgam checking plot from untransformed data (Figure 2C) shows that the higher the $\text{rel}\Delta\text{hs-cTnT}$, the more the $\hat{P}(\Delta\text{hs-cTnT}(\text{ng/L}) < \hat{\mu})$ deviates from 0.5, especially above 200% where $\hat{P}(\Delta\text{hs-cTnT}(\text{ng/L}) < \hat{\mu})$ is < 0.4 or > 0.6 . This is not the case for the OQN-based qgam checking plot (Figure 2D), which shows $\hat{P}(\Delta\text{hs-cTnT}(\text{ng/L}) < \hat{\mu})$ closer to

0.5 from -50 to 500% rel Δ hs-cTnT, thus demonstrating OQN as an effective way to optimize the qgam regression. Considering the OQN-based qgam, replacing the -50, -20, +20, +50% rel Δ hs-cTnT cut-offs from recommendations by the corresponding predicted abs Δ hs-cTnT (i.e., -90.0, -43.8, +19.9, 52.0 ng/L, Table 1) resulted in similar rates of MI, with only 0.06% diagnostic discrepancies ($P=0.68$), which demonstrates the quality of the qgam regression. Conversely, still according to the OQN-based qgam, a 70 ng/L abs Δ hs-cTnT cut-off (i.e., 5-fold the 99PURL, known to have >90% positive predictive value for acute type 1 MI [4]) would correspond to a 70.9% rel Δ hs-cTnT cut-off. In the same way, a 140 ng/L abs Δ hs-cTnT (e.g., 10 times the 99PURL cut-off used for CABG-related type 5 MI diagnosis [1]), would correspond to a 132.0% rel Δ hs-cTnT cut-off. Considering now the untransformed data-based qgam, the same 70 and 140 ng/L cut-offs correspond to 74.8 and 151.2% rel Δ TnT cut-offs respectively. Classifying patients above these corresponding rel Δ hs-cTnT cut-offs as positive, the theoretical proportions of MI were significantly different between the untransformed and OQN-transformed data based qgams (+0.53 and +1.07%, respectively; $P < 0.0001$). Given the most reliable checking profile (Figure 2D compared to 2C), these minor -but significant- differences in theoretical MI proportions suggest the OQN-based qgam as most efficient for the prediction of high abs Δ hs-cTnT cut-offs.

Discussion

Few studies have compared the absolute vs. relative changes in hs-cTnT from the standpoint of diagnostic cut-offs. Among these, Reichlin et al. found an area under receiver operating characteristic curve (AUROC) of 0.95 (abs Δ hs-cTnT) vs. 0.76 (rel Δ hs-cTnT) for the diagnosis of AMI, with an AUROC-derived cut-off for a 2-hour abs Δ hs-cTnT of 7 ng/L [2]. Mueller et al. found an AUROC of 0.90 (abs Δ hs-cTnT) vs. 0.75 with a ROC-optimized rise or fall of 9.2 ng/L to rule out NSTEMI in patients admitted at the emergency room (ER), and of 6.9 ng/L to rule out NSTEMI in ACS patients [3]. Irfan et al. also confirmed the higher accuracy of abs Δ hs-cTnT over rel Δ hs-cTnT for AMI diagnosis, either 1 hour (AUROC 0.93 vs. 0.67) or 2 hours after being admitted in the ER (AUROC 0.95 vs. 0.75), with ROC-optimized absolute cut-offs of 5 and 7 ng/L, respectively [9]. Furthermore, Biener et al. calculated a sensitivity of 95.6% and specificity of 57.4% for NSTEMI diagnosis for a rising abs Δ hs-cTnT ROC-optimized cut-off of 8.8 ng/L, but a lower sensitivity (82.2%) and specificity (56.8%) for a 20% rel Δ hs-cTnT. They therefore recommended to consider absolute rather than relative hs-cTnT changes -and rising rather than falling- to diagnose NSTEMI [10]. All these studies highlight the higher accuracy of abs Δ hs-cTnT, as specified in the 4UDMI [1]. One must, however, remember the main drawback of using the absolute change: the fact that it is assay dependent. This makes our study of interest since, to the best of our knowledge, the correspondence between absolute and relative change has never been specifically assessed for the Roche hs-cTnT. This work provides a new approach, the qgam

regression, optimized through an ordered-quantile transformation to take account for tail-heavy distributions so that the spline basis functions are evenly distributed across rel Δ hs-cTnT (i.e., to concentrate the spline basis functions where there was more data). This method could serve for further studies aiming at modelling relationships between biomarkers with highly skewed distributions, and/or for which outlier removal should be avoided.

Acknowledgements: The authors are grateful to Vincent Fitzpatrick for his English proofreading.

References

1. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, et al.; Executive Group on behalf of the Joint European Society of Cardiology (ESC)/American College of Cardiology (ACC)/American Heart Association (AHA)/World Heart Federation (WHF) Task Force for the Universal Definition of Myocardial Infarction. Fourth Universal Definition of Myocardial Infarction (2018). *J Am Coll Cardiol* 2018;72:2231-64. (PMID: 30153967)
2. Reichlin T, Irfan A, Twerenbold R, Reiter M, Hochholzer W, Burkhalter H, et al. Utility of absolute and relative changes in cardiac troponin concentrations in the early diagnosis of acute myocardial infarction. *Circulation* 2011;124:136-45. (PMID: 21709058)
3. Mueller M, Biener M, Vafaie M, Doerr S, Keller T, Blankenberg S, et al. Absolute and relative kinetic changes of high-sensitivity cardiac troponin T in acute coronary syndrome and in patients with increased troponin in the absence of acute coronary syndrome. *Clin Chem* 2012;58:209-18. (PMID: 22134520)
4. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al; ESC Scientific Document Group. 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J* 2020;42:1289-1367. (PMID: 32860058)
5. Monneret D, Fasiolo M, Bonnefont-Rousselot D. Relationships between renal function variations and relative changes in cardiac troponin T concentrations based on quantile generalized additive models (qgam). *Clin Chem Lab Med* 2021; 59:1115-25. (PMID: 33915606)
6. Monneret D, Gellerstedt M, Bonnefont-Rousselot D. Determination of age- and sex-specific 99th percentiles for high-sensitive troponin T from patients: an analytical imprecision- and partitioning-based approach. *Clin Chem Lab Med* 2018;56:685-96. (PMID: 29176015)
7. Fasiolo M, Wood SN, Zaffran M, Nedellec R, Goude Y. qgam: Bayesian non-parametric quantile regression modelling in R (2020). Available on <https://arxiv.org/pdf/2007.03303.pdf> (access April, 2021).
8. Fasiolo M, Wood SN, Zaffran M, Nedellec R, Goude Y. Fast calibrated additive quantile regression (2020). Available on <https://arxiv.org/pdf/1707.03307.pdf> (access April, 2021).
9. Irfan A, Reichlin T, Twerenbold R, Meister M, Moehring B, Wildi K, et al. Early diagnosis of myocardial infarction using absolute and relative changes in cardiac troponin concentrations. *Am J Med* 2013;126:781-8. (PMID: 23870791)

10. Biener M, Mueller M, Vafaie M, Jaffe AS, Widera C, Katus HA, et al. Diagnostic performance of rising, falling, or rising and falling kinetic changes of high-sensitivity cardiac troponin T in an unselected emergency department population. *Eur Heart J Acute Cardiovasc Care* 2013;2:314-22. (PMID: 24338290)

Table 1. Corresponding cut-offs between absolute and relative $\Delta\text{hs-cTnT}$

No transformation			Ordered-quantile normalization		
Relative $\Delta\text{hs-cTnT}$ (%)	Absolute $\Delta\text{hs-cTnT}$ (ng/L)	(-ACL ; + ACL)	Relative $\Delta\text{hs-cTnT}$ (%)	Absolute $\Delta\text{hs-cTnT}$ (ng/L)	(-ACL ; + ACL)
-50	-93.7	(-100 ; -87.4)	-50	-90.0	(-96.3 ; -83.7)
-20	-36.8	(-42.6 ; -31.0)	-20	-43.8	(-49.6 ; -38.0)
-10	-17.9	(-23.0 ; -12.9)	-10	-14.0	(-19.1 ; -8.9)
-5	-9.0	(-13.7 ; -4.3)	-5	-8.1	(-13.0 ; -3.1)
0	-0.7	(-5.1 ; 3.8)	0	0	(-4.5 ; 4.5)
5	6.8	(2.6 ; 11.0)	5	6.1	(2.1 ; 10.1)
10	13.2	(9.1 ; 17.3)	10	10.6	(6.6 ; 14.6)
20	23.3	(19.2 ; 27.3)	20	19.9	(15.9 ; 23.9)
50	47.9	(43.5 ; 52.2)	50	52.0	(47.6 ; 56.5)
100	95.8	(90.2 ; 102)	100	105	(99.7 ; 111)
200	167	(160 ; 174)	200	179	(172 ; 185)
500	235	(227 ; 243)	500	271	(262 ; 280)
1,000	361	(351 ; 371)	1,000	450	(437 ; 463)
10,000	2,323	(2,269 ; 2,377)	10,000	2,050	(2,001 ; 2,098)

Legends

Figure 1.

Flowchart of the data selection process. Abbreviations: Δ hs-cTnT: high-sensitive cardiac troponin T variation; LIS: laboratory information system.

Figure 2.

Qgam plots of the absolute Δ hs-cTnT $=f$ (relative Δ hs-cTnT) regression determined using untransformed data (A) and ordered-quantile-normalized data (B) ($n=9,753$). The solid black curve and its gray area represent the predicted values and its two-standard deviations. The dashed black curves represent the predicted analytical change limit. The concentric gray contour lines represent the ten-by-ten percentiles of the nonparametric kernel density estimation. The qgam checking plots represent the proportion of absolute Δ hs-cTnT that falls below the fitted quantile ($\hat{P}(\Delta$ hs-cTnT(ng/L) $<\hat{\mu}$) for untransformed data (C) and ordered-quantile-normalized data (D), knowing that roughly 50% of absolute Δ hs-cTnT values would be expected to fall below the fitted median quantile. The horizontal black dashed line represents the median quantile, the dots are the $\hat{P}(\Delta$ hs-cTnT(%) $<\hat{\mu}$ for each bin of relative Δ hs-cTnT, and the black crosses are the 95% confidence intervals for median quantile. If the dots fall outside the confidence intervals, then $\hat{P}(\Delta$ hs-cTnT(%) $<\hat{\mu}$ might be deviating too much from the median quantile. Overall, the deviations from the theoretical median quantile are more reduced for normalized data-based qgam than for untransformed data-based ones. Abbreviations: $\hat{P}(\Delta$ hs-cTnT(ng/L) $<\hat{\mu}$): proportion of absolute Δ hs-cTnT that falls below the fitted median quantile; Δ hs-cTnT: high-sensitive cardiac troponin T variation; qgam: quantile generalized additive model.