



**HAL**  
open science

## **M-CSF-mediated macrophage development is dominantly inhibited by NOD2 signaling for replenishment of immunogenic dendritic cells**

Camille Chauvin, Daniel Alvarez-Simon, Paul Régnier, Katarina Radulovic, Olivier Boulard, Myriam Delacre, Nadine Waldschmitt, Laurent Peyrin-Biroulet, Jérôme Kluza, Guillaume Darrasse-Jèze, et al.

### ► To cite this version:

Camille Chauvin, Daniel Alvarez-Simon, Paul Régnier, Katarina Radulovic, Olivier Boulard, et al.. M-CSF-mediated macrophage development is dominantly inhibited by NOD2 signaling for replenishment of immunogenic dendritic cells. 6th European Congress of Immunology, Sep 2021, Belgrade, Serbia. 51, pp.307, P-0672. 10.1002/eji.202170200 . hal-03870668

**HAL Id: hal-03870668**

**<https://cnrs.hal.science/hal-03870668>**

Submitted on 24 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## POSTER PRESENTATIONS

## P-0672

**M-CSF-mediated macrophage development is dominantly inhibited by NOD2 signaling for replenishment of immunogenic dendritic cells**

Camille Chauvin<sup>1</sup>, Daniel Alvarez Simon<sup>2</sup>, Paul Régnier<sup>3</sup>, Katarina Radulovic<sup>4</sup>, Olivier Boulard<sup>5</sup>, Myriam Delacré<sup>6</sup>, Nadine Waldschmitt<sup>6</sup>, Laurent Peyrin Biroulet<sup>7</sup>, Jérôme Kluzza<sup>8</sup>, Guillaume Darrasse Jèze<sup>9</sup>, Mathias Chamaillard<sup>5</sup>, **Lionel Franz Poulin<sup>5</sup>**

<sup>1</sup>Institut Pasteur, Paris 75015, France

<sup>2</sup>Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019, F-59000 Lille

<sup>3</sup>Immunology-Immunopathology-Immunotherapy (i3) Laboratory, INSERM UMR-S 959, Sorbonne Université, 75005 Paris, France; Biotherapy Unit (CIC-BTI), Inflammation-Immunopathology-Biotherapy Department (DHU i2B), Groupe Hospitalier Pitié-Salpêtrière, Assistance Publique-Hôpitaux de Paris (AP-HP), 75013 Paris, France

<sup>4</sup>Unité de Recherche Clinique, Centre Hospitalier de Valenciennes, 59322 Valenciennes CEDEX, France

<sup>5</sup>Univ. Lille, Inserm, U1003, F-59019 Lille

<sup>6</sup>Chair of Nutrition and Immunology, School of Life Sciences, Technische Universität München, Freising-Weihenstephan, Germany

<sup>7</sup>Department of Gastroenterology, Nancy University Hospital, University of Lorraine, Vandœuvre-lès-Nancy, France

<sup>8</sup>University Lille, UMR9020-U1277 - CANTHER - Cancer Heterogeneity Plasticity and Resistance to Therapies, F-59000 Lille, France

<sup>9</sup>Immunology-Immunopathology-Immunotherapy (i3) Laboratory, INSERM UMR-S 959, Sorbonne Université, 75005 Paris, France; Université de Paris, Paris Descartes, Faculté de Médecine, Paris, France

Despite recent advances, it remains unclear whether monocyte-derived dendritic cells (moDCs) represent alternative context-dependent fate in the gut. We found here that Nod2-dependent sensing of bacteria lowers the ability of circulating monocytes to respond to M-CSF for generating moDCs. Such inhibitory effect on monocyte-to-macrophage transition was prevented upon blockade of TNF- $\alpha$ . Recognition of the gut microbiota by Nod2 was sufficient to promote the expansion of moDCs within the colonic mucosa. A competitive bone marrow transplant model further demonstrated that Nod2 promotes the conversion of monocytes into dendritic cells. Equally of importance, tumours with the highest transcript levels of NOD2 were associated with a favorable prognosis and characterized by an enrichment of a gene signature related to moDCs. This study implicates that Nod2-dependent sensing of the gut microbiota influences monocytic lineage commitment into dendritic cells, which sets the stage for future investigations to achieve accurate outcome prediction in colorectal cancer.

**Keywords:** Immune development, macrophage, dendritic cells, *in vivo* tumor models

## P-0673

**Ex situ heart perfusion and standard of care cold storage differentially affect the ischemic secretome of donor hearts in perfusates but not the reperfusion response in recipient plasma**

**Evgeny Chichelnitskiy<sup>1</sup>**, Bettina Wiegmann<sup>2</sup>, Lena Radomsky<sup>1</sup>, Nadine Ledwoch<sup>1</sup>, Fabio Ius<sup>2</sup>, Franziska Wandrer<sup>1</sup>, Jenny Kühne<sup>1</sup>, Kerstin Beushausen<sup>1</sup>, Jana Keil<sup>1</sup>, Sebastian Rojas Hernandez<sup>2</sup>, Wiebke Sommer<sup>2</sup>, Christian Kühn<sup>2</sup>, Igor Tudorache<sup>2</sup>, Murat Avsar<sup>2</sup>, Axel Haverich<sup>2</sup>, Gregor Warnecke<sup>4</sup>, Christine S. Falk<sup>3</sup>

<sup>1</sup>Institute of Transplant Immunology, Hannover Medical School, Germany

<sup>2</sup>Department of Cardiothoracic, Transplantation and Vascular Surgery, Hannover Medical School, Germany

<sup>3</sup>Institute of Transplant Immunology, Hannover Medical School, Germany, German Centre for Infection Research, DZIF, TTU-IICH Hannover-Braunschweig

<sup>4</sup>Clinic for heart surgery, University Hospital Heidelberg, Germany

Allograft preservation procedures may influence the donor organ status and in turn affect heart transplant (HTx) recipient. Here we aimed to compare the secretomes in recipient plasma and perfusates in patients whose hearts were either preserved using *ex situ* heart perfusion (ESH) or standard of care (SOC) cold static preservation in order to identify potential biomarker candidates for heart preservation. Using multiplex techniques, we measured 50 cytokines/chemokines in recipient plasma before (pre), after (T0), 24h and 3 weeks after HTx. Unsupervised cluster analyses identified top-10 plasma cytokines and chemokines clearly separating T0 from other time points after HTx and reflecting a reperfusion injury-specific pattern. Surprisingly, ESH or SOC heart preservation did not have a significant impact on these inflammatory plasma profiles at T0, T24 or 3 weeks. The two strongest discriminators separating T0 from other time points i.e. IFN- $\gamma$ , SCGF- $\beta$  were detected in both ESH and SOC recipients at comparable concentrations. In contrast, the preservation method clearly affected the cytokine/chemokine profile in perfusates highlighted by higher concentrations of pro- (IFN- $\gamma$ , CXCL10) and anti-inflammatory (IL-10, IL-1RA) mediators in ESH compared to SOC samples. Although ESH or SOC preservation did not affect the reperfusion response in plasma at T0 after HTx, normothermic oxygenated preservation of donor hearts was accompanied by secretion of pro- and anti-inflammatory cytokines, chemokines that may affect long-term functionality and longevity of the graft. With a better understanding of molecular changes during ESH, we expect to identify biomarker candidates for improved organ function pre HTx.

**Keywords:** Biomarkers, chemokines, cytokines and mediators, transplantation

## P-0676

**The role of NK cells in the resolution of malaria-associated acute respiratory distress syndrome**

**Emilie Pollenus<sup>1</sup>**, Thao Thy Pham<sup>1</sup>, Hendrik Possemiers<sup>1</sup>, Leen Vandermosten<sup>1</sup>, Sofie Knoops<sup>1</sup>, Patrick Matthys<sup>2</sup>, Ghislain Opendakker<sup>2</sup>, Philippe Van Den Steen<sup>1</sup>

<sup>1</sup>Laboratory of Immunoparasitology, Department of Microbiology, Immunology and Transplantation, Rega Institute for Medical Research, KU Leuven, University of Leuven, Belgium

<sup>2</sup>Laboratory of Immunobiology, Department of Microbiology, Immunology and Transplantation, Rega Institute for Medical Research, KU Leuven, University of Leuven, Belgium

Malaria is a global health disease with >400 000 deaths each year, caused by complications such as malaria-associated acute respiratory distress syndrome (MA-ARDS). Despite efficient parasite killing with antimalarial drugs, 15% of patients with complications still die. This shows the need to study resolution and wound healing mechanisms involved in the recovery from these complications. Disease resolution is coordinated by several leukocyte subtypes. Here, we investigate the role of NK cells in the disease resolution of MA-ARDS. C57BL/6 mice were infected with *Plasmodium berghei* NK65 (*PbNK65*), resulting in the development of MA-ARDS on day 8 post infection. On this day, antimalarial treatment with artesunate and chloroquine was started for 5 days. To study the role of NK cells, NK cells were depleted by injection of anti-NK1.1. Depletion of NK cells did not affect the development of MA-ARDS, but resulted in a decreased survival during resolution. In particular, only 50% of the anti-NK1.1-treated mice recovered from MA-ARDS upon anti-malarial treatment, compared to >80% without depletion. Interestingly, the resolution of alveolar edema occurred as efficiently in the NK cell-depleted mice that could be rescued compared to the non-depleted mice. The treatment of *PbNK65*-infected C57BL/6 mice with antimalarial drugs serves as a good model to study the resolution of MA-ARDS. NK cells are not involved during the development of MA-ARDS, but are critical during the resolution process upon anti-malarial treatment.

**Keywords:** Innate host defence, NK cells, parasite infections, tissue damage and repair