



The triplet state of deprotonated mesoionic carbene Breslow intermediates is thermally accessible: Distal difunctionalization of aldehydes

Zengyu Zhang, Shiqing Huang, Chen-Yuan Li, Liang-Liang Zhao, Wei Liu, Mohand Melaimi, Guy Bertrand, Xiaoyu Yan

► To cite this version:

Zengyu Zhang, Shiqing Huang, Chen-Yuan Li, Liang-Liang Zhao, Wei Liu, et al.. The triplet state of deprotonated mesoionic carbene Breslow intermediates is thermally accessible: Distal difunctionalization of aldehydes. *Chem Catalysis*, 2022, 2 (12), pp.3517-3527. 10.1016/j.checat.2022.09.047 . hal-03872960

HAL Id: hal-03872960

<https://cnrs.hal.science/hal-03872960>

Submitted on 27 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The triplet state of deprotonated mesoionic carbene Breslow intermediates is thermally accessible: Distal difunctionalization of aldehydes

Zengyu Zhang,¹ Shiqing Huang,¹ Chen-Yuan Li,¹ Liang-Liang Zhao,¹ Wei Liu,¹ Mohand Melaimi,² Guy Bertrand,^{2*} and Xiaoyu Yan^{1*}

¹Key Laboratory of Advanced Light Conversion Materials and Biophotonics, Department of Chemistry, Renmin University of China, Beijing 100872, China.

²UCSD-CNR Joint Research Laboratory (IRL3555), Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA 92093-0358, United States.

★

Correspondence: gbertrand@ucsd.edu

★Correspondence: yanxy@ruc.edu.cn

INTRODUCTION

N-heterocyclic carbenes (NHCs), especially thiazol-2-ylidenes **A**¹ and 1,2,4-triazol-5-ylidenes **B**,² have long been known as organocatalysts for aldehyde transformations, such as the benzoin condensation, Stetter reaction, cycloadditions, Claisen rearrangement, etc.^{3, 4, 5, 6, 7} (Fig 1a). These reactions occur via electron-pair-transfer processes through the so-called Breslow Intermediates (**BI**s).^{8,9,10,11,12} The role of the **BI** in single electron transfer (SET) pathways was also established a long time ago in biosynthesis,¹³ and expanded in the last decade to NHC-organocatalysis.^{14, 15, 16, 17} Among the important differences between the electron-pair-transfer and the SET pathways is that the first relies on the **BI**, whereas the second involves the deprotonated **BI** (**BI**⁻).^{18,19,20,21,22,23,24} Also noteworthy is the recent development of NHC/photoredox dual catalysis²⁵ involving acyl azolium salts **BI**⁺,^{26,27,28,29,30,31,32,33,34,35,36} as well as few reports^{37,38,39} dealing with their excited state (**BI**^{+*}), both of which being the doubly oxidized deprotonated **BI** (Fig 1b). Obviously, the latter processes are based on the oxidizing power of the **BI**⁺ and the **BI**^{+*}, whereas

NHC-organocatalysis via SET reaction depends on the reducing ability of **BI**[•] in the ground state. Herein we report that the excited triplet state of **BI**[•] (**BI**^{•*}) derived from 1,2,3-triazol-5-ylidenes (mesoionic carbenes, MIC) **D**^{40,41} is thermally accessible, allowing for the catalytic distal difunctionalization of aryl aldehydes. At the formyl side, an oxidative esterification reaction takes place, concomitant with reduction of the aryl side. Here, three types of chemical transformation are demonstrated, namely hydro(deutero)defunctionalization, hydrogenation and reductive alkylation. (Fig. 1c).




Figure 1 A) Carbenes considered in this study; B) Breslow Intermediate (**BI**), deprotonated Breslow Intermediate (**BI**[•]), and acyl azolium intermediate in the ground state (**BI**⁺) and the excited state (**BI**^{•*}); C) mesoionic carbene (**D**)-catalyzed distal difunctionalization of aryl aldehydes via the excited state of deprotonated Breslow Intermediate (**(D)BI**^{•*}).

RESULTS AND DISCUSSION

To gain further insight into the electronic properties of **BIs** and **BI**[•]s derived from **A**, **B**, **C**⁴² and MICs **D**, we performed density functional theory (DFT) calculations at the PWPB95-D3/def2QZVPP//B3LYP-D3(BJ)/6-31G** + SMD (THF) level of theory. The adiabatic singlet-triplet gaps for **BIs** derived from classical NHCs **A-C** are greater

than 37.0 kcal/mol, whereas for the mesoionic carbene (**D**)**BI**,⁴³ it is only 23.7 kcal/mol (Fig. 2). The same trend is observed for the singlet-triplet gap of the corresponding deprotonated **BIs** (**BI**⁻s). They vary from 26.1 to 28.7 kcal/mol for NHCs **A-C**, but the gap is strikingly as low as 4.0 kcal/mol for the mesoionic analogue (**D**)**BI**⁻. The small S/T gap for (**D**)**BI** and (**D**)**BI**⁻ is due to the polarized exocyclic CC bond,⁴³ which in the singlet state leads to a repulsion between the negatively charged carbon and adjacent oxygen. The SOMO of (**D**)**BI**⁻ has contributions from the π orbital of the triazole ring, the carbonyl and, interestingly, the phenyl ring. The SOMO-1 is mainly located on the π^* orbital of the triazole ring and on the carbonyl group. From these computational results, we were confident that the very low singlet-triplet gap predicted for (**D**)**BI**⁻ would allow access to the triplet form (**D**)**BI**⁻* without photoexcitation. Moreover, since the SOMO is at least partially located on the phenyl group of the aldehyde moiety, it implied that the ring has a radical anion character. Thus, we envisioned that if a leaving group is present on the aryl ring, it could be eliminated, affording a biradical (**D**)**BR**⁻*. DFT calculations predict that such compounds would have a singlet ground state 14.5 kcal/mol lower in energy than the triplet and should therefore be regarded as the zwitterionic compound (**D**)**ZW**, which should allow for distal difunctionalization of aryl aldehydes.




Figure 2 A) Singlet/triplet gap of **BI**s and **BI⁻**s derived from carbenes **A-D**; B) SOMO and SOMO-1 of **(D)BI***, isovalue is 0.05; C) Postulated reaction sequence leading to the formation of the zwitterionic compound **(D)ZW**, allowing for difunctionalization of aryl aldehydes.

We capitalized on the potent leaving group ability of sulfonyl groups through single-electron reduction ⁴⁴ to check our hypothesis. We first reacted 4-sulfonylbenzaldehyde **1a** with methanol in the presence of 10 mol% of a 1,2,3-triazolium salt as precatalyst, using a base to generate *in situ* the corresponding mesoionic carbene **D** and to deprotonate the ensuing Breslow Intermediate. With 4,5-unsubstituted triazolium **Cat_a** and **Cat_b**, methyl benzoate **2aa** was formed in 42 and 38% yield, respectively (Figure 3, entries 1-2). The leaving group was quantitatively recovered as benzenesulfinic acid (PhSO₂H). The yield in **2aa** significantly increased to 65 and 74% by using ester substituted triazolium salts **Cat_c** and **Cat_d**, (entries 3-4), which have proven to be the best catalysts for the previously reported MIC-catalyzed arylacylation of alkynes via SET.⁴⁵ Classical-NHCs were less efficient; only very low yields were obtained (entries 5-7). Base and solvent screening showed *t*-BuOK and *t*-BuOMe (MTBE) to be the best choices (entries 8-12). Reducing the amounts of MeOH or decreasing the reaction temperature gave lower yields (entries 13-15). A control experiment in the dark gave the same yield, which revealed that this reaction did not proceed via photoexcitation (entries 16). Note that the transformation of the 4-sulfonylbenzaldehyde **1a** into methyl benzoate **2aa**, involved a reductive desulfonylation at the aryl group and an oxidative esterification at formyl carbon, and is therefore overall redox-neutral.




Figure 3 Optimization of the reductive desulfonylation and oxidative esterification.


^aReaction carried out in 0.3 mmol scale. Yields were determined by ¹H spectroscopy with CH₂Br₂ as internal standard. Isolated yield is given in parentheses. ^b1.0 equiv

MeOH was used. ^c3.0 equiv MeOH was used. ^dTemperature was changed from -15 °C to 0 °C. ^eThe reaction was performed in dark.


The scope of the MIC-catalyzed hydrodefunctionalization reaction was investigated under the optimized conditions. Treatment of **1a** with CH₃OD and CD₃OD gave deuterated products **2aa-D₁** and **2aa-D₄**, respectively. Note that the only partial deuteration at the phenyl ring is due to the presence of the formyl-H and MIC⁺-H, which are exchangeable. Different nucleophiles were next employed. The yields decreased as the chain length and steric hindrance of the alcohols increase. Primary alcohols gave benzoate **2aa-2ae** in 30-71% yields. For secondary alcohols, only cyclopropanol gave the corresponding benzoate **2af** in 37% yield, while other alcohols gave trace products. When ethanethiol was employed with ethanol as proton source, *S*-ethyl benzothioate **2ag** could be isolated in 47% yield. We extended our investigations to sulfonyl-substituted aryl aldehydes **1b-1e**. When bromo, trifluoromethyl and methyl substituents were embedded on 4-sulfonylbenzaldehyde, using CH₃OD as deuterium source, the corresponding deuterodesulfonylation products **2b-D₁**–**2d-D₁** were isolated in 43-80% yields with deuteration levels of 50-53% at the 4-position. 5-Sulfonyl-2-thiophenecarbaldehyde **1e** gave deuterated product **2e-D₁** in 73% isolated yield. Other sulfonyl groups, such as methylsulfonyl (**1f**) and aminosulfonyl (**1g**) were converted less efficiently, giving benzoate **2aa** in 30 and 42% yield, respectively. Interestingly, the sulfonyl group can be placed in a more distal position, as shown by the desulfonylation of **1h**, **1i** and **1j** which gave **2h** and **2i** in 27, 38 and 25 % yield, respectively.

The scope of the MIC-catalyzed hydrodefunctionalization and esterification of aryl aldehydes is not restricted to the sulfonyl functionality. Esters can also be used as the leaving group, as shown by the formation of the toluoate **2i** in 21% yield. Fluoride is also an attractive leaving group, since the reductive defluorination of trifluoromethylarenes is a versatile strategy for the construction of difluoromethylarenes.⁴⁶ When 4-trifluoromethyl substituted aryl aldehydes **1l** were employed, the corresponding 4-difluoromethylbenzoates **2l** was formed in 42%


isolated yields. The 4-deuterodifluoromethylbenzoate **2l-D₁** was obtained in 40% yield and 60% deuteration level, using CH₃OD as deuterium source. When the trifluoromethyl group was placed in the *ortho*-position, the corresponding 2-difluoromethylbenzoate **2m** was obtained in 22% yield. Interestingly, when two trifluoromethyl groups were placed on the *ortho*- and *para*-positions, reductive defluorination selectively occurred on the *ortho*-position, giving 2-difluoromethyl-4-trifluoromethylbenzoate **2n** in 38% yield. Using CH₃OD as deuterium source, the deuterated analogue **2n-D₁** was isolated in 35% yield with 70% deuteration level.




Desulfonation with different NuH/NuD




Desulfonation from different aldehydes



Desulfonation from different sulfonyl leaving group



Desulfonation from more distal position



Different Leaving Groups:




Figure 4 Scope of the MIC-catalyzed hydrodefunctionalization and esterification of aryl aldehydes. Reactions were carried out in 0.3 mmol scale. ^b3 equiv. EtOH, 2.5 equiv ^tBuOK, and 1.0 equiv EtSH were used. Isolated yields are given. ^cTemperature was changed from -15 °C to 0 °C. ^d**Cat**_c (20 mol%) was used as catalyst, 90 °C for 5 h. ^e**Cat**_c (20 mol%) was used as catalyst, 50 °C for 5 h.

We were curious to test whether the MIC-catalyzed distal difunctionalization of aryl aldehydes could be extended to other types of reduction processes. We first chose the hydrogenation reaction of azo compounds (Fig. 5). Under the optimized conditions, in the presence of methanol as the NuH agent, 4-azobenzaldehyde **3a** was converted into the corresponding hydrazine **4a** in 90% yield. By using CD₃OD as deuterium source, the deuterated product **4a-D₃** was obtained in 87% yield with a high deuteration level of 95%. Ethanol can also act as the NuH, affording **4b**, which was isolated in 67% yield. Azo derivatives with different substituents **3c-3f** also gave hydrazine derivatives **4c-f** in 47-76% yields. The azo group can also be placed in *meta*- and even a more distal position, giving the target products **4g-4h** in 50-61% yields. Note that these results allow to rule out a two-electron pathway via *p*- or *o*-quinodimethane.⁴⁷ These hydrogenation reactions can also be extended to other unsaturated substrates. When 4-benzoylbenzaldehyde **5** was used, the benzoyl group was reduced to the α-hydroxybenzyl group while the formyl group was oxidized to an ester, giving **6** which was isolated in 40% yield. A dearomatization reaction was also observed for 9-anthraldehyde **7**, giving 9,10-dihydroanthracene-9-carboxylate **8** in 32% yield.^{48,49}




Figure 5 MIC-catalyzed hydrogenation reactions. Reactions carried out in 0.3 mmol scale. Isolated yields are given. ^aCD₃OD was used. ^bEtOH was used.

Lastly, we wanted to demonstrate that the process could allow for C-C bond formation, therefore adding complexity to aryl aldehydes (Fig. 6). To this end, we reacted 4-iodobenzaldehyde **9** under the optimized catalytic conditions with alkenes **10** in the presence of methanol. Gratifyingly, 2-vinylpyridine derivatives (**10a-c**), 2-vinylbenzothiazole (**10d**), and even methyl methacrylate (**10e**) reacted smoothly with 4-iodobenzaldehyde **9**, generating the desired products **11a-e** in 37-55% yields. Note

that in these cases, the second inversion of state could occur either just after the cleavage of the carbon-iodine bond, or after the addition to the alkene.




Figure 6 C-C bond formation, adding complexity to aryl aldehydes.

CONCLUSIONS

Photochemical reactions involving direct excitation of a substrate to its triplet state are well explored.⁵⁰ Recently, this area experienced a renaissance by exploiting a photosensitizer as a catalyst for energy transfer.^{51,52,53} This work shows that the combination of aryl aldehydes with carbenes leads to low S/T gap adducts (Breslow Intermediates and their deprotonated forms) due to the radical-stabilization effect of carbenes.⁵⁴ The S/T gap is especially small with mesoionic carbenes, which results in their low activation energy and viability of thermal excitation. The resulting triplet state of deprotonated Breslow Intermediates enables tandem distal functionalization of aryl aldehydes. The reactivity of these biradicals is significantly different from that of the aforementioned photo-excited triplet substrates, which have a high activation energy, high reactivity and limited reaction patterns (Fig. 7). Synthetic strategies, based on the thermal accessibility of triplet states of other carbene-derived intermediates, such as azolium enolates, dienolates and homoenolates, are under current investigation.




Figure 7 Carbenes, especially MICs, lead to adducts featuring low-lying triplet states.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Nos. 22171284 and 21602249), the Public Computing Cloud Platform, Renmin University of China, and the NSF (CHE-2153475)

Author contributions

X.Y. and G.B. conceived and designed the project and wrote the manuscript with the help of M.M., Z.Z., C.-Y.L., L.-L.Z., and W.L. conducted the experiments and analyzed the data. S.H. carried out the computational analyses. All authors provided comments on the experiments and the manuscript during its preparation.

Declaration of interest

The authors declare no competing interests.

- ¹ Arduengo III, A. J., Goerlich, J. R., and Marshall, W. J. (1997). A Stable Thiazol-2-ylidene and Its Dimer. *Liebigs Ann.* 365-374.
- ² Enders, D., Breuer, K., Raabe, G., Runsink, J., Teles, J. H., Melder, J. -P., Ebel, K., and Brode, S. (1995). Preparation, Structure, and Reactivity of 1,3,4-Triphenyl-4,5-dihydro-1*H*-1,2,4-triazol-5-ylidene, a New Stable Carbene. *Angew. Chem. Int. Ed.* 34, 1021–1023.
- ³ Bellotti, P., Koy, M., Hopkinson, M. N., and Glorius, F. (2021). Recent advances in the chemistry and applications of N-heterocyclic carbenes. *Nat. Rev. Chem.* 5, 711–725.

-
- ⁴ Pareek, M., Yernaidu, S., and Sunoj, R. B. (2021). Tale of the Breslow intermediate, a central player in N-heterocyclic carbene organocatalysis: then and now. *Chem. Sci.* *12*, 7973-7992.
- ⁵ Biju, A. T. (2019). *N-Heterocyclic Carbenes in Organocatalysis*. Wiley-VCH Verlag GmbH & Co. KGaA Weinheim, Germany.
- ⁶ Bugaut, X., and Glorius, F. (2012). Organocatalytic umpolung: N-heterocyclic carbenes and beyond. *Chem. Soc. Rev.* *41*, 3511–3522.
- ⁷ Flanigan, D. M.; Romanov-Michailidis, F., White, N. A., and Rovis, T. (2015). Organocatalytic Reactions Enabled by N-Heterocyclic Carbenes. *Chem. Rev.* *115*, 9307–9387.
- ⁸ Breslow, R. (1958). On the Mechanism of Thiamine Action. IV. Evidence from Studies on Model Systems. *J. Am. Chem. Soc.* *80*, 3719-3726.
- ⁹ Berkessel, A., Elfert, S., Etzenbach-Effers, K., and Teles, J. H. (2010). Aldehyde Umpolung by N-Heterocyclic Carbenes: NMR Characterization of the Breslow Intermediate in its Keto Form, and a Spiro-Dioxolane as the Resting State of the Catalytic System. *Angew. Chem. Int. Ed.* *49*, 7120 –7124.
- ¹⁰ Berkessel, A., Yatham, V. R., Elfert, S., and Neudörfl, J. -M. (2013). Characterization of the Key Intermediates of Carbene-Catalyzed Umpolung by NMR Spectroscopy and X-Ray Diffraction: Breslow Intermediates, Homoenolates, and Azolium Enolates. *Angew. Chem. Int. Ed.* *52*, 11158 –11162.
- ¹¹ Paul, M., Neudörfl, J. -M., and Berkessel, A. (2019). Breslow Intermediates from a Thiazolin-2-ylidene and Fluorinated Aldehydes: XRD and Solution-Phase NMR Spectroscopic Characterization. *Angew. Chem. Int. Ed.* *58*, 10596 –10600.
- ¹² Wessels, A., Klusmann, M., Breugst, M., Schlörer, N. E., and Berkessel, A. (2022). Formation of Breslow Intermediates from N-Heterocyclic Carbenes and Aldehydes Involves Autocatalysis by the Breslow Intermediate, and a Hemiacetal. *Angew. Chem. Int. Ed.* *61*, e202117682
- ¹³ Ragsdale, S.W. (2003). Pyruvate ferredoxin oxidoreductase and its radical intermediate. *Chem. Rev.* *103*, 2333-2346.

-
- ¹⁴ Guin, J., De Sarkar, S., Grimme, S., and Studer, A. (2008). Biomimetic Carbene-Catalyzed Oxidations of Aldehydes Using TEMPO. *Angew. Chem. Int. Ed.* **47**, 8727-8730.
- ¹⁵ Bay, A. V., and Scheidt, K. A. (2022). Single-electron carbene catalysis in redox processes. *Trends Chem.* **4**, 277-290.
- ¹⁶ Ishii, T., Nagao, K., and Ohmiya, H. (2020). Recent advances in N-heterocyclic carbene-based radical catalysis. *Chem. Sci.* **11**, 5630-5636.
- ¹⁷ Dai, L., and Ye, S. (2021). Recent advances in N-heterocyclic carbene-catalyzed radical reactions. *Chin. Chem. Lett.* **32**, 660-667.
- ¹⁸ Regnier, V., Romero, E. A., Molton, F., Jazzar, R., Bertrand, G., and Martin, D. (2019). What Are the Radical Intermediates in Oxidative N-Heterocyclic Carbene Organocatalysis? *J. Am. Chem. Soc.* **141**, 1109-1117.
- ¹⁹ Delfau, L., Nichilo, S., Molton, F., Broggi, J, Tomás-Mendivil, E., and Martin, D. (2021). Critical Assessment of the Reducing Ability of Breslow-type Derivatives and Implications for Carbene-Catalyzed Radical Reactions. *Angew. Chem. Int. Ed.* **60**, 26783–26789.
- ²⁰ Ishii, T., Kakeno, Y., Nagao, K., and Ohmiya, H. (2019). N-Heterocyclic Carbene-Catalyzed Decarboxylative Alkylation of Aldehydes. *J. Am. Chem. Soc.* **141**, 3854–3858.
- ²¹ Ishii, T., Ota, K., Nagao, K., and Ohmiya, H. (2019) N-Heterocyclic Carbene-Catalyzed Radical Relay Enabling Vicinal Alkylacylation of Alkenes. *J. Am. Chem. Soc.* **141**, 14073-14077.
- ²² Li, J. -L., Liu, Y. -Q., Zou, W. -L., Zeng, R., Zhang, X., Liu, Y., Han, B., Leng, H. -J., and Li, Q.-Z. (2020). Radical Acylfluoroalkylation of Olefins through N-Heterocyclic Carbene Organocatalysis. *Angew. Chem. Int. Ed.* **59**, 1863–1870.
- ²³ Kim, I., Im, H., Lee, H., and Hong, S. (2020). N-Heterocyclic carbene-catalyzed deaminative cross-coupling of aldehydes with Katritzky pyridinium salts. *Chem. Sci.* **11**, 3192–3197.

-
- ²⁴ Matsuki, Y., Ohnishi, N., Kakeno, Y., Takemoto, S., Ishii, T., Nagao, K., and Ohmiya, H. (2021). Aryl radical-mediated N-heterocyclic carbene catalysis. *Nat. Commun.* *12*, 3848.
- ²⁵ For a review see: Liu, J., Xing, X. -N., Huang, J. -H., Lu, L. -Q., and Xiao, W. -J. (2020). Light opens a new window for N-heterocyclic carbene catalysis. *Chem. Sci.* *11*, 10605.
- ²⁶ Bay, A. V., Fitzpatrick, K. P., Betori, R. C., and Scheidt, K. A. (2020). Combined Photoredox and Carbene Catalysis for the Synthesis of Ketones from Carboxylic Acids. *Angew. Chem. Int. Ed.* *59*, 9143–9148.
- ²⁷ Meng, Q., Döben, N., and Studer, A. (2020). Cooperative NHC and Photoredox Catalysis for the Synthesis of β -Trifluoromethylated Alkyl Aryl Ketones. *Angew. Chem. Int. Ed.* *59*, 19956–19960.
- ²⁸ Meng, Q.; Lezius, L., and Studer, A. (2021). Benzylic C–H Acylation by Cooperative NHC and Photoredox Catalysis. *Nat. Commun.* *12*, 2068.
- ²⁹ Liu, K., and Studer, A. (2021). Direct α -Acylation of Alkenes via N-Heterocyclic Carbene, Sulfinate and Photoredox Cooperative Triple Catalysis. *J. Am. Chem. Soc.* *143*, 4903–4909.
- ³⁰ Zuo, Z., Daniliuc, C. G., and Studer, A. (2021). Cooperative NHC/Photoredox Catalyzed Ring-Opening of Aryl Cyclopropanes to 1-Aroyloxy-3-Acylated Alkanes. *Angew. Chem. Int. Ed.* *60*, 25252–25257.
- ³¹ Yu, X., Meng, Q.-Y., Daniliuc, C. G., and Studer, A. (2022). Aroyl Fluorides as Bifunctional Reagents for Dearomatizing Fluoroarylation of Benzofurans. *J. Am. Chem. Soc.* *144*, 7072–7079.
- ³² Bay, A. V., Fitzpatrick, K. P., González-Montiel, G. A., Farah, A. O., Cheong, P. H., and Scheidt, K. A. (2021). Light-Driven Carbene Catalysis for the Synthesis of Aliphatic and α -Amino Ketones. *Angew. Chem. Int. Ed.* *60*, 17925–17931.
- ³³ Bayly, A. A., McDonald, B. R., Mrksich, M., and Scheidt, K. A. (2020). High-throughput photocapture approach for reaction discovery. *Proc. Natl. Acad. Sci. U. S. A.* *117*, 13261–13266.

-
- ³⁴ Ren, S.-C., Yang, X., Mondal, B., Mou, C., Tian, W., Jin, Z., and Chi, Y. R. (2022). Carbene and photocatalyst-catalyzed decarboxylative radical coupling of carboxylic acids and acyl imidazoles to form ketones. *Nat. Commun.* *13*, 2864.
- ³⁵ Bay, A. V., Farnam, E. J., and Scheidt, K. A. (2022). Synthesis of Cyclohexanones by a Tandem Photocatalyzed Annulation. *J. Am. Chem. Soc.* *144*, 7030–7037.
- ³⁶ Sato, Y., Goto, Y., Nakamura, K., Miyamoto, Y., Sumida, Y., and Ohmiya, H. (2021). Light-Driven N-Heterocyclic Carbene Catalysis Using Alkylborates. *ACS Catal.* *11*, 12886–12892.
- ³⁷ Ren, S. -C., Lv, W. -X., Yang, X., Yan, J. -L., Xu, J., Wang, F. -X., Hao, L., Chai, H., Jin, Z., and Chi, Y.R. (2021). Carbene-Catalyzed Alkylation of Carboxylic Esters via Direct Photoexcitation of Acyl Azolium Intermediates. *ACS Catal.* *11*, 2925–2934.
- ³⁸ Huang, H., Dai, Q. -S., Leng, H. -J., Li, Q. -Z., Yang, S. -L., Tao, Y. -M., Zhang, X., Qi, T., and Li, J. -L. (2022). Suzuki-type cross-coupling of alkyl trifluoroborates with acid fluoride enabled by NHC/photoredox dual catalysis. *Chem. Sci.* *13*, 2584–2590.
- ³⁹ Mavroskoufis, A., Rajes, K., Golz, P., Agrawal, A., Ruß, V., Götze, J. P., and Hopkinson, M. N. (2020). N-Heterocyclic Carbene Catalyzed Photoenolization/Diels–Alder Reaction of Acid Fluorides. *Angew. Chem. Int. Ed.* *59*, 3190–3194.
- ⁴⁰ Guisado-Barrios, G., Bouffard, J., Donnadieu, B., and Bertrand, G. (2010). Crystalline 1*H*-1,2,3-Triazol-5-ylidenes: New Stable Mesoionic Carbenes (MICs). *Angew. Chem. Int. Ed.* *49*, 4759–4762.
- ⁴¹ Guisado-Barrios, G., Soleilhavoup, M., and Bertrand, G. (2018). 1*H*-1,2,3-Triazol-5-ylidenes: Readily Available Mesoionic Carbenes. *Acc. Chem. Res.* *51*, 3236–3244.
- ⁴² Arduengo III, A. J., Harlow, R. L., and Kline, M. (1991). A stable crystalline carbene. *J. Am. Chem. Soc.* *113*, 361–363.

-
- ⁴³ Liu, W., Zhao, L. -L., Melaimi, M., Cao, L., Xu, X., Bouffard, J., Bertrand, G., and Yan, X. (2019). Mesoionic Carbene (MIC)-Catalyzed H/D Exchange at Formyl Groups. *Chem* 5, 2484-2494.
- ⁴⁴ Chu, X. -Q., Ge, D., Cui, Y. -Y., Shen, Z. -L., and Li, C. -J. (2021). Desulfonation via Radical Process: Recent Developments in Organic Synthesis. *Chem. Rev.* 121, 12548–12680.
- ⁴⁵ Liu, W., Vianna, A., Zhang, Z., Huang, S., Huang, L., Melaimi, M., Bertrand, G., and Yan, X. (2021). Mesoionic Carbene-Breslow Intermediates as Super Electron Donors: Application to the Metal-Free Arylacylation of Alkenes. *Chem. Catalysis* 1, 196-206.
- ⁴⁶ Chen, K., Berg, N., Gschwind, R., and König, B. (2017). Selective Single C(sp³)–F Bond Cleavage in Trifluoromethylarenes: Merging Visible-Light Catalysis with Lewis Acid Activation. *J. Am. Chem. Soc.* 139, 18444-18447.
- ⁴⁷ Dai, L., and Ye, S. (2020). NHC-Catalyzed ε -Umpolung via p-Quinodimethanes and Its Nucleophilic Addition to Ketones. *ACS Catal.* 10, 994–998.
- ⁴⁸ Findlay, N. J., Park, S. R., Schoenebeck, F., Cahard, E., Zhou, S., Berlouis, L. E. A., Spicer, M. D., Tuttle, T., and Murphy, J. A. (2010) Reductions of Challenging Organic Substrates by a Nickel Complex of a Noninnocent Crown Carbene Ligand. *J. Am. Chem. Soc.* 132, 15462–15464.
- ⁴⁹ Chatterjee, A. and König, B. (2019) Birch-Type Photoreduction of Arenes and Heteroarenes by Sensitized Electron Transfer. *Angew. Chem. Int. Ed.* 58, 14289–14294.
- ⁵⁰ Hoffmann, N. (2008). Photochemical Reactions as Key Steps in Organic Synthesis. *108*, 1052-1103.
- ⁵¹ Strieth-Kalthoff, F., and Glorius, F. (2020). Triplet Energy Transfer Photocatalysis: Unlocking the Next Level. *Chem* 6, 1888–1903.
- ⁵² Zhou, Q.-Q., Zou, Y.-Q., Lu, L.-Q., and Xiao, W.-J. (2019). Visible-Light-Induced Organic Photochemical Reactions through Energy-Transfer Pathways. *Angew. Chem. Int. Ed.* 58, 1586–1604.

-
- ⁵³ Strieth-Kalthoff, F., James, M. J., Teders, M., Pitzer, L., and Glorius, F. (2018). Energy transfer catalysis mediated by visible light: principles, applications, directions. *Chem. Soc. Rev.* *47*, 7190-7202.
- ⁵⁴ Martin, C. D., Soleilhavoup, M., and Bertrand, G. (2013). Carbene-stabilized main group radicals and radical ions. *Chem. Sci.* *4*, 3020-3030.