



HAL
open science

Arabidopsis thaliana SHOOT MERISTEMLESS Substitutes for Medicago truncatula SINGLE LEAFLET1 to Form Complex Leaves and Petals

Véronique Pautot, Ana Berbel, Thibaud Cayla, Alexis Eschstruth, Bernard Adroher, Pascal Ratet, Francisco Madueño, Patrick Laufs

► **To cite this version:**

Véronique Pautot, Ana Berbel, Thibaud Cayla, Alexis Eschstruth, Bernard Adroher, et al.. Arabidopsis thaliana SHOOT MERISTEMLESS Substitutes for Medicago truncatula SINGLE LEAFLET1 to Form Complex Leaves and Petals. International Journal of Molecular Sciences, 2022, 23, 10.3390/ijms232214114 . hal-03875926

HAL Id: hal-03875926

<https://cnrs.hal.science/hal-03875926>

Submitted on 28 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Article

Arabidopsis thaliana SHOOT MERISTEMLESS Substitutes for *Medicago truncatula* SINGLE LEAFLET1 to Form Complex Leaves and Petals

Véronique Pautot ^{1,*}, Ana Berbel ², Thibaud Cayla ¹, Alexis Eschstruth ^{3,4}, Bernard Adroher ¹, Pascal Ratet ^{3,4}, Francisco Madueño ² and Patrick Laufs ¹

¹ Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), 78000 Versailles, France

² Instituto de Biología Molecular y Celular de Plantas, Consejo Superior de Investigaciones Científicas-Universidad Politécnica de Valencia, Campus de la Universidad Politécnica de Valencia, 46022 Valencia, Spain

³ Université Paris-Saclay, CNRS, INRAE, Université Evry, Institute of Plant Sciences Paris-Saclay (IPS2), 91405 Orsay, France

⁴ Université de Paris, Institute of Plant Sciences Paris-Saclay (IPS2), 91405 Orsay, France

* Correspondence: veronique.pautot@inrae.fr; Tel.: +33-(0)-1-30-83-30-58

Citation: Pautot, V.; Berbel, A.; Cayla, T.; Eschstruth, A.; Adroher, B.; Ratet, P.; Madueño, F.; Laufs, P. *Arabidopsis thaliana* SHOOT MERISTEMLESS Substitutes for *Medicago truncatula* SINGLE LEAFLET1 to Form Complex Leaves and Petals. *Int. J. Mol. Sci.* **2022**, *23*, 14114. <https://doi.org/10.3390/ijms232214114>

Academic Editor: Tomotsugu Koyama

Received: 14 October 2022

Accepted: 10 November 2022

Published: 15 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: LEAFY plant-specific transcription factors, which are key regulators of flower meristem identity and floral patterning, also contribute to meristem activity. Notably, in some legumes, *LFY* orthologs such as *Medicago truncatula* SINGLE LEAFLET (*SGL1*) are essential in maintaining an undifferentiated and proliferating fate required for leaflet formation. This function contrasts with most other species, in which leaf dissection depends on the reactivation of *KNOTTED-like* class I homeobox genes (*KNOX1*). *KNOX1* and *SGL1* genes appear to induce leaf complexity through conserved downstream genes such as the meristematic and boundary *CUP-SHAPED COTYLEDON* genes. Here, we compare in *M. truncatula* the function of *SGL1* with that of the *Arabidopsis thaliana* *KNOX1* gene, *SHOOT MERISTEMLESS* (*AtSTM*). Our data show that *AtSTM* can substitute for *SGL1* to form complex leaves when ectopically expressed in *M. truncatula*. The shared function between *AtSTM* and *SGL1* extended to the major contribution of *SGL1* during floral development as ectopic *AtSTM* expression could promote floral organ identity gene expression in *sgl1* flowers and restore sepal shape and petal formation. Together, our work reveals a function for *AtSTM* in floral organ identity and a higher level of interchangeability between meristematic and floral identity functions for the *AtSTM* and *SGL1* transcription factors than previously thought.

Keywords: *Medicago truncatula*; meristematic activity; flower development; SINGLE LEAFLET/LEAFY; SHOOT MERISTEMLESS transcription factors

1. Introduction

Meristems are essential for plant development, as they are required for the continuous growth and development that are distinguishing features of plants. Amongst all the different types of meristems, the shoot apical meristem (SAM) and the floral meristem (FM) share many features and have been well characterized. The class I *KNOTTED-like* homeobox (*KNOX1*) *SHOOT MERISTEMLESS* (*STM*) and *CUP-SHAPED COTYLEDON* (*CUC1*), *CUC2* and *CUC3* genes are essential regulators of meristem and boundary activities in *Arabidopsis thaliana* (*A. thaliana*) [1,2]. Boundaries are domains of restricted growth located between the meristem and initiating organ primordia or between two organs. These domains control organ separation, inflorescence architecture, organ abscission, fruit opening and leaf shape. Boundaries share overlapping features with meristems, and the regulation of both involves common factors [3]. *CUC* genes are required for SAM

initiation and establish boundaries together with *STM*, which is in turn required for SAM maintenance [4–11]. The three *A. thaliana* *CUC* genes, *CUC1*, *CUC2* and *CUC3*, share partially redundant roles, while also having specific functions. *CUC1* and *CUC2* but not *CUC3* transcripts are negatively regulated by *microRNA164* (*miR164*) [12,13].

LEAFY (*LFY*) is a key regulator of flower meristem identity and floral patterning [14–17]. *LFY* acts as a pioneer transcription factor and promotes chromatin accessibility to its target genes *APETALA1* (*AP1*) and *AGAMOUS* (*AG*) [18,19]. *LFY* also contributes to meristem function, particularly to the formation of floral meristems in *A. thaliana* [20–23]. *LFY* acts together with UNUSUAL FLOWER (*UFO*), an F-box protein, which is a substrate adaptor of CULLIN1–RING ubiquitin ligase complexes (*CRL1*) [24,25] to control meristem function and identity [17,26]. In flowers, both *LFY* and auxin transport contribute to proper positioning of sepal primordia through the regulation of *CUC2* expression [27]. Besides determining floral identity and patterning, *LFY* also contributes to the meristematic identity of floral or axillary meristems with several regulators [21,28–31]. Among them are PENNYWISE (*PNY*) and POUNDFOOLISH (*PNF*), two BEL1-like (*BELL*) homeodomain partners forming heterodimers with *STM* [32,33]. The *LFY* implication in axillary meristem emergence is mediated through REGULATOR OF AXILLARY MERISTEMS1 (*RAX1*), a MYB transcription factor [34,35], and through the repression of *ARABIDOPSIS RESPONSE REGULATOR* (*ARR7*), encoding a cytokinin signaling component [21,36].

Meristematic features can also be found outside *bone fide* meristems, such as in the leaves. This attribute is particularly obvious in compound leaves, in which leaflet formation requires a transient maintenance of a meristematic-like stage. Indeed, in most species with compound leaves, *KNOX1* gene down-regulation at leaf initiation is only transient and these genes are reactivated following leaf initiation, leading to leaflet formation [37–40]. In the inverted repeat-lacking (IRLC) clade of legume species, the formation of compound leaves requires the activity of the *LFY* orthologs called *UNIFOLIATA* (*UNI*) in pea (*P. sativum*) and *SINGLE LEAFLET1* (*SGL1*) in *Medicago truncatula* (*M. truncatula*) [41,42]. These *LFY* orthologs substitute for *KNOX1* expression, which is permanently excluded from the initiating leaf primordia [43,44]. *SGL1* is expressed in the entire SAM and highly expressed in developing leaves, where its prolonged expression is required for the formation of compound leaves [42,45]. However, ectopic expression of *KNOX1* genes in *M. sativa* and *M. truncatula* leaves can further increase leaf dissection [44,46], suggesting that these two *Medicago* species retain the capacity to respond to both *LFY* and *KNOX1* pathways. Consistent with a central role of *LFY* orthologs in IRLC legume leaf morphology, loss of function of the pea *UFO* ortholog, *STAMINA PISTILLOIDA* (*STP*), leads to leaf complexity reduction [47]. In contrast, in non-IRLC legumes such *Lotus japonicus* and soybean, *LFY* orthologs only play a minor role, and *KNOX1* proteins accumulate in leaves and are likely associated with compound leaf development [44,48].

In simple leaves, such as in *A. thaliana*, repression of *KNOX1* genes is permanent, limiting their meristematic features [49,50]. However, these leaves are still able to develop an increased complexity in response to ectopic expression of *KNOX1* genes [40,51–53] and to *UFO* [54].

The observation that depending on the species, *LFY* and *KNOX1* genes can similarly increase leaf complexity (through the formation of leaflets or serrations) and that some species are able to respond to both factors, suggests that both pathways may at least partially converge to control leaf development. *CUC* genes could be such a convergence point as both *KNOX1* and *LFY* pathways require the activity of *CUC2* to make compound leaves [55,56]. Similar to *CUC1/2* in *A. thaliana*, the expression of the *M. truncatula* *NO APICAL MERISTEM* (*MtNAM*) ortholog is regulated by *miR164* [57], and *MtNAM* is required to maintain boundaries both for cotyledon and leaflet separation besides its role in apical meristem initiation [58]. *SGL1* function in leaflet primordium initiation is epistatic to *MtNAM* activity and *MtNAM* RNAs levels are reduced in *sgl1* mutant [58], suggesting that *SGL1* acts upstream of *MtNAM* in this species.

Besides its function in leaves, SGL1 also plays a role in floral meristem identity. *M. truncatula* is a legume species developing compound inflorescences. Upon floral transition, the shoot apical meristem transforms into a primary inflorescence meristem (I1) and gives rise to a lateral secondary inflorescence meristem (I2), which produces a bract, one to three flowers and a spike [59–61]. In contrast to other flowering species which show a sequential floral ontogeny with successive formation of sepals, petals, stamens and carpels, each floral organ derives from a specific primordium; petals and sepals differentiate from common primordia in *M. truncatula* [59]. Thus, each floral meristem gives rise sequentially to five sepals and four common primordia, which further differentiate into five petals and ten stamens, and one carpel. The *Arabidopsis* floral organ identity genes are conserved in legumes [62]. Loss of function of *SGL1* leads to the reversion of common primordia into incomplete floral meristems, giving rise to sepals and carpels without petals and stamens [42]. This phenotype is related to a B function loss. Similar to *LFY* in *Arabidopsis* [63], *SGL1* acts synergistically with *MtPROLIFERATING INFLORESCENCE MERISTEM* (*MtPIM*), the *A. thaliana* ortholog of *APETALA1* (*AP1*) in *M. truncatula* to determine floral meristem identity [61,64]. *MtNMH7* and *MtTM6* are the *A. thaliana* *AP3*-like paralogs. *MtNMH7* determines petal identity whereas *MtTM6* controls stamen identity [65]. *MtPISTILLATA* (*MtPI*) and *MtNGL9* are the two *A. thaliana* *PI*-like paralogs, with *MtPI* functioning as the master regulator of B function [66,67]. The *M. truncatula* genome harbors two redundant *MtAG* members, *MtAGa* and *MtAGb*, which specify stamen and carpel identity and floral meristem determinacy [68,69]. Recently, a novel regulator of inflorescence development and floral organ identity was identified in *M. truncatula*: the *AGAMOUS*-like *FLOWERS* (*AGLF*) gene, which encodes a MYB domain protein that promotes the C floral identity function besides repressing A and B functions [69,70].

Here, we further compared the meristematic activity of *SGL1* (*LFY*) and *AtSTM* (*KNOX1*) using *M. truncatula* compound leaf as a model system. We first showed that *AtSTM* could substitute for *SGL1* to form complex leaves. We next tested whether *AtSTM* could also substitute for *SGL1*'s role during floral development. Indeed, *AtSTM* expression could restore petal formation in *sgl1* flowers, revealing that *AtSTM* could substitute for *SGL1* function to specify petal identity and promote floral organ identity gene expression. Therefore, our data reveal a high level of interchangeability between *SGL1* and *KNOX1* activities in *M. truncatula* that extends beyond the generally accepted meristematic function to the determination of the identity and growth of the flower perianth.


2. Results

2.1. *AtSTM* Substitutes for *SGL1* in *M. truncatula* to Form Compound Leaves

The *M. truncatula* genome harbors two *MtSTM*-like genes, *MtKNOX1* and *MtKNOX6*, and a previous report describes in vitro plantlets overexpressing the *MtSTM*-like genes, *MtKNOX1* and *MtKNOX6*, in *M. truncatula* [46]. However, only the vegetative phenotype was described, as the phenotype of *MtKNOX1* and *MtKNOX6* overexpressors was extremely severe. Therefore, to overcome such strong phenotypes, we thought to use a *KNOX* gene from a heterologous system. *AtSTM* shares 62.8% amino acid identity with *MtKNOX1* and 64.86% with *MtKNOX6*, and in addition to modifying leaf shape when ectopically expressed, *AtSTM* also has an established role in *Arabidopsis* floral identity [5,6,32,52,71,72]. Thus, we selected *AtSTM* to be expressed in *M. truncatula* and to explore its potential more widely; we expressed it under two different promoters by generating the *p35S:AtSTM* and *pSGL1:AtSTM* constructs that we first introduced in wild-type plants (see Section 4 and Supplemental Figure S1).

Transgenic lines expressing high levels of *AtSTM* presented a severe phenotype and were not viable in the greenhouse, similar to in vitro plantlets overexpressing *Mt-KNOX1*-like genes [46] (Figure S2). Only transgenic plants with low levels of *AtSTM* expression could be investigated (Figures 1 and S3). The overall development of these lines was quite normal, although their fertility was reduced. In wild-type *M. truncatula*, the juvenile first

leaf is simple, while adult later leaves are trifoliate and composed of a terminal leaflet with two lateral leaflets (Figure 1A–D). Ectopic *AtSTM* under the *p35S* or the *pSGL1* promoters occasionally led to the formation of an additional leaflet fused to the terminal leaflet of adult leaves (Figures 1F,H and S3). Quantitative analyses were performed using the *p35S:AtSTM* line (Figure 1Q). The wild-type first leaves (rank 1) were simple, while the majority of adult leaves (ranks 2 to 5) were trifoliate (only 4 out of 72 leaves had more than three leaflets). The *p35S:AtSTM* sequences seldom led to complex leaves, as only 8 out of 72 adult leaves (ranks 2 to 5) were more complex (Figure 1Q).



Q




Figure 1. Ectopic expression of *AtSTM* rescues the *sgl1* leaf phenotype. Phenotype of juvenile (L1) and adult leaves (rank L2, L3, L4) of 5-week-old plants. (A–D) R108 control line. Juvenile leaves are

simple, while adult leaves are trifoliate and composed of a terminal leaflet plus two lateral leaflets. The petiole (p) and the rachis (r) are indicated. (E–H) *p35S:AtSTM*, a transgenic line expressing *AtSTM* under the *p35S* promoter producing a L2 and a L4 heart-shaped adult leaves with an ectopic leaflet fused to the terminal leaflet (arrowheads). This phenotype was occasionally observed. Leaflet margins are serrated. (I–L) *sgl1* line, showing simple juvenile (L1) and adult leaves (L2–L4). (M–P) *p35S:AtSTM sgl1* line, showing trifoliate L3 and L4 leaves similar to wild-type. (Q) Quantification of the leaflet number. Four-week-old plants were analyzed (n = 18 plants per genotype). Average \pm SD are shown. Lowercase letters indicate significant differences between genotypes at each leaf rank (one-way ANOVA with Tukey's post hoc test; $p \leq 0.001$). Bars = 5 mm.

We then tested whether *AtSTM* expression is sufficient to rescue the *sgl1* leaf phenotype (see Section 4). In the *sgl1* mutant, the majority of leaves are simple (Figure 1I–L). All rank 5 leaves were simple, but 12 out of 54 leaves (ranks 2 to 4) were bi- or trifoliate in the *sgl1* mutant (Figure 1Q). In contrast, in *p35S:AtSTM sgl1* plants, the majority of adult leaves were trifoliate as in wild-type (Figure 1B–D,O,P). The *p35S:AtSTM* construct restored almost systematically the capacity to form trifoliate leaves, with 52 out of 54 leaves (ranks 3 to 5) producing at least three leaflets (Figure 1Q). Therefore, we concluded that *AtSTM* can replace *SGL1* to promote leaflet formation.

To explore the developmental origin of the extra or rescued leaflets in the different backgrounds, we imaged by SEM young developing leaf primordia (Figure 2). As observed in wild-type apices, a pair of lateral leaflets and a terminal leaflet initiated in *AtSTM* transgenic lines during early leaf primordium development (Figure 2A,C). At stage S8, additional leaflets could form at the base of the terminal leaflet in *AtSTM* (arrows Figure 2D), which were not observed in the wild-type (Figure 2B) and therefore resulted from secondary morphogenesis. This indicates that the morphogenetic window during which leaflets can be initiated is extended following *AtSTM* expression. In *p35S:AtSTM sgl1* plants, the terminal primordium was surrounded by two lateral primordia (Figure 2G,H), already visible at early stages (S4), as seen in the wild-type (Figure 2A,G). Thus, leaflet restoration in *p35S:AtSTM sgl1* does not appear to rely on a late production of leaflets but a rescue of the normal developmental process with a restoration of early lateral leaflet initiation, as occurs in the wild-type.




Figure 2. SEM analysis of early stages of leaf development. (A,B) R108 wild-type control line. (A) SAM and a leaf primordia at S6 stage showing a terminal leaflet (TL) developing trichomes with one of the lateral leaflets (LL) and one stipule (ST). (B) S8 stage leaf primordia with one terminal leaflet folded on itself between the two lateral leaflets. (C,D) *p35S:AtSTM* line. (C) SAM with a typical S4 stage leaf primordia. (D) At S8, the *p35S:AtSTM* line has formed a new leaflet (NL) at the base of the terminal leaflet (arrow). Leaflet margins are dissected. (E,F) *sgl1* mutant. (E) SAM with a simple S5 leaf primordia (SL). (F) At S8, the leaflet is folded on itself. (G,H) *p35S:AtSTM sgl1* line.

(G) SAM with S4 leaf primordia that has formed lateral leaflets similar to wild-type. (H) At S8, the *p35S:AtSTM sgl1* leaf primordia was similar to wild-type with the terminal leaflet surrounded by two lateral leaflets. Bars = 100 μ m.

2.2. *AtSTM* Substitutes for *SGL1* in *M. truncatula* in Specifying Petal Formation

M. truncatula is a legume species developing compound inflorescences. The wild-type *M. truncatula* mature flower (Figure 3A–I) comprises four whorls consisting of a calyx formed by five sepals fused at their base (Figure 3B), a corolla containing three types of yellow petals, the standard or the vexillum at the adaxial position (Figure 3C), the keel formed by two fused petals at the abaxial position surrounded by two lateral petals and the alae or wings (Figure 3D–G). The third whorl consists of an independent stamen filament at the adaxial position, the vexillary stamen filament and nine stamen filaments fused into a staminal tube that surrounds a monocarpous gynoecium [59] (Figure 3H,I). The *sgl1* mutants produce inflorescences with cauliflower-like floral structures, containing incomplete floral meristems (FMs), elongated sepals and occasionally carpels [42] (Figure 3U). These cauliflowers do not produce petals nor stamens, similar to *lfy* mutants in *Ara-bidopsis* [73].

Wild-type plants for *SGL1* overexpressing *AtSTM* occasionally produced abnormal flowers showing fused organs and are characterized by an increase in petal identity with petaloid sepals and petaloid stamens (Figure 3K–S). Some petals showed alterations in shape or serrated margins (Figure 3L,M). These flowers occasionally produced two to three unfused carpels (Figure 3M–O). Flowers can show petaloid sepals (Figure 3N,S), petaloid stamens (Figure 3O,P) and petaloid carpels (Figure 3Q). The fertility was severely reduced, with some plants infertile. The fruits were small, with fewer discs and unbent spines compared with wild-type fruits (Figure 3T,J). These fruits contained a few seeds. The same phenotypes were occasionally observed in *pSGL1:AtSTM* flowers (Figure 3F,G). We then tested the effects of *p35S:AtSTM* on *sgl1* flower development. Surprisingly, the ectopic expression of *AtSTM* rescued sepal shape and petal formation in the *sgl1* mutant (Figure 3V–Y). Similar to wild-type flowers, *p35S:AtSTM sgl1* flowers formed a calyx with five sepals fused at their base (Figure 3B,W). Inside the calyx, the *p35S:AtSTM sgl1* flowers showed a cauliflower phenotype with incomplete FMs, producing a few sepals and a majority of petals or petals with sepal sectors. Petals were partially restored as some of them had a vexillum-like, wing-like or keel-like shape (Figure 3X,Y). Thus, when ectopically expressed, *AtSTM* restores petal formation in *sgl1*. These flowers did not form carpels, in contrast to *sgl1* flowers, suggesting a deficiency in C function (Figure 3). The majority of organs formed were petals, as one cauliflower flower from a *35S:AtSTM sgl1* line could produce up to 65 petals (Figure S4).




Figure 3. Ectopic expression of *AtSTM* in *M. truncatula* promotes petal identity. (A–J) R108 wild-type control line. (A) The wild-type flower showing the calyx (C) and the corolla containing 5 petals: the standard or vexillum (Vx), the keel (K) and two alae (A) or wings. (B–I) A wild-type dissected flower: (B) the calyx, formed by 5 fused sepals at their base. (C) The standard or vexillum, abaxial side. (D,E) The keel formed by two fused petals (arrows) surrounded by two lateral petals, the alae or wings, adaxial (D) and abaxial (E) sides, (F) a dissected wing. (G) A dissected keel petal. (H,I) A single carpel enclosed by a staminal tube comprising nine fused stamens plus one independent “vexillary” stamen at the adaxial position (I). (J) After fertilization, the carpel grows out to form a coiled fruit with spines. (K–T) *p35S:AtSTM* line. (K–Q) Phenotypes of *p35S:AtSTM* flowers. (L,M)

Petals were abnormal and can show dissected margins. (N) Flower showing petaloid sepals. (M–O,Q) Flowers forming two or three carpels (arrowheads), with some showing an unfused carpel (arrows). (O,P) Flowers showing petaloid stamens. (Q) A dissected flower (the corolla was removed) showing 3 carpels with one developing petaloid sectors. (R) Dissected petals, some of them showing sepal sectors. (S) A dissected calyx showing petaloid sepals. (T) Fruits were smaller with unbent spines. (U) *sgl1* inflorescence containing three flowers with a cauliflower-like morphology, *sgl1* flowers contain sepal and carpeloid structures and lack petals and stamens. (V–Y) A *p35S:AtSTM sgl1* flower showing petals. This flower contains inside incomplete FMs that produce mainly petals or petaloid sepals and a few sepals. (V) FMs are visible (*). (W) Bottom view showing the calyx (arrow), the sepal form is restored (see also Figure 4M), some other sepals are visible (arrowhead). Dissected petals: a vexillum-like petal (X), a keel-like (arrow Y) and wing-like petals (arrowhead Y). Bars = 2 mm, except for F, G, I, N and U, for which bars = 1 mm.

SEM analyses were performed to further characterize these flowers at early developmental stages. Figure 4A–D shows wild-type floral development. At stage 4, the wild-type floral meristem had formed five sepal primordia, four common primordia and a carpel primordium (Figure 4B). At late stage 5, the wild-type floral meristem displayed the complete set of floral organ primordia, with petal and stamen primordia deriving from the differentiation of common primordia (Figure 4D). Figure 4E–H shows the floral development of a *p35S:AtSTM* plant wild-type for *SGL1*. Figure 4F shows a late stage 5 *p35S:AtSTM* floral meristem. Based on sepal development, a delay in the formation of the inner floral organ primordia could be observed compared with the wild-type (Figure 4F,D). In contrast, Figure 4G shows a stage 5 floral meristem containing differentiated petals and stamen primordia and two carpel primordia, indicating that the delay in internal organ primordia differentiation is variable between flowers. Figure 4H shows a *p35S:AtSTM* flower developing two carpels. Similar to previous data [42,61], *sgl1* inflorescences showed multiple incomplete FMs, elongated sepals, defective common primordia and carpel primordia (Figure 4I–L). Sepal primordia further develop into elongated sepals and carpel primordia into a carpel-like structure. The cauliflower phenotype is caused by the iterative conversion of common primordia into incomplete floral meristems (Figure 4I,L). In *sgl1* mutants overexpressing *AtSTM* (Figure 4M–O), the sepal form was restored, suggesting that *AtSTM* could take over *SGL1* function for the control of sepal shape. The late stage 5 floral meristems showed a delay in the differentiation of other floral organ primordia, as observed in *p35S:AtSTM SGL1* plants (Figure 4N,F). Later, petals and sepals differentiated from these primordia (Figure 4O). Together, these observations show that expression of *AtSTM* partly restored normal early morphogenesis of *sgl1* flowers.




Figure 4. SEM analysis of flower development. (A–D) Flower development in wild-type R108 line. (A) An inflorescence showing floral meristems (FMs) at different stages, including a late S5 stage primordia and a spike (spk) at the base of the floral meristem. (B) S4 stage FM showing the abaxial sepal (Sab), the adaxial sepal (Sad), two lateral sepals (SI), four common primordia (CPab, CPad and two CPI) and one carpel primordia (Cp), Br (bract). (C) S5 stage FM showing anteseptal stamen (Sts), antepetal stamen (Stp) and petal (P) primordia. (D) A late S5 stage FM showing differentiated flower organ primordia, non-fused vexillary stamen (stp*), alae petals (A), vexillum (Vx), keel petals (K), carpel (C). (E–H) Flower development in *p35S:AtSTM* plants. (E) Inflorescence showing floral meristems at different stages. (F) Close-up of a late S5 stage FM showing the differentiation of sepal primordia (S) and the formation of bulges in the center (*). Other floral organ primordia are not differentiated. (G) Close-up of a late S5 FM showing two carpel primordia (C). (H) S8 stage flower with two differentiated carpels (C). (I–L) Flower development in *sgl1* plants. (I) *sgl1* inflorescence showing multiple incomplete FMs, elongated sepals (S) and S4 stage FM. (J) A close-up view of S4 stage *sgl1* FM showing defective common primordia (CP'). (K) S5 stage FM, CP's do not further differentiate. (L) Carpel primordium (C). (M–O) Flower development in *p35S:AtSTM sgl1* plants. (M) Inflorescence showing incomplete FMs at different stages. (N) Close-up of S4 and a late S5 stage incomplete FM showing the differentiated sepals and bulges in the center (*). Note that the sepal form is similar to wild-type sepal (see (D)). (O) S8 stage flower (the calyx has been removed) containing two FMs that differentiate sepals (S) and petals (P). An elongated sepal is visible. Bars: (A,E,I,M–O) = 100 μ m, (B–D,E,G,J,K) = 50 μ m, (L) = 250 μ m.

2.3. *AtSTM* Substitutes for *SGL1* to Promote Floral Organ Identity Gene Expression

To determine if *AtSTM* activates A and B functions to promote petal formation in *sgl1* flowers, we used in situ hybridization to analyze the expression pattern of floral organ identity genes in *p35S:AtSTM sgl1* flowers. We first investigated the expression of the A

class gene *MtPIM*, the *A. thaliana* ortholog of *AP1* in *M. truncatula*. *MtAP1* has a conserved role with orthologous genes and is required to specify floral meristem and floral organ identity [61,64]. In wild-type inflorescences, *MtAP1* transcripts localize to the floral meristem and bract (Figure 5A,B). In a stage 4 flower meristem, *MtAP1* expression was observed in sepal primordia and was restricted to the outer domain of the common primordia that further gives rise to sepals and petals and was absent from the inner part, which differentiates into stamens and carpel (Figure 5C) [61,64]. At later stages, *MtAP1* expression was maintained in sepals and petals (Figure 5D). Similar to the pattern described in [61], in *sgl1* flowers, *MtAP1* was expressed in the floral meristem and in the bract (Figure 5E). *MtAP1* was expressed uniformly in defective common primordia and in reiterated floral meristems (Figure 5F–H). At later stages, *MtAP1* expression localized to the outer incomplete floral meristem and disappeared from the central domain that further differentiates into carpels (Figure 5F,G). In *p35S:AtSTM sgl1* flowers, *MtAP1* was more widely expressed than in *sgl1* flowers, with *MtAP1* detected in reiterated floral meristems and in developing petals (I–K). Thus, in *p35S:AtSTM sgl1* flowers, *AtSTM* acts as a positive regulator of A function, contributing to enhanced petal identity.




Figure 5. *MtAP1* expression in wild-type, *sgl1* and *p35S:AtSTM sgl1* flowers. (A–D) R108 wild-type flowers. (A,B) *MtAP1* was expressed in the floral meristem (FM) and in the bract primordia (Br) and absent in secondary inflorescence meristem (I₂). (C) At stage 4, *MtAP1* was expressed in sepals (S). *MtAP1* was restricted to the outer part of the developing common primordia (CP), which will give rise to petal (P) and was absent in the inner part that will give rise to stamen (St). (D) At stage 6, *MtAP1* expression was maintained in sepals and petals. *MtAP1* was absent in carpels (C) and stamens. (E–H) *sgl1* flowers. (E) *MtAP1* was expressed in bract (Br) and floral meristem (FM). (F,G) *MtAP1* was uniformly expressed in defective common primordia (CP'), unlike in wild-type common primordia, *MtAP1* was absent in the inner part of the floral meristem where carpel will develop (C). (H) *MtAP1* was expressed in reiterated floral meristems deriving from floral primordia. (I–K) *p35S:AtSTM sgl1* flowers. *MtAP1* expression was detected in reiterated floral meristems (FM), in defective common primordia (CP') and in developing petals (P). Bars: (A–I) = 50 μ m, (J,K) = 100 μ m.

We then investigated the expression of the B class gene *MtPI*. In wild-type, *MtPI* transcripts were localized to common primordia cells and later restricted to petal and stamens (Figure 6A,B) and [66,67]. In the *sgl1* mutant, no *MtPI* expression was detected in defective common primordia, consistent with the phenotype of *sgl1* flowers, which lack petals and stamens (Figure 6C). In *sgl1* flowers overexpressing *AtSTM*, *MtPI* expression was detected in defective common primordia (Figure 6D,E inset-a). At a later stage, *MtPI* localized to the outer domain of the defective common primordia that further gives rise to petal-like organs (Figure 6E and inset-b). Later, *MtPI* is expressed in petal-like organs

(Figure 6D,E). Thus, *AtSTM* acts as a positive regulator of *MtPI* expression consistent with the restoration of petal identity.




Figure 6. *MtPI* expression in wild-type, *sgl1* and *p35S:AtSTM sgl1* flowers. (A,B) *MtPI* expression in R108 wild-type flowers. (A) *MtPI* expression was detected at stage 3 in cells of the common primordia (CP). (B) At later stages, *MtPI* expression was restricted to stamens (St) and petals (P). (C) *MtPI* was not detected in *sgl1* flowers. (D,E) *MtPI* expression in *p35S:AtSTM sgl1* flowers. (D) A flower developing sepals (S) and petals (P). *MtPI* expression was detected in defective common primordia (CP') and in petals (P). (E) A cauliflower showing several floral meristems and developing sepals (S) and petals (P). *MtPI* was detected in defective common primordia (CP') and in petals (P). Unlike in wild-type, in which the expression of *MtPI* is observed in the whole common primordia, the expression of *MtPI* was restricted to the periphery of the defective common primordia, which will give rise to petals (see details in a,b). (a,b) Close-ups of the areas marked in (E). Bars: (A,C,a,b) = 50 μ m, (B–E) = 100 μ m.

We further determined the expression of the *M. truncatula* ortholog of the *A. thaliana* C-class gene *AG*. *MtAGb* was used as a probe as its signal is stronger and it is more restricted than that of *MtAGa* [68]. In wild-type flowers, *MtAGb* expression was first detected at stage 2 in the central part of the floral meristem where the carpel will develop (Figure 7A). At stage 4, *MtAGb* expression was mainly localized to the inner domain of the common primordia that will further give rise to stamens and to carpel primordia (Figure 7B). At later stages, its expression was restricted to stamens, carpel and ovules (Figure 7C,D). In *sgl1* flowers, a weak signal was detected in floral meristems and defective common primordia and was absent in the L1 layer (Figure 7E–G). Later, its expression was detected in carpel-like structures and ovules (Figure 7H). In *sgl1* plants overexpressing *AtSTM*, *MtAGb* expression was detectable in only a few flowers (3 of 13). In these flowers, the signal was weak and restricted to a few cells in FM beneath the two outer most layers (Figure 7I).




Figure 7. *MtAGb* expression in wild-type, *sgl1* and *p35S:AtSTM sgl1* flowers. (A–D) *MtAGb* expression in wild-type flowers. (A) *MtAGb* expression was located in the center cells of the floral meristem (FM) at stage 2. (B) At stage 4, *MtAGb* expression was detected in the carpel primordia (C) and the half of the common primordia (CP) that will give rise to the stamens. (C) At stage 5, *MtAGb* expression was detected in stamen (St) and carpel (C) primordia. (D) In later stages, *MtAGb* expression was located in stamen (St), carpel (C) and developing ovules (Ov). (E–H) *MtAGb* expression in *sgl1* flowers. *MtAGb* was detected in FM (E, arrowhead) and in defective common primordia (CP', F arrowhead, G) and absent in the L1 layer. (H) In later stages, *MtAGb* expression was detected in carpel (C) and ovules (Ov). (I,J) *MtAGb* expression in *p35S:AtSTM sgl1* flowers. Expression was detected in 3 flowers out of 13. (I) An apex showing expression underneath the outermost cell layers in floral meristem (FM). (J) An apex showing no expression. Bars = 50 μ m.

3. Discussion

Here, we compared the activity of two transcription factors, AtSTM and SGL1, in *M. truncatula*. Our analysis is based on transgenic plants that were able to grow in a greenhouse and therefore expressed *AtSTM* at low levels. This allowed us to investigate the activity of *AtSTM* during flower development.

An increase in the leaflet number was only occasionally observed following *AtSTM* ectopic expression in wild-type *M. truncatula*. This limited effect of *AtSTM* could be linked to *AtSTM* expression levels in these lines, which were low. The additional leaflets were formed at the base of the terminal leaflet and resulted from a secondary morphogenesis. This suggests that AtSTM leads to additional leaflets through the extension of the meristematic activity, allowing more leaflets to emerge, and not from the division of the lateral leaflets into two structures. In *M. truncatula*, the terminal leaflet derives from the terminal zone where auxin maxima are located through the activity of *SMOOTH LEAF MARGIN1* (*SLM1*), the *PIN1* ortholog in *M. truncatula* [74]. Lateral leaflets result from the marginal blastozone activity and the formation of local auxin maxima that depend on SGL1 activity [74]. The tetrafoliate pattern seen in *AtSTM* transgenic lines likely results from a defect in auxin distribution in the terminal zone. This leaf patterning is also found in *M. truncatula* plants inactivated for *HEADLESS* (*HDL*) or *MtREVOLUTA1* (*MtREV1*), the putative orthologs of *A. thaliana* *WUSCHEL* and *REVOLUTA*, of which mutants are altered in auxin homeostasis [75,76]. The ectopic expression of *AtSTM* could rescue the formation of lateral leaflets in the *sgl1* mutant. These data show that *AtSTM* could substitute for *SGL1* via an independent pathway to form complex leaves. This suggests that AtSTM could bypass the requirement for SGL1 during the formation of compound leaves in *M. truncatula*, indicating shared functions between these proteins, a conclusion further reinforced by the study of the floral phenotype of *p35S:AtSTM sgl1* plants.

Our data revealed an unexpected effect of AtSTM on floral development, as AtSTM could induce petal identity. The effect of AtSTM on petal identity was moderately visible in an *SGL1* wild-type background, as only few chimeric petaloid floral organs were formed, but was dramatic in an *sgl1* mutant background. Indeed, all *p35S:AtSTM sgl1* flowers produced petals or petals with sepal sectors, while such organs were missing in *sgl1*. Although the increase in petal number could be in part due to the indeterminate state conferred by the *sgl1* mutation, it nevertheless indicates that AtSTM can restore petal formation in an *sgl1* mutant. The shape of *sgl1* sepals was also restored following *AtSTM* expression, showing that *AtSTM* could substitute for other functions of *SGL1* during flower development. The formation of petals in *p35S:AtSTM sgl1* was correlated with an activation of *MtAP1* and more notably of *MtPI* expression, suggesting that AtSTM could promote the expression of these floral organ identity genes to restore petals, and not through an indirect effect on floral meristem growth, for instance. Such a role for *KNOX1* genes in the promotion of B function was not yet reported in either *M. truncatula* nor in *A. thaliana* [44,46,72].

On the contrary, *p35S:AtSTM sgl1* flowers did not form stamens, and in contrast to *sgl1* cauliflowers, which developed carpels, *AtSTM sgl1* cauliflowers lacked carpels. *MtAGb* expression was only rarely detected in *p35S:AtSTM sgl1* cauliflowers, in agreement with the lack of carpel identity. Interestingly, the expression of *MtAGb* was systematically detected in floral meristems beneath the outermost layers in *sgl1* background. The localization and the low intensity of the *MtAGb* signal in *sgl1* cauliflowers suggest that *SGL1* influences *MtAGb* expression.

In *Arabidopsis*, a link for *AtSTM* with carpel identity was revealed with the analysis of plants compromised for AtSTM activity in line with *AtSTM* expression in flowers [5,6,71,73]. A more direct contribution to carpel identity was illustrated with the phenotype of *A. thaliana KNOX1* overexpressors showing homeotic conversion of ovules into pistils. However, *KNOX1* ectopic expression does not complement the *ag* mutant [52,72]. In line with these conclusions made in *Arabidopsis*, we observed that in *M. truncatula*, ectopic expression of *AtSTM* could not induce the C function in the absence of *SGL1* activity. It is possible that in *p35S:AtSTM sgl1* flowers, AG is playing a role related to floral meristem termination more than a function related to the specification of carpel identity.

The impact of *AtSTM* was more obvious both in leaves and flowers of the *sgl1* mutant compared with wild-type *SGL1* plants. This distinct impact could suggest that the STM pathway is more effective in the absence of *SGL1* activity. It is likely that *SGL1* acts in part through the *M. truncatula UFO* ortholog, as it does in *Arabidopsis* and other legumes. Indeed, in pea and in *Lotus japonicus* defective in *STAMINA PISTILLOIDA (STP)* or in *PROLIFERATING FLOWER ORGAN (PFO)*, the *A. thaliana UFO* orthologs lack petals and stamens and show a reduced carpel formation similar to *sgl1* flowers [47,77]. On the other hand, AtSTM was shown recently to function together in *A. thaliana* with AP1 to specify floral meristem identity in part via UFO [78]. This suggests that *SGL1* and STM pathways may converge on MtUFO and that a competition for UFO interaction or for targets shared between *SGL1* and AtSTM could be the basis for the higher effect of AtSTM in the absence of *SGL1*.

Our work shows that AtSTM substitutes for *SGL1* function in *M. truncatula* during both vegetative and reproductive development. A parallel has been proposed between compound leaflet primordia and common primordia formation. Both of these processes seem to require the maintenance of an indeterminate phase controlled by *SGL1* [61]. While in leaves, *SGL1* maintains the indeterminate state, in flowers, *SGL1* acts in opposite by promoting the formation of common primordia. The capacity for AtSTM to substitute for *SGL1* in both leaves and flowers underlines this parallel and the control of meristematic activity shared by these two transcription factors.

4. Materials and Methods

4.1. Plant Growth and Plant Material

M. truncatula plants were grown in a greenhouse or in growth chambers under long-day conditions (16 h light at 23 °C and 8 h dark at 15 °C). The wild-type (R108) and the *sgl1-1* mutant *M. truncatula* lines have been described [42].

The *pSGL1:GUS* reporter construct was generated as follows. A 2.7 kb fragment corresponding to the *SGL1* (Medtr3g098560) promoter sequence used in [42] (wild-type *M. truncatula* cv Jemalong) was amplified from the *M. truncatula* R108 ecotype using primers *pSGL1-for*, incorporating a *Bgl*III site, and *pSGL1-rev*, incorporating a *Bam*HI site. The promoter was cloned into *pCR Blunt II-TOPO* vector to create *pTOPO-pSGL1* and sequenced. The *pSGL1* promoter was moved into the binary vector *pCAMBIA 3301* in front of the β -glucuronidase (*GUS*) gene. For this, a *Bgl*III-*Bam*HI fragment containing the *SGL1* promoter was ligated into *pCAMBIA3301* cut with *Bam*HI and *Bgl*III to replace the 35S promoter.

The *pSGL1:HA-AtSTM* construct was generated as follows (*AtSTM*, AT1G62360). *pTOPOpSGL1* was cut with *Eco*R1-*Bam*HI to release the *pSGL1* promoter, which was cloned into the *pCAMBIA 3300* binary vector cut with *Eco*RI and *Bam*HI to create *pCAMBIA 3300 pSGL1*. The *alli2AtSTM* plasmid harboring the triple hemagglutinin (HA) tag-*AtSTM* fusion under the double enhanced cauliflower Mosaic Virus 35S promoter was used as a template to amplify the HA-*AtSTM* fusion using primers *AtSTM-for* and *AtSTM-rev* incorporating *Bam*HI and *Eco*RI sites, respectively. This fragment was ligated into the *pALC* vector (Syngenta Ltd., Jeolotts Hill, UK) cut with *Bam*HI and *Eco*RI. The *Bam*HI-*Xba*I fragment containing the HA-*AtSTM* fusion and the 35S terminator was cloned into *pCAMBIA 3300 pSGL1* to create *pCAMBIA pSGL1:HA-AtSTM 35S term*.

The *p35S:HA-AtSTM* construct was generated as follows. The *pSGL1* promoter sequence of the *pCAMBIA pSGL1:HA-AtSTM 35S term* was replaced with the 35S promoter sequence from *pCAMBIA 3301* using the *Bgl*III and *Bam*HI sites. The *pCAMBIA 3301* was cut with *Bam*HI and *Bgl*III to release the 35S promoter, and the *pCAMBIA 3300* containing the *pSGL1:HA-AtSTM* construct was cut with *Bgl*III and *Bam*HI to replace the *pSGL1* promoter with the 35S promoter to create *pCAMBIA p35S:HA-AtSTM 35S term*. *pSGL1-GUS*, *p35S:AtSTM* and *pSGL1:AtSTM* constructs were introduced into *A. tumefaciens* GV3101. The *pSGL1-GUS* construct was used to transform *M. truncatula* R108 wild-type plant, while *p35S:AtSTM* and *pSGL1:AtSTM* constructs were used to transform *M. truncatula* R108 plants heterozygous for the *sgl1-1* mutation. *M. truncatula* transgenic lines were created using a leaf disc protocol [79]. Transgenic calli were selected on media containing 3 mgL⁻¹ Basta (glufosinate-ammonium). Primers are listed in Table S1.

Four independent *pSGL1:GUS* transgenic lines were analyzed for *SGL1:GUS* activity. The *SGL1:GUS* activity was detected in meristem, vascular tissue and young leaves in R108 *M. truncatula* (Figure S1), which was similar to the activity of the *SGL1* promoter isolated from the JemalongA17 ecotype [42], and in axillary meristem, young floral buds and carpels (Figure S1).

Most of the transgenic plantlets expressing *AtSTM* were not viable when transferred to soil. RT-PCR were realized to compare the level of expression of *AtSTM* in transgenic lines. Total RNA was extracted from *AtSTM* transgenic lines expressing *p35S:AtSTM* (in vitro seedlings and transgenic plants grown in the greenhouse) using Tri reagent (Sigma-Aldrich, Saint-Quentin-Fallavier, France) and treated with DNase I (Invitrogen, Waltham, MA, USA) according to the manufacturer's instructions. *AtSTM* levels were monitored using *qAtSTM-F* and *qAtSTM-R* primers. Primers specific for the *M. truncatula* *UBIQUITIN* gene (Medtr3g091400) were used as an internal control [80]. Only transgenic plantlets expressing *AtSTM* at low levels were viable in the greenhouse. Four *p35S:AtSTM* independent lines and three *pSGL1:AtSTM* lines were obtained. Of these, two independent *p35S:AtSTM* lines and one *pSGL1:AtSTM* based on their phenotype were chosen for further characterization. These plants showed reduced fertility. Plants homozygous for

the *p35S:AtSTM* construct and heterozygous for *sgl1* were obtained and confirmed by PCR genotyping [42].

4.2. Phenotypic Observations

Leaves and flowers were observed under a binocular microscope (Nikon, SMZ1000) and imaged with a digital camera (ProgRes C10^{plus}). *M. truncatula* meristems showing GUS activity were dissected and photographed using a LeicaMZ12 dissecting microscope fitted with an AxioCam ICc5 digital camera.

4.3. Quantitative Analyses of Leaf Development

Progenies of *SGL1+/sgl1* (R108) and *p35S:AtSTM SGL1+/sgl1* lines were grown in a greenhouse. Four-week-old plants were used. The leaflet number was determined on R108 wild-type, *sgl1/sgl1*, *p35S:AtSTM SGL1+* and *p35S:AtSTM sgl1/sgl1* plants. Eighteen plants per genotype were analyzed.

4.4. Scanning Electron Microscopy (SEM)

Three to eight-week-old plants were dissected to observe leaf and flower primordia. The samples were imaged using SEC DESKTOP SEM (Scanning Electron Microscope, (SNE-1500M), SEC, Suwon, Korea) at an accelerating voltage of 15 kV.

4.5. In Situ Localization of GUS Activity and In Situ Hybridization

GUS staining and tissue embedding have been described in [81]. RNA in situ hybridization with digoxigenin-labeled probes was performed as previously described [82]. The RNA antisense and sense probes of *MtAP1* (Medtr8g066260) *MtPIM*, *MtPI* (Medtr3g088615) and *MtAGb* (Medtr8g087860) were generated using as cDNA templates a 426 bp fragment of *MtPIM* (282–707 from ATG), a 298 bp fragment of *MtPI* (504–801 from ATG) or a 215 bp fragment of *MtAGb* (558–773 from ATG), respectively, cloned into the pGEM-T Easy vector (Promega, Madison, WI, USA) and using the corresponding SP6 and T7 RNA polymerases in the vector for transcription. SP6 was used for transcription of RNA antisense probes and T7 for the sense. The in situ hybridization with control sense probes is presented in Figure S7.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijms232214114/s1>, Figure S1: *pSGL1:GUS* expression in *M. truncatula*, Figure S2: In vitro transgenic *M. truncatula* plantlets overexpressing *AtSTM*, Figure S3: Phenotype of a *pSGL1:AtSTM* transgenic line expressing *AtSTM* under the *pSGL1* promoter, Figure S4: Petal production following *AtSTM* expression in *sgl1* flowers, Figure S5: In situ hybridization with control sense probes, Table S1: List of primers.

Author Contributions: All authors made essential contributions to the project. V.P. performed most of the experiments; A.E. made the *Medicago truncatula* transgenics; T.C. performed the GUS assays, some leaf phenotypic characterization and some SEMs; B.A. provided technical assistance to V.P.; A.B. and F.M. performed the in situ analyses; P.L., P.R. and V.P. designed the research; V.P. and P.L. wrote the article. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by IJPB's Plant Observatory technological platforms. The IJPB benefits from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007). Work at F.M.'s lab was supported by the Spanish Ministerio de Ciencia Innovación y Universidades and FEDER (grants BIO2015-64307-R and PGC2018-099232-B-I00). Thibaud Cayla and Alexis Eschstruth were supported by the ANR-11-BSV2-0005 Charmful.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We thank Bruno Letarnec for help in the greenhouse. We are grateful to Shelley Hepworth and Nicolas Arnaud for critical reading.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Hepworth, S.R.; Pautot, V.A. Beyond the divide: Boundaries for patterning and stem cell regulation in Plants. *Front. Plant Sci.* **2015**, *6*, 1052. <https://doi.org/10.3389/fpls.2015.01052>.
- Maugarny-Calès, A.; Gonçalves, B.; Jouannic, S.; Melkonian, M.; Ka-Shu Wong, G.; Laufs, P. Apparition of the NAC transcription factors predates the emergence of land plants. *Mol. Plant* **2016**, *9*, 1345–1348. <https://doi.org/10.1016/j.molp.2016.05.016>.
- Žádníková, P.; Simon, R. How boundaries control plant development. *Curr. Opin. Plant Biol.* **2014**, *17*, 116–125. <https://doi.org/10.1016/j.pbi.2013.11.013>.
- Aida, M.; Ishida, T.; Tasaka, M. Shoot apical meristem and cotyledon formation during *Arabidopsis* embryogenesis: Interaction among the *CUP-SHAPED COTYLEDON* and *SHOOT MERISTEMLESS* Genes. *Development* **1999**, *126*, 1563–1570.
- Endrizzi, K.; Moussian, B.; Haecker, A.; Levin, J.Z.; Laux, T. The *SHOOT MERISTEMLESS* gene is required for maintenance of undifferentiated cells in *Arabidopsis* shoot and floral meristems and acts at a different regulatory level than the meristem genes *WUSCHEL* and *ZWILLE*. *Plant J.* **1996**, *10*, 967–979. <https://doi.org/10.1046/j.1365-313X.1996.10060967.x>.
- Long, J.A.; Moan, E.I.; Medford, J.I.; Barton, M.K. A Member of the KNOTTED class of homeodomain proteins encoded by the *STM* gene of *Arabidopsis*. *Nature* **1996**, *379*, 66–69. <https://doi.org/10.1038/379066a0>.
- Takada, S.; Hibara, K.; Ishida, T.; Tasaka, M. The *CUP-SHAPEDCOTYLEDON1* gene of *Arabidopsis* regulates shoot apical meristem formation. *Development* **2001**, *128*, 1127–1135.
- Belles-Boix, E.; Hamant, O.; Witiak, S.M.; Morin, H.; Traas, J.; Pautot, V. *KNAT6*: An *Arabidopsis* homeobox gene involved in meristem activity and organ separation. *Plant Cell* **2006**, *18*, 1900–1907. <https://doi.org/10.1105/tpc.106.041988>.
- Spinelli, S.V.; Martin, A.P.; Viola, I.L.; Gonzalez, D.H.; Palatnik, J.F. A Mechanistic link between *STM* and *CUC1* during *Arabidopsis* development. *Plant Physiol.* **2011**, *156*, 1894–1904. <https://doi.org/10.1104/pp.111.177709>.
- Scofield, S.; Murison, A.; Jones, A.; Fozard, J.; Aida, M.; Band, L.R.; Bennett, M.; Murray, J.A.H. Coordination of meristem and boundary functions by transcription factors in the *SHOOT MERISTEMLESS* Regulatory Network. *Development* **2018**, *145*, 157081. <https://doi.org/10.1242/dev.157081>.
- Aida, M.; Tsubakimoto, Y.; Shimizu, S.; Ogisu, H.; Kamiya, M.; Iwamoto, R.; Takeda, S.; Karim, M.; Mizutani, M.; Lenhard, M.; et al. Establishment of the embryonic shoot meristem involves activation of two classes of genes with opposing functions for meristem activities. *Int. J. Mol. Sci.* **2020**, *21*, 5864. <https://doi.org/10.3390/ijms21165864>.
- Laufs, P.; Peaucelle, A.; Morin, H.; Traas, J. MicroRNA Regulation of the *CUC* genes is required for boundary size control in *Arabidopsis* meristems. *Development* **2004**, *131*, 4311–4322. <https://doi.org/10.1242/dev.01320>.
- Mallory, A.C.; Reinhart, B.J.; Jones-Rhoades, M.W.; Tang, G.; Zamore, P.D.; Barton, M.K.; Bartel, D.P. MicroRNA Control of *PHABULOSA* in leaf development: Importance of pairing to the MicroRNA 5' Region. *EMBO J.* **2004**, *23*, 3356–3364. <https://doi.org/10.1038/sj.emboj.7600340>.
- Parcy, F.; Nilsson, O.; Busch, M.A.; Lee, I.; Weigel, D. A Genetic framework for floral patterning. *Nature* **1998**, *395*, 561–566. <https://doi.org/10.1038/26903>.
- Wagner, D.; Sablowski, R.W.M.; Meyerowitz, E.M. Transcriptional activation of *APETALA1* by *LEAFY*. *Science* **1999**, *285*, 582–584. <https://doi.org/10.1126/science.285.5427.582>.
- Lohmann, J.U.; Hong, R.L.; Hobe, M.; Busch, M.A.; Parcy, F.; Simon, R.; Weigel, D. A Molecular link between stem cell regulation and floral patterning in *Arabidopsis*. *Cell* **2001**, *105*, 793–803. [https://doi.org/10.1016/S0092-8674\(01\)00384-1](https://doi.org/10.1016/S0092-8674(01)00384-1).
- Chae, E.; Tan, Q.K.-G.; Hill, T.A.; Irish, V.F. An *Arabidopsis* F-Box Protein Acts as a transcriptional co-factor to regulate floral development. *Development* **2008**, *135*, 1235–1245. <https://doi.org/10.1242/dev.015842>.
- Jin, R.; Klasfeld, S.; Zhu, Y.; Fernandez Garcia, M.; Xiao, J.; Han, S.-K.; Konkol, A.; Wagner, D. *LEAFY* is a pioneer transcription factor and licenses cell reprogramming to floral fate. *Nat. Commun.* **2021**, *12*, 626. <https://doi.org/10.1038/s41467-020-20883-w>.
- Lai, X.; Blanc-Mathieu, R.; GrandVuillemin, L.; Huang, Y.; Stigliani, A.; Lucas, J.; Thévenon, E.; Loue-Manifel, J.; Turchi, L.; Daher, H.; et al. The *LEAFY* floral regulator displays pioneer transcription factor properties. *Mol. Plant* **2021**, *14*, 829–837. <https://doi.org/10.1016/j.molp.2021.03.004>.
- Moyroud, E.; Kusters, E.; Monniaux, M.; Koes, R.; Parcy, F. *LEAFY* blossoms. *Trends Plant Sci.* **2010**, *15*, 346–352. <https://doi.org/10.1016/j.tplants.2010.03.007>.
- Chahtane, H.; Vachon, G.; Le Masson, M.; Thévenon, E.; Pérignon, S.; Mihajlovic, N.; Kalinina, A.; Michard, R.; Moyroud, E.; Monniaux, M.; et al. A Variant of *LEAFY* reveals its capacity to stimulate meristem development by inducing *RAX1*. *Plant J.* **2013**, *74*, 678–689. <https://doi.org/10.1111/tpj.12156>.
- Li, W.; Zhou, Y.; Liu, X.; Yu, P.; Cohen, J.D.; Meyerowitz, E.M. *LEAFY* controls auxin response pathways in floral primordium formation. *Sci. Signal.* **2013**, *6*, ra23. <https://doi.org/10.1126/scisignal.2003937>.
- Yamaguchi, N.; Wu, M.-F.; Winter, C.M.; Berns, M.C.; Nole-Wilson, S.; Yamaguchi, A.; Coupland, G.; Krizek, B.A.; Wagner, D. A molecular framework for auxin-mediated initiation of flower primordia. *Dev. Cell* **2013**, *24*, 271–282. <https://doi.org/10.1016/j.devcel.2012.12.017>.
- Zhao, D. The *ASK1* gene regulates B function gene expression in cooperation with *UFO* and *LEAFY* in *Arabidopsis*. *Development* **2001**, *128*, 2735–2748.

25. Gagne, J.M.; Downes, B.P.; Shiu, S.-H.; Durski, A.M.; Vierstra, R.D. The F-Box subunit of the SCF E3 complex is encoded by a diverse superfamily of genes in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 11519–11524. <https://doi.org/10.1073/pnas.162339999>.
26. Risseeuw, E.; Venglat, P.; Xiang, D.; Komendant, K.; Daskalchuk, T.; Babic, V.; Crosby, W.; Datla, R. An activated form of UFO alters leaf development and produces ectopic floral and inflorescence meristems. *PLoS ONE* **2013**, *8*, e83807. <https://doi.org/10.1371/journal.pone.0083807>.
27. Yamaguchi, N.; Wu, M.-F.; Winter, C.; Wagner, D. LEAFY and polar auxin transport coordinately regulate *Arabidopsis* flower development. *Plants* **2014**, *3*, 251–265. <https://doi.org/10.3390/plants3020251>.
28. Running, M.P.; Fletcher, J.C.; Meyerowitz, E.M. The WIGGUM gene is required for proper regulation of floral meristem size in *Arabidopsis*. *Development* **1998**, *125*, 2545–2553.
29. Chen, Q.; Atkinson, A.; Otsuga, D.; Christensen, T.; Reynolds, L.; Drews, G.N. The *Arabidopsis* FILAMENTOUS FLOWER gene is required for flower formation. *Development* **1999**, *126*, 2715–2726. <https://doi.org/10.1242/dev.126.12.2715>.
30. Sawa, S.; Watanabe, K.; Goto, K.; Kanaya, E.; Morita, E.H.; Okada, K. FILAMENTOUS FLOWER, a meristem and organ identity gene of *Arabidopsis*, encodes a protein with a zinc finger and hmg-related domains. *Genes Dev.* **1999**, *13*, 1079–1088. <https://doi.org/10.1101/gad.13.9.1079>.
31. Norberg, M.; Holmlund, M.; Nilsson, O. The BLADE ON PETIOLE genes act redundantly to control the growth and development of lateral organs. *Development* **2005**, *132*, 2203–2213. <https://doi.org/10.1242/dev.01815>.
32. Kanrar, S.; Onguka, O.; Smith, H.M.S. *Arabidopsis* inflorescence architecture requires the activities of KNOX-BELL homeodomain heterodimers. *Planta* **2006**, *224*, 1163–1173. <https://doi.org/10.1007/s00425-006-0298-9>.
33. Kanrar, S.; Bhattacharya, M.; Arthur, B.; Courtier, J.; Smith, H.M.S. Regulatory networks that function to specify flower meristems require the function of homeobox genes PENNYWISE and POUND-FOOLISH in *Arabidopsis*. *Plant J.* **2008**, *54*, 924–937. <https://doi.org/10.1111/j.1365-313X.2008.03458.x>.
34. Keller, T.; Abbott, J.; Moritz, T.; Doerner, P. *Arabidopsis* REGULATOR OF AXILLARY MERISTEMS1 controls a leaf axil stem cell niche and modulates vegetative development. *Plant Cell* **2006**, *18*, 598–611. <https://doi.org/10.1105/tpc.105.038588>.
35. Müller, D.; Schmitz, G.; Theres, K. Blind homologous R2R3 myb genes control the pattern of lateral meristem initiation in *Arabidopsis*. *Plant Cell* **2006**, *18*, 586–597. <https://doi.org/10.1105/tpc.105.038745>.
36. Lee, D.J.; Park, J.Y.; Ku, S.-J.; Ha, Y.-M.; Kim, S.; Kim, M.D.; Oh, M.-Ho.; Kim, J. Genome-wide expression profiling of ARABIDOPSIS RESPONSE REGULATOR 7 (ARR7) overexpression in cytokinin response. *Mol. Genet. Genomics.* **2007**, *277*, 115–137. <https://doi.org/10.1007/s00438-006-0177>.
37. Hareven, D.; Gutfinger, T.; Parnis, A.; Eshed, Y.; Lifschitz, E. The making of a compound leaf: Genetic manipulation of leaf architecture in tomato. *Cell* **1996**, *84*, 735–744. [https://doi.org/10.1016/S0092-8674\(00\)81051-X](https://doi.org/10.1016/S0092-8674(00)81051-X).
38. Bharathan, G.; Goliber, T.E.; Moore, C.; Kessler, S.; Pham, T.; Sinha, N.R. Homologies in leaf form inferred from KNOXI gene expression during development. *Science* **2002**, *296*, 1858–1860. <https://doi.org/10.1126/science.1070343>.
39. Hay, A.; Tsiantis, M. The genetic basis for differences in leaf form between *Arabidopsis thaliana* and its wild relative *Cardamine hirsuta*. *Nat. Genet.* **2006**, *38*, 942–947. <https://doi.org/10.1038/ng1835>.
40. Shani, E.; Burko, Y.; Ben-Yaakov, L.; Berger, Y.; Amsellem, Z.; Goldshmidt, A.; Sharon, E.; Ori, N. Stage-specific regulation of *Solanum lycopersicum* leaf maturation by class 1 KNOTTED1-LIKE HOMEBOX proteins. *Plant Cell* **2009**, *21*, 3078–3092. <https://doi.org/10.1105/tpc.109.068148>.
41. Hofer, J.; Turner, L.; Hellens, R.; Ambrose, M.; Matthews, P.; Michael, A.; Ellis, N. UNIFOLIATA regulates leaf and flower morphogenesis in Pea. *Curr. Biol.* **1997**, *7*, 581–587. [https://doi.org/10.1016/S0960-9822\(06\)00257-0](https://doi.org/10.1016/S0960-9822(06)00257-0).
42. Wang, H.; Chen, J.; Wen, J.; Tadege, M.; Li, G.; Liu, Y.; Mysore, K.S.; Ratet, P.; Chen, R. Control of compound leaf development by FLORICAULA/LEAFY ortholog SINGLE LEAFLET1 in *Medicago truncatula*. *Plant Physiol.* **2008**, *146*, 1759–1772. <https://doi.org/10.1104/pp.108.117044>.
43. Hofer, J.; Gourlay, C.; Michael, A.; Ellis, T.H.N. Expression of a Class 1 *Knotted1*-like homeobox gene is down-regulated in pea compound leaf primordia. *Plant Mol. Biol.* **2001**, *45*, 387–398. <https://doi.org/10.1023/A:1010739812836>.
44. Champagne, C.E.M.; Goliber, T.E.; Wojciechowski, M.F.; Mei, R.W.; Townsley, B.T.; Wang, K.; Paz, M.M.; Geeta, R.; Sinha, N.R. Compound leaf development and evolution in the legumes. *Plant Cell* **2007**, *19*, 3369–3378. <https://doi.org/10.1105/tpc.107.052886>.
45. Busch, A.; Gleissberg, S. EcFLO, a FLORICAULA-like gene from *Eschscholzia californica* is expressed during organogenesis at the vegetative shoot apex. *Planta* **2003**, *217*, 841–848. <https://doi.org/10.1007/s00425-003-1046-z>.
46. Zhou, C.; Han, L.; Li, G.; Chai, M.; Fu, C.; Cheng, X.; Wen, J.; Tang, Y.; Wang, Z.-Y. STM/BP-like KNOXI is uncoupled from ARP in the regulation of compound leaf development in *Medicago truncatula*. *Plant Cell* **2014**, *26*, 1464–1479. <https://doi.org/10.1105/tpc.114.123885>.
47. Taylor, S.; Hofer, J.; Murfet, I. *Stamina pistilloida*, the pea ortholog of *Fim* and *UFO*, is required for normal development of flowers, inflorescences, and leaves. *Plant Cell* **2001**, *13*, 31–46.
48. Dong, Z.; Zhao, Z.; Liu, C.; Luo, J.; Yang, J.; Huang, W.; Hu, X.; Wang, T.L.; Luo, D. Floral patterning in *Lotus japonicus*. *Plant Physiol.* **2005**, *137*, 1272–1282. <https://doi.org/10.1104/pp.104.054288>.
49. Nikolov, L.A.; Runions, A.; Das Gupta, M.; Tsiantis, M. Leaf development and evolution. *Curr. Top. Dev. Biol.* **2019**, *131*, 109–139.

50. Challa, K.R.; Rath, M.; Sharma, A.N.; Bajpai, A.K.; Davuluri, S.; Acharya, K.K.; Nath, U. Active suppression of leaflet emergence as a mechanism of simple leaf development. *Nat. Plants* **2021**, *7*, 1264–1275. <https://doi.org/10.1038/s41477-021-00965-3>.
51. Lincoln, C.; Long, J. A *Knotted1*-like homeobox gene in *Arabidopsis* is expressed in the vegetative meristem and dramatically alters leaf morphology when overexpressed in transgenic plants. *Plant Cell* **1994**, *6*, 1859–1876.
52. Pautot, V.; Dockx, J.; Hamant, O.; Kronenberger, J.; Grandjean, O.; Jublot, D.; Traas, J. *KNAT2*: Evidence for a link between *Knotted*-like genes and carpel development. *Plant Cell* **2001**, *13*, 1719–1734.
53. Piazza, P.; Bailey, C.D.; Cartolano, M.; Krieger, J.; Cao, J.; Ossowski, S.; Schneeberger, K.; He, F.; de Meaux, J.; Hall, N.; et al. *Arabidopsis thaliana* leaf form evolved via loss of *KNOX* expression in leaves in association with a selective sweep. *Curr. Biol.* **2010**, *20*, 2223–2228. <https://doi.org/10.1016/j.cub.2010.11.037>.
54. Lee, I.; Wolfe, D.S.; Nilsson, O.; Weigel, D. A *LEAFY* co-regulator encoded by *UNUSUAL FLORAL ORGANS*. *Curr. Biol.* **1997**, *7*, 95–104. [https://doi.org/10.1016/S0960-9822\(06\)00053-4](https://doi.org/10.1016/S0960-9822(06)00053-4).
55. Blein, T.; Pulido, A.; Vialette-Guiraud, A.; Nikovics, K.; Morin, H.; Hay, A.; Johansen, I.E.; Tsiantis, M.; Laufs, P. A conserved molecular framework for compound leaf development. *Science* **2008**, *322*, 1835–1839. <https://doi.org/10.1126/science.1166168>.
56. Bilsborough, G.D.; Runions, A.; Barkoulas, M.; Jenkins, H.W.; Hasson, A.; Galinha, C.; Laufs, P.; Hay, A.; Prusinkiewicz, P.; Tsiantis, M. Model for the regulation of *Arabidopsis thaliana* leaf margin development. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 3424–3429. <https://doi.org/10.1073/pnas.1015162108>.
57. Vialette-Guiraud, A.C.M.; Chauvet, A.; Gutierrez-Mazariegos, J.; Eschstruth, A.; Ratet, P.; Scutt, C.P. A conserved role for the NAM/MiR164 developmental module reveals a common mechanism underlying carpel margin fusion in monocarpous and syncarpous eurosids. *Front. Plant Sci.* **2016**, *6*, 1239. <https://doi.org/10.3389/fpls.2015.01239>.
58. Cheng, X.; Peng, J.; Ma, J.; Tang, Y.; Chen, R.; Mysore, K.S.; Wen, J. NO APICAL MERISTEM (MtNAM) regulates floral organ identity and lateral organ separation in *Medicago truncatula*. *New Phytol.* **2012**, *195*, 71–84. <https://doi.org/10.1111/j.1469-8137.2012.04147.x>.
59. Benlloch, R.; Navarro, C.; Beltrán, J.; Cañas, L.A. Floral development of the model legume *Medicago truncatula*: Ontogeny studies as a tool to better characterize homeotic mutations. *Sex. Plant Reprod.* **2003**, *15*, 231–241. <https://doi.org/10.1007/s00497-002-0157-1>.
60. Benlloch, R.; Berbel, A.; Ali, L.; Gohari, G.; Millán, T.; Madueño, F. Genetic control of inflorescence architecture in legumes. *Front. Plant Sci.* **2015**, *6*, 543. <https://doi.org/10.3389/fpls.2015.00543>.
61. Cheng, X.; Li, G.; Tang, Y.; Wen, J. Dissection of genetic regulation of compound inflorescence development in *Medicago truncatula*. *Development* **2018**, *145*, 158766. <https://doi.org/10.1242/dev.158766>.
62. Hecht, V.; Foucher, F.; Ferrándiz, C.; Macknight, R.; Navarro, C.; Morin, J.; Vardy, M.E.; Ellis, N.; Beltrán, J.P.; Rameau, C.; et al. Conservation of *Arabidopsis* flowering genes in model legumes. *Plant Physiol.* **2005**, *137*, 1420–1434. <https://doi.org/10.1104/pp.104.057018>.
63. Weigel, D.; Alvarez, J.; Smyth, D.R.; Yanofsky, M.F.; Meyerowitz, E.M. *LEAFY* controls floral meristem identity in *Arabidopsis*. *Cell* **1992**, *69*, 843–859. [https://doi.org/10.1016/0092-8674\(92\)90295-N](https://doi.org/10.1016/0092-8674(92)90295-N).
64. Benlloch, R.; d’Erfurth, I.; Ferrandiz, C.; Cosson, V.; Beltrán, J.P.; Cañas, L.A.; Kondorosi, A.; Madueño, F.; Ratet, P. Isolation of *mtpim* proves Tnt1 a useful reverse genetics tool in *Medicago truncatula* and uncovers new aspects of AP1-like functions in legumes. *Plant Physiol.* **2006**, *142*, 972–983. <https://doi.org/10.1104/pp.106.083543>.
65. Roque, E.; Serwatowska, J.; Cruz Rochina, M.; Wen, J.; Mysore, K.S.; Yenush, L.; Beltrán, J.P.; Cañas, L.A. Functional specialization of duplicated *AP3*-like genes in *Medicago truncatula*. *Plant J.* **2013**, *73*, 663–675. <https://doi.org/10.1111/tpj.12068>.
66. Benlloch, R.; Roque, E.; Ferrándiz, C.; Cosson, V.; Caballero, T.; Penmetsa, R.V.; Beltrán, J.P.; Cañas, L.A.; Ratet, P.; Madueño, F. Analysis of B function in legumes: PISTILLATA proteins do not require the PI motif for floral organ development in *Medicago truncatula*. *Plant J.* **2009**, *60*, 102–111. <https://doi.org/10.1111/j.1365-313X.2009.03939.x>.
67. Roque, E.; Fares, M.A.; Yenush, L.; Rochina, M.C.; Wen, J.; Mysore, K.S.; Gómez-Mena, C.; Beltrán, J.P.; Cañas, L.A. Evolution by gene duplication of *Medicago truncatula* PISTILLATA-like transcription factors. *J. Exp. Bot.* **2016**, *67*, 1805–1817. <https://doi.org/10.1093/jxb/erv571>.
68. Serwatowska, J.; Roque, E.; Gómez-Mena, C.; Constantin, G.D.; Wen, J.; Mysore, K.S.; Lund, O.S.; Johansen, E.; Beltrán, J.P.; Cañas, L.A. Two euAGAMOUS genes control C-function in *Medicago truncatula*. *PLoS ONE* **2014**, *9*, e103770. <https://doi.org/10.1371/journal.pone.0103770>.
69. Zhu, B.; Li, H.; Hou, Y.; Zhang, P.; Xia, X.; Wang, N.; Wang, H.; Mysore, K.S.; Wen, J.; Pei, Y.; et al. AGAMOUS and TERMINAL FLOWER controls floral organ identity and inflorescence development in *Medicago truncatula*. *J. Integr. Plant Biol.* **2019**, *61*, 917–923. <https://doi.org/10.1111/jipb.12799>.
70. Zhao, Y.; Liu, R.; Xu, Y.; Wang, M.; Zhang, J.; Bai, M.; Han, C.; Xiang, F.; Wang, Z.-Y.; Mysore, K.S.; et al. AGLF provides C-function in floral organ identity through transcriptional regulation of AGAMOUS in *Medicago truncatula*. *Proc. Natl. Acad. Sci. USA* **2019**, *116*, 5176–5181. <https://doi.org/10.1073/pnas.1820468116>.
71. Clark, S.E.; Jacobsen, S.E.; Levin, J.Z.; Meyerowitz, E.M. The *CLAVATA* and *SHOOT MERISTEMLESS* loci competitively regulate meristem activity in *Arabidopsis*. *Development* **1996**, *122*, 1567–1575.
72. Scofield, S.; Dewitte, W.; Murray, J.A.H. The *KNOX* Gene *SHOOT MERISTEMLESS* is required for the development of reproductive meristematic tissues in *Arabidopsis*. *Plant J.* **2007**, *50*, 767–781. <https://doi.org/10.1111/j.1365-313X.2007.03095.x>.
73. Schultz, E.A.; Haughn, G.W. *LEAFY*, a homeotic gene that regulates inflorescence development in *Arabidopsis*. *Plant Cell* **1991**, *3*, 771–781.

74. Zhou, C.; Han, L.; Hou, C.; Metelli, A.; Qi, L.; Tadege, M.; Mysore, K.S.; Wang, Z.-Y. Developmental analysis of a *Medicago truncatula* *smooth leaf margin1* mutant reveals context-dependent effects on compound leaf development. *Plant Cell* **2011**, *23*, 2106–2124. <https://doi.org/10.1105/tpc.111.085464>.
75. Meng, Y.; Liu, H.; Wang, H.; Liu, Y.; Zhu, B.; Wang, Z.; Hou, Y.; Zhang, P.; Wen, J.; Yang, H.; et al. *HEADLESS*, a *WUSCHEL* homolog, uncovers novel aspects of shoot meristem regulation and leaf blade development in *Medicago truncatula*. *J. Exp. Bot.* **2019**, *70*, 149–163. <https://doi.org/10.1093/jxb/ery346>.
76. Wang, H.; Xu, Y.; Hong, L.; Zhang, X.; Wang, X.; Zhang, J.; Ding, Z.; Meng, Z.; Wang, Z.-Y.; Long, R.; et al. *HEADLESS* regulates auxin response and compound leaf morphogenesis in *Medicago truncatula*. *Front. Plant Sci.* **2019**, *10*, 1024. <https://doi.org/10.3389/fpls.2019.01024>.
77. Zhang, S.; Sandal, N.; Polowick, P.L.; Stiller, J.; Stougaard, J.; Fobert, P.R. *Proliferating Floral Organs (Pfo)*, a *Lotus japonicus* gene required for specifying floral meristem determinacy and organ identity, encodes an F-Box Protein. *Plant J.* **2003**, *33*, 607–619. <https://doi.org/10.1046/j.1365-313X.2003.01660.x>.
78. Roth, O.; Alvarez, J.P.; Levy, M.; Bowman, J.L.; Ori, N.; Shani, E. The *KNOXI* transcription factor *SHOOT MERISTEMLESS* regulates floral fate in *Arabidopsis*. *Plant Cell* **2018**, *30*, 1309–1321. <https://doi.org/10.1105/tpc.18.00222>.
79. Cosson, V.; Eschstruth, A.; Ratet, P. *Medicago truncatula* transformation using leaf explants. *Methods Mol. Biol.* **2015**, *1223*, 43–56. https://doi.org/10.1007/978-1-4939-1695-5_4.
80. Kakar, K.; Wandrey, M.; Czechowski, T.; Gaertner, T.; Scheible, W.-R.; Stitt, M.; Torres-Jerez, I.; Xiao, Y.; Redman, J.C.; Wu, H.C.; et al. A community resource for high-throughput quantitative RT-PCR analysis of transcription factor gene expression in *Medicago truncatula*. *Plant Methods* **2008**, *4*, 18. <https://doi.org/10.1186/1746-4811-4-18>.
81. Nikovics, K.; Blein, T.; Peaucelle, A.; Ishida, T.; Morin, H.; Aida, M.; Laufs, P. The balance between the *MIR164A* and *CUC2* genes controls leaf margin serration in *Arabidopsis*. *Plant Cell* **2006**, *18*, 2929–2945. <https://doi.org/10.1105/tpc.106.045617>.
82. Ferrándiz, C.; Liljegren, S.J.; Yanofsky, M.F. Negative Regulation of the *SHATTERPROOF* genes by *FRUITFULL* during *Arabidopsis* fruit development. *Science* **2000**, *289*, 436–438. <https://doi.org/10.1126/science.289.5478.436>.