



HAL
open science

**Modeling the Influence of Invasive Fish Species
Perccottus glenii (Dybowski, 1877) on the Distribution
of Newts in Eastern Europe, Exemplified by *Lissotriton
vulgaris* (Linnaeus, 1758) and Preserved *Triturus
crystatus* (Laurenti, 1768), Using a GIS Approach**

Oksana Nekrasova, Oleksii Marushchak, Mihails Pupins, Volodymyr Tytar,
Jean-Yves Georges, Kathrin Theissinger, Andris Čeirāns, Arturs Skute

► **To cite this version:**

Oksana Nekrasova, Oleksii Marushchak, Mihails Pupins, Volodymyr Tytar, Jean-Yves Georges, et al.. Modeling the Influence of Invasive Fish Species *Perccottus glenii* (Dybowski, 1877) on the Distribution of Newts in Eastern Europe, Exemplified by *Lissotriton vulgaris* (Linnaeus, 1758) and Preserved *Triturus cristatus* (Laurenti, 1768), Using a GIS Approach. *Biology and Life Sciences Forum*, 2022. hal-03846931

HAL Id: hal-03846931

<https://cnrs.hal.science/hal-03846931v1>

Submitted on 10 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Modeling the Influence of Invasive Fish Species *Percottus glenii* (Dybowski, 1877) on the Distribution of Newts in Eastern Europe, Exemplified by *Lissotriton vulgaris* (Linnaeus, 1758) and Preserved *Triturus cristatus* (Laurenti, 1768), Using a GIS Approach [†]

Oksana Nekrasova ^{1,2}, Oleksii Marushchak ^{1,*}, Mihails Pupins ², Volodymyr Tytar ¹, Jean-Yves Georges ³, Kathrin Theissingner ⁴, Andris Čeirāns ² and Arturs Skute ²

¹ I. I. Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, 01030 Kyiv, Ukraine; oneks22@gmail.com (O.N.); vtytar@gmail.com (V.T.)

² Department of Ecology, Institute of Life Sciences and Technologies, Daugavpils University, LV-5401 Daugavpils, Latvia; mihails.pupins@gmail.com (M.P.); andris.ceirans@lu.lv (A.Č.); arturs.skute@du.lv (A.S.)

³ Université de Strasbourg, CNRS, IPHC UMR7178, Strasbourg, France; jean-yves.georges@iphc.cnrs.fr

⁴ LOEWE Center for Translational Biodiversity Genomics, TBG - Senckenberg Nature Research Institute, Frankfurt, Germany; kathrin.theissingner@senckenberg.de

* Correspondence: ecopelobates@gmail.com; Tel.: +380-964882670

[†] Presented at the 2nd International Electronic Conference on Diversity (IECD 2022) – New Insights into the Biodiversity of Plants, Animals and Microbes, 15–30 March 2022; Available online: <https://sciforum.net/event/IECD2022>.

Citation: Nekrasova, O.; Marushchak, O.; Pupins, M.; Tytar, V.; Georges, J.-Y.; Theissingner, K.; Čeirāns, A.; Skute, A. Modeling the Influence of Invasive Fish Species *Percottus glenii* (Dybowski, 1877) on the Distribution of Newts in Eastern Europe, Exemplified by *Lissotriton vulgaris* (Linnaeus, 1758) and Preserved *Triturus cristatus* (Laurenti, 1768), Using a GIS Approach. *Biol. Life Sci. Forum* **2022**, *2*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor:

Published:

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Along with decrease in numbers among amphibians in Eastern Europe nowadays newts especially suffer from invasive predatory fish and turtles. Such aggressive predator as *P. glenii* has recently occupied water bodies important for newts' breeding. The fish eat eggs, larvae and even adult individuals. Using the ecological niche approach, it was showed that for built distribution models based on certain climatic scenario Coefficient of determination (R^2) between Chinese sleeper and preserved *T. cristatus* with *L. vulgaris* to 2090 will grow from 44% to 66% in Eastern Europe highlighting expanding of new territories by the fish as a real threat to biodiversity.

Keywords: climate change; GIS-analysis; modelling; invasive species; amphibians; Urodela; Europe; niche modelling

1. Introduction

Recently, a decrease in the numbers of amphibians in Eastern Europe was reported multiple times. Suitable habitats that are important for amphibians at certain stages of their lives shrink and degrade due to climate change [1,2], anthropogenic impact and the animals suffer from pollution and appearance of alien invasive species [2]. Alien invasive species often appear and adapt in new places far from their natural range for example as a result of uncontrolled release into the wild from the terrariums or being accidentally transported as a result of developing aquaculture [3–6]. Thus, the appearance of Chinese sleeper – *Percottus glenii* (Dybowski, 1877) in Europe in the water bodies of cities can also be explained by the deliberate release of amateur aquarists and fishermen in the early 20th century. Until now *P. glenii* is considered ornamental, and due to its ability to survive in any conditions, aggressiveness, flexible and wide range of food and ecological plasticity, this species has spread quite widely both in the North and in the South of Europe [7–11].

In Europe since 2016 Chinese sleeper is included in the list of Invasive Alien Species of Union concern (the Union list) [1].

According to the literature resources *P. glenii* is a great threat for local biota. It was reported, that its diet consists of macro- and micro crustaceans (Ostracoda, Copepoda, Cladocera, Isopoda, Amphipoda, Gastropoda), highly mobile invertebrates such as Coleoptera and Heteroptera [12]. Additionally, to direct preying this species also affects the feeding base for local inhabitants acting as a trophic competitor and a predator competing for the resources. Habitats heavily invaded by *P. glenii* are usually associated with lower fish species richness and diversity [13,14]. Moreover *P. glenii* can be a vector for transporting of several species of parasites, that can lead to significant damage in local balance of bioecology or aquaculture [15–17]. Also, by feeding on macro- and micro-invertebrates *P. glenii* effectively reduces the transport of nitrogen from aquatic to terrestrial environments [18].

Of all amphibian species newts are especially vulnerable and suffer from climate change, anthropogenic influence and, as a result, the emergence of new invasive predator species, diseases and habitat loss. The two most widely distributed species of newts in Eastern Europe—preserved in European Union crested newt *Triturus cristatus* (Laurenti, 1768) and smooth newt *Lissotriton vulgaris* (Linnaeus, 1758) are the two species that can demonstrate potential connection between decrease in their numbers and expansive moving of *P. glenii* to new territories in the investigated region.

Therefore, the purpose of our work was to identify the most potentially suitable habitats for these species in Eastern Europe, as well as to make a prediction about the threats and expansion of the *P. glenii* as aggressive predator on native species of newts.

2. Materials and Methods

Occurrence data on the studied species referred to the entire territory of Europe in order to increase the accuracy of models built, despite the fact that main focus of the investigation was to evaluate species potential future overlap in Eastern part of Europe. The data was collected from the original datasets [19], collection materials (I. I. Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, Kyiv; Department of Ecology, Institute of Life Sciences and Technologies, Daugavpils University, Daugavpils, Latvia), literature resources [7–10,20,21], GBIF databases [22–24]—all non-duplicate. To account for sampling bias, we used the nearest neighbor distance ('ntbox' package in R [25]) method to thin the data, where occurrence points that were ≤ 0.1 units away from each other were removed to avoid errors due to spatial autocorrelation. As a result, the number of points significantly decreased to total of 2096 for *L. vulgaris*—(127,715 initial records within Europe downloaded from GBIF), *T. cristatus*—2277 (132,121 initial records within Europe downloaded from GBIF), *P. glenii*—944 (842 initial records within Europe downloaded from GBIF).

Ecological niche modeling and species distribution modelling (SDM) methods have been used to determine the potential home range of invasive species in new environments (MaxEnt [26] with 25 replicates, DivaGis (Bioclim). We used 35 bioclimatic variables from the CliMond dataset [27,28]. Of 35 bioclimatic variables (V), highly correlated (>0.7) predictors were removed using the 'virtualspecies' package in R, resulting in a selection of 18 for 1975 (1970–2000), 2090 (2081–2100). The area under the receiver operating characteristic (ROC) curve (AUC) was used for assessing the discriminatory capacity of the models: $AUC > 0.85$ is considered excellent. We carried out separate modelling for each set of factors and for each species. Logistic output format was used to describe the relative probability of presence, which is a continuous habitat suitability (HS) range between 0 (unsuitable) and 1 (the most suitable). Using the Coefficient of determination, we assessed statistically significant to assess similarity ($p < 0.05$) between the predicted HS map obtained through Maxent for every single target species across the study area. GIS-modelling was accomplished using SagaGis, DivaGis, QGis [12]. Statistical processing of the obtained data was carried out using Statistica for Windows v.8.0.

3. Results

According to our own data and published information, in spring and in the first half of summer, newts are found mainly in small reservoirs, streams and ditches, where they spawn. However, after active expansion of *P. glenii* throughout Europe, this predator appeared in the same biotopes. Being less whimsical and more ecologically plastic, rather active predator *P. glenii* is capable of displacing amphibians sensitive to water quality. For example, according to our data, in the Volyn, Rivne, Zhytomyr and Kyiv regions of Ukraine during field expeditions in 2017–2021 newts were found only in water bodies where *P. glenii* was not registered.

The reduction in the number and sometimes the disappearance of newts' micropopulations in the south of the range have led to the fact, that they were included in the Red Book in some countries, for example *T. cristatus* in Ukraine, and *L. vulgaris* is now a contender to be listed in Red Book of Ukraine in the nearest future [12,24,29].

An analysis of the distribution of newts in Eastern Europe showed that the continental biographical zone has the highest priority in terms of potentially suitable territories. In the western part of their range, newts can "descend" into the steppe zone along the eco-corridors of large rivers, such as the Danube and the Dniester. *P. glenii* has a similar range, but more fragmented. According to our data, the ranges of newts will be greatly reduced by 2090, while the range of the fish, on the contrary, will increase. We have been observing the rapid spread of *P. glenii* since 2010 in the continental zones of Ukraine and in the east of Latvia. In most cases temporary water bodies are left uninvaded playing role of place, where many amphibians have time to breed, but, on the contrary, Chinese sleeper does not have time to breed and establish.

As a result of the modelling, it was revealed that the complex of factors associated with average temperature amplitudes for different periods and humidity has the greatest influence on the spatial distribution of the studied species and the Radiation of coldest quarter (27 V) factor complements this complex of factors and affects the humidity of habitats. Naturally, according to a number of factors, *P. glenii* as an aquatic animal differs from semi-terrestrial amphibians (Figure 1).




Figure 1. Graph of most contributing predictors (V (bio), %) for building SSDM 3 species of Eastern Europe: (A)—Percent contribution and (B)—Permutation importance; LV—*L. vulgaris*; TC—*T. cristatus*; PG—*P. glenii*; 1 V—Annual mean temperature (°C); 8 V—Mean temperature of wettest quarter

(°C); 18 V—Precipitation of warmest quarter (mm); 27 V—Radiation of coldest quarter ($W m^{-2}$); 34 V—Mean moisture index of warmest quarter.

Moreover, *P. glenii* appeared to be the most tolerant and resistant to the temperature regime with such factor as Annual mean temperature having its optimal values in the range from +2 to +12 °C (Figure 2).




Figure 2. Maxent model created using only the variable (1 V)—Annual mean temperature (°C, CliMond) from: (A)—*L. vulgaris*; (B)—*T. cristatus*; (C)—*P. glenii*; (D)—dependence graphs of grid SDM different species (HS, Coefficient of determination % R^2 , Saga): LV—*L. vulgaris*; TC—*T. cristatus*; PG—*P. glenii*.

Statistical analysis of the results showed high accuracy of the model. Despite the sufficient scatter and the existing accumulations of locality points in certain areas due to the characteristics of the type of range of the species under study, the data on test locations are in good agreement with the predicted dynamics calculated for the test data obtained from the Maxent distribution. The high area under the curve (AUC) > 0.85 confirms our assumptions.

Analyzing the obtained models, we came to the conclusion that the greatest similarity of SDM HS is achieved by newt species, Coefficient of determination—67.78% (Figure 2D, TC/LV) and this is explained by the fact that they can live together in similar biotopes. Although earlier, at the end of the 20th century, biotopic specialization and different calendar periods in reproduction were observed among amphibians. Recently, suitable reservoirs for amphibian breeding have become smaller, and breeding times have also shifted (phenology has shifted), and therefore it is common to see several species of amphibians breeding in the same reservoir and at the same period. This leads to the fact that

not all amphibians will be able to breed in such conditions, especially newts. According to GIS modeling data, the range of newts and their abundance will be greatly reduced by 2090, especially in the south. *T. cristatus* will be especially affected. As for the Chinese sleeper, the number and range of which will increase significantly and by 2090 potentially suitable sites will increase significantly and the Coefficient of determination will increase significantly—to 44% to 66% (Figure 2D, TC/PG and LV/PG) with *T. cristatus* and *L. vulgaris* respectively.

4. Discussion

The results of the GIS-analysis are supported in recent years by numerous field investigations when lakes, ponds and streams heavily invaded by *P. glenii* showed either no signs of *L. vulgaris* or *T. cristatus*, or only separate individuals were found even during the peak of breeding season. However, the Chinese sleeper can also serve as a food base for some species of reptiles, for example, *Natrix tessellata* (Laurenti, 1768) [30], rare and preserved in Europe *Emys orbicularis* (Linnaeus, 1758) [31] and *Bombina bombina* (Linnaeus, 1761), or the invasive exotic species *Trachemys scripta* (Thunberg in Schoepff, 1792) [4,25,26,32]. Moreover, in depleted anthropogenic urban ecosystems, where water quality does not allow native fish species to survive, this invasive species, like other aquarium fish, can occupy its ecological niche and act as a biological method of combating dangerous blood-sucking invasive insects [5,27,33].

5. Conclusions

As a result of our research, it was found that in the future, due to climatic and anthropogenic changes, two most widely distributed species of newts—*L. vulgaris* and preserved in European Union *T. cristatus*—will suffer, their range and abundance will decrease, and the expansion of such an invasive Chinese sleeper will intensify. Therefore, it is necessary to establish management plans for the protection of native amphibian species taking into account heavy influence of invasive species adequately taking their further expansion into strict control.

Author Contributions: Conceptualization, O.N., M.P., J.-Y.G. and V.T.; Data curation, O.N., M.P., A.S. and A.Č.; Formal analysis, O.N., V.T., O.M. and K.T.; Funding acquisition, M.P., A.S. and A.Č.; Investigation, O.N., M.P., O.M. and A.Č.; Methodology, O.N. and V.T.; Project administration, A.S., O.N., A.Č. and M.P.; Resources, O.N., M.P. and A.Č.; Software, O.N. and V.T.; Supervision, O.N., V.T. and M.P.; Validation, O.N., V.T., A.Č. and M.P.; Visualization, O.N., V.T. and M.P.; Writing—original draft and Writing—review & editing, O.N., V.T., M.P., A.Č., J.-Y.G., K.T., O.M. and A.S. All authors have read and agreed to the published version of the manuscript.

Funding: The research was partly funded by the project Emys-R <https://emysr.cnrs.fr> through the 2020-2021 Biodiversa & Water JPI joint call for research proposals, under the BiodivRestore ERA-Net CO-FUND programme, and with the funding organisations Agence Nationale de la Recherche (ANR, France), Bundesministerium für Bildung und Forschung (BMBF, Germany), State Education Development Agency (VIAA, Latvia), and National Science Center (NSC, Poland) and by the project "Ecological and socioeconomic thresholds as a basis for defining adaptive management triggers in Latvian pond aquaculture" (Izp-2021/1-0247). We thank the project "The mobile complex of Daugavpils University pond aquaculture scientific laboratories" (#16-00-F02201-000002) for the possibility of using the mobile complex of scientific laboratories for the research purposes and the project "Modern risks of degradation of ecosystems of Ukraine on the example of model zoocenoses: analysis of factors in terms of biological safety" (project #0122U000708).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are openly available in [*Triturus cristatus* (Laurenti, 1768) in GBIF Secretariat (2022); GBIF: Copenhagen, Denmark, 2022. <https://doi.org/10.15468/dl.wsmeiw> (accessed on 1 February 2022); *Lissotriton vulgaris* (Linnaeus, 1758) in GBIF Secretariat (2022); GBIF: Copenhagen, Denmark, 2022.

<https://doi.org/10.15468/dl.j52e2c> (accessed on 1 February 2022) and *Perccottus glenii* (Dybowski, 1877) in GBIF Secretariat (2022); GBIF: Copenhagen, Denmark, 2022. <https://doi.org/10.15468/dl.4g66a3> (accessed on 1 February 2022)]

Conflicts of Interest: The authors declare no conflict of interest.

References

- Invasive Alien Species of Union Concern. Available online: https://ec.europa.eu/environment/nature/pdf/IAS_brochure_species.pdf (accessed on 7 January 2022).
- Tytar, V.; Nekrasova, O.; Pupina, A.; Pupins, M.; Oskyrko, O. Long-term bioclimatic modelling the distribution of the fire-bellied toad, *Bombina bombina*, under the influence of global climate change. *Vest. Zool.* **2018**, *52*, 553–556.
- Nekrasova, O.; Marushchak, O.; Pupins, M.; Skute, A.; Tytar, V.; Čeirāns, A. Distribution and Potential Limiting Factors of the European Pond Turtle (*Emys orbicularis*) in Eastern Europe. *Diversity* **2021**, *13*, 280.
- Nekrasova, O.; Tytar, V.; Pupins, M.; Čeirāns, A. Range expansion of the alien red-eared slider *Trachemys scripta* (Reptilia, Testudines) in Eastern Europe, with special reference to Latvia and Ukraine. *Bioinvasions Rec.* **2022**, *11*, in press.
- Nekrasova, O.; Tytar, V.; Pupins, M.; Čeirāns, A.; Marushchak, O.; Skute, A. A GIS Modeling Study of the Distribution of Viviparous Invasive Alien Fish Species in Eastern Europe in Terms of Global Climate Change, as Exemplified by *Poecilia reticulata* Peters, 1859 and *Gambusia holbrooki* Girard, 1859. *Diversity* **2021**, *13*, 385.
- Kuybida, V.V.; Nekrasova, O.D.; Kutsokon, Y.K.; Lopatynska, V.V. Summer fish kills in the Kaniv Reservoir. *Hydrobiol. J.* **2019**, *55*, 103–106.
- Reshetnikov, A.N. Influence of introduced fish *Percottus glenii* (Odontobutidae, Pisces). *Zhurnal. Obs. Biol.* **2001**, *62*, 352–361.
- Reshetnikov, A.N. Does rotan *Percottus glenii* (Perciformes: Odontobutidae) eat the eggs of fish and amphibians? *J. Ichthyol.* **2008**, *48*, 336–344.
- Reshetnikov, A.N. Spatio-temporal dynamics of the expansion of rotan *Percottus glenii* from West-Ukrainian center of distribution and consequences for European freshwater ecosystems. *Aquat. Invasions* **2013**, *8*, 193–206.
- Reshetnikov, A.N.; Manteifel, Y.B. Newt-fish interactions in Moscow province: A new predatory fish colonizer, *Percottus glenii*, transforms matapopulations of newts, *Triturus vulgaris* and *T. cristatus*. *Adv. Amphib. Res. Former Sov. Union* **1997**, *2*, 1–12.
- Pupina, A.; Pupins, M.; Nekrasova, O.; Tytar, V.; Kozynenko, I.; Marushchak, O. Species distribution modelling: *Bombina bombina* (Linnaeus, 1761) and its important invasive threat *Percottus glenii* (Dybowski, 1877) in Latvia under global climate change. *J. Environ. Res. Eng. Manag.* **2018**, *74*, 79–86.
- Košco, J.; Manko, P.; Miklisová, D.; Košuthová, L. Feeding ecology of invasive *Percottus glenii* (Perciformes, Odontobutidae) in Slovakia. *Czech. J. Anim. Sci.* **2008**, *53*, 479–486.
- Košco, J.; Lusk, S.; Halacka, K.; Lusková, V. The expansion and occurrence of the Amur sleeper (*Percottus glenii*) in eastern Slovakia. *Folia Zool.* **2003**, *52*, 329–336.
- Kutsokon, I.; Tkachenko, M.; Bondarenko, O.; Pupins, M.; Snigirova, A.; Berezovska, V.; Ceirans, A.; Kvach, Y. The role of invasive Chinese sleeper *Percottus glenii* Dybowski, 1877 in the Ilgas Nature Reserve ecosystem: An example of a monospecific fish community. *Bioinvasions Rec.* **2021**, *10*, 396–410.
- Kvach, Y.; Kutsokon, I.; Roman, A.; Ceirans, A.; Pupins, M.; Kirjusina, M. Parasite acquisition by the invasive Chinese sleeper (*Percottus glenii* Dybowski, 1877) (Gobiiformes: Odontobutidae) in Latvia and Ukraine. *J. Appl. Ichthyol.* **2020**, *36*, 785–794.
- Jorgensen, A.; Sterud, E. The marine pathogenic genotype of *Spiroucleus barkhanus* from farmed salmonids redescribed as *Spiroucleus salmonicida* n. sp. *J. Eukaryot. Microbiol.* **2006**, *53*, 531–541.
- Williams, C.F.; Lloyd, D.; Poynton, S.L.; Jorgensen, A.; Millet, C.O.M.; Cable, J. *Spiroucleus* species: Economically-important fish pathogens and enigmatic single-celled eukaryotes. *J. Aquac. Res. Dev.* **2011**, *1*, S2.
- Reshetnikov, A.N. The introduced fish, rotan (*Percottus glenii*), depresses population of aquatic animals (macrovertebrates, amphibians, and fish). *Hydrobiologia* **2003**, *510*, 83–90.
- Nekrasova, O.D.; Tytar, V.M.; Kuybida, V.V. *GIS Modeling of Climate Change Vulnerability of Amphibians and Reptiles in Ukraine*; Schmalhauzen Institute of Zoology NAS: Kyiv, Ukraine, 2019; 204p.
- Kutsokon, I. The Chinese sleeper (*Percottus glenii* Dybowski, 1877) in Ukraine: New data on distribution. *J. Appl. Ichthyol.* **2017**, *33*, 1100–1107.
- Kvach, Y.; Kutsokon, Y. The non-indigenous fishes in the fauna of Ukraine: A potentia and actum. *Bioinvasions Rec.* **2017**, *6*, 269–279.
- GBIF. *Triturus cristatus* (Laurenti, 1768) in GBIF Secretariat (2022); GBIF: Copenhagen, Denmark, 2022. <https://doi.org/10.15468/dl.wsmejw>.
- GBIF. *Lissotriton vulgaris* (Linnaeus, 1758) in GBIF Secretariat (2022); GBIF: Copenhagen, Denmark, 2022. <https://doi.org/10.15468/dl.j52e2c>.
- GBIF. *Percottus glenii* (Dybowski, 1877) in GBIF Secretariat (2022); GBIF: Copenhagen, Denmark, 2022. <https://doi.org/10.15468/dl.4g66a3>.
- Osorio-Olvera, L.; Lira-Noriega, A.; Soberón, J.; Peterson, A.T.; Falconi, M.; Contreras-Díaz, R.G. nbox: An R package with graphical user interface for modelling and evaluating multidimensional ecological niches. *Methods Ecol. Evol.* **2020**, *11*, 1199–1206.

26. Phillips, S.J. A brief tutorial on Maxent. *ATT Res.* **2005**, *190*, 231–259. 260
27. Kriticos, D.J.; Jarošik, V.; Ota, N. Extending the suite of bioclim variables: A proposed registry system and case study using principal components analysis. *Methods Ecol. Evol.* **2014**, *5*, 956–960. 261
28. CliMond. Available online: <https://www.climond.org/> (accessed on 27 December 2020). 262
29. Nekrasova, O.; Smirnov, N.; Marushchak, O.; Korshunov, O.; Kotserzhynska, I. Change in the conservation status and location of the newt *Triturus cristatus* (Amphibia, Salamandridae) in Ukraine. *Mater. Red Book Ukraine. Fauna/Ser. Conserv. Biol. Ukr.* **2019**, *7*, 378–386. 263
30. Nekrasova, O.D.; Gavis, G.G.; Kuybida, V.V. Changes in the northern border of the home range of the dice snake, *Natrix tessellata* (Reptilia, Colubridae), in the Dnipro Basin (Ukraine). *Vest. Zool.* **2013**, *47*, 67–71. 264
31. Nekrasova, O.; Yanish, Y.; Tytar, V.; Pupins, M. GIS-modeling of the range shifts of the sub-fossil and extant European pond turtle (*Emys orbicularis*) in Eastern Europe in holocene. *Diversity* **2019**, *11*, 121. 265
32. Pupins, M.; Pupina, A. Reciprocal predation between preserved and invasive species: Adult *Bombina bombina* predate young whitebaits of alien fish *Percocottus glenii*. *Act. Biol. Univ. Daug.* **2018**, *18*, 217–223. 266
33. Kozynenko, I.I.; Tytar, V.M. Bioclimatic modeling of the European distribution of the invasive Asian tiger mosquito, *Aedes (Stegomyia) albopictus* (Skuse, 1895), with special reference to Ukraine. *Rep. Nat. Acad. Sc. Ukr.* **2020**, *3*, 88–93. 267

268
269
270
271
272
273
274