



HAL
open science

Modelling Keloids Dynamics: A Brief Review and New Mathematical Perspectives

Raluca Eftimie, Gwenaël Rolin, Olusegun Adebayo, Stéphane Urcun, Franz Chouly, Stéphane Pierre Alain Bordas

► **To cite this version:**

Raluca Eftimie, Gwenaël Rolin, Olusegun Adebayo, Stéphane Urcun, Franz Chouly, et al.. Modelling Keloids Dynamics: A Brief Review and New Mathematical Perspectives. 2022. hal-03897527

HAL Id: hal-03897527

<https://hal.science/hal-03897527>

Preprint submitted on 13 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modelling Keloids Dynamics: A Brief Review and New Mathematical Perspectives

R. Eftimie^{1,*}, G. Rolin^{2,3}, O. Adebayo¹, S. Urcun⁵, F. Chouly^{3,6,7} and S.P.A. Bordas⁵

¹Laboratoire de Mathématiques de Besançon, Université Bourgogne Franche-Comté, 25000, Besançon, France

²INSERM CIC-1431, CHU Besançon, 25000, Besançon, France

³INSERM, EFS BFC, UMR1098, RIGHT Interactions Greffon-Hôte-Tumeur/Ingénierie Cellulaire et Génique, Université Bourgogne Franche-Comté, F-25000, Besançon, France

⁴Institut de Mathématiques de Bourgogne, Université Bourgogne Franche-Comté, 21078, Dijon, France

⁵Institute for Computational Engineering, Faculty of Science, Technology and Communication, University of Luxembourg, Esch-sur-Alzette, Luxembourg

⁶Center for Mathematical Modelling and Department of Mathematical Engineering, University of Chile and IRL 2807 – CNRS, Santiago, Chile

⁷Departamento de Ingeniería Matemática, C^PMA, Universidad de Concepción, Casilla 160-C, Concepción, Chile

Correspondence*:

Corresponding Author: Raluca Eftimie
raluca.eftimie@univ-fcomte.fr

ABSTRACT

The keloids are fibroproliferative disorders described by an excessive growth of fibrotic tissue, which also invades adjacent areas (beyond the original wound borders). Since these disorders are specific to humans (no other animal species naturally develop keloid-like tissue), the experimental *in vivo/in vitro* research has not lead to significant advances in this field. One possible approach could be to combine *in vitro* human models with calibrated *in silico* mathematical approaches (i.e., models and simulations) to generate new testable biological hypotheses related to biological mechanisms and improved treatments. Since these combined approaches do not really exist for keloid disorders, in this brief review we start by summarising the biology of these disorders, then present various types of mathematical and computational approaches used for related disorders (i.e., wound healing and solid tumours), followed by a discussion of the very few mathematical and computational models published so far to study various inflammatory and mechanical aspects of the keloids. We conclude by discussing the open problems and opportunities offered in the context of keloid disorders by such combined *in vitro/in silico* approaches, and the need for multi-disciplinary research to enable clinical progress.

Keywords: keloid disorders, review, mathematical modelling and computational approaches, modelling cancer and wound healing, inflammation, biomechanics

1 INTRODUCTION

Since 2018, World Health Organization defines keloids as a member of a group of disorders “characterized by increased deposition of fibrous tissue in the skin and subcutaneous tissue” (WHO, 2022). Accordingly, keloids are considered not just pathological scars, but rather the outcome of an abnormal wound healing process with features similar to that of chronic inflammatory diseases and cancer (Tan et al., 2019). Clinically, keloids present as a fibrotic tissue that proliferates beyond the primary injured area and persist over time without natural regression. This fibrotic tissue also exhibits inter-lesional and intra-lesional heterogeneity (Limandjaja et al., 2020a). Keloids may cause pain and pruritus, and seriously affect patient quality of life when they are located in visible areas (Butler et al., 2008). Keloids may result from different skin injuries, including surgery, burns, trauma, piercing, and folliculitis, though their location appears mainly confined to chest, shoulders, neck and ears (Ogawa, 2011). Today, the lack of effective treatments clearly reflects our poor understanding of keloid pathogenesis, though keloid histopathology has been extensively described in the past decades (Jumper et al., 2015). Despite this, keloid aetiology and pathogenesis remain largely unknown.

Since keloid disorders are exclusive to humans, existent *in vivo* and *in vitro* studies that are based on different animals are not quite relevant to humans (as skin physiology, immunology and cell and tissue biomechanics are different between animals and humans (Limandjaja et al., 2020a)). Therefore, new *in vitro* human biological models are required to advance the research in this field (Suttho et al., 2017). These *in vitro* human models could be complemented by *in silico* approaches focused on mathematical models of these biological systems and computer simulations. These combined *in vitro/in silico* perspectives have become quite common for studies on wound healing and cancer evolution (disorders that have similar biological mechanisms as the keloids (Limandjaja et al., 2021; Tan et al., 2019)), but they are not usually used for keloid disorders.

The goal of this brief review is to present the complex problem of keloid disorders in the context of related disorders such as wound healing and tumour growth and spread, and to propose new multi-disciplinary approaches (that include mathematical modelling and simulations), which could bring new perspectives and solutions. To this end, we structure this review as follows. In Section 2 we discuss the biology of keloid disorders, with an emphasis on inflammatory and bio-mechanical aspects. In Section 3, we review a variety of mathematical models that have been developed over the last decades to describe and investigate wound healing, solid tumours, and keloid scars. In Section 4, we discuss some open problems in the context of modelling and computational approaches for these multi-scale disorders. We conclude in Section 5 with a summary of the main aspects reviewed in this study and some future perspectives.

2 BIOLOGICAL ASPECTS OF KELOID DISORDERS: INFLAMMATION AND BIOMECHANICS

The set of causes (aetiology) for keloid formation is still not fully understood. However, several studies have postulated different hypotheses on the pathogenesis of keloid formation (Limandjaja et al., 2020b; Bayat et al., 2005; Limandjaja et al., 2021). These hypotheses, which touch on inflammation and/or bio-mechanics (Limandjaja et al., 2020a), range from the keloid triad hypothesis (which encompasses three risk factors that could trigger the keloids, namely genetics, inflammation and external factors such as surgery), to the incomplete malignancy hypothesis (suggesting that the keloids could be a type of incomplete malignancy that has undergone only a few tumorigenic changes), the viral hypothesis (suggesting that the keloids could be triggered by a normally quiescent, unknown virus), the stiffness gap hypothesis (suggesting

that the keloids could be the result of an enlarged gap between the ECM stiffness and cellular stiffness), the immuno-nutritional hypothesis (suggesting that keloids could be the result of nutritional imbalance that promotes prolonged inflammation), or the metabolic hypothesis (suggesting that keloids could be the result of an increased metabolic activity of fibroblasts). We mentioned these different hypotheses here since some of them have been investigated through mathematical models, as discussed in Section 3 below. However, in the following we focus on the “keloid triad” that combines both intrinsic inflammatory aspects, as well as bio-mechanical external aspects (Agbenorkul et al., 1995), since we aim to emphasise the importance of understanding the connection between these two aspects.

Inflammatory aspects related to keloids. In the middle of the “keloid triad” there is the keloid fibroblast (KF) – see also Table 1 – a key cell submitted to both inflammatory and mechanical stimuli. In consequence, and compared to normal skin fibroblasts, KFs have an increased proliferation rate (Suttho et al., 2017; Shi et al., 2019) and a reduced apoptosis rate (Lu et al., 2007; Zhou et al., 2020). Moreover, KFs are more responsive to growth factors activation (i.e., by TGF- β 1 and PDGF) than their normal counterparts (Babu et al., 1992; Bettinger et al., 1996; Haisa et al., 1994), and this leads to higher production of type I and III collagens (the ratio I/III in keloid is 17:1 versus 5:1 in normal skin) and extracellular matrix (ECM) accumulation (Xue and Jackson, 2015). During wound healing, ECM degradation and remodelling is controlled by a family of enzymes called matrix metalloproteinases (MMPs). MMPs are synthesised by fibroblasts, keratinocytes and macrophages, and their activity is regulated by the tissue inhibitors of matrix metalloproteinase (TIMPs). Dysregulation of such MMP/TIMP equilibrium may underlie ECM over-accumulation. In keloids, the levels of MMP-1 and -13 (collagenase), MMP-2 (gelatinase) and TIMP-1 and -2 increase, whereas the level of MMP3 (stromelysin) decreases, compared to the respective levels in normal scars (Xue and Jackson, 2015; Ulrich et al., 2010; Aoki et al., 2014). The TGF- β 1/Smad signalling pathway is pivotal in the process of keloid fibrosis. TGF- β 1 regulates fibroblast-to-myofibroblast conversion, α -SMA expression and collagen synthesis during the development of fibrosis. TGF- β 1 is found in higher quantity in keloid than in normal scar (Desmoulière et al., 1993; Jagadeesan and Bayat, 2007; Carthy, 2018). Accordingly, a variety of therapeutic strategies (drugs, siRNA, shRNA, miRNA, etc) have been designed to target the TGF- β /Smad signalling pathway in order to interfere with fibroblast-mediated keloid progression (Zhang et al., 2020).

Other inflammatory cells involved in wound healing and aberrant scar formation, which impact also the formation and evolution of keloids, are the macrophages, T lymphocytes, mast cells and neutrophils (Limandjaja et al., 2020a; Wang et al., 2020); see also Table 1. All these inflammatory cells that have been found in higher numbers in keloids (Limandjaja et al., 2020a), are able to cross-talk with the KF all along the abnormal healing process via different cytokines (e.g., IL-4, IL-6, IL-10, IFN- γ), chemokines (MCP-1) and growth factors (e.g., TGF β , VEGF, bFGF) (Wang et al., 2020).

Bio-mechanical aspects related to keloids. All cells inside skin tissues are subjected to various mechanical forces that could be intrinsic (e.g., due to orientation of collagen fibres in the ECM, or due to cytoskeletal structures) or extrinsic (e.g., forces generated through muscle movement). These mechanical forces include stretching tension, shear force, scratch, compression, as well as hydrostatic and osmotic pressures (Ogawa, 2011). In (Ogawa, 2011; Ogawa et al., 2012) it was shown that keloids are more likely to develop in highly mobile anatomic regions subjected to skin stretching/contraction and with different stiffness/thickness characteristics, such as the anterior chest region, or the scapular regions. Moreover, the resulting keloid tissues have different mechanical properties compared to the healthy skin tissues: lower extensibility and higher rigidity for the keloids (Chambert et al., 2019). In addition, mechanical forces at the tissue level, can modulate inflammatory cellular behaviours via the activation of

Table 1. Some of the inflammatory cells and molecules most important in keloids, whose kinetics and interactions could be modelled mathematically.

Cells	Behaviour during keloid evolution	References
Fibroblasts	↑cell numbers, ↑proliferation, ↑migration, ↑metabolism; ↓apoptosis, ↑ECM degradation	(Limandjaja et al., 2020a)
Myofibroblasts	↑cell numbers	(Limandjaja et al., 2020a)
Keratinocytes	↑cell numbers	(Limandjaja et al., 2020a)
Macrophages	↑cell numbers (in particular M2 cells)	(Limandjaja et al., 2020a)
T cells	↑cell numbers	(Limandjaja et al., 2020a)
Mast cells	↑cell numbers	(Limandjaja et al., 2020a)
Molecules	Behaviour during keloid evolution	References
Cytokines	↑IL-6; ↑IL-8; ↑IL-18; ↑TNF- α ↓IL-10	(Shan et al., 2022; Wang et al., 2020)
Growth factors	↑TGF- β ; ↑PDGF; ↑VEGF	(Wang et al., 2020; Desmoulière et al., 1993; Niessen et al., 1999; Fujiwara et al., 2005)
Chemokines and their receptors	↑CCR1; ↓CCR7; ↑CXCL9; ↑MCP-1	(Shan et al., 2022; Wang et al., 2020)
Collagen	↑type-I:type-III ratio	(Abergel et al., 1985)

mechanoreceptors on fibroblasts and immune cells, which can further increase cells' roles on scar formation and spread (Huang and Ogawa, 2022). Some of the most studied cellular mechanoreceptors are the integrins (at cell-ECM junctions) and cadherins (at cell-cell junctions), which are known to play an important role in wound healing and fibrosis (Kuehlmann et al., 2020). In response to mechanoreceptors activation, the mechanical signal is transferred and interpreted inside the cell through a mechanotransduction process (that transforms mechanical inputs into biochemical signals), which then leads to the modulation of molecular signalling pathways, such as the FAK-ERK-MCP-1 pathway (Kuehlmann et al., 2020) or the TGF- β /Smad pathway mentioned above (Huang and Ogawa, 2022; Kuehlmann et al., 2020). Since the mechanotransduction process has a big impact on cell proliferation, survival, differentiation or motility, a recent study in (Zhou et al., 2022) hypothesised an important role for the mechanical microenvironment on the macrophage dysfunction in the context of keloid formation. In particular, it was suggested that tissue stretching combined with ECM stiffness can perturb the balance between M1 and M2 macrophages, which could then lead to sustained and concomitant pro- and anti-inflammatory signalling (Li et al., 2017; Xu et al., 2020).

Before discussing the mathematical approaches applied (or those that could be applied) to modelling keloids disorders, we mention briefly some of the differences and similarities between the keloids and the scars (either normal or abnormal), and between the keloids and the cancers. For more in-depth comparisons we refer the reader to the references cited below. This brief discussion below will help us understand the applicability of mathematical modelling approaches for wound healing, presented in Section 3.1, and those for cancer growth and invasion of surrounding tissue presented in Section 3.2, in the context of keloid research. We also note here that the cellular and biochemical processes involved in wound healing are similar to those involved in tumour development and invasion, and for this reason Dvorak (1986) suggested that tumours are “wounds that will not heal”. Also for this reason, the type and structure of the mathematical models for wound healing and cancer discussed in Sections 3.1 and 3.2 will be quite similar.

Differences and similarities between keloids and scars. Since different studies investigate the keloids from the perspective of abnormal scars (Limandjaja et al., 2021), in the following we mention briefly some of the differences and similarities between the keloids and the hypertrophic scars (the two types of abnormal scars); for a more detailed comparison see (Limandjaja et al., 2021).

- *Similarities.* Both hypertrophic and keloid scars are fibroproliferative disorders characterised by excessive and disorganised production ECM (in particular increased collagen type-I production), as a result of abnormal proliferation and differentiation of fibroblasts (Wang et al., 2020; Limandjaja et al., 2021). In addition, both types of scars are characterised by different inflammatory immune cells that infiltrate the tissue (mast cells, macrophages, T cells, dendritic cells), increased epidermal thickness, increased presence of myofibroblasts, increased TGF- β and VEGF levels (Limandjaja et al., 2021). They are also characterised by increased levels of some matrix metalloproteinases, such as the MMP-2, and increased levels of MMP inhibitors TIMP-1 and TIMP-2 (Limandjaja et al., 2021). For more similarities see (Limandjaja et al., 2021).
- *Differences.* There are various differences between the scars (in particular hypertrophic scars) and keloids in terms of clinical presentation, epidemiology, histologic findings (Ghazawi et al., 2018). For example, the hypertrophic scars do not exceed the original injury site, while the keloid scars grow and invade adjacent tissue, beyond the original injury site. Moreover, while the hypertrophic scars could regress spontaneously over time, this does not happen for the keloids (Wang et al., 2020; Bran et al., 2009). Also, the hypertrophic scars showed lower levels of hypoxia, as opposed to the keloids that are characterised by increased hypoxia (Vincent et al., 2008). Moreover, as mentioned above, the keloids usually develop at certain anatomical sites (e.g., chest, shoulder; characterised by certain mechanical properties) and are not associated with skin contracture, while the hypertrophic scars occur more commonly at joints and are associated with skin contracture. In regard to the keloid matrix, the ratio of type-I to type-III collagen is much higher (17:1) compared to the same ratio in normal scars (6:1) (Tan et al., 2019), or even hypertrophic scars dominated by type-III collagen (Bran et al., 2009). For more differences see (Limandjaja et al., 2021; Bran et al., 2009).

Differences and similarities between keloids and cancers. Since various studies mentioned the importance of investigating the keloids from the perspective of benign cancers (Tan et al., 2019; Kimura et al., 2014), in the following we briefly discuss this aspect by focusing on their differences and similarities. To start this discussion, we note that the word “cancer” appears to be coined by Hippocrates 2400 years ago, who compared the disease to a crab that adheres to its surroundings with its claws (Hajdu, 2011; Lonardo et al., 2015). The word “keloid” seems to have been coined by Alibert (1806) and/or Addison (1854), who described the resemblance of this disease with the crab’s claws, while acknowledging its tissue-invasive similarities with cancer.

- *Similarities.* Both cancers and keloids show hyper-proliferative cell behaviours, and cell invasion into the adjacent surrounding healthy tissue (in patterns that resemble crabs’ claws), which are fuelled by alterations in cytokine/growth factors levels Tan et al. (2019). Neither cancers nor keloids can regress spontaneously, and their surgical excision is followed by a relapse. Same as tumours, the keloids are vascularised through angiogenesis Tan et al. (2019). Moreover, both keloid fibroblasts and cancer cells seem to use similar bioenergetics based mainly on glycolysis (as opposed to non-keloid fibroblasts that use mainly oxidative phosphorylation) (Vincent et al., 2008). In fact, due to their similar clinical presentations, in some cases the keloids have been misdiagnosed as skin tumours, while in other cases the tumours have been misdiagnosed as keloids (Kimura et al., 2014). In addition to the clinical similarities between keloids and some specific skin tumours (e.g., dermato

fibrosarcoma protuberans, keloidal dermatofibroma (Kanitakis, 2013), keloidal basal cell carcinoma and keloidal atypical fibroxanthom (Tongdee et al., 2016)) a few studies reported that abnormal levels of cytokines and (keloid-promoting) growth factors that are associated with cancers, might induce also keloids development, leading to the question of whether the keloids could be seen as a para-neoplastic phenomenon (He et al., 2011). This question is also supported by the fact that the keloids can exhibit to a large extent the hallmarks of cancer (Tan et al., 2019): sustained proliferative signalling, evasion of growth suppressors, angiogenesis, resisting cell death. Moreover, the keloids can respond to a certain extent to cancer treatments such as radiotherapy, chemotherapy and targeted immunotherapy and virotherapy (Kim, 2021; Tan et al., 2019).

- *Differences.* The main differences between keloids and cancers are that (i) the keloids cannot occur spontaneously and cannot metastasise (He et al., 2011), and (ii) they are specific to humans, no other animal species developing naturally keloid tissues (Marttala et al., 2016). In addition to this there are the similarities between the keloids and scars mentioned above, which only emphasise the differences between the keloids and cancers and contribute to the usual classification of keloids as abnormal scars and not as neoplasms (Tan et al., 2019).

3 CURRENT MATHEMATICAL MODELLING APPROACHES




Figure 1. Keloid dynamics can be investigated at different spatial scales: (a) *microscopic level*: focus on the kinetics of individual cells, or interactions between individual cells via cytokines/growth factors; (b) *mesoscopic level*: focus on the evolution of densities of cells, and their interactions with gradients of cytokines/growth factors; (c) *macroscopic level*: focus on the skin tissue, and how the mechanical forces on the skin impact the evolution of the whole keloid.

Since the biological processes involved in keloid disorders are similar to those characterising benign cancers and wound healing, in the following two subsections we start by briefly summarising the current mathematical approaches used to model normal and/or hypertrophic scars (see Section 3.1), and the approaches used to model solid tumours (see Section 3.2). Then, we discuss briefly the mathematical and computational models used to describe the keloids dynamics (see Section 3.3).

Types of modelling approaches. The modelling approaches used to describe collective cell behaviours at different scales can be classified as (see also Figure 1): (a) *discrete (agent-based) models* that focus on individual cells; (b) *continuum models* that focus on densities of cells; (c) *hybrid models* that connect the dynamics of individual cells with continuum gradients of cytokines/growth factors or continuum densities of extracellular matrix or other cell populations; (d) *computational models* that focus mostly on cell and tissue bio-mechanics. The discrete, continuum and hybrid models are usually described by ordinary differential equations (ODEs), partial differential equations (PDEs) or integro-differential equations (IDEs) for the changes in the position of cells (or densities of cells), their differentiation, proliferation, death, etc., which are then simulated numerically. Some discrete models (e.g., cellular automata models) avoid the use of equations for changes in cell positions/velocities by focusing on the computational rules that govern such changes in one cell following interactions with other cells, or due to cell death, proliferation, etc. These cell behavioural rules are then implemented directly into the numerical code, and so they can be considered as computational models. Other computational models consider fixed cell densities inside tissues and focus on mechanical forces acting on these tissues (forces usually described by constitutive equations).

Due to the huge amount of mathematical and computational modelling of both wound healing and cancers (for some reviews we refer the reader to Arciero and Swingdon (2013); Buganza Tepole and Kuhl (2013); Flegg et al. (2015); Jorgensen and Sanders (2016); Sherratt and Dallon (2002); Thackham et al. (2009); Ziraldo et al. (2013), as well as to Altrock et al. (2015); Bekisz and Geris (2020); Chen et al. (2020); Eftimie et al. (2011); Enderling and Chaplain (2014); Yin et al. (2019) and the references therein), in the following two subsections we only mention briefly some of these studies, while emphasising the multi-scale aspects of these models as well as the bio-mechanical vs. immunological/inflammatory approaches considered. In each section, we split the discussion between mathematical/computational models for cell-level dynamics (i.e., models described by ODEs, PDEs for changes in the number or densities of cells) and mathematical/computational models for tissue-level dynamics (i.e., models usually described by constitutive equations for various mechanical forces acting on tissues).

REMARK 1. *We acknowledge the huge mathematical literature on collective cell movement, that focuses on the biological mechanisms behind cell-cell interactions and cell migration in development and disease; see, for example, the reviews in Buttenschön and Edelstein-Keshet (2020); Camley and Rappel (2017); Chen et al. (2020); Deutsch et al. (2020); Maini and Baker (2014) and the references therein. There are also even more studies on the collective movement of animals in ecological settings (Abaid et al., 2022). However, to keep this review focused, here we focus exclusively on models developed explicitly for wound healing, for cancer, and for keloids, and ignore related models investigating general collective movement of cells, or collective movement of cells in tissue development or other diseases.*

3.1 Modelling wound healing

Numerous mathematical and computational models have been developed since 1990s to explain the various phases of wound healing: from angiogenesis and inflammation, to wound closure and tissue remodelling. While many of these models focus only on the angiogenesis (Pettet et al., 1996; Gaffney et al., 2002) and inflammatory aspects related to wound healing (Mi et al., 2007; Stern et al., 2012; Ziraldo et al., 2013; Vodovotz et al., 2004; Waugh and Sherratt, 2006), others focus on generic interactions between cells and extracellular matrix (Sherratt and Murray, 1990; Dallon et al., 1999, 2000) and/or on the bio-mechanical aspects related to wound healing (Tranquillo and Murray, 1993; Bowden et al., 2016; Maroudas-Sacks and Zemel, 2018) without considering the impact of inflammation, and many other models combine both inflammatory and mechanical aspects (Cumming et al., 2010; Tepole and Kuhl, 2016; Koppenol et al.,

2017). Most of these models are continuum, but there are also a few discrete models (Mi et al., 2007; Stern et al., 2012) and some hybrid discrete-continuum models (Dallon et al., 1999; McDougall et al., 2006). Moreover, while most of these wound healing models are local, there are also a few non-local models (Webb, 2022) that incorporate the long-range (i.e., almost tissue-level) effects of mechanical forces acting on the cells (e.g., cell-cell adhesion forces).

In the following we discuss in more detail a few examples of older/newer models for wound healing, to emphasise that: (i) older models are simpler but also very relevant to current studies; and (ii) newer models encode inflammatory-mechanical interactions but these are still difficult to understand due to model complexity. However, for more pedagogical reviews of modelling wound healing we refer the reader to Arciero and Swingdon (2013); Flegg et al. (2015).

Cell-level dynamics. One of the earliest models for cell dynamics in wound healing was introduced by (Sherratt and Murray, 1990) to describe the changes in the density of epithelial cells as a result of linear/nonlinear cell migration and mitotic regeneration. Since a one-equation PDE model could not accurately reproduce the data about the decrease in wound radius with respect to time, the authors in (Sherratt and Murray, 1990) have then considered a two-equations PDE model for the changes in epithelial cell density in response to a generic mitosis-regulating chemical. Due to a lack of biological details related to wound healing, numerical simulations combined with experimental data on wound radius were then performed to investigate the role of the generic chemical: inhibitor or activator of cell mitosis. We note that given the very simple model, which involved very few parameters, the authors have estimated these parameters (e.g., linear and nonlinear diffusion coefficients, cell mitotic rate, half-life of the chemical) using available experimental data. They also estimated from the wave speed of epidermal wound healing.

A very recent mathematical study (Webb, 2022) also used a one-equation model for the collective movement of cells to investigate the impact of cell-cell adhesion and cells sensing radius on the incomplete wound closure. The cell-cell adhesion was implemented (using the same modelling framework introduced by Armstrong et al. (2006)) via non-local terms describing the tissue-level effects of these adhesive interactions on the surrounding cells. Numerical simulations performed with this nonlocal advection-diffusion-reaction model showed the existence of a critical value of the sensing radius and a critical interval for the adhesion force that ensured complete wound closure. Reducing the adhesion force below this interval allowed for wound closure for every value of the sensing radius. This theoretical study did not use data to parametrise the model.

More complex models have been derived to consider various inflammatory aspects involved in wound healing. For example, Stern et al. (2012) developed a discrete model for the movement of individual epithelial cells in response to the cross-talk between two signalling pathways EGFR and TGF- β . The cells were characterised by intracellular proteins and cell surface receptors (e.g. EGFR, TLR-4 integrins), and they can produce diffusible factors (e.g., EGF). A “scratch” was introduced into the model to create a wound edge, and cells at this edge were assumed to produce reactive oxygen species (ROS), damage associated molecular patterns (DAMPs) and adenosine triphosphate (ATP) which stimulated more epithelial cells. Note that this model could actually be considered a multiscale model, due to the dynamics of cells - at cell level - that is controlled by molecular-level processes. Numerical simulations proposed new testable hypotheses on the crosstalk between TGF- β and EGFR (with a possible point of regulation at the receptor tyrosine kinase level) that could lead to scratch healing. The simulation results that lead to this hypothesis depended on model parameters, which were “calibrated” in the sense that they were chosen from the plausible parameters to produce behaviours similar to the in vitro cell migration data and wound closure percentage.

Tissue-level dynamics. Bowden et al. (2016) considered a 10-equations model for growth and contraction in the dermal layer of the skin during wound healing. The model combines differential equations for changes in tissue deformation and radial stress, balance of linear and angular momentum, a constitutive stress-strain relationship for an incompressible hyperelastic material, as well as the time-evolution equation of a growth tensor. The boundary conditions modelled contraction forces as exerted by fibroblast migration (considered implicitly). Numerical results investigated the changes in stress and growth at various spatial positions, changes in wound radius, total tissue volume and tissue thickness as functions of time. Local sensitivity analysis was also performed to investigate the effect of four model parameters (radial, circumferential and axial growth, and shear elastic modulus) on tissue remodelling time, and identified the shear elastic modulus as the parameter with the greatest impact.

Multiscale cell-level and tissue-level dynamics. These types of models combine differential equations for changes in cell densities with constitutive equations for mechanical forces acting on the tissue. One of the earliest models on this topic was introduced by Tranquillo and Murray (1993), who considered a fibroblast cell population that can proliferate logistically, migrate randomly and can also undergo passive convection with the extracellular matrix (ECM), whenever ECM is which also undergoes convection. Finally, a constitutive equation described the mechanical force balance between traction forces exerted by the cells and forces associated with the properties of ECM. Since the behaviour of the model – showing an expansion of the wound boundary followed by relaxation to its initial position as fibroblast repopulate the wound – could not explain the features of a real wound (i.e., a first plateau phase in the wound area, followed by an exponential decrease), the authors incorporated implicitly inflammatory responses. To this end, they investigated numerically different scenarios and observed that an increased steady-state concentration of an inflammation-derived growth factor associated with the wound, that influences the traction force, the maximum ECM concentration or the cell chemotaxis, could lead to simulations qualitatively consistent with wound healing contraction. The final discussion in Tranquillo and Murray (1993) also emphasised that the simple incorporation of the inflammatory responses can create ambiguity in comparing the results since identical inflammatory mechanisms would not lead to the same chemotactic gradient in different wound geometries. Our discussion in Section 2 also emphasised the complexity of the inflammatory response in wound healing characterised by multiple types of immune cells, cytokines, growth factors, etc. (see also Table 1), which could lead to more ambiguity if simpler models are used.

A more detailed model was introduced by Koppenol et al. (2017) to investigate numerically the formation and subsequent regression of hypertrophic scar tissue following dermal wounding. The authors used a constitutive stress-strain equation for the explicit modelling of the dermal layer of the skin as a heterogeneous isotropic and compressible neo-Hookean solid. In addition, they used partial-differential equations for the spatio-temporal evolution of a generic fibroblast population, a myofibroblast population, and a generic signalling molecule produced and degraded by both fibroblasts and myofibroblasts. The collagen molecules are produced by both fibroblasts and myofibroblasts (and this production rate is enhanced by the signalling molecule), but collagen does not move as it is assumed to be attached instantly to the ECM. Finite element numerical simulations and their comparison with clinical measurements suggested that a relatively high apoptosis rate of myofibroblasts can lead to normal scar tissue, while a low apoptosis rate can lead to hypertrophic scar tissue. However, the results depended strongly on the parameter models: some chosen from the published biological and mathematical literature (even if those experimental studies were performed under different conditions), some resulting from other parameter values, and some guessed. Therefore the numerical results, however interesting and informative in terms of new possible experiments, were valid only for those parameter sets.

3.2 Modelling solid cancers

The development of mathematical models for solid cancers dates back to 1950's (Araujo and McElwain, 2004). While earlier models focused on the mechanisms explaining the growth of avascular tumour spheroids due to diffusion of nutrients in the environment (Thomlinson and Gray, 1955; Laird et al., 1965; Greenspan, 1972, 1976; Roose et al., 2007), later models focused also on the growth of vascular spheroids in response to angiogenesis (Chaplain et al., 2006), as well as on tissue invasion and metastasis (Anderson et al., 2000; Quaranta et al., 2008; Sfakianakis and Chaplain, 2021), interactions with the immune cells in the environment (Eftimie et al., 2011), or cancer response to treatments (Yankeelov et al., 2013). Some of these models are discrete (Macnamara, 2021; Metzcar et al., 2019), others are hybrid (Rejniak and Anderson, 2011), and many more models are continuum (Chaplain et al., 2006; Roose et al., 2007). Moreover, while the great majority of models developed until a decade ago focused on cancer dynamics at one scale (molecular, cell or tissue scale), the last decade has seen the development of numerous multiscale models for cancer invasion into the surrounding tissues (Quaranta et al., 2008; Stolarska et al., 2009; Diesboeck et al., 2011).

Again, as in the previous section, we discuss in more detail a few examples of older/newer models for cancer growth and invasion, to emphasise that: (i) older models are simpler (and so can be investigated analytically), but they are still relevant to current studies; and (ii) newer models encode inflammatory-mechanical interactions, but these are still difficult to understand due to model complexity (including also large numbers of parameters that need to be estimated). For many more models for tumour growth, invasion and interactions with other components of the tumour microenvironment (including stromal and inflammatory cells) we refer the reader to the reviews in Araujo and McElwain (2004); Chaplain et al. (2006); Anderson et al. (2000); Quaranta et al. (2008); Eftimie et al. (2011); Sfakianakis and Chaplain (2021); Yankeelov et al. (2013); Macnamara (2021); Stolarska et al. (2009); Diesboeck et al. (2011); Rejniak and Anderson (2011); Friedman (2004); Metzcar et al. (2019) and references therein.

Cell-level dynamics. The last few decades have seen the development of a large variety of mathematical models describing the temporal and spatio-temporal evolution of tumour cell populations, as well as its interactions with other cell densities: e.g., various immune cells involved in inflammation (Eftimie and Barelle, 2021; Eftimie et al., 2011), endothelial cells involved in angiogenesis (Chaplain et al., 2006), extracellular matrix components (Anderson et al., 2000). In the following we present two models, a discrete and a continuum model for immune responses to cancer.

An agent-based model for immune responses against colorectal tumours was introduced in Kather et al. (2017) to investigate the interactions between stem and non-stem tumour cells, and T lymphocytes. The immune cells that have killed successfully five time are assumed to lead to the increase in tumour stroma, which then reduces cell migration. To parametrise the model, some parameters were taken from previous studies, others were measured histologically, and the remaining parameters were set to some biologically plausible values. Then the numerical spatial distribution patterns of immune and tumour cells were compared visually with histological colorectal cancer tissue samples. Numerical simulations showed different types of tumour infiltrated with immune cells, and suggested a possible connection between immune cell numbers and stromal permeability: increase in stromal permeability without an increase in lymphocytes led to faster tumour progression.

The inflammatory immune responses to cancer do not include only lymphocytes but also many more types of immune cells, some triggering inflammation others reducing inflammation. In Kirshtein et al. (2020) the authors have derived a continuum model for immune responses to colorectal cancer (described

by ordinary differential equations), which considered not only the densities of cytotoxic T lymphocytes but also the densities of naive, helper and regulatory T cells, densities of macrophages, naive and activated dendritic cells, and various cytokines such as IFN- γ and TGF- β , carcinogenic cytokines (such as IL-6, IL17, IL-21) and immuno-suppressive agents (such as IL-10). To parametrise the model the authors used clinical data from the Cancer Genome Atlas (TCGA) project of Colon Adenocarcinoma, and estimated the fraction of each immune cell type in each tumour, tumour dimensions, densities of cancer cells and ratio of cancer to immune cells for different clusters of patients' data with similar immune patterns. Sensitivity analysis was then performed to investigate the impact of variability in model parameters on tumour-immune outcomes and identified cancer proliferation and death rates as the most sensitive parameters. Numerical simulations were used to investigate the temporal dynamics of cells in colorectal tumours, and confirmed different patterns of tumour growth based on the immune infiltration level. The simulations also showed that some immune-related parameters, such as the macrophage activation rates, have different effects on tumour growth depending on immune infiltration level.

Tissue-level dynamics. One of the earlier models for the dynamics (and deformation) of avascular tumour spheroids in response to available nutrients was introduced by Greenspan (1976). This moving-boundary mathematical model was described by an equation for the internal pressure of the spheroid in response to cell division and death, which then leads to the motion of cellular material. A second equation described the steady-state level of nutrients at the outer boundary of the tumour spheroid. In addition, it was assumed that (i) cell-cell adhesion led to a surface tension at the outer boundary of the spheroid proportional to the mean curvature of the boundary, tension which balanced the internal expanding pressure; (ii) cell mass is conserved; (iii) the rate of nutrient diffusion through the outer spheroid surface is equal to the rate of nutrient consumption inside the spheroid. Analytical results for this model investigated the stability conditions for the growing tumours, i.e., the conditions (on tumour spheroid radius, nutrient level,) for which the tumour preserves its “almost-spherical” shape under small environmental perturbations and does not desegregate into smaller cell colonies. No numerical simulations were performed for this simple model.

A more complex poro-mechanical model for the dynamics of a tumour spheroid as a result of internal tumour pressure was recently developed by Urcun et al. (2021). The model considered the tumour as a porous multiphase system, with the ECM representing a solid scaffold while the interstitial fluid and the tumour cells being modelled as fluid phases. Equations described the mass conservation of these three phases, a momentum conservation equation for the full system, and an advection-diffusion equation for the oxygen within the interstitial fluid (here the oxygen represented the nutrient that limits the growth of tumour cells). Some model parameters were quantified using an experimental protocol called the Cellular Capsule Technology (CCT), where multi-cellular tumour spheroids were cultured with spherical porous alginate capsules which work as mechanosensor. Other model parameters were taken from multiple previous numerical studies, and their averaged values were set as initial guesses for parameter identification. A sensitivity analysis (using Sobol's index) was performed to assess the sensitivity of the finite element solution to the parameters. Then some of the selected parameters were identified (via an optimisation algorithm) from the sensitivity profiles, and cross-validated on experimental results with thick and thin encapsulated multi-cellular tumour spheroids. Finally, numerical simulations were performed with the identified parameter set to investigate model dynamics in terms of oxygen profiles, necrotic fraction, interstitial fluid pressure, ECM displacement, as well as tumour cell pressure and saturation.

Multiscale cell-level and tissue-level dynamics. A hybrid model for cancer growth and spread was introduced in Suveges et al. (2021) to investigate the spatial structure of the extracellular matrix (ECM),

and in particular the density and orientation of collagen fibres, on the evolution of solid cancers. To this end, the authors combined an off-lattice agent based modelling framework *MultiCell-LF* (Kim et al., 2014) for individual tumour cell dynamics, with a multi-scale continuous framework (Shuttleworth and Trucu, 2019) for the dynamics of the two phases of the extracellular matrix (ECM) (i.e., the fibrous and non-fibrous phases). The coupling of the discrete model with the continuous model was realised through mechanical cell-matrix adhesive forces that act non-locally in a neighbourhood of each individual cells, and the re-arrangement of matrix fibres by the cancer cells. Note that the density and directionality of the ECM fibre phase (as well as the non-fibres phase) are specified across the whole spatial domain representing the tissue. Numerical simulations were performed with some parameter values taken from the literature, and other values guessed. The results showed different cell invasion patterns depending on the cell-cell and cell-matrix adhesive forces, cells sensing radius, as well as on the spatial structure and directionality of fibres in the ECM. The complexity and multiscale aspect of this particular model makes it difficult to be parametrised.

A different type of hybrid model was developed by Kim et al. (2007) to describe the growth of a tumour spheroid. The model considered a discrete representations of cells in the outer region of the spheroid, where the majority of growth and cell division occurred, and a continuum constitutive description of cells in the quiescent and necrotic regions of the tumour, and a continuum description of the extracellular matrix (i.e., surrounding gel). Finally, the dynamics of two nutrients, glucose and oxygen, was described by reaction-diffusion equations. The outside gel and the interior (quiescent and necrotic) tumour regions were assumed to be viscoelastic with different material properties, and their behaviour was described by constitutive equations. While these constitutive equations were solved using the finite element method with a triangular mesh, the reaction-diffusion equations for the nutrients were solved using an alternating-direction implicit finite-difference scheme on a regular grid. In regard to the discrete cells, they were assumed to be subjected to various forces caused by cell-cell and cell-ECM adhesion (which could deform the cells), could move with a specific velocity, and could undergo proliferation depending on the nutrient and stress levels. The connection between the regions with discrete cells and continuum cell fields was computed numerically by first determining the location, magnitude and direction of force exerted by individual cells onto the boundary of the continuum fields. Then finite element interpolation functions were used to distribute point forces from individual cells (at the boundary of the continuum field) onto the nodes of the finite elements (that cover the continuum field). This nodal force distribution was used as a boundary condition for the stress distribution in the continuum continuum regions of the domain (Kim et al., 2007). The continuum regions were also updated through the generations of a new mesh, to account for discrete proliferating cells not receiving enough nutrients and thus becoming quiescent and entering the continuum field. Numerical simulations of this model investigated the temporal changes in the size of different regions of the tumour, as well as the impact of gel stiffness on tumour growth rates. Again, the model was not parametrised using experimental data.

3.3 Modelling keloids

Despite the numerous biological and clinical studies on keloids that have been published since 1800s (Alibert, 1806; Addison, 1854), very few mathematical/computational models have been proposed to focus exclusively on keloid formation and/or growth. We briefly mention all these models below.

Cell-level dynamics. The few models for the temporal and spatio-temporal evolution of cell densities focus mainly on the inflammatory aspects involved in keloid progression (Cobbold and Sherratt, 2000; Bianca, 2011; Bianca and Fermo, 2011). For example, starting with some experimental studies (Schäffer

et al., 1997a,b) on the role of nitric oxide (NO) in wound healing (NO being produced by both fibroblasts and macrophages), Cobbold and Sherratt (2000) proposed an ODE model for the temporal interactions between fibroblasts, collagen, oxygen, nitric oxide (for a simpler model), as well as macrophages, TGF- β and blood vessels (for a more complex model). Numerical simulations performed with both models suggested that keloid scarring could be the result of increased NO concentrations (as opposed to hypertrophic scarring that does not seem to be associated with increased NO). In addition, simulations suggested that reducing NO levels in conjunction with surgical excision might lead to normal scars. A different modelling approach was considered by Bianca (2011); Bianca and Fermo (2011), who used the kinetic theory of active particles (KTAP) to propose a model for keloid formation from the perspective of malignant cells. Therefore, their integro-differential model described the interactions between virus particles, immune cells, normal fibroblasts, keloid fibroblasts and malignant cells. All these cells were characterised by an activity variable that described the proliferation ability of fibroblasts, the aggressiveness of virus particles, the activation of immune cells and the progression ability of malignant cells. Numerical simulations were performed to investigate the behaviour of cell variables in this kinetic system in response to changes in various parameters, and with respect to the activity variables describing the phenotypes of these cells (Bianca, 2011; Bianca and Fermo, 2011).

Tissue-level dynamics. In addition to the above few mathematical models (focused on the dynamics of specific cells and aggregations of cells), there are also some computational models of keloids that focus mainly on the constitutive equations that describe the relations between different mechanical forces acting directly on the tissue, the tissue displacement, and the material properties of the tissue (see also the above discussion on computational models for the skin) (Akaishi et al., 2008; Chambert et al., 2012; Sutula et al., 2020; Marie et al., 2022). The computational domain, representing the skin formed of both healthy and keloid tissue (each being characterised by different mechanical parameters), is then discretised using a finite element approach. For example, in Akaishi et al. (2008) the authors used a finite element analysis to investigate the relationship between the keloid growth pattern and the stretching tension, and concluded that (i) keloid centers were regions of low tension while the keloid edges were subjected by high tension; (ii) keloid expansion occurred in the direction in which was pulled, and the crab's claw-shaped invasion pattern was the result of increased stretching tension; (iii) adhesion to subcutaneous hard tissue increased the tension in the keloid. In Chambert et al. (2012) the authors considered a computational model for the simulation of the skin and various stress fields in the vicinity of a keloid scar in the presternal area of the body, and used it to predict the likely directions of evolution of this keloid based on the mechanical stress field. In Sutula et al. (2020) the authors considered a 2D hyperelastic bi-material model of the keloid tissue and the surrounding healthy tissue. A finite element solver was then used to identify the mechanical parameters that characterise soft tissue based on *in vivo* experiments of uni-axial displacement, and the sensitivity of model parameters with respect to the displacement field and the reaction force measurements. Since the domain for all these tissue-level models for keloids is defined computationally (in contrast to the previous models for wounds and tumours where the domain/ECM was described by time-evolving equations), a particular aspect that needs to be investigated is the boundary between the keloid scar and the healthy skin, which might require the refinement of the finite element mesh to improve the accuracy of numerical prediction. Such an issue can be addressed using, for example, the Dual Weighted Residual (DWR) technique (Marie et al., 2022).

Cell-level and tissue-level dynamics. One of the earlier mathematical models that were derived to investigate fibro-proliferative wound healing disorders such as hypertrophic scars and keloids was proposed by Olsen et al. (1996). The model is described by partial differential equations for the spatio-temporal dynamics of fibroblasts, myofibroblasts, collagen and a generic growth factor. The cells were also assumed

to be passively convected by the strained tissue, and thus the equations for the changes in the densities of these cells (as a result of inflammation) were coupled with a conservation law for tissue displacement (as a result of mechanical forces). Numerical simulations were performed with some arbitrary parameter values in the context of normal wound healing and pathological response characterised by increased fibroblasts, myofibroblasts, growth factor and collagen values. Analytical bifurcation approaches were considered to investigate the changes in model steady states with respect to various parameters, and a travelling wave analysis was performed to calculate the minimum wave velocity.

4 OPEN PROBLEMS

Despite the various modelling approaches summarised in the above sections, there are still many open problems related to the modelling, as well as the analytical and numerical investigation of wounds/tumours/keloids. There are also open problems related to data availability and model parametrisation. Before mentioning briefly some of these open problems in Sections 4.1 and 4.2 below, we need to return to the models discussed above and to notice that while the simpler earlier models (for wound healing and cancer) focused also on analytical results, the later more complex models focused almost exclusively on numerical results. The complexity of these newer multiscale models makes it sometimes very difficult to investigate them analytically (in terms of pattern formation, bifurcations, travelling waves, etc...). Also the discrete/hybrid models are investigated only computationally, since the analytical approaches to study them still represent open problems. However, since this review did not present in more detail the various analytical approaches considered so far to investigate the various discrete/hybrid/continuum models for wounds, cancers and keloids, we prefer to focus our discussion on open problems only on some issues related to computational and data collection and assimilation approaches.

4.1 Numerical issues.

Continuum models. Regarding the numerical resolution of tissue-level constitutive equations as well as many PDEs for cell-level dynamics, the finite element method (FEM) is the way most frequently used. FEM implies, however, specific challenges: the widely varying skin-scar or skin-tumour stiffness leads to large displacement field gradients at the interface, knowing this interface evolves in time (Duddu et al., 2008). Moreover computing the velocity field along the interface may be challenging because of difficulties in resolving the gradient fields across the interface (Bordas and Duflot, 2007; Bordas et al., 2008):

- The gradients of the unknown fields (strain field, chemical gradients, etc.) are discontinuous across the scar/skin or tumour/tissue interface because the physical properties of the keloid are different from that of the skin.
- Field gradients can be large at the interface between the skin and the scar.
- The skin/scar interface evolves in time (Bui et al., 2017, 2018; Duprez et al., 2020).
- Computing the velocity field along the interface may be challenging because of difficulties in resolving the gradient fields across the interface (Bordas and Duflot, 2007; Bordas et al., 2008).

One approach that has been used to address some of these issues is the goal oriented mesh adaptation to improve efficiently the accuracy of the prediction (Duprez et al., 2020; Marie et al., 2022). The very recent study in Marie et al. (2022) has shown the feasibility of goal oriented mesh adaptation to improve the accuracy of the prediction, in the context of numerical simulations of mechanical responses in a keloid scar.

Discrete & hybrid models. The main challenge of these models is to simulate very large numbers of cells. It is known that a 1cm^3 of tissue can contain between $10^8 - 10^9$ cells, depending on cell diameter (Monte, 2009). Despite the huge increase in computational power over the last few years, simulating numerically the behaviour of $10^8 - 10^9$ cells, that can interact linearly/non-linearly with other cells in a 3D environment, and which might be characterised by some internal variables (describing, for example, their phenotype or activation state), is still computational prohibitive. In fact, this computational issue led to the development of hybrid models that combined the detailed dynamics of individual cells with some coarse-grained representation of other cells or of the environment (e.g., concentrations of chemicals, densities of extracellular matrix components) (Kim et al., 2007).

Another issue with some of the discrete models is that the lattice-based models are at risk with potential grid-based artefacts (Metzcar et al., 2019). For example, the number/position of neighbours with whom a cell can interact, could lead to different results regarding the evolution of an aggregation of cells (fibroblasts, cancer cells, etc.).

4.2 Multiscale data collection and model parametrisation

Despite the multitude of mathematical models for wound healing and cancer, that can be applied (perhaps in slightly modified ways) to investigate keloid dynamics, one of the main challenges of these models is the number of parameters, sometimes modelling dynamics across multiple scales (molecular, cellular and tissue scales), and limited data sets. Even if one manages to estimate some of these parameters in a multiscale model using different data sets (usually obtained through different experiments on different types of cells), there is still the problem of investigating the uncertainty in model outcomes to parameters across multiple scales. Renardy et al. (2019) presented an overview of various single-scale sensitivity analysis methods and approaches that could be used for multiscale models (i.e., all-in-one sensitivity approaches, intra/inter-compartmental sensitivity approaches and hierarchical sensitivity), while discussing the strong and weak points of each approach.

The latest experimental developments towards the use of microfluidics and lab-on-a-chip technologies to establish microenvironments to study wound healing (Monfared et al., 2021), cancer (Liu et al., 2021) and other diseases such as keloids, will generate much more single-scale and multi-scale data. These technologies enable researchers to vary local molecular, cellular and environmental parameters in a controlled way, which can then be replicated by the mathematical and computational modelling approaches to match the experimental settings through the development of “digital twins” (DT) (Eftimie et al., 2022). The current push for the development of DT for precision medicine (Venkatesh et al., 2022) will impact also the research for keloids (especially due to the lack of appropriate murine models for keloids). This will eventually allow the use of real-time single-scale and multi-scale keloids data to parametrise multiscale models for keloid dynamics. Nevertheless, the computational challenges discussed above will have to be solved before the development of such DT.

5 SUMMARY AND CONCLUSIONS

Keloids are unique human disorders, as no other animal species can develop naturally such keloid scars. This particular aspect has impacted the lack of current treatments for these keloid disorders. However, since the keloids display many molecular and cellular characteristics similar to cancers and wound healing, it is normal to look at this other two disorders when trying to understand keloid evolution. Moreover, if *in vitro* and/or *in vivo* experiments are not possible, one can still use *in silico* approaches to propose and test various biological hypotheses.

The goal of this review was to present the state-of-the art in modelling and simulations of keloid disorders. To this end, we started with a review of the biology of these disorders (while emphasising the similarities and differences at molecular, cellular and tissue scales) between keloids and wound healing and cancer. Moreover, since there is a huge amount of theoretical (mathematical and computational) studies for both wound healing and cancer, it was normal to consider such theoretical approaches to drive keloid research further. Thus, in Sections 3.1 and 3.2 we started by listing a multitude of studies for wound healing and cancer, and then we mentioned in more detail some of the models developed in these studies. In our discussion we aimed to present both older and newer models, so we can (i) emphasise the simplicity of older models vs. the complexity of newer models; (ii) emphasise the lack of multiscale data to parametrise all these models. Then, in Section 3.3 we presented briefly most of the mathematical/computational models we found on keloids. The literature review we did in this section showed that at this moment there are more mathematical/computational approaches to investigate tissue-level keloids dynamics, compared to cell-level dynamics. This is in contrast to the modelling studies on wounds and cancers, where cell-level dynamics dominates. This trend can be probably explained by the relatively few *in vitro* experiments and almost a lack of *in vivo* experiments on keloids, which can be used to parametrise mathematical models for cell-level dynamics. We believe that clinical progress with this keloid disorder can be ensured only through the development of new single-scale cell-level mathematical models and/or multi-scale cell-level/tissue-level mathematical models for keloids, combined with the generation of experimental data to parametrise these models.

CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

This review article was originally conceived and structured by RE. All authors contributed to the writing of this manuscript.

FUNDING

RE and OA acknowledge funding from a French Agence Nationale de Recherche (ANR) grant number ANR-21-CE45-0025-01; SPAB and SU acknowledge funding from a Luxembourg National Research Fund (FNR) grant number INTER/ANR/21/16399490; G.R. acknowledges funding from an ANR grant number ANR-21-CE45-0025-03; FC's work is partially supported by the I-Site BFC project NAANoD, the EIPHI Graduate School (contract ANR-17-EURE-0002) and the ANR project S-KELOID (ANR-21-CE45-0025-04). FC is also grateful to the Center for Mathematical Modelling grant FB20005.

REFERENCES

- Abaid, N., Eftimie, R., Hutt, A., and van Veen, L. (2022). Editorial: Modelling collective motion across scales. *Front. Appl. Math. Stat.* 8, 932364
- Abergel, R., Pizzurro, D., Meeker, C., Lask, G., Matsuoka, L., Minor, R., et al. (1985). Biochemical composition of the connective tissue in keloids and analysis of collagen metabolism in keloid fibroblast cultures. *The Journal of Investigative Dermatology* 84, 384–390
- Addison, T. (1854). On the keloid of Alibert and on true keloid. *Med. Chir. Trans.* 37, 27–47

- Agbenorkul, P., Kus, H., and Myczkowski, T. (1995). The keloid triad hypothesis (kht): a basis for keloid etiopathogenesis and clues for prevention. *Eur. J. Plast. Surg.* 18, 301–304
- Akaishi, S., Akimoto, M., Ogawa, R., and Hyakusoku, H. (2008). The relationship between keloid growth pattern and stretching tension: visual analysis using the finite element method. *Ann. Plast. Surg.* 60, 445–451
- Alibert, J. (1806). *Description des maladies de la peau observées à l'Hôpital Saint-Louis* (Paris: Chez Barrois l'ainé et fils)
- Altrock, P., Liu, L., and Michor, F. (2015). The mathematics of cancer: integrating quantitative models. *Nat Rev Cancer* 15, 730–745
- Anderson, A., Chaplain, M., Newman, E., Steele, R., and Thompson, A. (2000). Mathematical modelling of tumour invasion and metastasis. *Computational and Mathematical Methods in Medicine* 2, 129–154
- Aoki, M., Miyake, K., Ogawa, R., Dohi, T., Akaishi, S., Hyakusoku, H., et al. (2014). siRNA knockdown of tissue inhibitor of metalloproteinase-1 in keloid fibroblasts leads to degradation of collagen type I. *J. Invest. Dermatol.* 134, 818–826
- Araujo, R. and McElwain, D. (2004). A history of the study of solid tumour growth: the contribution of mathematical modelling. *Bull. Math. Biol.* 66, 1039–1091
- Arciero, J. and Swingdon, D. (2013). Equation-based models of wound healing and collective cell migration. In *Complex Systems and Computational Biology Approaches to Acute Inflammation*, eds. Y. Vodovotz and G. An. (New York: Springer), chap. 11. 185–207
- Armstrong, N., Painter, K., and Sherratt, J. (2006). A continuum approach to modelling cell-cell adhesion. *J. Theor. Biol.* 243, 98–113
- Babu, M., Diegelmann, R., and Oliver, N. (1992). Keloid fibroblasts exhibit an altered response to $\text{tgf-}\beta$. *J. Invest. Dermatol.* 99, 650–655
- Bayat, A., Arscott, G., Ollier, W., Grouther, D. M., and Ferguson, M. (2005). Keloid disease: clinical relevance of single versus multiple site scars. *British Journal of Plastic Surgery* 58, 28–37
- Bekisz, S. and Geris, L. (2020). Cancer modeling: from mechanistic to data-driven approaches, and from fundamental insights to clinical applications. *J. Computational Science* 46, 101198
- Bettinger, D., Yager, D., Diegelmann, R., and Cohen, I. (1996). The effect of $\text{TGF-}\beta$ on keloid fibroblast proliferation and collagen synthesis. *Plast. Reconstr. Surg.* 98, 827–833
- Bianca, C. (2011). Mathematical modeling for keloid formation triggered by virus: Malignant effects and immune system competition. *Mathematical Models and Methods in Applied Sciences.* 21, 389–419
- Bianca, C. and Fermo, L. (2011). Bifurcation diagrams for the moments of a kinetic type model of keloid-immune system competition. *Computers and Mathematics with Applications* 61, 277–288
- Bordas, S. and Duflot, M. (2007). Derivative recovery and a posteriori error estimate for extended finite elements. *Computer Methods in Applied Mechanics and Engineering* 196, 3381–3399
- Bordas, S., Duflot, M., and Le, P. (2008). A simple error estimator for extended finite elements. *Communications in Numerical Methods in Engineering* 24, 961–971
- Bowden, L., Byrne, H., Maini, P., and Moulton, D. (2016). A morphoelastic model for dermal wound closure. *Biomech Model Mechanobiol.* 15, 663 – 681
- Bran, G., Goessler, U., Hormann, K., Riedel, F., and Sadick, H. (2009). Keloids: current concepts of pathogenesis (review). *International Journal of Molecular Medicine* 24, 283–293
- Buganza Tepole, A. and Kuhl, E. (2013). Systems-based approaches toward wound healing. *Pediatric Research* 73, 553–563. doi:<https://doi.org/10.1038/pr.2013.3>

- Bui, H. P., Tomar, S., Courtecuisse, H., Audette, M., Cotin, S., and Bordas, S. P. (2018). Controlling the error on target motion through real-time mesh adaptation: Applications to deep brain stimulation. *International journal for numerical methods in biomedical engineering* 34, e2958
- Bui, H. P., Tomar, S., Courtecuisse, H., Cotin, S., and Bordas, S. P. (2017). Real-time error control for surgical simulation. *IEEE Transactions on Biomedical Engineering* 65, 596–607
- Butler, P., Longaker, M., and Yang, G. (2008). Current progress in keloid research and treatment. *J. Am. Coll Surg.* 206, 731–741
- Buttenschön, A. and Edelstein-Keshet, L. (2020). Bridging from single to collective cell migration: a review of models and links to experiments. *PLOS Computational Biology* 16, e1008411
- Camley, B. and Rappel, W.-J. (2017). Physical models of collective cell motility: from cell to tissue. *J. Phys. D Appl. Phys.* 50, 113002
- Carthy, J. (2018). TGF β signaling and the control of myofibroblast differentiation: Implications for chronic inflammatory disorders. *J. Cell Physiol.* 233, 98–106
- Chambert, J., Jacquet, E., and Remache, D. (2012). Numerical analysis of keloid scar in the presternal area. *Computer Methods in Biomechanics and Biomedical Engineering* 15, 23–24
- Chambert, J., Lihoreau, T., Joly, S., Chatelain, B., Sandoz, P., Humbert, P., et al. (2019). Multimodal investigation of a keloid scar by combining mechanical tests in vivo with diverse imaging techniques. *J. Mech. Behav. Biomed. Mat.* 99, 206–215
- Chaplain, M., McDougall, S., and Anderson, A. (2006). Mathematical modeling of tumor-induced angiogenesis. *Annu Rev Biomed Eng* 8, 233–257
- Chen, L., Painter, K., Surulescu, C., and Zhigun, A. (2020). Mathematical models for cell migration: a non-local perspective. *Phil. Trans. R. Soc. B* 375
- Cobbold, C. and Sherratt, J. (2000). Mathematical modelling of nitric oxide activity in wound healing can explain keloid and hypertrophic scarring. *Journal of Theoretical Biology* 204, 257–288. doi:<https://doi.org/10.1006/jtbi.2000.2012>
- Cumming, B. D., McElwain, D. L., and Upton, Z. (2010). A mathematical model of wound healing and subsequent scarring. *Journal of the Royal Society Interface* 7, 19–34
- Dallon, J. C., Sherratt, J. A., and Maini, P. (1999). Mathematical modelling of extracellular matrix dynamics using discrete cells: fiber orientation and tissue regeneration. *Journal of Theoretical Biology* 199, 449–471
- Dallon, J. C., Sherratt, J. A., Maini, P., and Ferguson, M. (2000). Biological implications of discrete mathematical model for collagen deposition and alignment in dermal wound repair. *IMA J. Math. Appl. Med. Biol.* 17, 379–393
- Desmoulière, A., Geinoz, A., Gabbiani, F., and Gabbiani, G. (1993). Transforming growth factor- β 1 induces α -smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. *J. Cell. Biol.* 122, 103–111
- Deutsch, A., Friedl, P., Preziosi, L., and Theraulaz, G. (2020). Multi-scale analysis and modelling of collective migration in biological systems. *Phil. Trans. R. Soc. B* 375, 20190377
- Diesboeck, T., Wang, Z., Macklin, P., and Cristini, V. (2011). Multiscale cancer modelling. *Annu. Rev. Biomed. Eng.* 13, 127–155
- Duddu, R., Bordas, S., Chopp, D., and Moran, B. (2008). A combined extended finite element and level set method for biofilm growth. *International Journal for Numerical Methods in Engineering* 74, 848–870
- Duprez, M., Bordas, S. P. A., Bucki, M., Bui, H. P., Chouly, F., Lleras, V., et al. (2020). Quantifying discretization errors for soft tissue simulation in computer assisted surgery: A preliminary study. *Applied Mathematical Modelling* 77, 709–723

- Dvorak, H. (1986). Tumors: wounds that do not heal: similarities between tumor stroma generation and wound healing. *N. Engl. J. Med.* 315, 1650–1659
- Eftimie, R. and Barelle, C. (2021). Mathematical investigation of innate immune responses to lung cancer: the role of macrophages with mixed phenotypes. *J. Theor. Biol.* 524, 110739
- Eftimie, R., Bramson, J., and Earn, D. (2011). Interactions between the immune system and cancer: a brief review of non-spatial mathematical models. *Bull. Math. Biol.* 73, 2–32
- Eftimie, R., Mavrodin, A., and Bordas, S. (2022). From digital control to digital twins in medicine: a brief review and future perspectives. *Advances in Applied Mechanics* 56, 1–49
- Enderling, H. and Chaplain, M. (2014). Mathematical modelling of tumour growth and treatment. *Current Pharmaceutical Design* 20, 4934–4940
- Flegg, J., Menon, S., Maini, P., and McElwain, D. (2015). On the mathematical modeling of wound healing angiogenesis in skin as a reaction-transport process. *Frontiers in Physiology* 6, 1–17
- Friedman, A. (2004). A hierarchy of cancer models and their mathematical challenges. *DCDS-B* 4, 147–159
- Fujiwara, M., Muragaki, Y., and Ooshima, A. (2005). Upregulation of transforming growth factor-beta1 and vascular endothelial growth factor in cultured keloid fibroblasts: relevance to angiogenic activity. *Arch. Dermatol. Res.* 297, 161–169
- Gaffney, E. A., Pugh, K., Maini, P. K., and Arnold, F. (2002). Investigating a simple model of cutaneous wound healing angiogenesis. *J. Math. Biol.* 45, 337–374
- Ghazawi, F., and M.S. Gilardino, R. Z., Sasseville, D., and Jafarian, F. (2018). Insights into the pathophysiology of hypertrophic scars and keloids: How do they differ? *Advances in Skin & Wound Care* 31, 582 – 594
- Greenspan, H. (1972). Models for the growth of a solid tumour by diffusion. *Studies in Applied Mathematics* 51, 317–340
- Greenspan, H. (1976). On the growth and stability of cell cultures and solid tumours. *J. Theor. Biol.* 56, 229–242
- Haisa, M., Okochi, H., and Grotendorst, G. (1994). Elevated levels of PDGF alpha receptors in keloid fibroblasts contribute to an enhanced response to PDGF. *J. Invest. Dermatol.* 103, 560–563
- Hajdu, S. (2011). A note from history: landmarks in history of cancer, part 1. *Cancer* 117, 1097–1102
- He, Y., Merin, M., Sharon, V., and Maverakis, E. (2011). Eruptive keloids associated with breast cancer: a paraneoplastic phenomenon? *Acta Derm Venereol* 91, 480–481
- Huang, C. and Ogawa, R. (2022). Roles of inflammasomes in keloids and hypertrophic scars – lessons learned from chronic diabetic wounds and skin fibrosis. *Int J Mol Sci.* 23, 6820
- Jagadeesan, J. and Bayat, A. (2007). Transforming growth factor beta ($\text{tgf}\beta$) and keloid disease. *Int. J. Surg. Lond. Engl.* 5, 278–285
- Jorgensen, S. and Sanders, J. (2016). Mathematical models of wound healing and closure: a comprehensive review. *Med. Biol. Eng. Comput.* 54, 1297–1316
- Jumper, N., Paus, R., and Bayat, A. (2015). Functional histopathology of keloid disease. *Histol. Histopathol.* 30
- Kanitakis, J. (2013). Keloidal dermatofibroma: report of a rare dermatofibroma variant in a young white woman. *Am J Dermatopathol.* 35, 400–401
- Kather, J., Poleszczuk, J., Suarez-Carmona, M., Krisam, J., Charoentong, P., Valous, N., et al. (2017). In silico modeling of immunotherapy and stroma-targeting therapies in human colorectal cancer. *Cancer Res.* 77, 6442–6452

- Kim, M., Reed, D., and Rejniak, K. (2014). The formation of tight tumour clusters affects the efficacy of cell cycle inhibitors: a hybrid model study. *J. Theor. Biol.* 352, 31–50
- Kim, S. (2021). Management of keloid scars: noninvasive and invasive treatments. *Arch. Plast. Surg.* 48, 149–157
- Kim, Y., Stolarska, M. A., and Othmer, H. G. (2007). A hybrid model for tumor spheroid growth in vitro i: theoretical development and early results. *Mathematical Models and Methods in Applied Sciences* 17, 1773–1798. doi:10.1142/S0218202507002479
- Kimura, K., Inadomi, T., Yamauchi, W., Yoshida, Y., Kashimura, T., and Terui, T. (2014). Dermatofibrosarcoma protuberans on the chest with a variety of clinical features masquerading as a keloid: is the disease really protuberant? *Ann Dermatol.* 26, 643–645
- Kirshtein, A., Akbarinejad, S., Hao, W., Le, T., Su, S., Aronow, R., et al. (2020). Data driven mathematical model of colon cancer progression. *J. Clin. Med.* 9, 3947
- Koppenol, D., Vermolen, F., Niessen, F., van Zuijlen, P., and Vuijk, K. (2017). A mathematical model for the simulation of the formation and the subsequent regression of hypertrophic scar tissue after dermal wounding. *Biomech Model Mechanobiol.* 16, 15–32
- Kuehlmann, B., Bonham, C., Zucal, I., Prantl, L., and Gurtner, G. (2020). Mechanotransduction in wound healing and fibrosis. *J. Clin. Med.* 9, 1423
- Laird, A., Tyler, S., and Barton, A. (1965). Dynamics of normal growth. *Growth* 21, 233–248
- Li, X., Wang, Y., Yuan, B., Yang, H., and Qiao, L. (2017). Status of M1 and M2 type macrophages in keloid. *Int J Clin Exp Pathol* 10, 11098–11105
- Limandjaja, G., Nielsen, F., Scheper, R., and Gibbs, S. (2020a). The keloid disorder: heterogeneity, histopathology, mechanisms and models. *Front. Cell Dev. Biol.* 8, 360
- Limandjaja, G., Niessen, F., Scheper, R., and Gibbs, S. (2020b). The keloid disorder: Heterogeneity, histopathology, mechanisms and models. *Frontiers in Cell and Developmental Biology* 8
- Limandjaja, G., Niessen, F., Scheper, R., and S.Gibbs (2021). Hypertrophic scars and keloids: Overview of the evidence and practical guide for differentiating between these abnormal scars. *Exp. Dermatology* 30, 146–161
- Liu, X., Fang, J., Huang, S., Wu, X., Xie, X., Wang, J., et al. (2021). Tumour-on-a-chip: from bioinspired design to biomedical application. *Microsystems & Nanoengineering* 7, 50
- Lonardo, A. D., Nasi, S., and Pulciani, S. (2015). Cancer: we should not forget the past. *J. Cancer* 6, 29–39
- Lu, F., Gao, J., Ogawa, R., Hyakusoku, H., and Ou, C. (2007). Biological differences between fibroblasts derived from peripheral and central areas of keloid tissues. *Plast. Reconstr. Surg.* 120, 625–630
- Macnamara, C. (2021). Biomechanical modelling of cancer: agent-based force-based models of solid tumours within the context of the tumour microenvironment. *Comput. Syst. Oncol.* 1, e1018
- Maini, P. and Baker, R. (2014). Modelling collective cell motion in biology. *Advances in Applied Mathematics* , 1–11
- Marie, N., Lejeune, A., Chouly, F., Chambert, J., and Jacquet, E. (2022). Dwr error estimator for the biomechanics of the skin with a keloid scar. FigShare repository (DOI:10.6084/m9.figshare.19372280.v1)
- Maroudas-Sacks, Y. and Zemel, A. (2018). theoretical analysis of stress distribution and cell polarisation surrounding a model wound. *Biophysical Journal* 115, 398–410
- Marttala, J., Andrews, J., Rosenbloom, J., and Uitto, J. (2016). Keloids: animal models and pathological equivalents to study tissue fibrosis. *Matrix Biology* 51, 47–54

- McDougall, S., Dallon, J., Sherratt, J. A., and Maini, P. (2006). Fibroblast migration and collagen deposition during dermal wound healing: mathematical modelling and clinical implications. *Phil. Trans. R. Soc. A* 364, 1385–1405
- Metzcar, J., Wang, Y., Heiland, R., and Macklin, P. (2019). A review of cell-based computational modelling in cancer biology. *Clin. Cancer Inform.* 3, 1–13
- Mi, Q., Riviere, B., Clermont, G., Steed, D., and Vodovotz, Y. (2007). Agent-based model of inflammation and wound healing: insights into diabetic foot ulcer pathology and the role of transforming growth factor-beta1. *Wound Repair Regen.* 15, 671–682
- Monfared, G. S., P.Ertl, and Rothbauer, M. (2021). Microfluidic and lab-on-a-chip systems for cutaneous wound healing studies. *Pharmaceutics* 13, 793
- Monte, U. D. (2009). Does the cell number 10^9 still really fit one gram of tumour tissue? *Cell Cycle* 8, 505–506
- Niessen, F., Spauwen, P., and Schalkwijk, J. (1999). On the nature of hypertrophic scars and keloids: a review. *Plast. Reconstr. Surg.* 104, 1435–1458
- Ogawa, R. (2011). Mechanobiology of scarring. *Wound Repair Regen* 19, s2–s9
- Ogawa, R., Okai, K., Tokumura, F., Mori, K., Ohmori, Y., and H. Hyakusoku, C. H., et al. (2012). The relationship between skin stretching/contraction and pathologic scarring: the important role of mechanical forces in keloid generation. *Wound Repair Regen.* 20, 149–157
- Olsen, L., Sherratt, J., and Maini, P. (1996). A mathematical model for fibro-proliferative wound healing disorders. *Bull. Math. Biol.* 58, 787–808
- Pettet, G., Chaplain, M., McElwain, D., and Byrne, H. (1996). On the role of angiogenesis in wound healing. *Proc. R. Soc. Lond. B* 263, 1487–1493
- Quaranta, V., Rejniak, K., Gerlee, P., and Anderson, A. (2008). Invasion emerges from cancer cell adaptation to competitive microenvironments: quantitative predictions from multiscale mathematical models. *Semin. Cancer Biol.* 18, 338–348
- Rejniak, K. and Anderson, A. (2011). Hybrid models of tumour growth. *Wiley Interdiscip. Rev. Syst. Biol. Med.* 3, 115–125
- Renardy, M., Hult, C., Evans, S., Linderman, J., and Kirschner, D. (2019). Global sensitivity analysis of biological multiscale models. *Current Opinion in Biomedical Engineering* 11, 109–116
- Roose, T., Chapman, S., and Maini, P. (2007). Mathematical models of avascular tumour growth. *SIAM Review.* 49, 179–208
- Schäffer, M. R., Efron, P. A., Thornton, F. J., Klingel, K., Gross, S. S., and Barbul, A. (1997a). Nitric oxide, an autocrine regulator of wound fibroblast synthetic function. *The Journal of Immunology* 158, 2375–2381
- Schäffer, M. R., Tantry, U., van Wesep, R. A., and Barbul, A. (1997b). Nitric oxide metabolism in wounds. *Journal of Surgical Research* 71, 25–31. doi:10.1006/jsre.1997.5137
- Sfakianakis, N. and Chaplain, M. (2021). Mathematical modelling of cancer invasion: a review. In *Methods of Mathematical Oncology. MMDS 2020. Springer Proceedings in Mathematics & Statistics*, eds. T. Suzuki, C. Poignard, M. Chaplain, and V. Quaranta. vol. 370, 153–172
- Shan, M., Liu, H., Song, K., Liu, S., Hao, Y., and Wang, Y. (2022). Immune-related gene expression in skin, inflamed and keloid tissue from patients with keloids. *Oncology Letters* 23, 72
- Sherratt, J. and Dallon, J. (2002). Theoretical models of wound healing: past successes and future challenges. *C.R. Biologies* 325, 557–564
- Sherratt, J. A. and Murray, J. (1990). Models of epidermal wound healing. *Proc. R. Soc. Lond B.* 241, 29–36

- Shi, C., Zhao, Y.-P., Ge, P., and Huang, G.-B. (2019). Therapeutic effect of interleukin-10 in keloid fibroblasts by suppression of $\text{tgf-}\beta/\text{smad}$ pathway. *Eur. Rev. Med. Pharmacol. Sci.* 23, 9085–9092
- Shuttleworth, R. and Trucu, D. (2019). Multiscale modelling of fibres dynamics and cell adhesion within moving boundary cancer invasion. *Bull. Math. Biol.* 81, 2176–2219
- Stern, J., Christley, S., Zaborina, O., Alverdy, J., and An, G. (2012). Integration of TGF- β and EGFR based signaling pathways using an agent based model of epithelial restitution. *Wound Repair Regen.* 20, 862–871
- Stolarska, M. A., Kim, Y., and Othmer, H. G. (2009). Multi-scale models of cell and tissue dynamics. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 367, 3525–3553. doi:10.1098/rsta.2009.0095
- Suttho, D., Mankhetkorn, S., Binda, D., Pazart, L., Humbert, P., and Rolin, G. (2017). 3d modelling of keloid scars in vitro by cell and tissue engineering. *Arch. Dermatol. Res.* 309, 55–62
- Sutula, D., Elouneq, A., Sensale, M., Chouly, F., Chambert, J., Lejeune, A., et al. (2020). An open source pipeline for design of experiments for hyperelastic models of the skin with applications to keloids. *Journal of the mechanical behavior of biomedical materials* 112, 103999
- Suveges, S., Chamseddine, I., Rejniak, K., Eftimie, R., and Trucu, D. (2021). Collective cell migration in a fibrous environment: a hybrid multiscale modelling approach. *Front. Appl. Math. Stat.* , 680029
- Tan, S., Khumalo, N., and Bayat, A. (2019). Understanding keloid pathobiology from a quasi-neoplastic perspective: Less of a scar and more of a chronic inflammatory disease with cancer-like tendencies. *Frontiers in Immunology* 10, 1664–3224
- Tepole, A. B. and Kuhl, E. (2016). Computational modeling of chemo-bio-mechanical coupling: a systems-biology approach toward wound healing. *Computer Methods in Biomechanics and Biomedical Engineering* 19, 13–30
- Thackham, J., McElwain, D. S., and Turner, I. (2009). Computational approaches to solving equations arising from wound healing. *Bull. Math. Biol.* 71, 211–246
- Thomlinson, R. and Gray, L. (1955). The histological structure of some human lung cancers and the possible implications for radiotherapy. *Br. J. Cancer* 9, 539–549
- Tongdee, E., Touloei, K., Shitabata, P., Shareef, S., and Maranda, E. (2016). Keloidal atypical fibroxanthoma: case and review of the literature. *Case Rep Dermatol.* 8, 156–163
- Tranquillo, R. and Murray, J. (1993). Mechanistic model of wound contraction. *Journal of Surgical Research* 55, 233–247
- Ulrich, D., Ulrich, F., Unglaub, F., Piatkowski, A., and Pallua, N. (2010). Matrix metalloproteinases and tissue inhibitors of metalloproteinases in patients with different types of scars and keloids. *J. Plast. Reconstr. Aesthet. Surg.* 63, 1015–1021
- Urcun, S., Rohan, P.-Y., Skalli, W., Nassoy, P., Bordas, S. P. A., and Sciumè, G. (2021). Digital twinning of cellular capsule technology: Emerging outcomes from the perspective of porous media mechanics. *PLOS ONE* 16, 1–30. doi:10.1371/journal.pone.0254512
- Venkatesh, K., Raza, M., and Kvedar, J. (2022). Heath digital twins as tools for precision medicine: consideration for computation, implementation, and regulation. *npj Digital Medicine* 5, 150
- Vincent, A., Phan, T., Mukhopadhyay, A., Lim, H., Halliwell, B., and Wong, K. (2008). Human skin keloid fibroblasts display bioenergetics of cancer cells. *J Invest Dermatol.* 128, 702–709
- Vodovotz, Y., Clermont, G., Chow, C., and An, G. (2004). Mathematical models of acute inflammatory response. *Curr Opin Crit Care* 10, 383
- Wang, Z.-C., Zhao, W.-Y., Cao, Y., Liu, Y.-Q., Sun, Q., Shi, P., et al. (2020). The roles of inflammation in keloid and hypertrophic scars. *Front. Immunol.* 11, 603187

- Waugh, H. and Sherratt, J. (2006). Macrophage dynamics in diabetic wound healing. *Bull. Math. Biol.* 68, 197–207
- Webb, G. (2022). The force of cell-cell adhesion in determining the outcome in a nonlocal advection diffusion model of wound healing. *Mathematical Biosciences and Engineering* 19, 8689–8704
- [Dataset] WHO, I. (2022). Icd-11 for mortality and morbidity statistics
- Xu, X., Gu, S., Huang, X., Ren, J., Gu, Y., Wei, C., et al. (2020). The role of macrophages in the formation of hypertrophic scars and keloids. *Burns & Trauma* 8. doi:10.1093/burnst/tkaa006
- Xue, M. and Jackson, C. (2015). Extracellular matrix reorganization during wound healing and its impact on abnormal scarring. *Adv. Wound Care* 4, 119–136
- Yankeelov, T., Atuegwu, N., Hormuth, D., Weis, J., Barnes, S., Miga, M., et al. (2013). Clinically relevant modelling of tumour growth and treatment response. *Sci. Transl. Med.* 5, 187ps9
- Yin, A., Moes, D., van Hasselt, J., Swen, J., and Guchelaar, H.-J. (2019). A review of mathematical models for tumour dynamics and treatment resistance evolution of solid tumours. *CPT Pharmacometrics Syst. Pharmacol.* 8, 720–737
- Zhang, T., Wang, X.-F., Wang, Z.-C., Lou, D., Fang, Q.-Q., Hu, Y.-Y., et al. (2020). Current potential therapeutic strategies targeting the $\text{tgf-}\beta/\text{smad}$ signaling pathway to attenuate keloid and hypertrophic scar formation. *Biomed. Pharmacother. Biomedecine Pharmacother.* 129, 110287
- Zhou, B., Gao, Z., Liu, W., Wu, X., and Wang, W. (2022). Important role of mechanical microenvironment on macrophage dysfunction during keloid pathogenesis. *Experimental Dermatology* 31, 375–380. doi:https://doi.org/10.1111/exd.14473
- Zhou, Y., Sun, Y., Hou, W., Ma, L., Tao, Y., Li, D., et al. (2020). The JAK2/STAT3 pathway inhibitor, AG490, suppresses the abnormal behavior of keloid fibroblasts in vitro. *Int. J. Mol. Med.* 46, 191–200
- Zirardo, C., Mi, Q., An, G., and Vodovotz, Y. (2013). Computational modeling of inflammation in wound healing. *Adv Wound Care (New Rochelle)* 2, 527–537