



**HAL**  
open science

## Macromolecular organic matter in samples of the asteroid (162173) Ryugu

Hikaru Yabuta, George D Cody, Cécile Engrand, Yoko Kebukawa, Bradley de Gregorio, Lydie Bonal, Laurent Remusat, Rhonda Stroud, Eric Quirico, Larry Nittler, et al.

► **To cite this version:**

Hikaru Yabuta, George D Cody, Cécile Engrand, Yoko Kebukawa, Bradley de Gregorio, et al.. Macromolecular organic matter in samples of the asteroid (162173) Ryugu. *Science*, 2023, 379 (6634), pp.eabn9057. 10.1126/science.abn9057. hal-04034418

**HAL Id: hal-04034418**

**<https://cnrs.hal.science/hal-04034418v1>**

Submitted on 21 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

## Macromolecular organic matter in samples of the asteroid (162173) Ryugu

Hikaru Yabuta<sup>1\*§</sup>, George D. Cody<sup>2</sup>, Cécile Engrand<sup>3</sup>, Yoko Kebukawa<sup>4</sup>, Bradley De Gregorio<sup>5</sup>,  
Lydie Bonal<sup>6</sup>, Laurent Remusat<sup>7</sup>, Rhonda Stroud<sup>5#</sup>, Eric Quirico<sup>6</sup>, Larry Nittler<sup>2¶</sup>, Minako  
Hashiguchi<sup>8</sup>, Mutsumi Komatsu<sup>9, 10</sup>, Taiga Okumura<sup>11</sup>, Jérémie Mathurin<sup>12</sup>, Emmanuel Dartois<sup>13</sup>,  
5 Jean Duprat<sup>7</sup>, Yoshio Takahashi<sup>11, 14</sup>, Yasuo Takeichi<sup>14</sup>, David Kilcoyne<sup>15</sup>, Shohei Yamashita<sup>14</sup>,  
Alexandre Dazzi<sup>12</sup>, Ariane Deniset-Besseau<sup>12</sup>, Scott Sandford<sup>16</sup>, Zita Martins<sup>17</sup>, Yusuke  
Tamenori<sup>18</sup>, Takuji Ohigashi<sup>19</sup>, Hiroki Suga<sup>18</sup>, Daisuke Wakabayashi<sup>14</sup>, Maximilien Verdier-  
Paoletti<sup>7</sup>, Smail Mostefaoui<sup>7</sup>, Gilles Montagnac<sup>20</sup>, Jens Barosch<sup>2</sup>, Kanami Kamide<sup>1</sup>, Miho  
Shigenaka<sup>1</sup>, Laure Bejach<sup>3</sup>, Megumi Matsumoto<sup>21</sup>, Yuma Enokido<sup>21</sup>, Takaaki Noguchi<sup>22</sup>,  
10 Hisayoshi Yurimoto<sup>23</sup>, Tomoki Nakamura<sup>21</sup>, Ryuji Okazaki<sup>24</sup>, Hiroshi Naraoka<sup>24</sup>, Kanako  
Sakamoto<sup>25</sup>, Harold C. Connolly Jr.<sup>26</sup>, Dante S. Lauretta<sup>27</sup>, Masanao Abe<sup>25, 28</sup>, Tatsuaki Okada<sup>25</sup>,  
<sup>29</sup>, Toru Yada<sup>25</sup>, Masahiro Nishimura<sup>25</sup>, Kasumi Yogata<sup>25</sup>, Aiko Nakato<sup>25</sup>, Miwa Yoshitake<sup>25</sup>,  
Ayako Iwamae<sup>30</sup>, Shizuhō Furuya<sup>11</sup>, Kentaro Hatakeda<sup>30</sup>, Akiko Miyazaki<sup>25</sup>, Hiromichi  
Soejima<sup>30</sup>, Yuya Hitomi<sup>30</sup>, Kazuya Kumagai<sup>30</sup>, Tomohiro Usui<sup>25</sup>, Tasuku Hayashi<sup>25</sup>, Daiki  
15 Yamamoto<sup>25</sup>, Ryota Fukai<sup>25</sup>, Seiji Sugita<sup>11, 31</sup>, Kohei Kitazato<sup>32</sup>, Naru Hirata<sup>32</sup>, Rie Honda<sup>33</sup>,  
Tomokatsu Morota<sup>11</sup>, Eri Tatsumi<sup>34</sup>, Naoya Sakatani<sup>35</sup>, Noriyuki Namiki<sup>28, 36</sup>, Koji Matsumoto<sup>28</sup>,  
<sup>36</sup>, Rina Noguchi<sup>37</sup>, Koji Wada<sup>31</sup>, Hiroki Senshu<sup>31</sup>, Kazunori Ogawa<sup>38</sup>, Yasuhiro Yokota<sup>25</sup>,  
Yoshiaki Ishihara<sup>38</sup>, Yuri Shimaki<sup>25</sup>, Manabu Yamada<sup>31</sup>, Chikatoshi Honda<sup>32</sup>, Tatsuhiro  
Michikami<sup>39</sup>, Moe Matsuoka<sup>40</sup>, Naoyuki Hirata<sup>41</sup>, Masahiko Arakawa<sup>41</sup>, Chisato Okamoto<sup>41†</sup>,  
20 Masateru Ishiguro<sup>42</sup>, Ralf Jaumann<sup>43</sup>, Jean-Pierre Bibring<sup>44</sup>, Matthias Grott<sup>45</sup>, Stefan Schröder<sup>46</sup>,  
Katharina Otto<sup>1, 45</sup>, Cedric Pilorget<sup>44, 47</sup>, Nicole Schmitz<sup>45</sup>, Jens Biele<sup>48</sup>, Tra-Mi Ho<sup>49</sup>, Aurélie  
Moussi-Soffys<sup>50</sup>, Akira Miura<sup>25</sup>, Hirotomo Noda<sup>28, 36</sup>, Tetsuya Yamada<sup>25</sup>, Keisuke Yoshihara<sup>25</sup>,  
Kosuke Kawahara<sup>25</sup>, Hitoshi Ikeda<sup>25</sup>, Yukio Yamamoto<sup>25</sup>, Kei Shirai<sup>41</sup>, Shota Kikuchi<sup>31, 36</sup>,  
Naoko Ogawa<sup>25</sup>, Hiroshi Takeuchi<sup>25, 28</sup>, Go Ono<sup>51</sup>, Yuya Mimasu<sup>25</sup>, Kent Yoshikawa<sup>51</sup>, Yuto  
25 Takei<sup>25</sup>, Atsushi Fujii<sup>25</sup>, Yu-ichi Iijima<sup>25†</sup>, Satoru Nakazawa<sup>25</sup>, Satoshi Hosoda<sup>25</sup>, Takahiro  
Iwata<sup>25, 28</sup>, Masahiko Hayakawa<sup>25</sup>, Hirotaka Sawada<sup>38</sup>, Hajime Yano<sup>25, 28</sup>, Ryudo Tsukizaki<sup>25</sup>,  
Masanobu Ozaki<sup>25, 28</sup>, Fuyuto Terui<sup>52</sup>, Satoshi Tanaka<sup>25, 28</sup>, Masaki Fujimoto<sup>25</sup>, Makoto  
Yoshikawa<sup>25, 28</sup>, Takanao Saiki<sup>25</sup>, Shogo Tachibana<sup>11, 25</sup>, Sei-ichiro Watanabe<sup>8</sup>, Yuichi Tsuda<sup>25</sup>

<sup>1</sup>Department of Earth and Planetary Systems Science, Hiroshima University, Hiroshima 739-  
30 8526, Japan.

<sup>2</sup>Earth and Planets Laboratory, Carnegie Institution for Science, Washington, DC 20015,  
USA.

<sup>3</sup>Laboratoire de Physique des 2 Infinis Irène Joliot-Curie, Université Paris-Saclay, Centre  
National de la Recherche Scientifique, 91405 Orsay, France.

<sup>4</sup>Faculty of Engineering, Yokohama National University, Yokohama 240-8501, Japan.

<sup>5</sup>Materials Science and Technology Division, US Naval Research Laboratory, Washington,  
DC 20375, USA.

<sup>6</sup>Institut de Planétologie et d'Astrophysique, Université Grenoble Alpes, 38000 Grenoble,  
France.

<sup>7</sup>Institut de Mineralogie, Physique des Matériaux et Cosmochimie, Museum National  
d'Histoire Naturelle, Centre National de la Recherche Scientifique, Sorbonne Université,  
40 75231 Paris, France.

<sup>8</sup>Graduate School of Environmental Studies, Nagoya University, Nagoya 464-8601, Japan.

<sup>9</sup>Center for University-Wide Education, Saitama Prefectural University, Saitama 343-8540, Japan.

<sup>10</sup>Department of Earth Sciences, Waseda University, Tokyo 169-8050, Japan.

5 <sup>11</sup>Department of Earth and Planetary Science, The University of Tokyo, Tokyo 113-0033, Japan.

<sup>12</sup>Institut Chimie Physique, Université Paris-Saclay, Centre National de la Recherche Scientifique, 91405 Orsay, France.

10 <sup>13</sup>Institut des Sciences Moléculaires d'Orsay, Université Paris-Saclay, Centre National de la Recherche Scientifique, 91405 Orsay, France.

<sup>14</sup>Institute of Materials Structure Science, High Energy Accelerator Research Organization, Tsukuba 305-0801, Japan.

<sup>15</sup>Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley 94720-8229, USA.

15 <sup>16</sup>NASA Ames Research Center, Moffett Field 94035-1000, USA.

<sup>17</sup>Centro de Química Estrutural, Institute of Molecular Sciences and Department of Chemical Engineering, Instituto Superior Técnico, Universidade de Lisboa, Lisboa 1049-001, Portugal.

<sup>18</sup>Japan Synchrotron Radiation Research Institute, Hyogo 679-5198, Japan.

<sup>19</sup>Institute for Molecular Science, UVSOR Synchrotron Facility, Okazaki 444-8585, Japan.

20 <sup>20</sup>École normale supérieure de Lyon, University Lyon 1, 69342 Lyon, France.

<sup>21</sup>Department of Earth Science, Tohoku University, Sendai 980-8578, Japan.

<sup>22</sup>Department of Earth and Planetary Sciences, Kyoto University, Kyoto 606-8502, Japan.

<sup>23</sup>Department of Earth and Planetary Sciences, Hokkaido University, Sapporo 060-0810, Japan.

25 <sup>24</sup>Department of Earth and Planetary Sciences, Kyushu University, Fukuoka 819-0395, Japan.

<sup>25</sup>Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency (JAXA), Sagami-hara 252-5210, Japan.

<sup>26</sup>Department of Geology, School of Earth and Environment, Rowan University, Glassboro 08028, USA.

30 <sup>27</sup>Lunar and Planetary Laboratory, University of Arizona, Tucson 85721, USA.

<sup>28</sup>School of Physical Sciences, The Graduate University for Advanced Studies, Hayama 240-0193, Japan.

<sup>29</sup>Department of Chemistry, The University of Tokyo, Tokyo 113-0033, Japan.

<sup>30</sup>Marine Works Japan Ltd., Yokosuka 237-0063, Japan.

35 <sup>31</sup>Planetary Exploration Research Center, Chiba Institute of Technology, Narashino 275-0016, Japan.

<sup>32</sup>Aizu Research Cluster for Space Science, University of Aizu, Aizu-Wakamatsu 965-8580, Japan.

<sup>33</sup>Center of Data Science, Ehime University, Matsuyama 790-8577, Japan.

<sup>34</sup>Instituto de Astrofísica de Canarias, University of La Laguna, Tenerife E-38205, Spain.

5 <sup>35</sup>Department of Physics, Rikkyo University, Tokyo 171-8501, Japan.

<sup>36</sup>Research of Interior Structure and Evolution of Solar System Bodies, National Astronomical Observatory of Japan, Mitaka 181-8588, Japan.

<sup>37</sup>Faculty of Science, Niigata University, Niigata 950-2181, Japan.

<sup>38</sup>JAXA Space Exploration Center, JAXA, Sagamihara 252-5210, Japan.

10 <sup>39</sup>Faculty of Engineering, Kindai University, Higashi-Hiroshima 739-2116, Japan.

<sup>40</sup>Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology, Tsukuba 305-8567, Japan.

<sup>41</sup>Department of Planetology, Kobe University, Kobe 657-8501, Japan.

15 <sup>42</sup>Department of Physics and Astronomy, Seoul National University, Seoul 08826, Republic of Korea.

<sup>43</sup>Freie Universität Berlin, Institute of Geological Sciences, Planetary Sciences and Remote Sensing, Berlin 12249, Germany.

<sup>44</sup>Institut d'Astrophysique Spatiale, Université Paris-Saclay, Centre National de la Recherche Scientifique, 91405 Orsay, France.

20 <sup>45</sup>Institute of Planetary Research, German Aerospace Center, Berlin 12489, Germany.

<sup>46</sup>Department of Computer Science, Electrical and Space Engineering, Luleå University of Technology, 98128 Kiruna, Sweden.

<sup>47</sup>Institut Universitaire de France, 75005 Paris, France.

<sup>48</sup>Microgravity User Support Centre, German Aerospace Center, Cologne 53147, Germany.

25 <sup>49</sup>Institute of Space Systems, German Aerospace Center, Bremen 28359, Germany.

<sup>50</sup>Centre National d'Études Spatiales, 31401 Toulouse, France.

<sup>51</sup>Research and Development Directorate, Japan Aerospace Exploration Agency (JAXA), Sagamihara 252-5210, Japan.

30 <sup>52</sup>Department of Mechanical Engineering, Kanagawa Institute of Technology, Atsugi 243-0292, Japan.

\*Corresponding author. Email: [hyabuta@hiroshima-u.ac.jp](mailto:hyabuta@hiroshima-u.ac.jp)

† Deceased

35 §Current address: International Institute for Sustainability with Knotted Chiral Meta Matter, Hiroshima University, Hiroshima 739-8526, Japan.

#Current address: Buseck Center for Meteorite Studies, Arizona State University, Tempe AZ 85827, USA.

<sup>†</sup>Current address: School of Earth and Space Exploration, Arizona State University, Tempe AZ 85827, USA.

5 **Samples of the carbonaceous asteroid (162173) Ryugu were collected and brought to Earth by the Hayabusa2 spacecraft. We investigate the macromolecular organic matter in Ryugu samples, finding they contain aromatic and aliphatic carbon, ketone and carboxyl functional groups. The spectroscopic features of the organic matter are consistent with those in chemically primitive carbonaceous chondrite meteorites that experienced parent body aqueous alteration (reactions with liquid water). The morphology of the organic carbon includes nanoglobules and diffuse carbon associated with phyllosilicate and carbonate minerals. Deuterium and/or nitrogen-15 enrichments indicate the organic matter formed in a cold molecular cloud or the pre-solar nebula. The diversity of the organic matter indicates variable levels of aqueous alteration on Ryugu's parent body.**

15 Organic compounds in asteroids and comets were produced and modified within the pre-solar molecular cloud, the protoplanetary disk, during formation of planetesimals in the early Solar System, and their subsequent evolution. Delivery of extraterrestrial organic compounds might have contributed to the habitability of terrestrial planets, including the Earth. Analysis of pristine samples collected from primitive small bodies (asteroids and comets) could provide information on how organic compounds were formed and modified in space, and which organic compounds were supplied to the early Earth. Macromolecular organic matter, a dark, complex, acid-insoluble organic matter (IOM), accounts for most carbon in primitive carbonaceous chondrite meteorites.

25 The Hayabusa2 asteroid sample return mission visited the carbonaceous (C-type) asteroid (162173) Ryugu. The mission goals included investigating the origin and evolution of organic compounds in the early Solar System (1). Remote sensing and lander observations showed that Ryugu is a dark rubble-pile asteroid that contains hydrated minerals on its surface (2-4) and indicated a relationship between C-type asteroids and carbonaceous chondrites (3, 4). The average albedo of Ryugu is consistent with the thermally metamorphosed subgroups of the CI1 and CM2 meteorites (3, 4). CI1 meteorites are Ivuna-type carbonaceous chondrites of petrologic type 1 (extensively altered by aqueous fluids on the parent asteroid), and CM2 meteorites are Mighei-type carbonaceous chondrites of petrologic type 2 (moderately aqueously altered). Spectral and thermal inertia variations indicated that Ryugu consists of boulders that experienced different degrees of thermal and space weathering processes (5-7), derived from a much larger parent body, and potentially additional materials from different asteroids (8).

35 The Hayabusa2 spacecraft collected the surface material from two touchdown sites on Ryugu and returned them to Earth on 2020 December 6. The Ryugu samples exhibit near-infrared absorption features due to OH and carbonate/organic C-H bonds, at 2.7 and 3.4  $\mu\text{m}$  respectively (9, 10), indicating that the Ryugu samples are similar to CI carbonaceous chondrites (9).

40 We sought to determine the distributions and chemical characteristics of macromolecular organic matter in the Ryugu samples. We therefore measured the elemental, isotopic, and functional group compositions, structures, and textures of organic macromolecules from the Ryugu samples. The analytical procedures included micro-Fourier transform infrared (FTIR) spectroscopy, micro-Raman spectroscopy, synchrotron-based scanning transmission X-ray microscopy (STXM), X-ray absorption near edge structure (XANES), scanning transmission

electron microscopy (STEM) coupled with electron energy loss spectroscopy (EELS) and energy dispersive X-ray spectroscopy (EDS), atomic force microscope based infrared (AFM-IR) spectroscopy, and nanometer-scale secondary ion mass spectrometry (NanoSIMS) (11). The analytical workflow (fig. S1) was designed to optimize the use of these complementary techniques.

The samples we used were selected aggregates stored in collection chamber A (from the first touchdown) and collection chamber C (from the second touchdown) of the spacecraft sample catcher (12) (table S1). We studied i) intact grains (taken from aggregates designated A0108 and C0109), ranging from 200 to 900  $\mu\text{m}$  in size per particle; and ii) insoluble carbonaceous residues (fig. S2) isolated by acid treatment of Ryugu aggregates (designated A0106 and C0107). Each of these samples was split into several sub-samples for analysis with different techniques (table S1).

### Structural properties of macromolecular organic matter

To characterize the macromolecular structures of organic matter in Ryugu, we applied micro-Raman spectroscopy. Two peaks, identified as the D- ( $\sim 1350\text{ cm}^{-1}$ ) and G-bands ( $\sim 1580\text{ cm}^{-1}$ ) of polyaromatic molecular structures (13), are present in the Raman spectra of A0108 and C0109 (Fig. 1A and fig. S3A). The spectral features are broad, indicating lattice disorder in the organic macromolecules, and are superimposed on a fluorescence background. The numerical values of the derived spectral parameters of the D- and G-bands, such as their full widths at half maximum ( $\text{FWHM}_{\text{D}}$  and  $\text{FWHM}_{\text{G}}$ , respectively), peak positions ( $\omega_{\text{D}}$ ,  $\omega_{\text{G}}$ ), and intensity ratio ( $I_{\text{D}}/I_{\text{G}}$ ), are similar for all grains examined from the two aggregates (Figs. 1B-D and figs. S3B-D).

The macromolecular structures of organic matter in meteorites reflect the thermal histories of the meteorite parent bodies (13-16). To evaluate the thermal history of Ryugu, we compared the Raman parameters measured from the Ryugu samples to those measured from meteorites (Figs. 1B-D and figs. S3B-D). The closest matches to Ryugu are the primitive CI1 and CM2 carbonaceous chondrites. The Ryugu samples are distinct from carbonaceous chondrites of petrologic type 3 (thermally metamorphosed) (Fig. 1B) and from the thermally metamorphosed subgroup of CM2, such as the Jbilet Winselwan meteorite (Fig. 1C, D), and other ungrouped carbonaceous chondrites of petrologic type 2 (C2), such as the Wisconsin Range (WIS) 91600 and Pecora Escarpment (PCA) 02012 meteorites (Fig. 1C, D). This indicates that the Ryugu samples A0108 and C0109 did not experience long duration radiogenic thermal metamorphism on their parent bodies, as petrologic type 3 chondrites did (17), or impact-induced short duration heating, as experienced by some petrologic type 2 chondrites (18).

### Functional group compositions

We used micro-FTIR spectroscopy to characterize the organic molecules and minerals. The FTIR spectra of the Ryugu grains show bands due to organic aliphatic C-H stretching ( $3000$  to  $2800\text{ cm}^{-1}$ ,  $3.33$  to  $3.57\text{ }\mu\text{m}$ ), aromatic C=C stretching ( $\sim 1600\text{ cm}^{-1}$ ,  $\sim 6.25\text{ }\mu\text{m}$ ), and carbonyl C=O stretching modes ( $\sim 1700\text{ cm}^{-1}$ ,  $\sim 5.88\text{ }\mu\text{m}$ ), as well as bands due to mineral Si-O stretching ( $\sim 1000\text{ cm}^{-1}$ ,  $\sim 10.00\text{ }\mu\text{m}$ ), structural OH stretching of phyllosilicates ( $\sim 3680\text{ cm}^{-1}$ ,  $\sim 2.72\text{ }\mu\text{m}$ ), and the  $\nu_3$  stretching mode of carbonates ( $\sim 1435\text{ cm}^{-1}$ ,  $\sim 6.97\text{ }\mu\text{m}$ ) (Fig. 2A). The spectra also contain bands from interlayer water, of varying intensities, contributing at  $\sim 3300\text{ cm}^{-1}$  ( $\sim 3.03\text{ }\mu\text{m}$ , stretching) and  $1640\text{ cm}^{-1}$  ( $6.10\text{ }\mu\text{m}$ , bending). Part of this water is intrinsic to the Ryugu grains, and part can be attributed to water adsorbed on the grains under atmospheric conditions.

These absorption bands are commonly observed in unheated, aqueously altered carbonaceous chondrites (19, 20), while organic features are weaker in thermally metamorphosed CM chondrites, such as the Jbilet Winselwan meteorite (Fig. 2A). The spectral shape of the OH band in the Ryugu samples is characteristic of Mg-rich phyllosilicates, which have been observed in CI chondrites as saponite and serpentine (21, 22). There were large spectral heterogeneities among the Ryugu grains, but there is no obvious difference in the spectral variations observed between samples from chambers A and C. Compared to CI chondrites, the sulfate S=O stretching band ( $\sim 1100$  to  $1200\text{ cm}^{-1}$ ,  $\sim 9.09$  to  $8.33\text{ }\mu\text{m}$ ) is absent from the spectra of Ryugu samples. The absence of sulfates is consistent with other elemental and mineralogical measurements of Ryugu samples (23, 24). Sulfate can be produced by oxidation of sulfides during terrestrial weathering of the meteorites (25), so the lack of sulfate indicates that the Ryugu samples are pristine (23).

The shapes of the FTIR spectra of the grains are consistent with the reflectance spectra of Ryugu's surface acquired by the Hayabusa2 spacecraft (4). The OH band and aliphatic C-H band features in our absorption spectra are similar to reflectance spectra of other Ryugu samples (9), but we find lower intensities of aliphatic C-H peaks.

The FTIR spectra of insoluble carbonaceous residues obtained from our Ryugu samples show similar functional groups to those of the intact Ryugu grains (Fig. 2B). The aliphatic C-H stretching band ( $3000$  to  $2800\text{ cm}^{-1}$ ,  $3.33$  to  $3.57\text{ }\mu\text{m}$ ) from the carbonaceous residue from Ryugu is more intense than those of IOM from meteorites (26, 27). The peak intensity ratios of  $\text{CH}_2$  to  $\text{CH}_3$  ( $I_{\text{CH}_2}/I_{\text{CH}_3}$ ) of the Ryugu residues are 1.9, while those of IOMs in Murchison and Ivuna meteorites are 1.2 and 1.3, respectively. Because  $I_{\text{CH}_2}/I_{\text{CH}_3}$  ratios are proportional to the molar ratios of  $\text{CH}_2$  to  $\text{CH}_3$  ( $\text{CH}_2/\text{CH}_3$ ), we infer the  $\text{CH}_2/\text{CH}_3$  ratios of the residue, which are higher than those of meteoritic IOMs. This could indicate that Ryugu's organic matter contains longer aliphatic chains, or aliphatic chains with a higher degree of cross-linking. The Ryugu carbonaceous residue also exhibits an absorption band of C=O ( $\sim 1670\text{ cm}^{-1}$ ,  $\sim 5.99\text{ }\mu\text{m}$ ) which is not seen in meteoritic IOMs. We assign this C=O band to unsaturated ketones, aldehydes or amides.

## Chemical and morphological variations

### *Macromolecular diversity*

We used synchrotron-based STXM, with spatial resolution of 30 to 50 nm, to produce elemental X-ray maps and XANES spectra. Carbon X-ray maps (Figs. 3A, B) show discrete grains of organic material with sizes of  $\sim 200$  nm. X-ray absorption by carbon atoms is also present in the phyllosilicate matrix, at low levels but nearly ubiquitously (Fig. 3B). Carbon-XANES spectra of the discrete grains show three major peaks, due to aromatic carbon (C=C, 285 eV), aromatic ketone (C=C-C=O, 286.7 eV), and carboxyl (COOH, 288.5 eV) functional groups (Fig. 3E), which have been observed in primitive extraterrestrial carbonaceous matter, such as carbonaceous chondrites (28, 29) and their extracted IOMs (30), interplanetary dust particles (IDPs) (31), dust particles from Comet Wild 2 (32), and Antarctic micrometeorites (AMMs) (33). Some spectra of fine-grained matrices in the Ryugu samples contain an additional peak at 290.4 eV, which corresponds to a  $1s-\pi^*$  transition of carbonate groups, e.g., in calcite and other carbonate minerals. However, our matrix XANES spectra containing this feature lack extended X-ray absorption fine structure (EXAFS) features at higher energies (294-304 eV), indicating that the carbonate is not in a crystalline structure, so is more likely to be molecular carbonate. A similar carbonate feature has previously been reported in clay-bound carbon from extensively

hydrated carbonaceous chondrites, Ivuna (CI1), Orgueil (CI1), and Tagish Lake (C2) (34), and in diffuse carbon (organic matter) in the fine-grained matrices of Renazzo (CR2; a Renazzo-type carbonaceous chondrite of petrologic type 2), Murchison (CM2), and Orgueil (28).

We classify the STXM spectra of Ryugu organic matter based on spectral shape similarities into four representative types: i) Highly aromatic (~25% of individual carbon grains), ii) Aromatic (~35% of individual grains), iii) IOM-like (~40% of individual grains), and iv) Diffuse carbon associated with a molecular carbonate peak (Fig. 3E). Aromatic spectra show higher ratios of aromatic carbon to aromatic ketone than IOM-like spectra, while highly aromatic spectra show a broader peak for aromatic carbon, indicating increased diversity of aromatic structures. The frequency distributions of these classes were similar for both chamber A and C samples. We find a relationship between the morphology of organic matter and XANES spectral shape, with particulate and nanoglobular regions having more frequent aromatic or highly-aromatic XANES spectra, while organic matter dispersed in the matrix more commonly has IOM-like or diffuse carbon spectra. These observations indicate that the molecular functional groups present are influenced by aqueous processing on the asteroid parent body. We did not find any evidence of long-duration thermal metamorphism, such as the  $1s-\sigma^*$  exciton (291.6 eV) peak of graphite or other graphitized carbon materials (35).

Similar characteristics are found for the insoluble carbonaceous residues from Ryugu. Average C-XANES spectra have IOM-like spectral shapes, but with more prominent aromatic C=C, ketone, and carboxyl peaks, compared to IOM from Orgueil and Murchison (Fig. 3E). Several hollow and solid organic nanoglobules are apparent in the X-ray absorption images (Figs. 3C, D), which we confirmed using TEM imaging (Figs. 4A, B). Their XANES spectra were either IOM-like, aromatic, or highly-aromatic (Fig. 3E). The 290.4 eV carbonate feature was not observed in the insoluble carbonaceous residue, perhaps because this organic phase was incorporated into phyllosilicate interlayers, so was removed or destroyed during the acid extraction process.

Nitrogen-XANES spectra (fig. S4A) of the Ryugu intact grains and insoluble residue did not show clear absorption peaks, indicating low abundances of N-rich Ryugu organics. This is consistent with carbonaceous chondrites, where nitrogen-rich particles are only occasionally observed (32). Oxygen-XANES spectra (fig. S4B) of Ryugu organic matter often contain a peak at ~531.3eV, corresponding to carbonyl C=O bonds in the ketone and carboxyl functional groups (36). The peak intensity of the carbonyl absorption, relative to the main oxygen  $1s-\sigma^*$  peak in Ryugu, is similar to that of aqueously altered carbonaceous chondrites. We cannot determine whether these carbonyl functional groups are also present in the phyllosilicate-bound diffuse organic matter, because its oxygen-XANES spectrum is dominated by the surrounding phyllosilicate.

### *Nanoscale morphologies*

We performed TEM and STEM-EELS-EDS analysis (Fig. 4) on ultrathin sections of particles and carbonaceous residues, including from some of the same sections we analyzed previously with STXM. The two most abundant organic microstructures are nanoglobules (Figs. 4A, B), and diffuse carbon mixed with phyllosilicates, i.e, clay-bound carbon (Figs 4C, E). Other microstructures include dense, irregularly shaped particles (Fig. 4B), diffuse organic matter trapped in vesicles in carbonate grains (Figs. 4D, F), and organic matter coatings on sulfide grains. The presence of organic matter associated with Mg-rich phyllosilicates and carbonates, which likely formed through aqueous alteration (24), implies that much of the organic material



was altered by low temperature aqueous processing on Ryugu's parent body. However, we also find nanodiamonds associated with amorphous organic carbon, which probably formed in the interstellar medium or the Solar System's protoplanetary nebula (37, 38). The insoluble carbonaceous residues contain large nanoglobules, and fluffy or porous material, comprising demineralized grain coatings, intergranular material and small nanoglobules. These microstructures are present in samples from both chambers A and C; they are consistent in size and shape with those in CI and CM chondrites (28, 34).

Organic nanoglobules have previously been found in early Solar System materials, such as primitive carbonaceous chondrites (39-42), dust particles from Comet Wild 2 (32), IDPs (43), and AMMs (33, 44). The nanoglobules in Ryugu grains occur in solid and hollow form, typically 50 to 500 nm in diameter, with a few reaching 2000 nm, consistent with nanoglobules in other extraterrestrial materials. The carbon EELS spectra of many of the Ryugu nanoglobules are dominated by an aromatic carbon peak at around 285 eV (Fig. 4G), corresponding to the aromatic-rich carbon-XANES spectra (Fig. 3E). Other prominent peaks are due to aromatic ketone (286.7 eV) and carboxyl (288.5 eV). The fluffy material in the insoluble carbonaceous residues shows these three peaks in varying intensities. The presence of both aromatic- and IOM-like nanoglobules of various sizes indicates that Ryugu experienced heterogeneous aqueous alteration on its parent body. Correlated STEM-NanoSIMS measurements of a 2000-nm nanoglobule (particle A0108-37; figs. S6 and S7) show high abundances of aromatic carbon, and isotopically anomalous H and N, indicating preservation of material from the interstellar medium or protoplanetary nebula.

The widespread diffuse carbon mixed into phyllosilicates could have been formed from soluble molecules intercalated into clays, through oxidation during aqueous alteration (34). Alternatively, the clay-bound diffuse carbon might have been released by hydrolysis of macromolecular organic material, during the flow of aqueous fluids (28). The EELS spectra show peaks of either aromatic ketones or aliphatic carbon (287 eV) and carbonate ( $\text{CO}_3$  at 290.4 eV), along with aromatic carbon (285 eV) (Fig. 4G). These EELS data are consistent with the carbon-XANES spectrum of the diffuse carbon (Figs. 3B, E). The relative intensities of the 287 eV and 285 eV peaks vary, compared to the 290.4 eV peak. This is consistent with clay-bound organics in CI chondrites (34), but sometimes is a better match to the ungrouped C2 chondrite Tagish Lake (34). In all cases, Mg is detected in the EDS spectra from the clay-bound organic matter (Fig. 4F), however we cannot determine whether the Mg is associated with the carbonate. We find no evidence for crystalline Mg carbonate in the clay-bound organics, but the  $\text{CO}_3$  feature could be associated with either molecular  $\text{CO}_3$ , or  $\text{MgCO}_3$ , or a combination of these.

Pockets of organic matter occur in vesicles inside carbonates (Fig. 4D). A STEM-EDS image (Fig. 4F) shows this vesicle-bound diffuse organic matter is associated with phyllosilicates and particles of sulfide and Fe-Ni with size of ~10 nm, embedded in a larger grain of calcite. This morphology could indicate carbonate formation from diffuse organic matter. The lack of distinct particle boundaries for the organic matter could indicate it is soluble, and therefore potentially lost during our sample preparation, which used ultra-microtomy with a water bath.

We obtained AFM-IR maps of organic inclusions and nanoglobule-like matter at lateral resolutions of 25 and 50 nm. AFM-IR measurements map specific vibrational modes; we used the C=O and C=C modes. Samples from chambers A and C were analyzed using AFM-IR, both in tapping and contact modes. Figs. 5A and 5H show combined maps of carbonyl C=O ( $1720 \text{ cm}^{-1}$ ,  $5.81 \mu\text{m}$ ), aromatic C=C ( $1600 \text{ cm}^{-1}$ ,  $6.25 \mu\text{m}$ ) and Si-O ( $1020 \text{ cm}^{-1}$ ,  $9.80 \mu\text{m}$ ) modes. The diffuse organic component within the phyllosilicate matrix is evident in both samples. The maps

show small (up to ~100 nm) organic nanoglobule-like inclusions in the chamber A sample (Fig. 5A) and organic inclusions in the chamber C samples (Fig. 5H). The abundances of carbonyl and C=C vary between the organic inclusions. These AFM-IR results are consistent with, and complimentary to, those from STEM-EELS-EDS and STXM-XANES discussed above.

5

## Elemental abundances

The elemental abundances of macromolecular organic matter in meteorites are highly heterogeneous, both between meteorite groups and at the micrometer scale, reflecting the complex histories of these materials (45, 46). We used NanoSIMS to map the H, C, N, O and S elemental abundances of the Ryugu samples.

10

The ratios of O/C, N/C, and S/C in the acid-insoluble carbonaceous residues from Ryugu were estimated from NanoSIMS elemental maps of the major isotopes ( $^{16}\text{O}$ ,  $^{12}\text{C}$ ,  $^{14}\text{N}$ ,  $^{32}\text{S}$ ) and were compared between samples taken from chambers A and C. The bulk O/C ratio ( $0.12 \pm 0.03$ ) in chamber A was consistent with those of CI (0.15 to 0.18), CM (0.11 to 0.23), and CR chondrites (0.11 to 0.22) (45). The bulk O/C ratio ( $0.04 \pm 0.01$ ) in chamber C was three times lower than those values. We cannot determine whether this difference is due to intrinsic differences between the two sampling sites, or heterogeneity within the samples. There are smaller differences in the bulk ratios N/C ( $0.035 \pm 0.006$  (A0106);  $0.021 \pm 0.001$  (C0107)) and S/C ( $0.032 \pm 0.001$  (A0106);  $0.025 \pm 0.001$  (C0107)) between the two chambers, which are within the ranges of those for CI, CM, and CR chondrites ( $0.026 < \text{N/C} < 0.039$  and  $0.02 < \text{S/C} < 0.06$ ) (45).

15

20

We also used STEM-EDS to measure N, O, and S abundances, relative to C, at the nano- to micro-meter scale. An EDS spectrum from a  $\sim 1 \mu\text{m}^2$  region of an ultrathin section of carbonaceous residue from chamber A is shown in Fig. 4H. The average compositions measured for carbonaceous residues are  $\text{C}_{100}\text{N}_1\text{O}_{10}\text{S}_{0.8}$  for chamber A and  $\text{C}_{100}\text{N}_2\text{O}_{10}\text{S}_{0.8}$  for chamber C. The insoluble carbonaceous residues contain some nanoparticulate minerals, including chromite, with varying Al, Mg and Fe, and sulfides with variable Fe, Ni, and (more rarely) CuS. These particles were excluded from the STEM-EDS measurements when  $> 4$  nm in size. These nitrogen abundances are slightly lower than the bulk N/C ratio estimated using NanoSIMS, and consistent with the weak N-XANES absorption of the Ryugu grains.

25

30

## Isotopic compositions

NanoSIMS was also used to measure the isotopic compositions of hydrogen, carbon, and nitrogen. The resulting maps (Fig. 6A-C) show bulk enrichments of D ( $^2\text{H}$ ) and/or  $^{15}\text{N}$  in most analyzed Ryugu particles from aggregates A0108 and C0109, with a high degree of isotopic heterogeneity at the micrometer scale. The NanoSIMS H measurements are dominated by H from the organic matter, but also include H in phyllosilicates and possible contamination by terrestrial water. The isotopic compositions are expressed in delta notation:  $\delta R \equiv (R_{\text{sample}}/R_{\text{standard}} - 1) \times 1000$ , where  $R_{\text{sample}}$  is the isotopic ratio of sample and  $R_{\text{standard}}$  is the ratio of a terrestrial standard. The average measured H isotopic compositions ( $\delta\text{D}$ ) of particles range from  $+254 \pm 52\text{‰}$  to  $+490 \pm 100\text{‰}$  (table S3), consistent with the  $\delta\text{D} = +252 \pm 13\text{‰}$  measured in an analysis of other Ryugu grains with a different technique (47). There is no obvious difference between grains from chambers A and C. The bulk N isotopic compositions  $\delta^{15}\text{N} = +39 \pm 5$  to  $+43 \pm 4\text{‰}$  are also within the range reported for other Ryugu grains, measured with different methods [ $\delta^{15}\text{N}$

35

40

=  $+43.0 \pm 9.0\%$  (47) and  $\delta^{15}\text{N} = 0$  to  $+20\%$  (48)]. Figure S8 shows that the bulk  $\delta\text{D}$  and  $\delta^{15}\text{N}$  of the Ryugu grains are between the bulk values of CI chondrites [ $\delta\text{D} = +170$  to  $+300\%$ ,  $\delta^{15}\text{N} = +39$  to  $+52\%$  (49)] and IOM in CI chondrites [ $\delta\text{D} = \sim +975\%$ ,  $\delta^{15}\text{N} = +31\%$  (45)].

Isotopic ratios determined for individual sub- $\mu\text{m}$  to  $\mu\text{m}$  sized carbonaceous grains within the NanoSIMS maps are similar to those of carbonaceous chondrites and IDPs (43, 46). Most grains are consistent with the bulk average, within the uncertainties, while a small fraction of outliers have D and/or  $^{15}\text{N}$  enrichments or depletions (Fig. 6D-F), which we term hot-spots and cold-spots respectively. These have a similar range of compositions to those seen in CI (46) and CM (50) chondrites, but smaller than that seen in CR chondrites (29). There is no correlation between  $\delta\text{D}$  and  $\delta^{15}\text{N}$  for the hot- or cold-spots and a wide range of compositions occur over small spatial scales (Fig. 6B and C). The origin(s) of the H and N isotopic anomalies in meteorites are debated; current consensus models propose that they reflect isotopic fractionation at low temperatures in interstellar clouds or the outer protoplanetary nebula (51-54).

The NanoSIMS measurements show that the bulk Ryugu particles, and almost all carbon-rich particles within them, have  $\delta^{13}\text{C}$  values within the range of organic matter in carbonaceous chondrites [ $\delta^{13}\text{C} = -35$  to  $-5\%$  (45)]. However,  $\sim 0.5\%$  of the carbon-rich particles have  $^{13}\text{C}$  enrichments or depletions (Fig. 6G),  $>2\sigma$  above or below the bulk values. These particles have a range of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values similar to those seen in organic grains with anomalous carbon isotopes in primitive CM chondrites. For the Ryugu samples,  $^{13}\text{C}$ -rich particles have a wider range of  $\delta^{15}\text{N}$  values than  $^{13}\text{C}$ -poor ones, likely reflecting different origins. Two very small ( $\leq 150$  nm) regions have much higher  $^{13}\text{C}$  enrichments:  $\delta^{13}\text{C} = 700$  and  $4800\%$ , indicating they are pre-solar grains that formed in the outflows of previous generations of stars (55).

The acid-insoluble carbonaceous residues from the Ryugu samples showed bulk D-enrichments ( $\delta\text{D} = +306 \pm 42\%$  (A0106);  $+440 \pm 52\%$  (C0107)) (Fig. 7I), lower than the bulk hydrogen compositions of the IOM in CI chondrites ( $\delta\text{D} = +972\%$  for Orgueil,  $+978\%$  for Ivuna) and CM chondrites ( $+639\% < \delta\text{D} < +893\%$ ) (45). The distributions of D-rich hotspots (distribution modes  $\delta\text{D} = +1030\%$  (A0106) and  $+1374\%$  (C0107)) were consistent with CI and CM chondrites, but not CR chondrites (56).

The bulk N isotopic compositions of the insoluble carbonaceous residues from Ryugu samples ( $\delta^{15}\text{N} = +17.4 \pm 1.9\%$  (A0106);  $+30 \pm 4.3\%$  (C0107)) (Fig. 7J) are consistent with those of IOM in CI chondrites ( $\delta^{15}\text{N} = +30.7\%$  for Orgueil;  $+31.9\%$  for Ivuna) (45). Delta- $^{15}\text{N}$  was more heterogeneous in the sample from chamber C than chamber A. The  $^{15}\text{N}$ -rich hot-spots (distribution modes  $\delta^{15}\text{N} = +241\%$  (A0106) and  $+348\%$  (C0107)) were within the range of CI and CM carbonaceous chondrites (57).  $^{15}\text{N}$ -depleted cold-spots ( $\delta^{15}\text{N} = -100$  to  $-380\%$ ) were detected.  $^{15}\text{N}$  cold-spots have previously been reported in the matrices of carbonaceous chondrites (57-59) and IDPs (43), and interpreted as indicating organic grains that have been partially equilibrated with proto-solar or interstellar  $\text{N}_2$  gas (43).

### Comparison with D-type asteroids and Comets

We compare the organic matter of Ryugu (a C-type asteroid) with other primitive small bodies in the early Solar System. Of the carbonaceous chondrites, the Tagish Lake meteorite is thought to be related to dark (D-type) asteroids, which are mainly located in the outer regions of the asteroid belt and among the trojan asteroids of Jupiter (60). Tagish Lake contains organic and mineralogical variations between different specimens that reflect variable degrees of alteration

on the same parent body (61, 62). The ratios D/H, H/C, and aliphatic carbon to aromatic carbon, decrease systematically with increasing alteration of the Tagish Lake meteorite samples (61).

We infer that the organic matter in the Ryugu samples has not been heated to temperatures higher than 200°C, based on the similarity of its chemical features with primitive carbonaceous CI and CM chondrites. Like the IOM from Orgueil (63) and Murchison (64), the organic macromolecules in the Ryugu samples likely contain polyaromatic structures which are composed of small numbers of aromatic rings with short, cross-linked, aliphatic chains and various oxygen- and nitrogen bearing functional groups. This macromolecular structure is distinct from graphitic or glassy carbon produced by heating: we did not find any evidence of highly conjugated sp<sup>2</sup> carbon. Laboratory experiments have shown that aliphatic carbon in IOM from primitive carbonaceous chondrites is reduced after hydrous heating at 300°C for 6 days (65). Therefore, the higher abundance of aliphatic carbon in organic residue of Ryugu, than in that of primitive carbonaceous chondrites, indicates that the Ryugu grains did not experience short heating equivalent to the experimental conditions.

The distribution of organic functional groups present in the Ryugu samples is unlike aqueously altered carbonaceous chondrites, for which organic particles and nanoglobules show predominantly IOM-like XANES spectra, with only occasional highly aromatic grains (28, 30, 66). Ryugu samples contain abundant aromatic-rich particles and nanoglobules, a high abundance of IOM-like diffuse carbon in matrix, and molecular carbonate associated with phyllosilicates. Progressive aqueous alteration in situ causes i) an increase in the amount of diffuse organic matter associated with matrix phyllosilicates, ii) an increase in the proportion of aromatic and oxygen-bearing functional groups in discrete organic particles and nanoglobules, and iii) an increase in the variety of XANES spectral features. Based on these trends, much of the internal spectral variation in the Ryugu samples, including carbon grains with an aromatic XANES spectral shape (Fig. 3E), likely developed through redistribution of organic matter and formation of additional organic molecules during aqueous alteration on the parent body, rather than pre-accretionary diversity inherited from the proto-solar molecular cloud. Carbonaceous grains and nanoglobules with highly-aromatic XANES spectra (not aromatic spectra; Fig 3E) provide a possible exception to this proposal. These highly-aromatic grains are more likely to represent original accreted materials, because similar XANES spectra have been found in unaltered, petrologic type 3, carbonaceous chondrites (30) and in cometary dust particles (32). This is consistent with our observation of a large highly-aromatic nanoglobule containing D and <sup>15</sup>N isotopic anomalies (figs. S6 and S7).

The δD distributions of the carbon-rich particles in the Ryugu grains are consistent with those in the Tagish Lake meteorite (56), CI and CM chondrites (46, 57). The δD distributions of the Ryugu insoluble carbonaceous residues from chambers A and C are within the ranges of IOMs in the most and least altered fragments of Tagish Lake, respectively (61). The hydrogen isotopic variations among the different lithologies of Tagish Lake has previously been explained by depletion of D via hydrogen isotopic exchange or hydrolysis of D-rich structures of IOM during various degrees of planetesimal hydrothermal alteration (65, 67). In contrast, the δD distributions of our Ryugu samples are lower than those of less altered CR chondrites (56), the most primitive class of IDPs (68), and comet-derived AMMs (33, 69). We therefore propose that Ryugu organic matter is a product of heterogeneous aqueous processing, which occurred on both C- and D-type asteroids, of the common primordial materials formed at an earlier stage of the solar nebula. The δ<sup>15</sup>N distributions of the intact grains and insoluble residues from our Ryugu samples were within the ranges of carbon-rich particles (39, 56) and IOM (bulk) (61) in Tagish

Lake and Comet Wild 2 (32) as well as CI and CM chondrites. However, they are lower than the  $\delta^{15}\text{N}$  distributions of anhydrous IDPs (43, 68) and the IOM from CR chondrite (56). Nitrogen isotopic compositions are less affected than hydrogen isotopes by modification through parent body processes (45, 57, 61, 66), and their variations indicate mixing of different isotopic components from different precursors (53).

Nitrogen content can be an indicator of chemical evolution in the early Solar System. The measured N abundance ( $\text{N/C} = 0.01$  to  $0.035$ ) of the insoluble carbonaceous residue in Ryugu samples is similar to primitive carbonaceous chondrites (45), and consistent with the total bulk N abundance ( $\sim 0.15\text{wt}\%$ ) (47). The N/C ratio of IOM from Tagish Lake ( $\sim 0.042$ ) is higher than those of the Ryugu samples. In contrast, some cometary materials, such as dust particles from Comet Wild 2, anhydrous AMMs, and ultra-carbonaceous AMMs (UCAMMs), contain N-rich organics [ $\text{N/C} = 0.07$  to  $0.24$  (32, 44, 70)]. Those are composed of a variety of nitrogen-bearing functional groups, such as imine, nitrile and/or heterocyclic N, or amide (32, 33, 44, 70, 71). The N contents of Comet 67P/Churyumov-Gerasimenko particles were heterogeneous ( $\text{N/C} = 0.018$  to  $0.06$ ) (72) but lower than the N/C ratio of the Sun ( $0.3 \pm 0.1$ ) (73). The depletion of N in that comet might be due to the presence of ammonium salts (74, 75). These variations in N/C between different Solar System objects could arise from partitioning between gas and solid phases in the cold interstellar cloud or outer solar nebula, or exposure to warmer inner Solar System during perihelion passage (74), rather than processing in the interiors of planetesimals. Experiments simulating parent body aqueous alteration show the N/C ratios in meteoritic IOM are not substantially modified (65, 76). We therefore conclude that the N abundance in Ryugu macromolecular organic matter is intrinsic, not the product of extensive parent body processing.

## Conclusions

Our analysis of Ryugu samples indicate a direct link between macromolecular organic matter in C-type asteroids and that in primitive carbonaceous chondrites. The observed similarities and variations in molecular, isotopic, and morphological compositions between the Ryugu samples and other Solar System materials indicate a continuum of source material in the solar nebula, which was incorporated into C-type asteroids, D-type asteroids and comets in the early Solar System. Macromolecular organic matter in the surface grains of asteroid Ryugu reflects various degrees of parent body aqueous alteration and localized preservation of inherited nebular or molecular cloud history. The highly variable nature of the material indicates that the Ryugu organic matter is likely derived from material that was not subjected to long-term space weathering, and was only recently exposed to the asteroid surface. The macromolecular organic matter shows no record of high temperature impact heating of Ryugu, despite the asteroid being a rubble-pile body formed from impact debris, and geomorphological evidence that it experienced subsequent impact cratering (2, 3). This observation is consistent with the homogeneous reflectance spectra of Ryugu's surface (3).

Ryugu materials are much darker than primitive carbonaceous chondrites, despite having similar carbon contents (23, 47), so some other factor (or factors) must determine the albedo. Possibilities include the macromolecular organic matter mixed with phyllosilicates (this study) or nano-phase sulfides (24), both of which have high light absorption efficiency owing to their small grain sizes. Such complex aromatic macromolecules in C- and/or D-type asteroids could have supplied organic material, and possibly prebiotic molecules (47), to Earth.

## References and Notes

1. S. Tachibana, M. Abe, M. Arakawa, M. Fujimoto, Y. Iijima, M. Ishiguro, K. Kitazato, N. Kobayashi, N. Namiki, T. Okada, R. Okazaki, H. Sawada, S. Sugita, Y. Takano, S. Tanaka, S. Watanabe, M. Yoshikawa, H. Kuninaka, Hayabusa2: Scientific importance of samples returned from C-type near-Earth asteroid (162173) 1999 JU3. *Geochem. J.* 48, 571–587 (2014). DOI: 10.2343/geochemj.2.0350
2. S. Watanabe, M. Hirabayashi, N. Hirata, N. Hirata, R. Noguchi, Y. Shimaki, H. Ikeda, E. Tatsumi, M. Yoshikawa, S. Kikuchi, H. Yabuta, T. Nakamura, S. Tachibana, Y. Ishihara, T. Morota, K. Kitazato, N. Sakatani, K. Matsumoto, K. Wada, H. Senshu, C. Honda, T. Michikami, H. Takeuchi, T. Kouyama, R. Honda, S. Kameda, T. Fuse, H. Miyamoto, G. Komatsu, S. Sugita, T. Okada, N. Namiki, M. Arakawa, M. Ishiguro, M. Abe, R. Gaskell, E. Palmer, O. S. Barnouin, P. Michel, A. S. French, J. W. McMahon, D. J. Scheeres, P. A. Abell, Y. Yamamoto, S. Tanaka, K. Shirai, M. Matsuoka, M. Yamada, Y. Yokota, H. Suzuki, K. Yoshioka, Y. Cho, S. Tanaka, N. Nishikawa, T. Sugiyama, H. Kikuchi, R. Hemmi, T. Yamaguchi, N. Ogawa, G. Ono, Y. Mimasu, K. Yoshikawa, T. Takahashi, Y. Takei, A. Fujii, C. Hirose, T. Iwata, M. Hayakawa, S. Hosoda, O. Mori, H. Sawada, T. Shimada, S. Soldini, H. Yano, R. Tsukizaki, M. Ozaki, Y. Iijima, K. Ogawa, M. Fujimoto, T.-M. Ho, A. Moussi, R. Jaumann, J.-P. Bibring, C. Krause, F. Terui, T. Saiki, S. Nakazawa, Y. Tsuda, Hayabusa2 arrives at the carbonaceous asteroid 162173 Ryugu-A spinning top-shaped rubble pile. *Science* 364, 268–272 (2019). DOI: 10.1126/science.aav8032
3. S. Sugita, R. Honda, T. Morota, S. Kameda, H. Sawada, E. Tatsumi, M. Yamada, C. Honda, Y. Yokota, T. Kouyama, N. Sakatani, K. Ogawa, H. Suzuki, T. Okada, N. Namiki, S. Tanaka, Y. Iijima, K. Yoshioka, M. Hayakawa, Y. Cho, M. Matsuoka, N. Hirata, N. Hirata, H. Miyamoto, D. Domingue, M. Hirabayashi, T. Nakamura, T. Hiroi, T. Michikami, P. Michel, R.-L. Ballouz, O. S. Barnouin, C. M. Ernst, S. E. Schröder, H. Kikuchi, R. Hemmi, G. Komatsu, T. Fukuhara, M. Taguchi, T. Arai, H. Senshu, H. Demura, Y. Ogawa, Y. Shimaki, T. Sekiguchi, T. G. Müller, A. Hagermann, T. Mizuno, H. Noda, K. Matsumoto, R. Yamada, Y. Ishihara, H. Ikeda, H. Araki, K. Yamamoto, S. Abe, F. Yoshida, A. Higuchi, S. Sasaki, S. Oshigami, S. Tsuruta, K. Asari, S. Tazawa, M. Shizugami, J. Kimura, T. Otsubo, H. Yabuta, S. Hasegawa, M. Ishiguro, S. Tachibana, E. Palmer, R. Gaskell, L. Le Corre, R. Jaumann, K. Otto, N. Schmitz, P. A. Abell, M. A. Barucci, M. E. Zolensky, F. Vilas, F. Thuillet, C. Sugimoto, N. Takaki, Y. Suzuki, H. Kamiyoshihara, M. Okada, K. Nagata, M. Fujimoto, M. Yoshikawa, Y. Yamamoto, K. Shirai, R. Noguchi, N. Ogawa, F. Terui, S. Kikuchi, T. Yamaguchi, Y. Oki, Y. Takao, H. Takeuchi, G. Ono, Y. Mimasu, K. Yoshikawa, T. Takahashi, Y. Takei, A. Fujii, C. Hirose, S. Nakazawa, S. Hosoda, O. Mori, T. Shimada, S. Soldini, T. Iwata, M. Abe, H. Yano, R. Tsukizaki, M. Ozaki, K. Nishiyama, T. Saiki, S. Watanabe, Y. Tsuda, The geomorphology, color, and thermal properties of Ryugu: Implications for parent-body processes. *Science* 364, eaaw0422 (2019). DOI: 10.1126/science.aaw0422
4. K. Kitazato, R. E. Milliken, T. Iwata, M. Abe, M. Ohtake, S. Matsuura, T. Arai, Y. Nakauchi, T. Nakamura, M. Matsuoka, H. Senshu, N. Hirata, T. Hiroi, C. Pilorget, R. Brunetto, F. Poulet, L. Riu, J. P. Bibring, D. Takir, D. L. Domingue, F. Vilas, M. A. Barucci, D. Perna, E. Palomba, A. Galiano, K. Tsumura, T. Osawa, M. Komatsu, A. Nakato, T. Arai, N. Takato, T. Matsunaga, Y. Takagi, K. Matsumoto, T. Kouyama, Y. Yokota, E. Tatsumi, N.

- 5 Sakatani, Y. Yamamoto, T. Okada, S. Sugita, R. Honda, T. Morota, S. Kameda, H. Sawada, C. Honda, M. Yamada, H. Suzuki, K. Yoshioka, M. Hayakawa, K. Ogawa, Y. Cho, K. Shirai, Y. Shimaki, N. Hirata, A. Yamaguchi, N. Ogawa, F. Terui, T. Yamaguchi, Y. Takei, T. Saiki, S. Nakazawa, S. Tanaka, M. Yoshikawa, S. Watanabe, Y. Tsuda, The surface composition of asteroid 162173 Ryugu from Hayabusa2 near-infrared spectroscopy. *Science* 364, 272–275 (2019). DOI: 10.1126/science.aav7432
- 10 5. T. Morota, S. Sugita, Y. Cho, M. Kanamaru, E. Tatsumi, N. Sakatani, R. Honda, N. Hirata, H. Kikuchi, M. Yamada, Y. Yokota, S. Kameda, M. Matsuoka, H. Sawada, C. Honda, T. Kouyama, K. Ogawa, H. Suzuki, K. Yoshioka, M. Hayakawa, N. Hirata, M. Hirabayashi, H. Miyamoto, T. Michikami, T. Hiroi, R. Hemmi, O. S. Barnouin, C. M. Ernst, K. Kitazato, T. Nakamura, L. Riu, H. Senshu, H. Kobayashi, S. Sasaki, G. Komatsu, N. Tanabe, Y. Fujii, T. Irie, M. Suemitsu, N. Takaki, C. Sugimoto, K. Yumoto, M. Ishida, H. Kato, K. Moroi, D. Domingue, P. Michel, C. Pilorget, T. Iwata, M. Abe, M. Ohtake, Y. Nakauchi, K. Tsumura, H. Yabuta, Y. Ishihara, R. Noguchi, K. Matsumoto, A. Miura, N. Namiki, S. Tachibana, M. Arakawa, H. Ikeda, K. Wada, T. Mizuno, C. Hirose, S. Hosoda, O. Mori, T. Shimada, S. Soldini, R. Tsukizaki, H. Yano, M. Ozaki, H. Takeuchi, Y. Yamamoto, T. Okada, Y. Shimaki, K. Shirai, Y. Iijima, H. Noda, S. Kikuchi, T. Yamaguchi, N. Ogawa, G. Ono, Y. Mimasu, K. Yoshikawa, T. Takahashi, Y. Takei, A. Fujii, S. Nakazawa, F. Terui, S. Tanaka, M. Yoshikawa, T. Saiki, S. Watanabe, Y. Tsuda, Sample collection from asteroid (162173) Ryugu by Hayabusa2: Implications for surface evolution, *Science* 368, 654–659 (2020). DOI: 10.1126/science.aaz6306
- 15 6. N. Sakatani, S. Tanaka, T. Okada, T. Fukuhara, L. Riu, S. Sugita, R. Honda, T. Morota, S. Kameda, Y. Yokota, E. Tatsumi, K. Yumoto, N. Hirata, A. Miura, T. Kouyama, H. Senshu, Y. Shimaki, T. Arai, J. Takita, H. Demura, T. Sekiguchi, T. G. Müller, A. Hagermann, J. Biele, M. Grott, M. Hamm, M. Delbo, W. Neumann, M. Taguchi, Y. Ogawa, T. Matsunaga, T. Wada, S. Hasegawa, J. Helbert, N. Hirata, R. Noguchi, M. Yamada, H. Suzuki, C. Honda, K. Ogawa, M. Hayakawa, K. Yoshioka, M. Matsuoka, Y. Cho, H. Sawada, K. Kitazato, T. Iwata, M. Abe, M. Ohtake, S. Matsuura, K. Matsumoto, H. Noda, Y. Ishihara, K. Yamamoto, A. Higuchi, N. Namiki, G. Ono, T. Saiki, H. Imamura, Y. Takagi, H. Yano, K. Shirai, C. Okamoto, S. Nakazawa, Y. Iijima, M. Arakawa, K. Wada, T. Kadono, K. Ishibashi, F. Terui, S. Kikuchi, T. Yamaguchi, N. Ogawa, Y. Mimasu, K. Yoshikawa, T. Takahashi, Y. Takei, A. Fujii, H. Takeuchi, Y. Yamamoto, C. Hirose, S. Hosoda, O. Mori, T. Shimada, S. Soldini, R. Tsukizaki, M. Ozaki, S. Tachibana, H. Ikeda, M. Ishiguro, H. Yabuta, M. Yoshikawa, S. Watanabe, Y. Tsuda, Anomalously porous boulders on (162173) Ryugu as primordial materials from its parent body, *Nature Astronomy* 5, 766–774 (2021). DOI: 10.1038/s41550-021-01371-7
- 20 7. E. Tatsumi, N. Sakatani, L. Riu, M. Matsuoka, R. Honda, T. Morota, S. Kameda, T. Nakamura, M. Zolensky, R. Brunetto, T. Hiroi, S. Sasaki, S. Watanabe, S. Tanaka, J. Takita, C. Pilorget, J. de León, M. Popescu, J. Luis Rizos, J. Licandro, E. Palomba, D. Domingue, F. Vilas, H. Campins, Y. Cho, K. Yoshioka, H. Sawada, Y. Yokota, M. Hayakawa, M. Yamada, T. Kouyama, H. Suzuki, C. Honda, K. Ogawa, K. Kitazato, N. Hirata, N. Hirata, Y. Tsuda, M. Yoshikawa, T. Saiki, F. Terui, S. Nakazawa, Y. Takei, H. Takeuchi, Y. Yamamoto, T. Okada, Y. Shimaki, K. Shirai, S. Sugita, Spectrally blue hydrated parent body of asteroid (162173) Ryugu, *Nature Communications* 12, 5837 (2021). DOI: <https://doi.org/10.1038/s41467-021-26071-8>
- 25 30 35 40 45

8. E. Tatsumi, C. Sugimoto, L. Riu, S. Sugita, T. Nakamura, T. Hiroi, T. Morota, M. Popescu, T. Michikami, K. Kitazato, M. Matsuoka, S. Kameda, R. Honda, M. Yamada, N. Sakatani, T. Kouyama, Y. Yokota, C. Honda, H. Suzuki, Y. Cho, K. Ogawa, M. Hayakawa, H. Sawada, K. Yoshioka, C. Pilorget, M. Ishida, D. Domingue, N. Hirata, S. Sasaki, J. de León, M. A. Barucci, P. Michel, M. Suemitsu, T. Saiki, S. Tanaka, F. Terui, S. Nakazawa, S. Kikuchi, T. Yamaguchi, N. Ogawa, G. Ono, Y. Mimasu, K. Yoshikawa, T. Takahashi, Y. Takei, A. Fujii, Y. Yamamoto, T. Okada, C. Hirose, S. Hosoda, O. Mori, T. Shimada, S. Soldini, R. Tsukizaki, T. Mizuno, T. Iwata, H. Yano, M. Ozaki, M. Abe, M. Ohtake, N. Namiki, S. Tachibana, M. Arakawa, H. Ikeda, M. Ishiguro, K. Wada, H. Yabuta, H. Takeuchi, Y. Shimaki, K. Shirai, N. Hirata, Y. Iijima, Y. Tsuda, S. Watanabe, M. Yoshikawa, Collisional history of Ryugu's parent body from bright surface boulders, *Nature Astronomy* 5, 39–45 (2021). DOI: 10.1038/s41550-020-1179-z
9. T. Yada, M. Abe, T. Okada, A. Nakato, K. Yogata, A. Miyazaki, K. Hatakeda, K. Kumagai, M. Nishimura, Y. Hitomi, H. Soejima, M. Yoshitake, A. Iwamae, S. Furuya, M. Uesugi, Y. Karouji, T. Usui, T. Hayashi, D. Yamamoto, R. Fukai, S. Sugita, Y. Cho, K. Yumoto, Y. Yabe, J.-P. Bibring, C. Pilorget, V. Hamm, R. Brunetto, L. Riu, L. Lourit, D. Loizeau, G. Lequertier, A. Moussi-Soffys, S. Tachibana, H. Sawada, R. Okazaki, Y. Takano, K. Sakamoto, Y. N. Miura, H. Yano, T. R. Ireland, T. Yamada, M. Fujimoto, K. Kitazato, N. Namiki, M. Arakawa, N. Hirata, H. Yurimoto, T. Nakamura, T. Noguchi, H. Yabuta, H. Naraoka, M. Ito, E. Nakamura, K. Uesugi, K. Kobayashi, T. Michikami, H. Kikuchi, N. Hirata, Y. Ishihara, K. Matsumoto, H. Noda, R. Noguchi, Y. Shimaki, K. Shirai, K. Ogawa, K. Wada, H. Senshu, Y. Yamamoto, T. Morota, R. Honda, C. Honda, Y. Yokota, M. Matsuoka, N. Sakatani, E. Tatsumi, A. Miura, M. Yamada, A. Fujii, C. Hirose, S. Hosoda, H. Ikeda, T. Iwata, S. Kikuchi, Y. Mimasu, O. Mori, N. Ogawa, G. Ono, T. Shimada, S. Soldini, T. Takahashi, Y. Takei, H. Takeuchi, R. Tsukizaki, K. Yoshikawa, F. Terui, S. Nakazawa, S. Tanaka, T. Saiki, M. Yoshikawa, S.-I. Watanabe, Y. Tsuda, Preliminary analysis of the Hayabusa2 samples returned from C-type asteroid Ryugu, *Nature Astronomy*, (2021). DOI: 10.1038/s41550-021-01550-6
10. C. Pilorget, T. Okada, V. Hamm, R. Brunetto, T. Yada, D. Loizeau, L. Riu, T. Usui, A. Moussi-Soffys, K. Hatakeda, A. Nakato, K. Yogata, M. Abe, A. Aléon-Toppani, J. Carter, M. Chaigneau, B. Crane, B. Gondet, K. Kumagai, Y. Langevin, C. Lantz, T. Le Pivert-Jolivet, G. Lequertier, L. Lourit, A. Miyazaki, M. Nishimura, F. Poulet, M. Arakawa, N. Hirata, K. Kitazato, S. Nakazawa, N. Namiki, T. Saiki, S. Sugita, S. Tachibana, S. Tanaka, M. Yoshikawa, Y. Tsuda, S. Watanabe & J.-P. Bibring, First compositional analysis of Ryugu samples by the MicrOmega hyperspectral microscope, *Nature Astronomy*, (2021). DOI: 10.1038/s41550-021-01549-z
11. Materials and methods are available as supplementary materials.
12. S. Tachibana, H. Sawada, R. Okazaki, Y. Takano, K. Sakamoto, Y. N. Miura, C. Okamoto, H. Yano, S. Yamanouchi, P. Michel, Y. Zhang, S. Schwartz, F. Thuillet, H. Yurimoto, T. Nakamura, T. Noguchi, H. Yabuta, H. Naraoka, A. Tsuchiyama, N. Imae, K. Kurosawa, A. M. Nakamura, K. Ogawa, S. Sugita, T. Morota, R. Honda, S. Kameda, E. Tatsumi, Y. Cho, K. Yoshioka, Y. Yokota, M. Hayakawa, M. Matsuoka, N. Sakatani, M. Yamada, T. Kouyama, H. Suzuki, C. Honda, T. Yoshimitsu, T. Kubota, H. Demura, T. Yada, M. Nishimura, K. Yogata, A. Nakato, M. Yoshitake, A. I. Suzuki, S. Furuya, K. Hatakeda, A. Miyazaki, K. Kumagai, T. Okada, M. Abe, T. Usui, T. R. Ireland, M. Fujimoto, T. Yamada, M. Arakawa, H. C. Connolly Jr., A. Fujii, S. Hasegawa, N. Hirata, N. Hirata, C. Hirose, S.



- Hosoda, Y. Iijima, H. Ikeda, M. Ishiguro, Y. Ishihara, T. Iwata, S. Kikuchi, K. Kitazato, D. S. Lauretta, G. Libourel, B. Marty, K. Matsumoto, T. Michikami, Y. Mimasu, A. Miura, O. Mori, K. Nakamura-Messenger, N. Namiki, A. N. Nguyen, L. R. Nittler, H. Noda, R. Noguchi, N. Ogawa, G. Ono, M. Ozaki, H. Senshu, T. Shimada, Y. Shimaki, K. Shirai, S. Soldini, T. Takahashi, Y. Takei, H. Takeuchi, R. Tsukizaki, K. Wada, Y. Yamamoto, K. Yoshikawa, K. Yumoto, M. E. Zolensky, S. Nakazawa, F. Terui, S. Tanaka, T. Saiki, M. Yoshikawa, S. Watanabe, Y. Tsuda, Pebbles and sand on asteroid (162173) Ryugu: In situ observation and particles returned to Earth. *Science* 375, 1011-1016 (2022). DOI: 10.1126/science.abj8624
13. L. Bonal, E. Quirico, M. Bourot-Denise, G. Montagnac, Determination of the petrologic type of CV3 chondrites by Raman spectroscopy of included organic matter. *Geochimica Cosmochimica Acta* 70, 1849-1863 (2006). DOI: 10.1016/j.gca.2005.12.004
14. H. Busemann, C. M. O'D. Alexander, L. R. Nittler, Characterization of insoluble organic matter in primitive meteorites by microRaman spectroscopy, *Meteoritics & Planetary Science* 42, 1387-1416 (2007). DOI: 10.1111/j.1945-5100.2007.tb00581.x
15. L. Bonal, E. Quirico, L. Flandinet, G. Montagnac, Thermal history of type 3 chondrites from the Antarctic meteorite collection determined by Raman spectroscopy of their polyaromatic carbonaceous matter, *Geochimica Cosmochimica Acta* 189, 312-337 (2016). DOI: 10.1016/j.gca.2016.06.017
16. E. Quirico, L. Bonal, P. Beck, C.M.O'D. Alexander, H. Yabuta, T. Nakamura, A. Nakato, L. Flandinet, G. Montagnac, P. Schmitt-Koppling, C.D.K. Herd, Prevalence and nature of heating processes in CM and C2-ungrouped chondrites as revealed by insoluble organic matter, *Geochimica Cosmochimica Acta* 241, 17-37 (2018). DOI: 10.1016/j.gca.2018.08.029
17. G.R. Huss, A.E. Rubin, J.N. Grossman, Thermal Metamorphism in Chondrites, *Meteorites and the Early Solar System II* (Eds. D. S. Lauretta and H. Y. McSween), pp. 567-586. (2006).
18. T. Nakamura, Yamato 793321 CM chondrite: Dehydrated regolith material of a hydrous asteroid, *Earth and Planetary Science Letters* 242, 26-38 (2006). DOI: 10.1016/j.epsl.2005.11.040
19. T. Osawa, H. Kagi, T. Nakamura, T. Noguchi, Infrared spectroscopic taxonomy for carbonaceous chondrites from speciation of hydrous components, *Meteoritics and Planetary Science* 40, 71-86. DOI: 10.1111/j.1945-5100.2005.tb00365.x
20. Y. Kebukawa, C. M. O'D. Alexander, G.D. Cody, Comparison of FT-IR spectra of bulk and acid insoluble organic matter in chondritic meteorites: An implication for missing carbon during demineralization, *Meteoritics & Planetary Science* 54, 1632-1641 (2019). DOI: 10.1111/maps.13302
21. P. Beck, E. Quirico, G. Montes-Hernandez, L. Bonal, J. Bollard, F.-R.Orthous-Daunay, K. T. Howard, B. Schmitt, O. Brissaud, F. Deschamps, B. Wunder, S. Guillot, Hydrous mineralogy of CM and CI chondrites from infrared spectroscopy and their relationship with low albedo asteroids, *Geochimica et Cosmochimica Acta* 74, 4881-4892. DOI: 10.1016/j.gca.2010.05.020

22. D. Takir, J. P. Emery, H. Y. Mewsween Jr., C. A. Hibbitts, R. N. Clark, N. Pearson, A. Wang, Nature and degree of aqueous alteration in CM and CI carbonaceous chondrites, *Meteoritics & Planetary Science* 48, 1618-1637. DOI: 10.1111/maps.12171
23. T. Yokoyama, K. Nagashima, I. Nakai, E. D. Young, Y. Abe, J. Aléon, C. M. O'D. Alexander, S. Amari, Y. Amelin, K. Bajo, M. Bizzarro, A. Bouvier, R. W. Carlson, M. Chaussidon, B.-G. Choi, N. Dauphas, A. M. Davis, T. D. Rocco, W. Fujiya, R. Fukai, I. Gautam, M. K. Haba, Y. Hibiya, H. Hidaka, H. Homma, P. Hoppe, G. R. Huss, K. Ichida, T. Iizuka, T. R. Ireland, A. Ishikawa, M. Ito, S. Itoh, N. Kawasaki, N. T. Kita, K. Kitajima, T. Kleine, S. Komatani, A. N. Krot, M.-C. Liu, Y. Masuda, K. D. McKeegan, M. Morita, K. Motomura, F. Moynier, A. Nguyen, L. Nittler, M. Onose, A. Pack, C. Park, L. Piani, L. Qin, S. S. Russell, N. Sakamoto, M. Schönbächler, L. Tafla, H. Tang, K. Terada, Y. Terada, T. Usui, S. Wada, M. Wadhwa, R. J. Walker, K. Yamashita, Q.-Z. Yin, S. Yoneda, H. Yui, A.-C. Zhang, H. C. Connolly, Jr., D. S. Lauretta, T. Nakamura, H. Naraoka, T. Noguchi, R. Okazaki, K. Sakamoto, H. Yabuta, M. Abe, M. Arakawa, A. Fujii, M. Hayakawa, N. Hirata, N. Hirata, R. Honda, C. Honda, S. Hosoda, Y. Iijima, H. Ikeda, M. Ishiguro, Y. Ishihara, T. Iwata, K. Kawahara, S. Kikuchi, K. Kitazato, K. Matsumoto, M. Matsuoka, T. Michikami, Y. Mimasu, A. Miura, T. Morota, S. Nakazawa, N. Namiki, H. Noda, R. Noguchi, N. Ogawa, K. Ogawa, T. Okada, C. Okamoto, G. Ono, M. Ozaki, T. Saiki, N. Sakatani, H. Sawada, H. Senshu, Y. Shimaki, K. Shirai, S. Sugita, Y. Takei, H. Takeuchi, S. Tanaka, E. Tatsumi, F. Terui, Y. Tsuda, R. Tsukizaki, K. Wada, S. Watanabe, M. Yamada, T. Yamada, Y. Yamamoto, H. Yano, Y. Yokota, K. Yoshihara, M. Yoshikawa, K. Yoshikawa, S. Furuya, K. Hatakeda, T. Hayashi, Y. Hitomi, K. Kumagai, A. Miyazaki, A. Nakato, M. Nishimura, H. Soejima, A. Suzuki, T. Yada, D. Yamamoto, K. Yogata, M. Yoshitake, S. Tachibana, H. Yurimoto, Samples returned from the asteroid Ryugu are similar to Ivuna-type carbonaceous meteorites, *Science*, this issue. DOI: 10.1126/science.abn7850
24. T. Nakamura, M. Matsumoto, K. Amano, Y. Enokido, M. E. Zolensky, T. Mikouchi, H. Genda, S. Tanaka, M. Y. Zolotov, K. Kurosawa, S. Wakita, R. Hyodo, H. Nagano, D. Nakashima, Y. Takahashi, Y. Fujioka, M. Kikuri, E. Kagawa, M. Matsuoka, A. J. Brearley, A. Tsuchiyama, M. Uesugi, J. Matsuno, Y. Kimura, M. Sato, R. E. Milliken, E. Tatsumi, S. Sugita, T. Hiroi, K. Kitazato, D. Brownlee, D. J. Joswiak, M. Takahashi, K. Ninomiya, T. Takahashi, T. Osawa, K. Terada, F. E. Brenker, B. J. Tkalcec, L. Vincze, R. Brunetto, A. Aléon-Toppani, Q. H. S. Chan, M. Roskosz, J.-C. Viennet, P. Beck, E. E. Alp, T. Michikami, Y. Nagaashi, T. Tsuji, Y. Ino, J. Martinez, J. Han, A. Dolocan, R. J. Bodnar, M. Tanaka, H. Yoshida, K. Sugiyama, A. J. King, K. Fukushi, H. Suga, S. Yamashita, T. Kawai, K. Inoue, A. Nakato, T. Noguchi, F. Vilas, A. R. Hendrix, C. Jaramillo, D. L. Domingue, G. Dominguez, Z. Gainsforth, C. Engrand, J. Duprat, S. S. Russell, E. Bonato, C. Ma, T. Kawamoto, T. Wada, S. Watanabe, R. Endo, S. Enju, L. Riu, S. Rubino, P. Tack, S. Takeshita, Y. Takeichi, A. Takeuchi, A. Takigawa, D. Takir, T. Tanigaki, A. Taniguchi, K. Tsukamoto, T. Yagi, S. Yamada, K. Yamamoto, Y. Yamashita, M. Yasutake, K. Uesugi, I. Umegaki, I. Chiu, T. Ishizaki, S. Okumura, E. Palomba, C. Pilorget, S. M. Potin, A. Alasli, S. Anada, Y. Araki, N. Sakatani, C. Schultz, O. Sekizawa, S. D. Sitzman, K. Sugiura, M. Sun, E. Dartois, E. De Pauw, Z. Dionnet, Z. Djouadi, G. Falkenberg, R. Fujita, T. Fukuma, I. R. Gearba, K. Hagiya, M. Y. Hu, T. Kato, T. Kawamura, M. Kimura, M. K. Kubo, F. Langenhorst, C. Lantz, B. Lavina, M. Lindner, J. Zhao, B. Vekemans, D. Baklouti, B. Bazi, F. Borondics, S. Nagasawa, G. Nishiyama, K. Nitta, J. Mathurin, T. Matsumoto, I. Mitsukawa, H. Miura, A. Miyake, Y. Miyake, H. Yurimoto, R. Okazaki, H. Yabuta, H. Naraoka, K. Sakamoto, S. Tachibana, H. C. Connolly Jr., D. S. Lauretta, M. Yoshitake, M.

- Yoshikawa, K. Yoshikawa, K. Yoshihara, Y. Yokota, K. Yogata, H. Yano, Y. Yamamoto, D. Yamamoto, M. Yamada, T. Yamada, T. Yada, K. Wada, T. Usui, R. Tsukizaki, F. Terui, H. Takeuchi, Y. Takei, A. Iwamae, H. Soejima, K. Shirai, Y. Shimaki, H. Senshu, H. Sawada, T. Saiki, M. Ozaki, G. Ono, T. Okada, N. Ogawa, K. Ogawa, R. Noguchi, H. Noda, M.  
5 Nishimura, N. Namiki, S. Nakazawa, T. Morota, A. Miyazaki, A. Miura, Y. Mimasu, K. Matsumoto, K. Kumagai, T. Kouyama, S. Kikuchi, K. Kawahara, S. Kameda, T. Iwata, Y. Ishihara, M. Ishiguro, H. Ikeda, S. Hosoda, R. Honda, C. Honda, Y. Hitomi, N. Hirata, N. Hirata, T. Hayashi, M. Hayakawa, K. Hatakeda, S. Furuya, R. Fukai, A. Fujii, Y. Cho, M. Arakawa, M. Abe, S. Watanabe, Y. Tsuda, Formation and evolution of carbonaceous asteroid  
10 Ryugu: Direct evidence from returned samples, *Science*, this issue. DOI: 10.1126/science.abn8671
25. M. Gounelle, M. E. Zolensky, The Orgueil meteorite: 150 years of history, *Meteoritics & Planetary Science* 49, 1769-1794 (2014). DOI: 10.1111/maps.12351
- 15 26. Y. Kebukawa, C.M.O'D. Alexander, G. D. Cody, Compositional diversity in insoluble organic matter in type 1, 2 and 3 chondrites as detected by infrared spectroscopy, *Geochimica et Cosmochimica Acta* 75, 3530–3541 (2011). DOI:10.1016/j.gca.2011.03.037
27. F.-R. Orthous-Daunay, E. Quirico, P. Beck, O. Brissaud, E. Dartois, T. Pino, B. Schmitt, Mid-infrared study of the molecular structure variability of insoluble organic matter from primitive chondrites, *Icarus* 223, 534-543 (2013). DOI: 10.1016/j.icarus.2013.01.003
- 20 28. C. Le Guillou, S. Bernard, A. J. Brearley, L. Remusat, Evolution of organic matter in Orgueil, Murchison and Renazzo during parent body aqueous alteration: In situ investigations, *Geochimica et Cosmochimica Acta* 131, 368–392 (2014). DOI: 10.1016/j.gca.2013.11.020
- 25 29. L. R. Nittler, R. M. Stroud, J. M. Trigo-Rodríguez, B. T. De Gregorio, C. M. O'D. Alexander, J. Davidson, C. E. Moyano-Camero, S. Tanbakouei, A cometary building block in a primitive asteroidal meteorite, *Nature Astronomy* 3, 659–666 (2019). DOI: 10.1038/s41550-019-0737-8
- 30 30. B. T. De Gregorio, R. M. Stroud, L. R. Nittler, C. M.O'D. Alexander, N. D. Bassim, G. D. Cody, A. L. D. Kilcoyne, S. A. Sandford, S. N. Milam, M. Nuevo, T. J. Zega, Isotopic and chemical variation of organic nanoglobules in primitive meteorites, *Meteoritics & Planetary Science* 48, 904-928 (2013). DOI: 10.1111/maps.12109
31. L. P. Keller, S. Messenger, G. J. Flynn, S. Clemett, S. Wirick, C. Jacobsen, The nature of molecular cloud material in interplanetary dust, *Geochimica et Cosmochimica Acta* 68, 2577–2589 (2004). DOI: <https://doi.org/10.2465/jmps.101.122>
- 35 32. B. T. De Gregorio, R. M. Stroud, L. R. Nittler, C. M.O'D. Alexander, A. L. D. Kilcoyne, T. J. Zega, Isotopic anomalies in organic nanoglobules from Comet 81P/Wild 2: Comparison to Murchison nanoglobules and isotopic anomalies induced in terrestrial organics by electron irradiation, *Geochimica et Cosmochimica Acta* 74, 4454–4470 (2010). DOI:10.1016/j.gca.2010.05.010
- 40 33. T. Noguchi, H. Yabuta, S. Itoh, N. Sakamoto, T. Mitsunari, A. Okubo, R. Okazaki, T. Nakamura, S. Tachibana, K. Terada, M. Ebihara, N. Imae, M. Kimura, H. Nagahara, Variation of mineralogy and organic material during the early stages of aqueous activity recorded in Antarctic micrometeorites, *Geochimica et Cosmochimica Acta* 208, 119–144 (2017). DOI: 10.1016/j.gca.2017.03.034

34. L. A. J. Garvie and P. R. Buseck, Prebiotic carbon in clays from Orgueil and Ivuna (CI), and Tagish Lake (C2 ungrouped) meteorites, *Meteoritics & Planetary Science* 42, 2111-2117 (2007), DOI: 10.1111/j.1945-5100.2007.tb01011.x
- 5 35. G. D. Cody, C.M.O'D. Alexander, H. Yabuta, A. L. D. Kilcoyne, T. Araki, H. Ade, P. Dera, M. Fogel, B. Militzer, B. O. Mysen, Organic thermometry for chondritic parent bodies, *Earth and Planetary Science Letters* 272, 446–455 (2008). DOI:10.1016/j.epsl.2008.05.008
36. C.M.O'D. Alexander, G.D. Cody, B.T. De Gregorio, L.R. Nittler, R.M. Stroud, The nature, origin and modification of insoluble organic matter in chondrites, the major source of Earth's C and N, *Chemie der Erde* 77, 227–256 (2017). DOI: 10.1016/j.chemer.2017.01.007
- 10 37. D. D. Clayton, Origin of heavy xanon in meteoritic diamonds, *Astrophysical Journal* 340, 613-619 (1989).
38. Z. R. Dai, J. P. Bradley, D. J. Joswiak, D. E. Brownlee, H. G. M. Hill, M. J. Genge, Possible in situ formation of meteoritic nanodiamonds in the early Solar System, *Nature* 418, 157–159 (2002). DOI: 10.1038/nature00897
- 15 39. K. Nakamura-Messenger, S. Messenger, L. P. Keller, S. J. Clemett, M. E. Zolensky, Organic Globules in the Tagish Lake Meteorite: Remnants of the Protosolar Disk, *Science* 314, 1439-1442 (2006). DOI: 10.1126/science.1132175
40. T. Matsumoto, A. Tsuchiyama, K. Nakamura-Messenger, T. Nakano, K. Uesugi, A. Takeuchi, M. E. Zolensky, *Geochimica et Cosmochimica Acta* 116, 84–95 (2013). DOI: 20 10.1016/j.gca.2012.05.007
41. M. Hashiguchi, S. Kobayashi, H. Yurimoto, In situ observation of D-rich carbonaceous globules embedded in NWA 801 CR2 chondrite, *Geochimica et Cosmochimica Acta* 122, 306–323 (2013). DOI: 10.1016/j.gca.2013.08.007
- 25 42. C. Le Guillou, A. Brearley, Relationships between organics, water and early stages of aqueous alteration in the pristine CR3.0 chondrite MET 00426, *Geochimica et Cosmochimica Acta* 131, 344–367 (2014). DOI: 10.1016/j.gca.2013.10.024
43. G. Matrajt, S. Messenger, D. Brownlee, D. Joswiak, Diverse forms of primordial organic matter identified in interplanetary dust particles, *Meteoritics & Planetary Science* 47, 525–549 (2012). DOI: 10.1111/j.1945-5100.2011.01310.x
- 30 44. H. Yabuta, T. Noguchi, S. Itoh, T. Nakamura, A. Miyake, S. Tsujimoto, N. Ohashi, N. Sakamoto, M. Hashiguchi, K-I Abe, A. Okubo, A. L. D. Kilcoyne, S. Tachibana, R. Okazaki, K. Terada, M. Ebihara, H. Nagahara, Formation of an ultracarbonaceous Antarctic micrometeorite through minimal aqueous alteration in a small porous icy body, *Geochimica et Cosmochimica Acta* 214, 172–190 (2017). DOI: 10.1016/j.gca.2017.06.047
- 35 45. C.M.O'D. Alexander, M. Fogel, H. Yabuta, G.D. Cody, The origin and evolution of chondrites recorded in the elemental and isotopic compositions of their macromolecular organic matter, *Geochimica et Cosmochimica Acta* 71, 4380–4403 (2007). DOI: 10.1016/j.gca.2007.06.052
- 40 46. L. Remusat, Y. Guan, Y. Wang, J. M. Eiler, Accretion and preservation of D-rich organic particles in carbonaceous chondrites: Evidence for important transport in the early Solar System nebula, *The Astrophysical Journal*, 713, 1048–1058 (2010). DOI: 10.1088/0004-637X/713/2/1048

47. H. Naraoka, Y. Takano, J. P. Dworkin, Y. Oba, K. Hamase, A. Furusho, N. O. Ogawa, M. Hashiguchi, K. Fukushima, D. Aoki, P. Schmitt-Kopplin, J. C. Aponte, E. T. Parker, D. P. Glavin, H. L. McLain, J. E. Elsila, H. V. Graham, J. M. Eiler, F.-R. Orthous-Daunay, C. Wolters, J. Isa, V. Vuitton, R. Thissen, S. Sakai, T. Yoshimura, T. Koga, N. Ohkouchi, Y. Chikaraishi, H. Sugahara, H. Mita, Y. Furukawa, N. Hertkorn, A. Ruf, H. Yurimoto, T. Nakamura, T. Noguchi, R. Okazaki, H. Yabuta, K. Sakamoto, S. Tachibana, H. C. Connolly, Jr., D. S. Lauretta, M. Abe, T. Yada, M. Nishimura, K. Yogata, A. Nakato, M. Yoshitake, A. Suzuki, A. Miyazaki, S. Furuya, K. Hatakeda, H. Soejima, Y. Hitomi, K. Kumagai, T. Usui, T. Hayashi, D. Yamamoto, R. Fukai, K. Kitazato, S. Sugita, N. Namiki, M. Arakawa, H. Ikeda, M. Ishiguro, N. Hirata, K. Wada, Y. Ishihara, R. Noguchi, T. Morota, N. Sakatani, K. Matsumoto, H. Senshu, R. Honda, E. Tatsumi, Y. Yokota, C. Honda, T. Michikami, M. Matsuoka, A. Miura, H. Noda, T. Yamada, K. Yoshihara, K. Kawahara, M. Ozaki, Y. Iijima, H. Yano, M. Hayakawa, T. Iwata, R. Tsukizaki, H. Sawada, S. Hosoda, K. Ogawa, C. Okamoto, N. Hirata, K. Shirai, Y. Shimaki, M. Yamada, T. Okada, Y. Yamamoto, H. Takeuchi, A. Fujii, Y. Takei, K. Yoshikawa, Y. Mimasu, G. Ono, N. Ogawa, S. Kikuchi, S. Nakazawa, F. Terui, S. Tanaka, T. Saiki, M. Yoshikawa, S. Watanabe and Y. Tsuda, Soluble organic molecules in samples of the carbonaceous asteroid (162173) Ryugu, *Science*, this issue. DOI: 10.1126/science.abn9033
48. R. Okazaki, B. Marty, H. Busemann, K. Hashizume, J. D. Gilmour, A. Meshik, T. Yada, F. Kitajima, M. W. Broadley, D. Byrne, E. Füri, M. E. I. Riebe, D. Krietsch, C. Maden, A. Ishida, P. Clay, S. A. Crowther, L. Fawcett, T. Lawton, O. Pravdivseva, Y. N. Miura, J. Park, K. Bajo, Y. Takano, K. Yamada, S. Kawagucci, Y. Matsui, M. Yamamoto, K. Righter, S. Sakai, N. Iwata, N. Shirai, S. Sekimoto, M. Inagaki, M. Ebihara, R. Yokochi, K. Nishiizumi, K. Nagao, J. Ik Lee, A. Kano, M. W. Caffee, R. Uemura, T. Nakamura, H. Naraoka, T. Noguchi, H. Yabuta, H. Yurimoto, S. Tachibana, H. Sawada, K. Sakamoto, M. Abe, M. Arakawa, A. Fujii, M. Hayakawa, N. Hirata, N. Hirata, R. Honda, C. Honda, S. Hosoda, Y. Iijima, H. Ikeda, M. Ishiguro, Y. Ishihara, T. Iwata, K. Kawahara, S. Kikuchi, K. Kitazato, K. Matsumoto, M. Matsuoka, T. Michikami, Y. Mimasu, A. Miura, T. Morota, S. Nakazawa, N. Namiki, H. Noda, R. Noguchi, N. Ogawa, K. Ogawa, T. Okada, C. Okamoto, G. Ono, M. Ozaki, T. Saiki, N. Sakatani, H. Senshu, Y. Shimaki, K. Shirai, S. Sugita, Y. Takei, H. Takeuchi, S. Tanaka, E. Tatsumi, F. Terui, R. Tsukizaki, K. Wada, M. Yamada, T. Yamada, Y. Yamamoto, H. Yano, Y. Yokota, K. Yoshihara, M. Yoshikawa, K. Yoshikawa, S. Furuya, K. Hatakeda, T. Hayashi, Y. Hitomi, K. Kumagai, A. Miyazaki, A. Nakato, M. Nishimura, H. Soejima, A. Iwamae, D. Yamamoto, K. Yogata, M. Yoshitake, R. Fukai, T. Usui, H. C. Connolly Jr., D. Lauretta, S. Watanabe, and Y. Tsuda, Noble gases and nitrogen in samples of asteroid Ryugu record its volatile sources and recent surface evolution, *Science*, this issue. DOI: 10.1126/science.abo0431
49. J. F. Kerridge, Carbon, hydrogen and nitrogen in carbonaceous chondrites: Abundances and isotopic compositions in bulk samples, *Geochimica et Cosmochimica Acta* 49, 1707-1714 (1985). DOI: 10.1016/0016-7037(85)90141-3
50. L. R. Nittler, C. M. O'D. Alexander, M. J. Verdier-Paoletti, paper presented at the 84th Annual Meeting of the Meteoritical Society, Chicago, Illinois, August 15-21, 2021. LPI Contribution No. 2609.
51. Y. Aikawa, G. J. van Zadelhoff, E. F. van Dishoeck, E. Herbst, Warm molecular layers in protoplanetary disks, *Astron. Astrophys.* 386, 622-632 (2002). DOI: 10.1051/0004-6361:20020037

52. N. Watanabe, A. Kouchi, Ice surface reactions: A key to chemical evolution in space. *Progress in Surface Science*, 83, 439–489 (2008). DOI: 10.1016/j.progsurf.2008.10.001
53. J. Aléon, Multiple origins of nitrogen isotopic anomalies in meteorites and comets, *The Astrophysical Journal*, 722, 1342–1351 (2010). DOI: 10.1088/0004-637X/722/2/1342
- 5 54. R. Terzieva, E. Herbst, The possibility of nitrogen isotopic fractionation in interstellar clouds, *Monthly Notices of the Royal Astronomical Society* 317, 563-568 (2000). DOI: 10.1046/j.1365-8711.2000.03618.x
55. E. K. Zinner, Presolar Grains, In: *Meteorites, Comets, and Planets* (Ed. A. M. Davis), *Treatise on Geochemistry Vol. 1*, 2005, Elsevier, pp. 17-39.
- 10 56. H. Busemann, A. F. Young, C. M. O'D. Alexander, P. Hoppe, S. Mukhopadhyay, L. R. Nittler, Interstellar Chemistry Recorded in Organic Matter from Primitive Meteorites, *Science* 312, 727-730 (2006). DOI: 10.1126/science.1123878
57. L. Remusat, J. -Y. Bonnet, S. Bernard, A. Buch, E. Quirico, Molecular and isotopic behavior of insoluble organic matter of the Orgueil meteorite upon heating, *Geochimica et*  
15 *Cosmochimica Acta* 263, 235 - 247 (2019). DOI: 10.1016/j.gca.2019.07.013
58. C. Floss, F. Stadermann, Auger nanoprobe analysis of presolar ferromagnesian silicate grains from primitive CR chondrites QUE 99177 and MET 00426, *Geochimica et Cosmochimica Acta* 73, 2415-2440 (2009). DOI: 10.1016/j.gca.2009.01.033
59. C. Vollmer, J. Leitner, D. Kepaptsoglou, Q. M. Ramasse, A. J. King, P. F. Schofield, A. Bischoff, T. Araki, P. Hoppe, A primordial <sup>15</sup>N-depleted organic component detected within  
20 the carbonaceous chondrite Maribo, *Scientific Reports* 10, 20251 (2020). DOI: 10.1038/s41598-020-77190-z
60. T. Hiroi, M. E. Zolensky, C. M. Pieters, The Tagish Lake meteorite: A Possible Sample from a D-Type Asteroid, *Science* 293, 2234-2236 (2001). DOI: 10.1126/science.1063734
- 25 61. C. D. K. Herd, A. Blinova, D. N. Simkus, Y. Huang, R. Tarozo, C. M. O'D. Alexander, F. Gyngard, L. R. Nittler, G. D. Cody, M. L. Fogel, Y. Kebukawa, A. L. D. Kilcoyne, R. W. Hilts, G. F. Slater, D. P. Glavin, J. P. Dworkin, M. P. Callahan, J. E. Elsila, B. T. De Gregorio, R. M. Stroud, Origin and Evolution of Prebiotic Organic Matter As Inferred from the Tagish Lake Meteorite, *Science* 332, 1304-1307 (2011). DOI: 10.1126/science.1203290
- 30 62. Gilmour, C. M., Herd, C. D. K., and Beck, P., 2019. Water abundance in the Tagish Lake meteorite from TGA and IR spectroscopy: Evaluation of aqueous alteration. *Meteorit. Planet. Sci.* DOI: 10.1111/maps.13362.
63. A. Gardinier, S. Derenne, F. Robert, F. Behar, C. Largeau, J. Maquet, Solid state CP/MAS <sup>13</sup>C NMR of the insoluble organic matter of the Orgueil and Murchison meteorites:  
35 quantitative study, *Earth and Planetary Science Letters* 184, 9-21 (2000). DOI: 10.1016/S0012-821X(00)00300-9
64. G. D. Cody, C. M. O'D. Alexander, F. Tera, Solid-state (<sup>1</sup>H and <sup>13</sup>C) nuclear magnetic resonance spectroscopy of insoluble organic residue in the Murchison meteorite: A self-consistent quantitative analysis, *Geochimica et Cosmochimica Acta*, 66, 1851–1865, (2002).  
40 DOI: 10.1016/S0016-7037(01)00888-2
65. H. Yabuta, L. B. Williams, G. D. Cody, C. M. O'D. Alexander, S. Pizzarello, The insoluble carbonaceous material of CM chondrites: A possible source of discrete organic compounds

under hydrothermal conditions, *Meteoritics & Planetary Science* 42, 37–48 (2007). DOI: 10.1111/j.1945-5100.2007.tb00216.x

- 5 66. V. Vinogradoff, C. Le Guillou, S. Bernard, L. Binet, P. Cartigny, A.J. Brearley, L. Remusat, Paris vs. Murchison: Impact of hydrothermal alteration on organic matter in CM chondrites, *Geochimica et Cosmochimica Acta* 212, 234–252 (2017). DOI: 10.1016/j.gca.2017.06.009
67. Y. Oba and H. Naraoka, Elemental and isotope behavior of macromolecular organic matter from CM chondrites during hydrous pyrolysis, *Meteoritics & Planetary Science* 44, 943–953 (2009). DOI: 10.1111/j.1945-5100.2009.tb00779.x
- 10 68. S. Messenger, Identification of molecular-cloud material in interplanetary dust particles, *Nature* 404, 968-971 (2000). DOI: 10.1038/35010053
69. J. Duprat, E. Dobrica, C. Engrand, J. Ale'ón, Y. Marrocchi, S. Mostefaoui, A. Meibom, H. Leroux, J.-N. Rouzaud, M. Gounelle, F. Robert, Extreme deuterium excesses in ultracarbonaceous micrometeorites from central Antarctic snow, *Science* 328, 742–745 (2010). DOI: 10.1126/science.1184832
- 15 70. G. D. Cody, H. Ade, C. M. O'D. Alexander, T. Araki, A. Butterworth, H. Fleckensteins, G. Flynn, M. K. Gilles, C. Jacobsen, A. L. D. Kilcoyne, K. Messenger, S. A. Sandford, T. Tyliczszak, A. J. Westphal, S. Wirick, H. Yabuta, Quantitative organic and light-element analysis of comet 81P/Wild 2 particles using C-, N-, and O- $\mu$ -XANES, *Meteoritics & Planetary Science* 43, 353–365 (2008). DOI: 10.1111/j.1945-5100.2008.tb00627.x
- 20 71. E. Dartois, C. Engrand, R. Brunetto, J. Duprat, T. Pino, E. Quirico, L. Remusat, N. Bardin, G. Briani, S. Mostefaoui, G. Morinaud, B. Crane, N. Szwec, L. Delauche, F. Jamme, Ch. Sandt, P. Dumas, Ultracarbonaceous Antarctic micrometeorites, probing the Solar System beyond the nitrogen snow-line. *Icarus* 224, 243–252 (2013). Doi: 10.1016/j.icarus.2013.03.002
- 25 72. N. Fray, A. Bardyn, H. Cottin, D. Baklouti, C. Briois, C. Engrand, H. Fischer, K. Hornung, R. Isnard, Y. Langevin, H. Lehto, L. Le Roy, E. M. Mellado, S. Merouane, P. Modica, F.-R. Orthous-Daunay, J. Paquette, J. Rynö, R. Schulz, J. Silén, S. Siljeström, O. Stenzel, L. Thirkell, K. Varmuza, B. Zaprudin, J. Kissel, M. Hilchenbach, Nitrogen-to-carbon atomic ratio measured by COSIMA in the particles of comet 67P/Churyumov–Gerasimenko, *Monthly Notices of the Royal Astronomical Society* 469, S506–S516, (2017). DOI: 10.1093/mnras/stx2002
- 30 73. K. Lodders, Solar system abundances of the elements, *Principles and Perspectives in Cosmochemistry* (Eds. A. Goswami and B. E. Reddy), pp. 379–417. (2010).
- 35 74. O. Poch, I. Istiqomah, E. Quirico, P. Beck, B. Schmitt, P. Theulé, A. Faure, P. Hily-Blant, L. Bonal, A. Raponi, M. Ciarniello, B. Rousseau, S. Potin, O. Brissaud, L. Flandinet, G. Filacchione, A. Pommerol, N. Thomas, D. Kappel, V. Mennella, L. Moroz, V. Vinogradoff, G. Arnold, S. Erard, D. Bockelée-Morvan, C. Leyrat, F. Capaccioni, M. C. De Sanctis, A. Longobardo, F. Mancarella, E. Palomba, F. Tosi, Ammonium salts are a reservoir of nitrogen on a cometary nucleus and possibly on some asteroids, *Science* 367, eaaw7462, (2020). DOI: 10.1126/science.aaw7462
- 40 75. K. Altwegg, H. Balsiger, N. Hänni, M. Rubin, M. Schuhmann, I. Schroeder, T. Sémon, S. Wampfler, J.-J. Berthelier, C. Briois, M. Combi, T. I. Gombosi, H. Cottin, J. De Keyser, F. Dhooghe, B. Fiethe, S. A. Fuselier, Evidence of ammonium salts in comet 67P as explanation

for the nitrogen depletion in cometary comae, *Nature Astronomy* 4, 533–540 (2020). DOI: 10.1038/s41550-019-0991-9

76. H. Naraoka, H. Mita, M. Komiya, S. Yoneda, H. Kojima, A. Shimoyama, A chemical sequence of macromolecular organic matter in the CM chondrites, *Meteoritics & Planetary Science* 39, 401–406 (2004). DOI: 10.1111/j.1945-5100.2004.tb00101.x
77. N. D. Bassim, B. T. De Gregorio, A.L.D. Kilcoyne, K. Scott, T. Chou, S. Wirick, G. Cody, R.M. Stroud, Minimizing damage during FIB sample preparation of soft materials, *Journal of Microscopy*, 245, 288–301 (2012). DOI: 10.1111/j.1365-2818.2011.03570.x
78. Y. Takeichi, N. Inami, H. Suga, C. Miyamoto, T. Ueno, K. Mase, Y. Takahashi, K. Ono, Design and performance of a compact scanning transmission X-ray microscope at the Photon Factory, *Review of Scientific Instruments* 87, 013704. (2016). DOI: 10.1063/1.4940409
79. A.P. Hitchcock, aXis 2000 - Analysis of X-ray Images and Spectra. <http://unicorn.mcmaster.ca/aXis2000.html>
80. M. Lerotic, R. Mak, S. Wirick, F. Meirer, C. Jacobsen, MANTiS: a program for the analysis of X-ray spectromicroscopy data, *Journal of Synchrotron Radiation* 21, 1206–1212 (2014). DOI:10.1107/S1600577514013964
81. J. Mathurin, E. Dartois, T. Pino, C. Engrand, J. Duprat, A. Deniset-Besseau, F. Borondics, C. Sandt, A. Dazzi, Nanometre-scale infrared chemical imaging of organic matter in ultra-carbonaceous Antarctic micrometeorites (UCAMMs), *Astronomy & Astrophysics* 622, A160 (2018). DOI: 10.1051/0004-6361/201833957
82. A. Thomen, F. Robert, L. Remusat, Determination of the nitrogen abundance in organic materials by NanoSIMS quantitative imaging, *Journal of Analytical Atomic Spectrometry* 29, 512–519 (2014). DOI: 10.1039/c3ja50313e
83. L. Nittler, L'image: Larry's Image Program. <https://sites.google.com/carnegiescience.edu/limagesoftware/>
84. J. Davidson, H. Busemann, L.R. Nittler, C. M. Alexander, F.-R. Orthous-Daunay, I. A. Franchi, P. Hoppe, Abundances of presolar silicon carbide grains in primitive meteorites determined by NanoSIMS, *Geochimica et Cosmochimica Acta* 139, 248 – 266 (2014). DOI: 10.1016/j.gca.2014.04.026
85. C. M. O'D. Alexander, R. Bowden, M. L. Fogel, K. T. Howard, C. D. K. Herd, L. R. Nittler, The provenances of asteroids, and their contributions to the volatile inventories of the terrestrial planets, *Science* 337, 721–723 (2012). DOI: 10.1126/science.1223474

**Acknowledgments:** We thank the anonymous reviewers and the editors for their helpful comments. The Hayabusa2 spacecraft was developed and built under the leadership of Japan Aerospace Exploration Agency (JAXA), with contributions from the German Aerospace Center (DLR) and the Centre National d'Études Spatiales (CNES), and in collaboration with NASA, Nagoya Univ., Univ. of Tokyo, National Astronomical Observatory of Japan (NAOJ), Univ. of Aizu, Kobe Univ., and other universities, institutes, and companies in Japan. We thank the many engineers, including N. Inaba at JAXA and T. Masuda, S. Yasuda, K. Matsushima, and T. Ohshima at NEC Corp. for their dedicated work on the Hayabusa2 mission.



5 The STXM data were acquired at beam line 19A of the Photon Factory, High Energy Accelerator Research Organization, and at beam line 5.3.2.2 of the Advanced Light Source (ALS), Lawrence Berkeley National Laboratory. Use of the ALS was supported by the Department of Energy (DOE) (contract no. DE-AC02-05CH11231). The synchrotron FTIR data were acquired at the SMIS beamline of SOLEIL. The NanoSIMS facility at Muséum National d'Histoire Naturelle (MNHN) in Paris was established by funds from the CNRS, Région Ile de France, Ministère délégué à l'Enseignement supérieur et à la Recherche, and the Muséum National d'Histoire Naturelle. The FIB facility at the University of Tokyo is supported by the Advanced Characterization Nanotechnology Platform in the Nanotechnology Platform Project sponsored by 10 the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan.

We thank N. Sugimura at Waseda University, Y. Fujioka at Tohoku University, C. Sandt. and F. Borondics at SOLEIL for technical support. We thank S. Kawakami at Gifu Shotoku University, T. Fagan at Waseda University, M. E. Zolensky at NASA Johnson Space Center, MNHN, and B. Fectay and C. Bidaut for providing the meteorite samples used in this study.

15 **Funding:** H.Yab was supported by KAKENHI from the Japan Society for the Promotion of Science (JSPS) (grant nos. JP20H05846, 19H01954, 18H04461, 20H04615) and the Astrobiology Center Program of National Institutes of Natural Sciences (NINS) (Grant Number AB312007, AB022001 and AB032004). Y.Taka, Y.Take, Y.K., H.Suga, S.Y., and D.W. were supported by KAKENHI from JSPS (grant no. JP17H06458). H.Yab, Y.K., M.Hash. and H.Suga were supported by the JSPS Core-to-Core program “International Network of Planetary Sciences”.

L.Bo. and E.Q. acknowledge funding from the Centre National d'Etudes Spatiales (CNES-France).

25 L.R. acknowledges funding from the European Research Council through the consolidator grant HYDROMA (grant agreement No. 819587).

C.E., J.D., E.D., and J.M. acknowledge funding from CNES (MIAMI2), DIM-ACAV+ (C3E), LabEx P2IO, PCMI, and ANR (COMETOR ANR-18-CE31-0011).

30 R.M.S and B.D.G acknowledge funding from NASA Awards 80HQTR20T0014 and 80HQTR20T0050.

L.R.N. acknowledges funding from NASA Grants NNX16AK72G and 80NSSC20K0340.

A.Da., A.De. and J.M. acknowledge funding from Paris Ile-de-France Region-DIM “Matériaux anciens et patrimoniaux.”

35 Centro de Química Estrutural (Z.M.) acknowledges the financial support of Fundação para a Ciência e Tecnologia (FCT) (grants UIDB/00100/2020, UIDP/00100/2020 and LA/P/0056/2020).

#### **Author contributions:**

40 H.Yab. led the research and wrote the paper with contributions from L.R.N., B.D.G., R.M.S., R.L., L.B., Y.K. and G.D.C.

Sample preparation for the research: H.Yab., B.D.G., T.O., M.Hash, C. E., L.Bo., M.Matsum., Y.E., K.K., M.S.;

Raman analyses: L.Bo, M.K., E.Q., G.M., Y.K.;

FTIR analyses: Y.K., E.Q., E.D., L.Bo., C.E., J.D.;

5 STXM analyses: B.D.G., H.Yab., Y.K., G.C., D.K., S.S., R.S., Y.Taka, T.O., H.Suga, Y.Take, S.Y., D.W., Y.Tam;

STEM-EELS-EDS analyses: R.M.S.;

AFM-IR analyses: J.M., A.Da., A.De., E.D., C.E., J. D., L.Be;

NanoSIMS analyses: L.R., L.R.N., M.V.-P., S.M., J.B.

10 Interpretation and writing: H.Yab., L.R.N., R.M.S., B.D.G., R.L., L.B. Y.K, E.D., J.M., E.Q., M.K., G.D.C., C. E., S.S., T.O., M.Hash., S.Tachi.

Administration of the initial analysis: S.Tachi., H.Yur, T.Nak, T.Nog, R.O., H.Yab., H.Na, K.S., H.C.Jr., D.S.L.

15 Sample curation: M.Abe, T.O., T.Yada, M.N., K.Y., A.N., M.Y., A.I., A.Miya, S.F., K.Hata, H.So, Y.H., K.Ku, T.U., T.H., D.Y., and R.F.

20 Sample collection at Ryugu: Y.Tsu., T.S., F.T., S.Tanak, M.O., R.T., H.Yan., H.Sawa., M.Haya., T.I., S.H, Y-II., A.F., Y.Takei, K.Yoshikawa, Y.M., G.O., H.T., N.O., S.K., K.S., Y.Y., H.I., K.K., K.Yoshihara, T.Y., H.No., A.Miur., S.Su, K.Ki., N.H., R.H., T.Mo., E.T., T.O., N.S., N.N., K.M., R.N., K.W., H.Se, K.O, Y.Y.,Y.I., Y.S., M.Y., C.H., T.Mi, M.Matsuo, N.H., M.A., C.O., M.I., H.Yan, S.Tachi., R.O., S.W., T.Nak, H.Yab., R.J., J.-P. B., M.G., S.Sc, K.O., C.P., N.S., J.B., T.-M. H., and A.Mous.

Project management: Y.Tsu., S.W., M.Y., M.F., and S.N.

All authors discussed the results and commented on the manuscript.

25 **Competing interests:** The authors declare that they have no competing interests.

### **Data and materials availability:**

30 All images used in this study and machine-readable data files are available at the JAXA Data Archives and Transmission System (DARTS) at [https://data.darts.isas.jaxa.jp/pub/hayabusa2/paper/sample/Yabuta\\_2022/](https://data.darts.isas.jaxa.jp/pub/hayabusa2/paper/sample/Yabuta_2022/) . Data for all Hayabusa2 samples and other data from the mission are available at the DARTS archive at <https://www.darts.isas.jaxa.jp/curation/hayabusa2> and <https://www.darts.isas.jaxa.jp/planet/project/hayabusa2/>, respectively. The samples of Ryugu are curated by the JAXA Astromaterials Science Research Group; distribution for analysis is through an Announcement of Opportunity available at <https://jaxa-ryugu-sample-ao.net> . The samples we used are listed in Table S1.

35

## Supplementary Materials

Materials and Methods

Supplementary Text

Figs. S1 to S8

5 Tables S1 to S3

References (77–85)

10 **Fig. 1. Raman spectra and spectral parameters: comparison between Ryugu samples and chondrites.** (A): Average Raman spectra of Ryugu grains from Chamber A (samples A0108-6, -10, and -18) (in grey) and Chamber C (samples C0109-5, -9, and -12) (in black), compared to the meteorites Murchison (CM, green) and Orgueil (CI, yellow). All spectra were acquired in the same analytical conditions (11). (B-D): Average spectral parameters (error bars show standard deviations) determined from the Raman spectra: (B)  $\text{FWHM}_G$  as a function of  $\omega_G$ ; (C)  $\text{FWHM}_D$  as a function of  $I_D/I_G$ ; (D)  $\omega_G$  as a function of  $\omega_D$ . Data are for individual grains from Chamber A (open circles) and Chamber C (black filled circles) compared to petrologic type 1 (Orgueil, orange diamond), unheated petrologic type 2 chondrites (green filled diamonds), heated petrologic type 2 chondrites (open blue diamonds) and petrologic type 3 chondrites (filled grey diamonds) (11). Petrologic type 2 chondrites were classified (16) as unheated (e.g., Murchison, Nogoya, and Tarda) or heated (e.g., Jbilet Winselwan and WIS 91600).

25 **Fig. 2. Micro-FTIR spectra of Ryugu samples compared to chondrites.** (A) Infrared transmission spectra (11) of six Ryugu grains from the aggregates A0108 and C0109 (black), heated CM chondrite (Jbilet Winselwan) (red), unheated CM Murchison (pink), and two CI chondrites Orgueil (light blue) and Ivuna (dark blue). All the spectra were baseline-corrected using spline curves and normalized by the peak height of the band at  $\sim 1000 \text{ cm}^{-1}$  ( $\sim 10.00 \mu\text{m}$ ). Dashed lines indicate identified bands: silicate OH and SiO at  $3685$  to  $3675 \text{ cm}^{-1}$  ( $2.71$  to  $2.72 \mu\text{m}$ ) and  $\sim 1000 \text{ cm}^{-1}$  ( $\sim 10.00 \mu\text{m}$ ), aliphatic C-H bands at  $2960 \text{ cm}^{-1}$  ( $3.38 \mu\text{m}$ ,  $\text{CH}_3$  asymmetric stretching),  $2930$  to  $2925 \text{ cm}^{-1}$  ( $3.41$  to  $3.42 \mu\text{m}$ ,  $\text{CH}_2$  asymmetric stretching),  $2855$  to  $2850 \text{ cm}^{-1}$  ( $3.50$  to  $3.51 \mu\text{m}$ ,  $\text{CH}_3$  and  $\text{CH}_2$  symmetric stretching),  $1460 \text{ cm}^{-1}$  ( $6.85 \mu\text{m}$ ) and  $1380 \text{ cm}^{-1}$  ( $7.25 \mu\text{m}$ , bending), other organic features at  $1705$  to  $1690 \text{ cm}^{-1}$  ( $5.87$  to  $5.92 \mu\text{m}$ , C=O), and  $\sim 1600 \text{ cm}^{-1}$  ( $\sim 6.25 \mu\text{m}$ , aromatic with some water bending mode contribution). Some spectra show a peak at  $\sim 1430 \text{ cm}^{-1}$  ( $\sim 6.99 \mu\text{m}$ ) due to carbonates. A broad water stretching band at  $\sim 3400 \text{ cm}^{-1}$  ( $\sim 2.94 \mu\text{m}$ ) is observed, which is weaker in samples that was measured at  $60^\circ\text{C}$  or higher temperatures. The peaks at  $2360 \text{ cm}^{-1}$  ( $4.24 \mu\text{m}$ ) are due to atmospheric  $\text{CO}_2$ . The A0108-5 and C0109-1 Ryugu grains, and Murchison and Ivuna meteorites were measured at  $60^\circ\text{C}$  under  $\text{N}_2$  flow. The Jbilet Winselwan meteorite was measured at  $80^\circ\text{C}$  under  $\text{N}_2$  flow. The A0108-10 and C0109-12 Ryugu grains were measured at  $80^\circ\text{C}$  under vacuum. The Orgueil meteorite was measured at  $130^\circ\text{C}$  under vacuum. (B) Same as panel A, but for insoluble carbonaceous residues from the Ryugu samples (A0106 and C0107), compared to IOM from the heated CM meteorite Y-793321 (16), the CM Murchison (27), and CI meteorite Ivuna (27). The spectra were the averages of main fractions of A0106 and C0107, respectively, after baseline-correction using

spline curves and normalization by the peak height of the aromatic C=C band at  $\sim 1600\text{ cm}^{-1}$  ( $\sim 6.25\ \mu\text{m}$ ).

**Fig. 3. STXM elemental maps and carbon-XANES spectra of Ryugu samples.** STXM greyscale images of FIB sections extracted from the Ryugu grains (A) A0108-3, observed at 530 eV and (B) A0108-11, observed at 390 eV (11). In both panels, color overlays on both FIB sections are false-color maps of X-ray absorptions due to aromatic C (cyan), carbonate functional groups (magenta), and calcite minerals (yellow). Dashed circles and boxes indicate regions measured for panel E. (C) STXM image at 290 eV of insoluble carbonaceous residue from Ryugu sample C0107-18. The orange box shows the region in panel D. (D) X-ray absorption map of aromatic carbon (cyan), ketones (magenta), and carboxyl (yellow) functional groups. Circled features are solid nanoglobules except for feature 18f, which is a cluster of typical Ryugu IOM. (E) Carbon-XANES spectra for carbonaceous grains and matrix regions identified in panels A, B and D. IOM from the meteorites Orgueil and Murchison (black) are shown for comparison (11).

**Fig. 4. STEM-EELS-EDS analysis of Ryugu samples.** All scale bars are 500 nm. (A) False color bright field TEM image of the ultrathin section of C0109-11, indicating nanoglobules (magenta), Mg-phyllsilicates (blue), FeS (orange). (B) bright field TEM image of insoluble organic residue from a microtome slice of sample C0107, showing hollow and solid nanoglobules and fluffy materials. (C) High angle annular dark field (HAADF) STEM image of a FIB section of A0108-11 (region 11h in Fig. 3B) showing two magnetite grains (labelled Mag) adjacent to carbon-bearing Mg-rich phyllosilicates (Mg-phy) with Fe, Ni sulfide nanoparticles (Sulf). (D) HAADF STEM image of a vesicle in the large calcite grain from a FIB section of sample A0108-3 (region 3a in Fig. 3A). (E) EDS element map of carbon (magenta), oxygen (cyan) and sulfur (yellow) of the area shown in (C). (F) EDS element map of carbon (magenta), calcium (green), oxygen (yellow) and sulfur (cyan) of the vesicle shown in (D) showing diffuse organic matter, Fe,Ni-sulfides and Ca- sulfate. (G) Carbon EEL spectra from molecular carbonate incorporated into phyllosilicates, i.e., clay-bound carbon (C0109-11), an aromatic nanoglobule (A0108-39), an IOM-like nanoglobule (C0109-11), and nanodiamonds (A0108-8). Spectra were acquired at 0.02 eV per channel, and are smoothed to 0.4 eV resolution, after power law background subtraction and arbitrary scaling. (H) EDS spectra from molecular carbonate clay-bound carbon (C0109-11, green) and an insoluble organic residue (A0106-9, magenta). The asterisk (\*) indicates a Cu background peak from the sample support and microscope.

**Fig. 5. AFM-IR analysis of intact Ryugu grains A0108 and C0109.** Composite AFM-IR images of the intact Ryugu grains (A) A0108-15 in tapping mode,  $2 \times 2\ \mu\text{m}$  and (H) C0109-4, in contact mode,  $3 \times 3\ \mu\text{m}$ . Colors indicate the C=O ( $1720\text{ cm}^{-1}$ ,  $5.81\ \mu\text{m}$ , red), C=C ( $1600\text{ cm}^{-1}$ ,  $6.25\ \mu\text{m}$ , blue) and Si-O ( $1020\text{ cm}^{-1}$ ,  $9.80\ \mu\text{m}$ , green) peaks. Each image has been normalized to its maximum peak value. Organic matter is widespread throughout each sample. In panel A, small organic globules are visible (red-brown) surrounded by a dominant phyllosilicate. The magenta box indicates the inclusion used for panel D. (B) AFM topographic map of the  $2 \times 2\ \mu\text{m}$

area in panel A. (C) Visible light image ( $75 \times 75 \mu\text{m}$ ) showing the location of the AFM-IR map (green square). (D) AFM-IR spectrum (red) of the globule indicated in panel A and shown in the inset, and background spectrum (green) taken in a region 100 nm away. The AFM-IR signal is shown in mV, which is proportional to optical depth ( $I/I_0$ ). The C=C and C=O absorptions spectral region is also shown with a  $\times 5$  factor for better visibility. (E-G) Synchrotron FTIR maps of the same  $75 \times 75 \mu\text{m}$  area as panel C, of (G) the C=O ( $1720 \text{ cm}^{-1}$ ,  $5.81 \mu\text{m}$ , red), (F) C=C ( $1600 \text{ cm}^{-1}$ ,  $6.25 \mu\text{m}$ , blue) and (E) Si-O ( $1020 \text{ cm}^{-1}$ ,  $9.80 \mu\text{m}$ , green) peaks. Data were acquired with a  $6 \times 6 \mu\text{m}$  beam size, sampled with  $3 \mu\text{m}$  steps. Color bars show the integrated optical depths calculated for each band. (H) Organic matter in the Ryugu grain C0109-4 appears as red-purple inclusions standing out from the dominant phyllosilicate signal. Arrows indicate the positions used for the spectra in panel K. (I) AFM topographic map of the  $3 \times 3 \mu\text{m}$  area. (J) Visible image ( $51 \times 51 \mu\text{m}$ ) showing the location of the AFM-IR map (green square). (K) AFM-IR spectra of a C=O poor region (labelled 1 in panel H) and organic inclusions (labelled 2), compared to the average synchrotron FTIR spectrum. Red, blue and green tick marks indicate spectral positions corresponding to the AFM-IR images combined in panel H. (L-N) Same as panels E-G, but for the  $51 \times 51 \mu\text{m}$  region shown in panel J. To aid color-blind readers, another version of this figure with alternatives colors is provided in fig. S5.

**Fig. 6. NanoSIMS analysis of the intact Ryugu grains A0108 and C0109.** (A-C) NanoSIMS images of a  $\sim 3 \times 3 \text{ mm}$  area of a microtome slice from Ryugu grain A0108-11. (A) map of  $^{16}\text{O}$  (cyan) and  $^{12}\text{C}$  (red). White outlines indicate four carbon-rich particles embedded in the silicate matrix. (B) Hydrogen and (C) nitrogen isotope abundances of the same region. The ratios vary between the carbonaceous inclusions. (D-F) H and N- isotopic ratios for individual carbon-rich particles (regions of interest, ROIs) within Ryugu grains. Most ROIs are consistent (within the uncertainties) with the bulk averages; they are represented by box-and-whisker plots where the box size represents the inner 50% of data around the median (interquartile range, IQR) and the whiskers indicate the  $\pm 1.5$  IQR range beyond that. Dots indicate outliers beyond those ranges (hot-spots and cold-spots) from A0108 (blue) and C0109 (orange). In panel D, white dots indicate the grains shown in panels A-C and the error bar is  $1\sigma$ . The distributions of H and N isotope ratios are consistent between panel D and those in E and F (acquired in different laboratories) and between the two Ryugu samples. The pink shaded region indicates the ranges of H and N hot-spots and cold-spots in primitive CM chondrite meteorites (50) and the green box indicates the ranges seen in CI chondrite Orgueil (46). The Ryugu outliers span a similar range to the CI and CM chondrites. (G)  $\delta^{15}\text{N}$  as a function of  $\delta^{13}\text{C}$  for the Ryugu grain A0109. The grey points indicate all carbon-rich ROIs, while blue circles are outliers ( $>2\sigma$  away from bulk average). The pink region is the same as in panel D. The Ryugu outliers span a similar range to that seen in CM chondrites. The dashed vertical line indicates the bulk  $\delta^{13}\text{C}$  of CI chondrite IOM.

**Fig. 7. NanoSIMS analysis of insoluble carbonaceous residues from Ryugu grains A0106 and C0107.** Hydrogen (A,B) and Carbon (C,D) maps for insoluble carbonaceous residues of A0106 and C0107, respectively, acquired at the same locations. The corresponding H (E, F) and N (G, H) isotope ratio maps show numerous micron-sized D-rich and  $^{15}\text{N}$ -rich hot-spots. The carbon and hydrogen images indicate that these hot-spots consist of organic carbon. (I) Hydrogen

and (J) nitrogen box plot diagrams showing the distributions of isotope ratios in the insoluble carbonaceous residues A0106-8 (orange) and C0107-14 (blue). Open grey circles show hot-spots (or cold-spots), micron-sized areas with isotope ratios much higher (or lower) than the average composition. Large orange and blue circles indicate the bulk average value for each residue. Horizontal dashed lines indicate the mode of each distribution of hot-spots. Data for the IOM of CI (light blue), CM (green), CR (red) and the Tagish Lake (yellow) chondrites are shown for comparison. Filled circles are bulk values (45); for Tagish Lake subsamples 5b (less altered) and 11v (altered) (61) are shown separately. Upward pointing arrows (some extend beyond the plot) indicate the range of hot-spots reported for those meteorites (46, 56, 57). For Orgueil, the downward arrow indicates the extent of <sup>15</sup>N-depleted cold-spots.