



## Quantitative Analysis of NH<sub>2</sub>\* in Premixed Ammonia-Air Flames

Alka Karan, Guillaume Dayma, Christian Chauveau, Fabien Halter

### ► To cite this version:

Alka Karan, Guillaume Dayma, Christian Chauveau, Fabien Halter. Quantitative Analysis of NH<sub>2</sub>\* in Premixed Ammonia-Air Flames. 11th European Combustion Meeting (ECM 2023), Apr 2023, Rouen, France. . hal-04085381

HAL Id: hal-04085381

<https://cnrs.hal.science/hal-04085381>

Submitted on 28 Apr 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Quantitative Analysis of NH<sub>2</sub>\* in Premixed Ammonia-Air Flames

Alka KARAN<sup>I,II</sup>, Guillaume DAYMA<sup>I,II</sup>, Christian CHAUVEAU<sup>II</sup>, Fabien HALTER<sup>I,II</sup>

<sup>I</sup>Université d'Orléans, Château de la Source, 45100 Orléans, France

<sup>II</sup>CNRS – ICARE, 1C Avenue de la Recherche Scientifique, 45071 Orléans Cedex 2, France


The interest in studying ammonia flames arises from the need to curb carbon emissions and to solve the energy crisis arising from the depletion of fossil reserves.

## Why chemiluminescence?

- A cheap and non-intrusive diagnostic tool
- It provides various information on the combustion process, like assessing flame ratio and understanding flame topology.

## Why NH<sub>2</sub>\* chemiluminescence?

- NH<sub>2</sub>\* chemiluminescence is used to determine the position of the maximum HRR in premixed ammonia-air flames.
- The relative intensity of NH<sub>2</sub>\* can give information on NOx formation





## Why quantitative analysis?

Possibility to have reference data for the chemiluminescent part of the species to model the total excited species.



## Procedure

- NH<sub>2</sub>\* chemiluminescence data at 632 nm for ambient conditions.
- Calibration to quantify the data.

## Chemiluminescence set-up



Determining the position of NH<sub>2</sub>\* within the flame





Normalized NH<sub>2</sub>\* distribution along the flame for different  $\varphi$

## Calibration process


### Step 1: Optics Calibration

NH<sub>2</sub>\* filter response and camera response against a fixed light source



### Step 2: Tungsten lamp calibration

- Irradiation data as a function of wavelength
- Filter and camera response against the tungsten lamp



### Step 3: Converting intensity to concentration<sup>1</sup>


$$\frac{I_{\text{species,Abel}}}{I_c} = \frac{1}{4\pi} [\text{species}] N_A \eta_\lambda \tau_1 \Delta t_1 \int_{\lambda 1}^{\lambda 2} \frac{S_c(\lambda)}{h\nu} \eta_\lambda d\lambda \tau_2 \Delta t_2$$

### Step 4: Adapting to the configuration

- Irradiation values were provided at a distance of 50 cm
- Values adapted to fit the experimental set-up distance

## Results

- On moving from fresh gases towards burnt gases, NH<sub>2</sub>\* profiles for a low stretch can be plotted.
- It is seen that the thickness of NH<sub>2</sub>\* increases with the equivalence ratio.
- Preferential diffusion of lighter species towards the base of the flame is noted for leaner flames.
- Future work includes accommodating these species in any existing chemical kinetic schemes.



NH<sub>2</sub>\* concentrations for different  $\varphi$  plotted for minimum stretch values

## Acknowledgement

Support from the CAPRYSSES project (ANR-11-LABX-006-01) funded by the PIA (Programme d'Investissement d'Avenir) is gratefully acknowledged.

ICARE-CNRS and Université d'Orléans are acknowledged as well.

## Reference

<sup>1</sup>Y. Liu, J. Tan, M. Wan, L. Zhang, X. Yao, Quantitative measurement of OH\* and CH\* chemiluminescence in jet diffusion flames, ACS Omega, (2020), 15922–30

