

### Citizen science approach for Galactic Cosmic Radiation and SEP monitoring at commercial flight altitude

Francois Trompier

#### ▶ To cite this version:

Francois Trompier. Citizen science approach for Galactic Cosmic Radiation and SEP monitoring at commercial flight altitude. European Radiation Protection Week 2022 (ERPW 2022), Oct 2022, Estrosil, Portugal. , 10.13140/RG.2.2.32620.03208. hal-04096070

#### HAL Id: hal-04096070 https://cnrs.hal.science/hal-04096070

Submitted on 12 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License




# **COSMIC UNAR** : Citizen science approach for **Galactic Cosmic Radiation and SEP** monitoring at commercial flight altitude

The Solar flare hunt is onl F. Trompier<sup>1</sup>, G. Darley<sup>1</sup>, N. Berthelot<sup>2</sup>, JF. Bottollier-Depois<sup>1</sup>, C. Briand<sup>3</sup>, A. Buffler<sup>4</sup>, S. Cnudde<sup>3</sup>, C. Damon<sup>1</sup>, N. Fuller<sup>3</sup>, T. Hutton<sup>4</sup>, E. Jarvie<sup>4</sup>, K.-L. Klein<sup>3</sup>, V. Lejeune<sup>1</sup>

### Scientific context

Intense solar energy proton events with an effect in the atmosphere are sporadic and rare but can generate a high dose rate at flight altitude. Very few such events have been measured, making it impossible to validate the dose estimation model. This project aims to provide measured data to the scientific community through a citizen science approach.






Figure 1. Solar flares.

### 06:39 08:21 09:48 NO:51 NY:42 NO.98 NO:55 NO:41

### Temps universel

**Figure 2.** Measured dose rate due to the galactic component of cosmic rays compared with additional contribution from SEP: the dose is doubled on the same flight.

### Project partners 3 1) **RSN bservatorre** INSTITUT de Paris open DE RADIOPROTECTION ET DE SÛRETÉ NUCLÉAIRE radiation $\begin{array}{c|c} \textbf{Department of} \\ \textbf{PHYSICS} \end{array} M_{|} e_{|} A_{|} S_{|} U_{|} R_{|} e \end{array}$

# How to participate?

Contact via OpenRadiation website or cosmic@irsn.fr to participate in:

- **Detector development or characterization;**
- Improving website and apps;
- Data analysis and measuring; and
- Proposing new actions (balloon flights,..).






Figure 3. Measurement campaigns with balloon flights (crédit Planète sciences).

Figure 5.

## Data collected and calibration aspects

Data are collected through the Openradiation




- website and apps or any other detectors with a data logger (Safecast, GQ family).
- A dedicated mode of measurements in airplanes is now available in the OR app.
- Specific calibration coefficients for CR have been derived from in-flight measurements by comparison with reference detectors (Hawk TEPC) or with calculation (CARI-7).



- 225 flights measured, 57195 measurements.
- All measurements were made under normal space weather conditions, but the solar maximum is expected to be reached in 2025, with an increased probability of intense solar flares.


**Figure 4.** In-flight measurements with OR app.



OR kit detectors.

### **Devices characterized**

- Devices can be connected by BT to the OpenRadiation app (openradiation) kit, Safecast, Rium GM, Atomtag, Pocketgeiger, Polimaster).
- **Detection principles: GM tube or photodiode.**





Océan


retique

Ocean

ecifique.

Alternative approaches

Due to the cost of the devices, evaluations with webcams or CMOS from smartphone cameras are being studied to democratise the measurement, or with scintillators for the measurement of secondary neutrons.



**Figure 8.** Screenshots of CR detection in CMOS sensor of smartphone camera.



Figure 9. Prototype of the compact neutron detector based on scintillator from MeASURe.