

Protocol for emergency dosimetry based on EPR spectrometry on nails

Francois Trompier, Alexander Romanyukha, Lucile Rémy-Kornak, Céline Calas, Blaise LeBlanc, Isabelle Clairand, Chad Mitchell, Harold Swartz

► To cite this version:

Francois Trompier, Alexander Romanyukha, Lucile Rémy-Kornak, Céline Calas, Blaise LeBlanc, et al.. Protocol for emergency dosimetry based on EPR spectrometry on nails. 7th International Symposium on EPR Dosimetry and Applications and the 2nd International Conference on Biodosimetry, Jul 2006, Bethesda, United States. Elsevier, Radiation Measurements, 42 (6-7), pp.1085, 2007, 10.13140/RG.2.2.18565.73443. hal-04103726

HAL Id: hal-04103726 https://cnrs.hal.science/hal-04103726

Submitted on 23 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Protocol for emergency dosimetry based on EPR spectrometry on nails

In case of an **accident** involving a **large** number of victims, most of dosimetric techniques are limited by the necessary short delay of the answer or the invasive sampling. In this specific case, it is very important in a first time and as quick as possible to sort population according to severity of the exposure. the Nevertheless, EPR dosimetry on materials easily sampled on the victims (hairs or nails) may be a pertinent tool. In this context, the objective of this work was to study the EPR properties of nails to propose a first protocol for fast triage of population.



Dose estimation Protocol

The protocol is divided in different stages. A first estimation is done very quickly using averaged value of main characteristics (fading constant, radiation sensitivity, amplitude of background signal) previously determined on a large set of samples. At each next stage, the determination of the real values of the different constants for each sample allows to **increase the accuracy** of the dose estimation.





F. Trompier⁽¹⁾, A. Romanyukha⁽²⁾, L. Remy-Kornak⁽¹⁾, C. Calas⁽¹⁾, B. LeBlanc⁽²⁾, I. Clairand⁽¹⁾, C. Mitchell⁽²⁾, H. Swartz⁽³⁾ ⁽¹⁾ IRSN, ⁽²⁾USUHS, ⁽³⁾ Dartmouth Medical School Contact : francois.trompier@irsn.fr

EPR Dose estimation

Dose= (S_{total} - S_{native-}S_{MIS}).F(t, T°, H).A with $S_{total} = S_{RIS} + S_{native} + S_{MIS}$

F : fading correction factor depending on time elapsed between irradiation and measurement \mathbf{t} , temperature \mathbf{T}° and humidity **H** and **A**: dose coefficient

Limitation and Elimination of MIS

Determination of S_{total} , dose estimation with mean value of S_{native} , A and F

Determination of real value of S_{native}

Determination of real value of **A**

Estimation of sample humidity (H): improvment of estimation of F

This study showed that it was possible, starting from EPR measurement of nails to carry out a population triage according to the severity of the exposure