



Chemiluminescence in Premixed Ammonia-Air Flames

Alka Karan, Guillaume Dayma, Christian Chauveau, Fabien Halter

► To cite this version:

Alka Karan, Guillaume Dayma, Christian Chauveau, Fabien Halter. Chemiluminescence in Premixed Ammonia-Air Flames. Topic-Oriented Technical Meeting "Chemical energy carriers for long-term storage and long-distance transport of renewable energies" (TOTeM 49), Jun 2023, Mulhouse, France.
hal-04159133

HAL Id: hal-04159133

<https://cnrs.hal.science/hal-04159133>

Submitted on 11 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chemiluminescence in Premixed Ammonia-Air Flames

Alka KARAN^{I,II}, Guillaume DAYMA^{I,II}, Christian CHAUVEAU^{II}, Fabien HALTER^{I,II}


^IUniversité d'Orléans, Château de la Source, 45100 Orléans, France

^{II}CNRS – ICARE, 1C Avenue de la Recherche Scientifique, 45071 Orléans Cedex 2, France

Ammonia combustion is of great interest as it is a zero-carbon fuel and has a high octane rating. However, the flame properties have not been studied well. The objective of this work is to use an economic diagnostic tool to obtain useful data.


Chemiluminescence is a cheap and non-intrusive diagnostic tool that is widely used to obtain a large range of information pertaining to flame behaviour. In ammonia combustion, NH_2^* , NH^* , and OH^* have been identified as species of relevance.

Heat release rate marker




NH^* and NH_2^* have been identified as the HRR marker pure ammonia flames on a Bunsen burner configuration.

Flame reactivity



Change of intensity in NH_2^* profiles on a Bunsen burner configuration indicating local change of reactivity.

NO emissions



Relative NO* emissions vs corresponding proportional OH* emissions for two different equivalence ratios indicating OH* predictions for relative NO quantities


KURO Camera Objective lens Filter

Chemiluminescence set-up


NH_2^* - 632 nm

NH^* - 337 nm









OH^* - 310 nm



On superimposing NH^* and NH_2^* inverse Abel transformed image with the DEHS droplets evaporating at 525 K, the position of the species can be determined within the flame front



NH_2^* sub-mechanism: proposed reactions



Experiment vs simulation

The proposed sub-mechanism can predict the width of the profiles quite well. The peak values depend on the base model. There is still room for improvement. Future work includes proposing sub-mechanism for other species.

Acknowledgement

Support from the CAPRYSES project (ANR-11-LABX-006-01) funded by the PIA (Programme d'Investissement d'Avenir) is gratefully acknowledged.