



HAL
open science

TMED10 mediates the loading of neosynthesised Sonic Hedgehog in COPII vesicles for efficient secretion and signalling

Yonis Bare, Tamás Matusek, Sophie Vriz, Maika S Deffieu, Pascal Théron, Raphael Gaudin

► **To cite this version:**

Yonis Bare, Tamás Matusek, Sophie Vriz, Maika S Deffieu, Pascal Théron, et al.. TMED10 mediates the loading of neosynthesised Sonic Hedgehog in COPII vesicles for efficient secretion and signalling. Cellular and Molecular Life Sciences, 2023, 80 (9), pp.266. 10.1007/s00018-023-04918-1 . hal-04261574

HAL Id: hal-04261574

<https://cnrs.hal.science/hal-04261574>

Submitted on 27 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

TMED10 mediates the loading of neosynthesised Sonic Hedgehog in COPII vesicles for efficient secretion and signalling

Yonis Bare^{1,2}, Tamas Matusek³, Sophie Vriz^{4,5}, Maika S. Deffieu^{1,2}, Pascal Therond³, Raphael Gaudin^{1,2}

5 #

¹ Institut de Recherche en infectiologie de Montpellier (IRIM) CNRS, 34293 Montpellier, France.

² Université de Montpellier, 34090 Montpellier, France.

³ Université Côte d'Azur, UMR7277 CNRS, Inserm 1091, Institut de Biologie de Valrose (iBV), Parc
10 Valrose, 06108 Nice, France.

⁴ Laboratoire des biomolécules (LBM), Département de chimie, École Normale Supérieure, PSL
University, Sorbonne Université, CNRS, 75005 Paris, France

⁵ Faculty of Science, Université de Paris, Paris, France.

15 #Corresponding author:

Raphael Gaudin

1919 Route de Mende

34293 Montpellier, France

Tel: +33 4 34 35 94 78

20 Email: raphael.gaudin@irim.cnrs.fr

Keywords: Biosynthetic pathway; p24 proteins; ERES; membrane trafficking; endoplasmic reticulum

Running title: TMED10-mediated secretion of SHH

25 **Abstract**

The morphogen Sonic Hedgehog (SHH) plays an important role in coordinating embryonic development. Short- and long-range SHH signalling occurs through a variety of membrane-associated and membrane-free forms. However, the molecular mechanisms that govern the early events of the trafficking of neosynthesised SHH in mammalian cells are still poorly understood. Here, we employed the retention using selective hooks (RUSH) system to show that newly-synthesised SHH is trafficked through the classical biosynthetic secretory pathway, using TMED10 as an ER cargo receptor for efficient ER-to-Golgi transport and Rab6 vesicles for Golgi-to-cell surface trafficking. TMED10 and SHH colocalized at ER exit sites (ERES), and TMED10 depletion significantly delays SHH loading onto ERES and subsequent exit leading to significant SHH release defects. Finally, we utilised the *Drosophila* wing imaginal disc model to demonstrate that the homologue of TMED10, *Baiser (Bai)*, participates in Hedgehog (Hh) secretion and signalling *in vivo*. In conclusion, our work highlights the role of TMED10 in cargo-specific egress from the ER and sheds light on novel important partners of neosynthesised SHH secretion with potential impact on embryonic development.

40

Introduction

Intercellular communication plays an essential role in directing cell fate during embryonic development (Wolpert 2016). The Hedgehog signalling pathway illustrates very well this statement as the Hedgehog (Hh) family proteins have been shown to play pivotal roles such as acting as a morphogen, embryonic patterning during development, and regeneration of adult tissues (Dahmane and Ruiz i Altaba 1999; Palma et al. 2005; Straface et al. 2009; Briscoe and Théron 2013; Álvarez-Buylla and Ihrie 2014; Groves, Placzek, and Fletcher 2020). Abnormal Hh expression and signalling has been shown to be responsible for cerebellar hypoplasia, and also for numerous cancers, including medulloblastoma and glioblastoma (Corrales et al. 2006; Rubin and de Sauvage 2006; Melamed et al. 2018). In metazoans, Hh has been shown to mediate cell differentiation and proliferation in the central nervous system (Yam and Charron 2013; Hill et al. 2019; Brady and Vaccarino 2021). In addition, Sonic Hedgehog (SHH), one of three Hh isoforms expressed in mammals, plays a pleiotropic and pivotal role in human physiopathology, such as holoprosencephaly and Pallister-Hall syndrome (Maity, Fuse, and Beachy 2005; McClelland, Li, and Rosenblum 2022). However, the mechanisms governing the secretion of neosynthesized Hh family proteins remains incompletely understood.

The biosynthetic secretory pathway is a tightly regulated process to ensure precise coordinated trafficking of cargoes to their destined compartments. Newly synthesised proteins are transported into the endoplasmic reticulum (ER) through a protein translocon and undergo protein quality control through various molecular chaperones. Proteins are then sorted and concentrated into ER exit sites (ERES) by the coat protein complex II (COPII), and bud off as ER-derived vesicles (Zanetti et al. 2012; Hutchings and Zanetti 2019). These vesicles fuse with other vesicular-tubular structures in the ER-Golgi intermediate compartment (ERGIC) for microtubule-dependent sorting to the Golgi complex (Saraste and Marie 2018; Peotter et al. 2019). At the Golgi complex, cargoes are packaged and sorted to be delivered to various subcellular compartments, such as endosomes, lysosomes and the plasma membrane (PM), with the help of the recruitment of small Rab GTPases, including Rab6 and Rab7 (Goud, Liu, and Storrie 2018; Borchers, Langemeyer, and Ungermann 2021; Deffieu et al. 2021).

SHH is first synthesised as a full-length 45 kDa precursor in the cytoplasm, with an N-terminal ER signal sequence. Upon entering into the ER, the signal peptide is cleaved, revealing a cysteine residue as the extreme N-terminal residue. The C-terminal domain of SHH contains an auto-processing catalytic domain, which mediates the self-cleavage of the precursor protein into two halves. The SHH-C protein (26 kDa) is degraded in the ER. In contrast, the SHH-N fragment possesses all signalling

activity of SHH that is post-translationally modified with a cholesterol molecule covalently bound to the C-terminus of SHH-N and a palmitoyl-CoA on the N-terminal cysteine residue of SHH-N (Porter, Young, and Beachy 1996; Chamoun et al. 2001; Buglino and Resh 2008). These modifications are critical for the signalling activity of the resulting dual-lipidated SHH-N (19 kDa) that represents the mature, biologically active molecule that exits the ER and is subsequently delivered via the secretory pathway to the cell surface (Maity, Fuse, and Beachy 2005; Resh 2021). Given the importance of these post-translational modifications, it is particularly difficult to monitor the sequential steps governing the subcellular transport of neosynthesised SHH using conventional N- and/or C-terminal tagging strategies.

The further processing/delivery of mature SHH requires the activity of several small GTPases. In polarized cells, it is established, that Hh is first introduced to the plasma membrane, and then endocytosed in a clathrin dependent manner to Rab5 positive early endosomes (D'Angelo et al. 2015; Pizette et al. 2021; Gore et al. 2021; Sandoval et al. 2022). Subsequently, SHH is sorted in a Rab8/Rab11 dependent way to its final site of release. Importantly however, these studies were performed in the context of apicobasal secretion, while the secretion of SHH in non-polarized cells remains to be determined. Moreover, the biosynthetic secretory pathway often represents a marginal fraction of the secretory pool at steady-state, convolved with the recycling and endo/exocytic trafficking routes. Hence, it is essential to develop tools to study the biosynthetic secretory pathway independently of the other SHH trafficking routes.

The Retention Using Selective Hook (RUSH) system allows for retention of a given protein in the ER and synchronously release upon biotin addition to follow its trafficking pathway in real time (Boncompain et al. 2012). Here, we adapted the SHH-RUSH system that we previously developed (Thauvin et al. 2022) to study the secretion of SHH from its neo-synthesis in the ER up to the plasma membrane (PM) and secretion into the extracellular media. We found that SHH-RUSH is functional, retaining signalling activity and is transported from the ER to the Golgi up to the plasma membrane within 45 minutes. We highlight that SHH exits the Golgi in vesicles harbouring Rab6, a small GTPase protein required for SHH to reach the PM. Furthermore, we report that TMED10, a member of the p24 family of transmembrane proteins known to interact with COPII machinery, is necessary for efficient ER-to-Golgi transport of SHH, and that silencing of TMED10 leads to decreased SHH secretion. This was confirmed in *Drosophila*, as depletion of *Baiser* (*Bai*; TMED10 homologue), leads to reduced extracellular Hh levels in producing cells, partial inhibition of Hh-dependent imaginal disc outgrowth and affects both short- and long-range Hh activity.

Results

Characterisation of the SHH-RUSH system

110 In order to investigate the SHH biosynthetic secretory pathway, we tagged murine SHH with a streptavidin-binding peptide (SBP) and an mCherry fluorescent protein to produce the SHH-RUSH reporter protein (**Fig. 1A**; Thauvin et al. 2022). The SBP and mCherry tags were introduced internally to preserve the N-terminal palmitate and C-terminal cholesterol modifications in order to track the biologically-active form of SHH (Chamberlain et al. 2008). This modification conditionally traps SHH in the ER in cells expressing Streptavidin-KDEL, a “hook” that retain SBP-expressing proteins in the
115 lumen of the ER. This system allows to synchronise SHH-RUSH release and transport along the secretory pathway upon the addition of biotin (**Fig. 1B**).

To verify that SHH-RUSH still undergoes the necessary processing for SHH secretion, we performed western blotting alongside lysates from the human astroglial cell line SVG-A either mock-transfected or transfected with a plasmid encoding untagged SHH or SHH-RUSH. Immunoblotting revealed little
120 to no degradation of SHH proteins and demonstrates that the SHH-RUSH construct expresses the precursor form of SHH and undergoes processing similar to untagged SHH, while wild-type cells do not express endogenous SHH (**Fig. 1C**). SHH protein secretion from cell culture supernatant by ELISA showed a strong increase upon incubating cells with biotin-rich media (**Fig. 1D**). Of note, SHH was also detected in the supernatant of cells that did not receive biotin treatment, which is likely due to
125 some SHH-RUSH not being retained by the hook. This residual secreted SHH was poorly able to induce *Gli1* RNA expression, suggesting that the leaky forms of SHH might not be fully processed into an active form and/or loaded onto biologically active carriers, such as extracellular vesicles (EVs). Importantly, biotin addition induced secretion of SHH in an active state, able to induce *Gli1* RNA expression in NIH 3T3 fibroblasts almost as efficiently as wild-type non-tagged *Shh* does (**Fig. 1E-F**).
130 Together, this data indicates that the fully matured and secreted SHH-RUSH protein is biologically functional.

To visualise the trafficking pathway of SHH-RUSH, we performed high-resolution confocal imaging on SHH-RUSH cells fixed at different timepoints after biotin addition. Before the addition of biotin, SHH-RUSH displays an ER distribution (**Fig. 1G, upper panel**), at 15 min post-biotin addition, SHH-RUSH is
135 enriched in the Golgi apparatus (**Fig. 1G, middle panel**), and at 45 min post-biotin addition, SHH-RUSH positive vesicles are trafficking towards the cell periphery and localise at the cell surface (**Fig. 1G, lower panel**). We confirmed that the cell population had relatively homogenous synchronization as most cells showed SHH-RUSH in the ER prior biotin addition, while expressing a Golgi-like signal 30 min post-biotin addition (**Supplemental Fig. S1A**). Further confocal analysis by co-staining with

140 markers against the ER (Calnexin), cis-Golgi (GM130), trans-Golgi (TGN46) and the cell periphery (Actin) demonstrated that SHH-RUSH was trafficked through the canonical biosynthetic secretory pathway (**Supplemental Fig. S1B**).

To better understand the trafficking of SHH-RUSH between subcellular compartments, the construct was co-transfected with the medial Golgi marker, Mannosidase-II-EGFP. Approximately 5 min after
145 biotin addition, SHH-RUSH was accumulating at dotted structures within the ER, which likely represent ER exit sites (ERES) (**Fig. 1H + Video S1**). An increase in SHH-RUSH intensity was observed at the Golgi at approximately 15-20 min post biotin addition (**Fig. 1H-I**). At 30 min post-biotin addition, a majority of SHH-RUSH had accumulated at the Golgi, and at 90 min post-biotin addition, SHH-RUSH had vacated the Golgi complex and was observed at the plasma membrane (**Video S2**).

150 Altogether, these results demonstrate that our SHH-RUSH reporter is a suitable system to study the secretion and signalling of neosynthesised active SHH.

SHH-RUSH follows the classical biosynthetic pathway

To further investigate the SHH secretory pathway, we looked at the role of Rab GTPases in SHH-RUSH trafficking to the cell surface, as they play important roles in the biosynthetic pathway. Rab6 is a
155 well-described marker of anterograde and retrograde vesicular transport toward and from the Golgi complex (Goud, Liu, and Storrie 2018). Rab7, a key regulator of late endosome maturation, has recently been shown to be involved as an intermediate compartment of some post-Golgi cargoes (Deffieu et al. 2021). By fixing cells transfected with SHH-RUSH at different timepoints after biotin addition, then performing immunostaining with different Rab GTPases (**Supplemental Fig. S2A**), we
160 observed a significant subset of post-Golgi SHH-RUSH vesicles colocalising with Rab6-positive vesicles at 45 min post-biotin addition (**Fig. 2A-B**). By contrast, Rab5- and Rab7-positive vesicles did not colocalise with SHH-RUSH at this time-point, suggesting that post-Golgi SHH-RUSH is not trafficking through the Rab7 secretory pathway and is predominantly transported post-Golgi through the Rab6 secretory pathway *en route* to the plasma membrane (**Supplemental Fig. S2A**). Rab11 has previously
165 been shown to be involved in secretion of endocytosed SHH at steady-state (Sandoval et al. 2022), but in our assay, investigating specifically neosynthesized SHH trafficking toward the plasma membrane, we could not detect significant enrichment of Rab11 in the SHH-containing vesicles at the investigated time points (**Supplemental Fig. S2B**). Of note, we were unable to investigate Rab8 localization as we lacked good anti-Rab8 antibodies.

170 To define the fate of Rab6-containing post-Golgi SHH-RUSH vesicles, total internal reflection fluorescence (TIRF) microscopy was performed on cells co-transfected with GFP-Rab6a and SHH-RUSH (**Fig. 2C (left panel) and Video S3**). At approximately 30 min post biotin addition, we observed

the arrival of SHH-RUSH in the evanescent TIRF field. We observed that GFP-Rab6a remained associated with SHH-RUSH vesicles until fusion with the plasma membrane (**Fig. 2C (right panels) and Video S4**).

To further support the evidence that SHH-RUSH is transported through Rab6-positive vesicles, we wanted to observe the effects of Rab6 depletion on the SHH secretory pathway. First, we used small interfering RNA (siRNA) against Rab6a or Rab7a, and validated their specificity at the mRNA and protein levels (**Supplemental Fig. S2C-D**). To quantitatively measure the kinetics of SHH-RUSH arriving at the cell surface, we performed a cell surface stain flow cytometry assay as previously described (Deffieu et al. 2021, see Materials & Methods). At specific timepoints following the addition of biotin, cells were incubated on ice to arrest membrane trafficking and SHH-RUSH localised at the surface of non-permeabilised cells was labelled using an anti-SHH antibody (**Fig. 2D**). In cells treated with siCTRL, SHH surface expression peaked at 45 min post-biotin addition. While the profile of Rab7a-depleted cells was similar to siCTRL, a significant decrease in surface SHH fluorescence was observed in siRab6a-treated cells (**Fig. 2E**). Our data indicate that, after exiting the Golgi, SHH is transported in Rab6-harboured vesicles *en route* to the cell surface in mammalian cells, a role that was never reported previously to our knowledge.

TMED10 regulates SHH transport through the biosynthetic pathway

Although we demonstrated that Rab6 is an important player for post-Golgi transport of SHH to the plasma membrane, the molecular mechanism underlying ER-to-Golgi trafficking of SHH remains obscure. The transmembrane emp24 domain (TMED) family of proteins are key players in the regulation of protein trafficking and secretion, exhibiting evolutionary conservation across eukaryotic species (Aber et al. 2019). Following heterodimer formation, TMED proteins oligomerize to form the p24 complex, which contains a single member from each of the four subfamilies (α , β , γ , δ) (Strating and Martens 2009). The p24 complex facilitates the interaction between soluble cargoes in the ER lumen and the cytosolic COPII machinery (Schuiki and Volchuk 2012; D'Arcangelo et al. 2015).

Due to its distribution and function at the ER-Golgi interface (Pastor-Cantizano et al. 2016), we next investigated the potential role of TMED10, also known as Tmp21 or p24 delta (p24 δ), in intracellular SHH trafficking. First, TMED10 protein expression was silenced in SVG-A cells, using siRNA (**Fig. 3A-B**). Using a SHH construct coupled to Nanoluciferase (SHH-NLuc; Petrov et al. 2020), we observed that TMED10-depleted cells secreted significantly less SHH than control cells, although the decrease was modest at steady state (**Fig. 3C**). We observed that the decreased secretion was only observed in TMED10-depleted cells transfected with SHH-NLuc, while other NLuc constructs were not significantly impacted by the silencing of TMED10 (**Supplemental Fig. S3A**).

These experiments were performed at steady state, and the contribution of neosynthesised SHH (and thus TMED10) in this system may be masked by the large amount of SHH that would be already synthesised and recycled. To obtain more information on the role of TMED10 in the intracellular trafficking of neosynthesised SHH, we went back to our SHH-RUSH system and measured the amount of SHH at the cell surface by flow cytometry as in **Fig 2E**. We observed a significant decrease in SHH surface levels in siTMED10-treated cells compared to control cells at early timepoints, while the delay was compensated at later timepoints (**Fig. 3D**). This decrease was even more severe than the one observed for Rab6a, underlying the importance of TMED10 during SHH secretion. As a control, we also showed that TMED10 silencing decreased significantly the secretion of a neosynthesized GPI-anchored protein using the RUSH system (**Supplemental Fig. S4**), consistent with previous report (Tashima et al. 2021). Interestingly, in stark contrast to SHH-RUSH and GPI-RUSH, the trafficking of TfR-RUSH to the cell surface was not perturbed in TMED10-depleted cells (**Fig. 3E**), highlighting that TMED10 is important for the biosynthetic pathway secretion of a specific subset of cargoes.

Next, to better define at which stage TMED10 could intervene in SHH trafficking, we labelled the cis-Golgi (GM130) and the trans-Golgi (TGN46) compartments at different timepoints after biotin addition. Colocalisation analyses revealed that silencing TMED10 led to significant decrease in colocalisation between SHH-RUSH and GM130 at 15 min post biotin addition, and with TGN46 at 45 min post biotin addition compared to control cells (**Fig. 3F-G and Supplemental Fig. S3B-E**), suggesting that SHH-RUSH trafficking is delayed in reaching and passing through the Golgi stacks.

Due to a lack of reliable antibodies against endogenous TMED10, we decided to co-express GFP-tagged TMED10 (Blum and Lepier 2008) alongside SHH-RUSH in SVG-A cells to observe TMED10 dynamics in the context of SHH secretion (**Fig. 3E, left panel and Video S5**). We observed that a large fraction of TMED10 was enriched in a structure reminiscent of the Golgi complex, with a small subset of TMED10 present in distinct puncta reminiscent of ERES. These puncta were also positive for SHH-RUSH. At 60 s post-biotin addition, we observed double-positive puncta (GFP-TMED and SHH-RUSH) trafficking towards the perinuclear region together (**Fig. 3H, right panels and Video S6**).

Thus, our data indicate that TMED10 is an important regulator of SHH biosynthetic secretion and may play a role of ER-to-Golgi trafficking of SHH.

TMED10 associates with SHH-RUSH at ERES and aids in its exit from ER

TMED10 has been shown to play an important role in loading lipidated cargo proteins (such as GPI-anchored proteins and Wnt) in the ER into COPII vesicles *en route* to the cis Golgi (Tashima et al. 2021). As SHH is also a lipidated protein, and lacks the necessary transmembrane motifs to interact with the COPII machinery, we hypothesised that TMED10 could act as an ER cargo receptor for SHH.

To assess whether there was an association between SHH and TMED10, we setup a proximity ligation assay (PLA). Cells were co-transfected with SHH-RUSH and GFP-TMED10. The number of PLA puncta per cell (indicative of SHH and TMED10 proximity) was quantified through confocal imaging and automated image analysis (**Supplemental Fig. S5A**). At 0 min post-biotin addition, we saw no/few PLA puncta per cell, suggesting that SHH-RUSH is not in proximity with GFP-TMED10. However, at 5 min and 10 min post-biotin, we observed PLA puncta, suggesting that at these timepoints, TMED10 and SHH were in close proximity with each other, likely in the same subcellular structures. At 15 min post biotin, fewer PLA puncta were observed, while the majority of SHH-RUSH was localised to the Golgi. These results suggest that TMED10 may associate with SHH at pre-Golgi stages.

Our numerous attempts to co-immunoprecipitate TMED10 and SHH were inconclusive and irreproducible, which prevents us to definitely conclude on the potential interactions between the two proteins. However, the p24 complex is composed of four subfamilies, and the stoichiometry of p24 subfamily members in the p24 complex is key for efficient cargo transport. Hence it is possible that other partners interact more directly with SHH. On this line, we found that TMED2, another subfamily member of the p24 complex, is also required for efficient SHH arrival to the plasma membrane (**Supplemental Fig. S5B**). Moreover, TMED10 knock-down results in the slight upregulation of TMED7 mRNA (**Supplemental Fig. S5C**). These data suggest that the different p24 members are likely cooperating to allow SHH exit from the ER.

As TMED10 interacts with the inner COPII coat protein SEC24D (Lopez et al. 2020), we next investigated the dynamics of SHH-RUSH at ERES using live confocal time-lapse microscopy. Even before the addition of biotin, a significant subset of SHH-RUSH was localised at SEC24D-puncta (**Fig. 4A and associated Video S7**). Fluorescent intensity analysis of this structure containing both SHH-RUSH and GFP-SEC24D revealed that shortly after biotin addition, the intensity of SHH-RUSH dropped whereas the intensity of SEC24D remains the same (**Fig. 4B**), suggesting that SHH exited the ER through an ERES. Of note, given the recent advances on the structure of ERES (Weigel et al. 2021), super-resolution microscopy could help investigating further the sublocation of TMED10 and SHH in ERES.

To investigate the role of TMED10 in the ER exit of SHH-RUSH, SHH-RUSH expressing cells were treated with siCTRL or siTMED10 and immunolabelled with antibodies against SEC31, an outer COPII coat protein known to localise to ERES (Peotter et al. 2019). Before the addition of biotin, SHH-RUSH was distributed to the ER in both siCTRL and siTMED10 treated cells. Colocalisation analysis demonstrated that SHH-RUSH colocalised with SEC31 for a longer period of time in siTMED10 cells compared to siCTRL (**Fig. 4C-D**), suggesting that TMED10 depletion delayed the exit of SHH-RUSH

from ER exit sites. To assess whether TMED10 could help SHH to reach ERES, we measured the fluorescence intensity of SHH-RUSH at SEC31-positive puncta over time in siCTRL and siTMED10 cells and found that indeed, SHH-RUSH intensity at SEC31 puncta was significantly reduced in TMED10-
275 depleted cells compared to control cells (**Fig. 4E**).

Taken together, these results suggest that TMED10 promotes the recruitment of SHH to ERES, thereby favouring its trafficking towards the Golgi and subsequent secretory pathways.

Loss of Baiser function affects Hh secretion and signalling in vivo

Given that TMED10 depletion leads to a delay in neosynthesised SHH trafficking from the ER to the
280 cell surface and reduced SHH secretion from producing cells, we investigated whether similar phenotypes could be observed in an established *in vivo* model such as *Drosophila*.

Hh is known to play a vital role in axial patterning of the epithelial wing imaginal disc cells (WIDs) in *Drosophila* larvae (Briscoe and Théron 2013). The WID is mainly composed of a monolayer of two populations of epithelial cells (columnar cells), posterior cells expressing Hh and anterior cells
285 responding to Hh signalling. Hh induces differential responses in the first proximal 6-8 anterior cells located at the antero-posterior (A/P) border (Tabata and Kornberg 1994; Hatori and Kornberg 2020). This readout, and also the failure to respond, can be directly analysed by measuring the expression level of Hh target genes which give specific response to different levels of Hh concentration: short-range target genes, such as the transcription factor Engrailed (*En*) and the Hh receptor Patched (*Ptc*),
290 and long-range target genes, such as Decapentaplegic (*dpp*), the *Drosophila* homologue of TGF- β (Tanimoto et al. 2000; Torroja, Gorfinkiel, and Guerrero 2005).

First, we used a 'tester line' in which Hh is conditionally overexpressed in the posterior compartment of the WID, leading to distal ectopic expression of the long-range Hh target gene *dpp*, resulting in an outgrowth of the anterior compartment at the opposite site of the source of Hh (Matusek et al.
295 2014) and post-larvae lethality during pupal stage (**Fig. 5A,I-J**). The expression of two different RNAis targeting *bai*, the *Drosophila* homologue of TMED10, in the posterior cells induced a moderate rescue of Hh-induced pupal lethality, as well as *dpp*-dependent anterior outgrowth (**Fig. 5A**). Because the RNAi, which is expressed only in posterior cells, induces a non-cell autonomous effect on Hh-dependent activity in anterior cells, our data suggest that *bai* may play a role in Hh secretion *in vivo*.

300 To verify the role of *bai* in the establishment of Hh gradient activity, we examined the expression levels of Hh target genes in *bai* RNAi treated WIDs expressing a physiological level of Hh (**Fig. 5B-D**). We observed an increased number of cells expressing the high-level Hh target *En* in anterior Hh-receiving cells (3-4 rows, instead of 1-2 in the wild-type discs, **Fig. 5B, E**). We did not detect

significant changes in *Ptc* expression (**Fig. 5C, F**). In contrast, the range of *dpp-lacZ* expression covered
305 a smaller territory in posterior *bai*-depleted discs compared to wild-type (**Fig. 5D, G**). This change was
very apparent in cells close to the A/P border, where, since *En* is a transcriptional repressor for *dpp*,
the increased ectopic *En* expression is known to suppressed *dpp* transcription.

To determine whether *bai* depletion plays a direct role on Hh secretion, we stained extracellular Hh
and actin as a control, and found that within the same animal, the *bai* RNAi region had decreased
310 extracellular Hh compared to non-RNAi treated areas, while actin staining remained similar
regardless of the region observed (**Fig. 5H**). Together, the data suggests that *in vivo* depletion of *bai*
decreases Hh secretion, likely differentially affecting Hh targets, and perturbs the physiological Hh
gradient involved in WID morphogenesis.

Finally, ectopic distal *dpp-lacZ* patterns were observed in the WID of the UAS-Hh tester line together
315 with either control GFP RNAi or with RNAi targeting *bai* or *Dispatched* (*disp*; a strong regulator of Hh
secretion in both flies and mammals (Burke et al. 1999) (**Fig. 5I-J** and **Supplemental Fig. S6A-B**). In
control samples, ectopic *dpp-lacZ* was found at the distal tip of the outgrowing WID at 1- and 4-days
post induction. In contrast, we found that *bai* RNAi results in decreased distal *dpp-lacZ* expression
with a reduced outgrowth area at 4 days post-induction. This phenotype was very reminiscent of the
320 one observed for *disp* RNAi, used here as a positive control. These data suggest that *bai* regulates Hh
and is particularly important for long-range Hh signalling in flies.

Discussion

The present study aimed to provide a better understanding of the complex mechanisms governing
325 the intracellular transport of SHH by using a novel SHH-RUSH reporter system. Our results
demonstrate that the SHH-RUSH system enables efficient transport of SHH through the classical
biosynthetic secretory pathway while maintaining its full signalling capability. This system allowed to
identify two novel players involved in the trafficking of neosynthesised SHH: TMED10 and Rab6.
Moreover, we have uncovered TMED10 as a novel regulator of SHH biosynthetic secretory pathway.
330 We show that TMED10 mediates ER-to-Golgi translocation of SHH by contributing to its loading onto
ERES. Furthermore, TMED10 plays a significant role in Hh/SHH secretion and downstream signalling
in vitro and *in vivo*, highlighting its physiological relevance during embryonic development.

Recently, a suite of sophisticated tools has been developed to elucidate the intricate spatial and
temporal mechanisms underlying the secretory pathway. Early investigations into protein secretion
335 were conducted using the thermosensitive vesicular stomatitis virus glycoprotein (ts045VSV-G)

(Balch et al. 1994). However, to retain physiological conditions and monitor the trafficking of cargoes other than VSV-G, the RUSH system was developed (Boncompain et al. 2012). Here, the cells we used do not express endogenous SHH and thus the CRISPR approach could not be employed. The RUSH system may also perturb intracellular trafficking as the RUSH-tagged protein is released as a single
340 wave, potentially causing an overload of the system. Although this is important to take into account, we investigated SHH-RUSH kinetics in various cell lines (Thauvin *et al.*, 2022 and this study), testing its signalling activity, and confirmed our main findings with non-RUSH approaches, allowing us to further consolidate our conclusions. Nonetheless, it is imperative to exercise caution when utilizing the RUSH system to study the normal trafficking and function of cargoes, taking into account their
345 size and post-translational modifications (Barlowe and Helenius 2016). These findings suggest that our approach would be suitable for future investigations into other aspects of Hh/SHH biology.

We identified Rab6-positive carriers as a mean of transport of neosynthesised SHH from the Golgi apparatus to the cell surface in non-polarised mammalian cells. This finding is consistent with previous reports on the importance of Rab6 for the biosynthetic secretory pathway of other proteins
350 (Fourriere et al. 2019). Rab6 has also been implicated in cargo transport to other subcellular compartments, such as the recycling endosome, late endosomes/lysosomes, and Golgi stacks, highlighting its versatility in intracellular transport (Patwardhan et al. 2017; Dickson, Liu, and Storrie 2020; Huber et al. 2020). However, silencing of Rab6a induces only a mild (although significant) decrease of SHH arrival at the PM, which suggest either that other Rab6 isoforms are participating in
355 SHH secretion, or that other parallel pathways are at play. Future studies should be undertaken to explore the regulatory partners involved in post-Golgi transport of SHH. Such investigations would significantly deepen our understanding of the interwoven dynamics of the intracellular trafficking machinery that governs protein secretion.

In our study, the association of TMED10 with SHH at ERES provides insight into the delayed SHH
360 secretion observed upon TMED10 depletion. Of note however, given the recent advances on the structure of ERES (Weigel et al. 2021), super-resolution microscopy could help investigating further the sublocation of TMED10 and SHH in ERES. We found that the depletion of one p24 family member results in abnormal expression of other members and that TMED2 silencing also decreases SHH secretion. These observations argue in favour of the requirement for a mature p24 complex,
365 composed of the four subfamily members. Yet, only TMED10 was found in SHH-containing EVs (Coulter et al. 2018), suggesting that TMED10 could play additional roles that remain to be characterized.

The p24 family proteins have been extensively studied as cargo receptors for proteins such as GPI-anchored and Wnt proteins in mammals, flies and plants (Mañuel and Howard 2016; Bernat-Silvestre et al. 2020; Lopez et al. 2020). As lipidated cargo proteins have been identified as requiring p24 family proteins for efficient ER-to-Golgi trafficking, we hypothesize that such proteins utilise common mechanisms of ER egress. Recently, SURF4 was identified as another cargo receptor for SHH in the ER (Tang et al. 2022). The SURF4-SHH interaction was found in HeLa and 293T cells, while we used an astrocytic cell line to study SHH secretion as well as *in vivo* models. One can hypothesize that SURF4 and TMED10 are differentially recruited according to the producer cell type investigated, and thus, further work is needed to decipher the interplay of the various SHH partners in the ER. Whether SURF4 and TMED10 are redundant or playing different roles during SHH secretion remains unclear at this stage. SURF4 is a major player in the role of maintaining the architecture of the ERGIC and Golgi apparatus (Mitrovic et al. 2008), and thus, it may be possible that SURF4 has a broad effect on protein secretion. In contrast, TMED10 silencing had no effect on the TfR neosynthetic pathway in mammalian cells, suggesting that TMED10 is cargo-specific. Yet, the molecular determinants governing specificity remain to be identified. Interestingly, TMED10 was recently shown to be involved in SMO secretion, another component of SHH signalling cascade (Di Minin et al. 2022). It would be interesting to determine whether TMED10 plays a role on other components of the hedgehog pathway, as well as on Desert hedgehog and Indian hedgehog secretion. Finally, TMED10 has been involved in various neurodevelopmental defects (Vetrivel et al. 2008; Xie et al. 2014; Qiu et al. 2019; Kim, Scott, and Meador-Woodruff 2019; Shin et al. 2019) and found associated to SHH-containing extracellular vesicles in ART-EVs, a subtype of vesicles found in the cerebrospinal fluid of a glioblastoma patient (Coulter et al. 2018).

In this study, we highlight that *bai*/TMED10 participates in the secretion of Hh/SHH in *Drosophila* and in mammalian cells, and revealed its pivotal role during ER egress. *Bai* has been shown to be required for the secretion of other morphogens such as Wingless, the *Drosophila* homologue of Wnt (Port, Hausmann, and Basler 2011; Li et al. 2015), providing further evidence supporting our hypothesis on the role of TMEDs in selective cargo secretion. Hh is secreted in various forms: a form that is involved in short-range signalling (monitored through *En* staining), and a form more potent for long-range signalling mediated by extracellular vesicles (monitored through *Dpp* staining). We have often observed a tradeoff between the long-range and the short-range Hh pathway targets, in other words that defects in long-range Hh signalling was associated to increased short-range signaling (Ayers et al. 2010; Gore et al. 2021; Pizette et al. 2021). This is observed specifically when trafficking of Hh in its secreting cells is perturbed, suggesting that apical and basolateral trafficking of Hh in the polarized producing cells are not independent. Thus, the increased *En* staining observed may solely result from

a defective apical secretion induced by *bai* RNAi. This trade-off is extremely difficult to investigate because little is known about polarized trafficking in wing imaginal discs.

Intriguingly, the low, yet significant and consistent, impact of *bai* depletion in the WID outgrowth assays closely resembles the effect of TMED10 depletion that we observed on SHH secretion under steady-state conditions. As previously stated, steady-state condition is not an ideal system to decipher the biosynthetic secretory pathway, and thus, to further elucidate the role of *bai* in Hh secretion and downstream signalling, it will be beneficial to employ the RUSH system, which has recently been implemented in *Drosophila* (Glashauer et al. 2023).

In conclusion, our study lays the groundwork for future research on the intricate workings of SHH biology and showcases the potential of the SHH-RUSH reporter system as a potent instrument for delineating the molecular mechanisms underlying the intracellular trafficking and subsequent secretion of SHH. Notably, our discovery of the pivotal role of TMED10 as an important modulator of SHH secretion and its therapeutic potential as a target against abnormal SHH signalling should be further considered.

Methods

Cell culture and transfections

Human brain astroglial cell line SVG-A (Major et al. 1985) was cultured in high glucose Dulbecco's modified Eagle's medium (DMEM-GlutaMAX, Gibco), supplemented with 10% foetal bovine serum (FBS, Dominique Dutscher), 100 U/ml penicillin, and 100 µg/ml streptomycin (Gibco). Transient transfections were performed on 70% confluent cells using JetPRIME (Polyplus) according to the manufacturer's instructions. Briefly, a known concentration of DNA was diluted in JetPRIME buffer in a microcentrifuge tube and vortex. A volume of JetPRIME reagent was added to the solution at a 1:2 DNA/JetPRIME ratio. The tube was briefly vortexed, centrifuged and incubated for 10 min at room temperature (RT). The transfection mix was added dropwise to cells in complete growth media and incubated for 24 – 48 hrs at 37°C and 5% CO₂. For siRNA transfections, single siRNA sequences targeting a gene of interest were purchased from Ambion (ThermoFisher Scientific). 50,000 cells were seeded in 24-well plates, and the next day, 11 pmol of siRNA was transfected using Lipofectamine 3000 (ThermoFisher Scientific) following the manufacturer's instructions. For RUSH experiments, cells were maintained in avidin-rich media (2 µg/ml, Sigma) to prevent leakage of the reporter.

Plasmids

The SHH-RUSH construct was previously described (Thauvin et al. 2022). The GFP-Rab6A cDNA
435 construct cloned into the pBS vector under control of the weak promoter L30 was generated by the
Montpellier Genomic Collection (MGC). The TfR-RUSH construct (pStrepKDEL-TfR-SBP-EGFP) was a
kind gift from Dr. Franck Perez. The Nanoluciferase (NLuc) plasmid (pNL2.1) was purchased from
Promega. SHH-NLuc was a kind gift from Dr. Adrian Salic (Harvard Medical School) (Petrov et al.
2020). Hsp70-NLuc was a kind gift from Dr. Gregory Lavieu (Institut Curie) (Bonsergent et al. 2021).
440 GFP-TMED10 was a kind gift from Dr. Robert Blum (Uni. Wurzburg) (Blum and Lepier 2008).

Antibodies and reagents

For immunofluorescence, primary antibodies used were rabbit anti-Calnexin (1:500; Elabscience),
rabbit anti-Rab6 (1:1000; Cell Signalling Technology), rabbit anti-Rab7 (1:250; Cell Signalling
Technology), mouse anti-GM130 (1:1000; BD Biosciences), sheep anti-TGN46 (1:1000; Bio-Rad),
445 mouse anti-SEC31 (1:250; BD Biosciences), rabbit anti-SHH (1:250; Abcam), mouse anti-GFP (1:250;
Invitrogen), guinea-pig anti-Engrailed (1:100; Pizette et al. 2021), mouse anti-Ptc (1:400; kind gift
from P. Ingham), chicken anti- β -gal (1:1000; GeneTex) and rabbit anti-Hh (1:500; Matusek et al.
2014). Secondary antibodies used were Alexa Fluor 488 donkey anti-mouse PLUS (1:5000;
ThermoFisher Scientific), Alexa Fluor 488 donkey anti-rabbit PLUS (1:5000; ThermoFisher Scientific),
450 Alexa Fluor 647 donkey anti-rabbit PLUS (1:5000; ThermoFisher Scientific), Alexa Fluor 647 donkey
anti-sheep (1:1000; ThermoFisher Scientific), Cy3-conjugated donkey anti-chicken (1:200; Jackson
ImmunoResearch), goat anti-mouse Alexa 488 (1:500; Life Technologies), Cy3-conjugated anti-guinea
pig (1:200; Jackson ImmunoResearch), goat anti-rabbit Cy5 (1:200; Jackson ImmunoResearch), and
Rhodamine phalloidin (1:100; Sigma-Aldrich). Probes used for immunofluorescence were Duolink In
455 Situ PLA Probe donkey anti-rabbit PLUS, donkey anti-mouse MINUS and Detection Reagents Far-Red
(Sigma-Aldrich). DAPI (4,6-diamidino-2-phenylindole) (1:1000; Sigma-Aldrich) was used to stain the
nucleus. For flow cytometry, rabbit anti-SHH antibody (1:1000; Abcam) and Alexa Fluor 647 donkey
anti-rabbit PLUS (1:5000; ThermoFisher Scientific) were used. Antibodies used for immunoblotting
were rabbit anti-Rab6 (1:1000; Cell Signalling Technology), rabbit anti-Rab7 (1:1000; GeneTex), rabbit
460 anti-TMED10 (1:1000; Sigma-Aldrich), mouse anti-GAPDH (1:5000, GeneTex), mouse anti-GFP
(1:1000, Invitrogen), goat anti-SHH (1:1000, Santa Cruz) and rabbit anti-SHH (1:1000;
Abcam). Secondary antibodies used for immunoblotting were goat anti-mouse immunoglobulin G
(IgG) horseradish peroxidase (HRP) antibody (1:10,000; Jackson ImmunoResearch), goat anti-rabbit
IgG HRP antibody (1:10,000; Jackson ImmunoResearch) and donkey anti-goat IgG HRP antibody
465 (1:10,000; Invitrogen).

Retention using selective hook (RUSH) procedure

Cells were cultured with 2 µg/ml avidin (Sigma-Aldrich) 24 h prior experiments started to quench endogenous biotin from the media and/or produced by the cells. To release RUSH constructs, media was replaced with complete medium containing 40 µM biotin (Sigma-Aldrich) and incubated for the
470 indicated amount of time at 37°C and 5% CO₂. Cells were then placed on ice, and media was replaced with ice-cold phosphate-buffered saline (PBS) and cells were maintained on ice for 10 min.

Cell surface staining and flow cytometry

To measure the amount of protein at the cell surface, cells were incubated for 1 hr with a primary antibody, washed three times with ice-cold PBS then incubated for 30 min with a secondary antibody
475 coupled to Alexa Fluor 647 (abbreviated as AF-647 on the corresponding figures). Cells were then washed twice with cold PBS and detached with pre-warmed 0.5 mM EDTA solution. Cells were collected and centrifuged at 500 x *g* for 5 min at 4°C. Cell fixation was performed using 4% PFA for 15 min at RT, and after two washes, they were resuspended in FACS buffer (PBS, 0.5% BSA (w/v), 0.5 mM EDTA). Samples were run on a NovoCyte (ACEA Biosciences) flow cytometer equipped with 561-
480 and 640-nm lasers and four-filter set. Data analysis was performed using FlowJo software (FlowJo LLC, v10.8.1).

Nanoluciferase secretion assay

SVG-A cells were seeded in 6-well plates and transfected with plasmids encoding NLuc-tagged proteins. At 16-24 hrs post-transfection, cells were washed with PBS and media was replaced with
485 serum-free DMEM. After 24 hrs, cell culture supernatant was collected and centrifuged at 10,000 x *g* for 5 min. Cells were also trypsinised and pelleted at 500 x *g* for 3 min before lysis using NanoGlo Luciferase Assay Reagent (Promega). NLuc activity was measured on an Infinite200 PRO Plate Reader (TECAN) using the NanoGlo Luciferase Assay System kit (Promega) according to the manufacturer's instructions.

Immunofluorescence and proximity ligation assay

Cells were seeded onto 12 mm glass coverslips (EMS) in 24-well plates and cultured overnight in complete medium. For RUSH assays, cells were cultured in complete medium supplemented with avidin (2 µg/ml). At 16-24 hrs post-transfection, cells were incubated at indicated timepoints in complete medium containing 40 µM biotin. Cells were washed twice with 1X PBS and then fixed in
495 4% PFA for 15 min at RT. Cells were then simultaneously permeabilised and blocked for 15 min at RT with PBS containing 0.1 % Triton X-100 (v/v) (Sigma-Aldrich) and 0.5 % BSA (w/v) (Euromedex). Cells were subsequently incubated with primary antibodies diluted in perm/block buffer for 2 hr at RT in a humidified chamber. For immunofluorescence, cells were incubated for 1 hr at RT in a humidified

chamber with the corresponding secondary antibodies and DAPI diluted in the same buffer as the
500 primary antibodies. Secondary antibodies were washed three times for 5 min in PBS at RT and
coverslips were then mounted onto glass slides using Mowiol 4-88 (Sigma-Aldrich). For PLA, after
incubation with primary antibodies, cells were washed twice for 5 min with 1X Wash Buffer A at RT.
Cells were then incubated for 1 hr at 37°C in a pre-heated humidity chamber with Sigma PLUS and
MINUS PLA probes in the 1X Sigma dilution solution. Sigma probes were washed twice for 5 min with
505 1X Wash Buffer A. The ligation step was performed at 37°C for 30 min followed by two washes with
1X Wash Buffer A. Amplification was performed for 100 min at 37°C in dark conditions. Cells were
then washed twice at RT for 10 min each with 1X Wash Buffer B and finally for 1 min in 0.01X Wash
Buffer B. Coverslips were then mounted onto glass slides using Mowiol 4-88 (Sigma-Aldrich).

Image acquisition

510 For fixed imaging, confocal microscopy was performed using an Eclipse Ti inverted microscope
(Nikon) mounted with a CSU-X1 spinning disk head (Yokogawa), a back-illuminated EMCCD Andor
camera (Oxford Instruments), and a 100X (1.45 NA) oil objective lens controlled by the Andor iQ3
software (Oxford Instruments).

For live cell imaging, cells were seeded onto 25 mm glass coverslips (EMS) and transfected using
515 JetPRIME (Polyplus) according to the manufacturer's instructions. Coverslips with transfected cells
were then transferred into an Attofluor Cell Chamber (Invitrogen) and placed onto the microscope
stage and maintained in a dark atmosphere-controlled chamber at 37°C and 5% CO₂. Live-cell
confocal imaging was performed using either an inverted IX83 microscope (Olympus) coupled to a
CSU-W1 spinning disk unit (Yokogawa), a sCMOS Fusion camera (Hamamatsu) and a 60X (1.3 NA) oil
520 objective lens controlled by the CellSens software (Olympus), or an Eclipse Ti2 inverted microscope
(Nikon) coupled to the Andor Dragonfly spinning disk system (Oxford Instruments), an EMCCD Andor
camera (Oxford Instruments), and a 100X APO (1.45 NA) oil objective lens controlled by Andor Fusion
software (Oxford Instruments). For TIRF microscopy, live imaging was performed with an Eclipse Ti
inverted microscope (Nikon), a back-illuminated EMCCD camera (Evolve 512, Photometrics) and a
525 100X APO (1.49 NA) oil objective lens controlled by MetaMorph. The TIRF angle was chosen to obtain
a calculated evanescent field depth of <100 nm. Acquisition was performed from 30 min after biotin
addition.

Image analyses

Image processing was performed using either the Fiji upgrade of ImageJ or Imaris software v9.2
530 (Bitplane, Oxford Instruments). Quantifications for mean fluorescence intensities and colocalisation
measurements were performed using Imaris software v9.2 (Bitplane, Oxford Instruments). For

colocalisation analysis of the different Rab GTPases, cells were stained with a Golgi marker (Mannosidase-II-EGFP) and the Golgi staining was masked prior colocalization measurements to assess colocalization in non-Golgi area. The fluorescence intensities of SHH and given Rab proteins were then measured within each manually delimited cell.

Drosophila stocks and experimental conditions

Flies were cultured on standard *Drosophila* medium. In rescue experiments of Hh-dependent wing imaginal disc outgrowth and lethality, “tester line” virgins with the genotype of w¹¹¹⁸; tubGal80ts; hhGal4<UAS-Hh(M4)/TM6b-Tb (Matusek et al. 2014) were crossed to control w¹¹¹⁸ and *bai* RNAi lines (UAS-*bai*330020SH: Vienna *Drosophila* Resource Center, VDRC#330020, UAS-baiser100612KK: Vienna *Drosophila* Resource Center, VDRC#100612). Parents were transferred to new vials every 12 hours at 25°C. This allowed sufficient number of eggs to be laid without considerable overcrowding. Early 2nd instar larvae were shifted to 29°C to inactivate tubGal80ts thus inducing UAS-Hh(M4) expression and the UAS transgenes tested in the system. Following 3 days of incubation at 29°C, wandering 3rd instar larvae were dissected and their WIDs were scored into categories (with wild-type like, moderately outgrown, fully outgrown morphologies). To assess lethality rescue, in parallel, vials with the same progeny were kept at 25°C till hatching, and then scored for pharate adult viability (empty pupae). In 50% of the progeny Tb animals (represented by empty small circles on Figure 5A) do not contain the hhGal4<UASHh (M4) recombinant chromosome, expected to be viable, therefore serve as an internal control. Non-Tb animals carry the hhGal4<UASHh (M4) recombinant chromosome and either a neutral GFP RNAi (VDRC#60102) or *bai* RNAi-s. To overexpress *bai* RNAi in wild-type background, UAS-*bai*330020SH UAS-*bai*100612KK and were crossed to tubGal80ts<dpp-lacZ; hhGal4/TM6b-Tb virgins. As a control we crossed tubGal80ts<dpp-lacZ; hhGal4/TM6b-Tb virgins to w¹¹¹⁸ as well. Egg laying was allowed at 25°C and vials were then shifted up to 29°C until dissection.

Immunostaining and imaging of WIDs

Conventional and extracellular immunolabeling was performed as previously described (Matusek et al. 2014; D’Angelo et al. 2015). For extracellular Hh labelling without detergent but with initial fixation, discs were dissected in ice-cold PBS, and then fixed in 2% PFA (v/v) in PBS for 15 min at RT. After fixation, discs were rinsed in PBS three times for 5 min at RT, and then incubated with anti-Hh antibody for 1 hr in 2% BSA (w/v) in PBS. Discs were then rinsed three times in 0.3% Triton-X100 (v/v) in PBS and blocked again for an hour in 0.1% Triton X-100 (v/v), 1% BSA (w/v) in PBS. Following this step, anti-Ptc was added in the detergent-containing block solution and the samples were incubated at 4°C overnight. The next day, samples were washed three times for 10 minutes in 0.3% Triton-X100

565 (v/v) in PBS before incubation with secondary antibodies and phalloidin. For imaging, wing imaginal discs were mounted in VECTASHIELD mounting medium (Vector Laboratories). All confocal images were recorded with Leica TCS SP5 confocal laser scanning microscope. dpp-lacZ intensity was measured on single slices on images recorded in 1024x1024 resolution, with 40x objective (HCX PLAN APO; NA=1.3) at 1,3x zoom level, with 400hz speed, within a 200x100 pixel rectangle. For En and Ptc 570 range measurements, the same imaging conditions, and a 100x100 pixel square was used. Optical Z sections within the posterior compartment along the dorsoventral axis were taken at 1µm thickness in 1024x1024 resolution, with 40x objective (HCX PLAN APO; NA=1.3) at 3.52x zoom level, with 400hz speed. Images presented in the manuscript are a stack of 5 single sections. To image whole discs single confocal sections were collected with 20x objective (HCX PL APO CS 20.0x0.70 IMM UV 575 NA=0.7) without zoom at 1024x1024 resolution with 400Hz speed. To analyze ectopic dpp-lacZ patterns, expression of UAS-Hh (tester line) together with either control UAS-GFP RNAi (VDRC #60102), UAS-*disp* RNAi (Bloomington Drosophila Stock Center, BDSC#27247) or with UAS-*bai* RNAi (VDRC #100612) was induced for either 1 or 4 days.

Western blotting

580 Cells were seeded in 6-well plates in complete growth media. After 16-24 hrs post-transfection, cells were washed twice with ice-cold PBS and lysed with NP40 buffer [150 mM NaCl, 1% NP-40, 50 mM Tris (pH 8.0), and protease inhibitor (Promega)]. Cells were rotated on a carousel for 10 min at 4°C then centrifuged at 15,000 x *g* for 20 min at 4°C. The supernatant was collected and protein concentration was determined using the Pierce BCA Assay kit (ThermoFisher Scientific). A total of 20 585 µg of protein lysates was run on Bolt 4-12% Bis-Tris Mini Protein gels (Invitrogen) and proteins were transferred onto nitrocellulose membranes using the iBlot2 Transfer system (Invitrogen) according to the manufacturer's instructions. Membranes were blocked with 5% low-fat milk (w/v) in TBS-T [Tris-buffered saline (TBS) at pH 7.4 and 0.05% Tween20] for 30 min at RT prior to incubation with primary antibodies overnight at 4°C in TBS-T containing 0.5% milk. Secondary antibodies were incubated for 1 590 hr at RT in TBS-T containing 0.5% milk. After several washes with TBS-T, membranes were revealed by chemiluminescence using Clarity Western ECL Blotting Substrates (Bio-Rad) and images were acquired using the ChemiDoc Touch System (Bio-Rad). Relative intensities of western blot bands were quantified using ImageLab software (Bio-Rad, v6.0.1) using Volume Tools, global background subtraction and linear regression method.

595 **SHH-ELISA**

SVG-A cells transfected with SHH-RUSH were incubated in complete growth medium supplemented with 40 μ M biotin at indicated timepoints. The media was collected and centrifuged at 10,000 x g for 5 min and used in the SHH-ELISA kit (R&D Systems) according to the manufacturer's instructions.

SHH signalling and RT-qPCR

600 NIH 3T3 cells were seeded into 48-well plates and cultured in complete growth media until confluent. Cells were then serum-starved for 24 hrs then incubated with cell culture supernatants taken from SVG-A cells expressing SHH or SHH-RUSH in absence or presence of biotin. After 48 hrs, cells were lysed and prepared for RT-qPCR analysis. For all RT-qPCR experiments, cells were lysed and RNA immediately extracted using the Luna Cell Ready Lysis Module (New England Biolabs) according to
605 the manufacturer's instructions. RT-qPCR on extracted RNA was performed using the Luna One-Step RT-qPCR kit (New England Biolabs) and samples were analysed using a LightCycler 480 instrument (Roche) using the following thermal cycle: reverse transcription step at 55°C for 15 min, initial denaturation at 95°C for 2 min; then, 40 cycles were programmed consisting of 15 s of denaturation at 95°C followed by 30 s of annealing/extension at 60°C. Fluorescent data collection was performed
610 at the end of each cycle.

Oligonucleotides	Sequence (5' to 3')
Gli1 qPCR F (Ms) a	CCTTTCTTTGAGGTTGGGATGA
Gli1 qPCR R (Ms) a	TTGGATTGAACATGGCGTCT
Gli1 qPCR F (Ms) b	CTTCAAGGCCCAATACATGC
Gli1 qPCR R (Ms) b	GTGAATAGGACTTCCGACAGC
Gapdh qPCR F (Ms)	AATGGTGAAGGTCGGTGTG
Gapdh qPCR R (Ms)	GTGGAGTCATACTGGAACATGTAG
RAB6A qPCR F	CGTTTCCGTAGCCTCATTC
RAB6A qPCR R	ACATCACTTCTCTTTCTGTTC
RAB7A qPCR F	CTCACACACACACACACA
RAB7A qPCR R	GGGAAAGGGGGGGTAAAAAG
TMED10 qPCR F	GAGATGCGTGATACCAACGA
TMED10 qPCR R	TTCTTGGCCTTGAAGAAGCG
TMED9 qPCR F	GCGCTCTACTTTCACATCGG
TMED9 qPCR R	CACCTCCACAAACATGCCAA
TMED7 qPCR F	TTGGAGAAGACCCACCTTTG

TMED7 qPCR R	GCCCTATGCTAACCACCAGA
TMED2 qPCR F	GTAAAGCACGAACAGGAATACATG
TMED2 qPCR R	GTCATGGCAACTAGAACAAGAGC
GAPDH qPCR F	GAGTCAACGGATTTGGTCGT
GAPDH qPCR R	TTGATTTTGGAGGGATCTCG

Statistical analysis

All statistical analyses were performed using GraphPad Prism (v9.5.1). Student's t-test was performed for comparisons between two groups of data that seemed to be normally distributed with similar variances. Multiple group comparisons were performed using one-way ANOVA.

Acknowledgments

We acknowledge the MRI imaging facility, member of the national infrastructure France-BioImaging, for advice and training. We thank Drs. Adrian Salic, Franck Perez, Gregory Lavieu and Robert Blum for the gift of constructs used in this study. **Funding:** This work was supported by the Agence National de la Recherche (ANR-18-CE13-0003-01) to RG and PT. YB obtained a post-doctoral fellowship from Fondation pour la Recherche Medicale (FRM, SPF202110014043). **Author contributions:** Conceptualisation: YB, RG. Methodology: YB, TM, SV, PT, MSD, RG. Investigation: YB, TM, PT, RG. Analysis: YB, TM, PT, RG. Supervision: PT, RG. Writing – original draft: YB, RG. Writing – review & editing: YB, TM, SV, PT, MSD, RG. **Competing interest:** The authors declare that they have no competing interests.

Data and materials availability

All data are available in the main text, the supplementary materials or will be made available on reasonable request.

References

- Aber, Rachel, Wesley Chan, Sevane Mugisha, and Loydie A. Jerome-Majewska. 2019. 'Transmembrane Emp24 Domain Proteins in Development and Disease.' *Genetics Research* 101: e14. <https://doi.org/10.1017/S0016672319000090>.
- Álvarez-Buylla, Arturo, and Rebecca A. Ihrie. 2014. 'Sonic Hedgehog Signaling in the Postnatal Brain'. *Seminars in Cell & Developmental Biology* 33:105-11. <https://doi.org/10.1016/j.semcdb.2014.05.008>.

- 640 Ayers, Katie L., Armel Gallet, Laurence Staccini-Lavenant, and Pascal P. Thérond. 2010. 'The Long-Range Activity of Hedgehog Is Regulated in the Apical Extracellular Space by the Glypican Dally and the Hydrolase Notum'. *Developmental Cell* 18 (4): 605–20. <https://doi.org/10.1016/j.devcel.2010.02.015>.
- 645 Balch, William E., J. Michael McCaffery, Helen Plutner, and Marilyn Gist Farquhar. 1994. 'Vesicular Stomatitis Virus Glycoprotein Is Sorted and Concentrated during Export from the Endoplasmic Reticulum'. *Cell* 76 (5): 841–52. [https://doi.org/10.1016/0092-8674\(94\)90359-X](https://doi.org/10.1016/0092-8674(94)90359-X).
- Barlowe, Charles, and Ari Helenius. 2016. 'Cargo Capture and Bulk Flow in the Early Secretory Pathway'. *Annual Review of Cell and Developmental Biology* 32: 197–222. <https://doi.org/10.1146/annurev-cellbio-111315-125016>.
- 650 Bernat-Silvestre, César, Vanessa De Sousa Vieira, Judit Sanchez-Simarro, Noelia Pastor-Cantizano, Chris Hawes, María Jesús Marcote, and Fernando Aniento. 2020. 'P24 Family Proteins Are Involved in Transport to the Plasma Membrane of GPI-Anchored Proteins in Plants'. *Plant Physiology* 184 (3): 1333–47. <https://doi.org/10.1104/pp.20.00880>.
- Blum, Robert, and Alexandra Lepier. 2008. 'The Luminal Domain of P23 (Tmp21) Plays a Critical Role in P23 Cell Surface Trafficking'. *Traffic* 9 (9): 1530–50. <https://doi.org/10.1111/j.1600-0854.2008.00784.x>.
- 655 Boncompain, Gaele, Severine Divoux, Nelly Gareil, Helene de Forges, Aurianne Lescure, Lynda Latreche, Valentina Mercanti, Florence Jollivet, Graça Raposo, and Franck Perez. 2012. 'Synchronization of Secretory Protein Traffic in Populations of Cells'. *Nature Methods* 9 (5): 493–98. <https://doi.org/10.1038/nmeth.1928>.
- 660 Bonsergent, Emeline, Eleonora Grisard, Julian Buchrieser, Olivier Schwartz, Clotilde Théry, and Grégory Lavieau. 2021. 'Quantitative Characterization of Extracellular Vesicle Uptake and Content Delivery within Mammalian Cells.' *Nature Communications* 12 (1): 1864. <https://doi.org/10.1038/s41467-021-22126-y>.
- 665 Borchers, Ann-Christin, Lars Langemeyer, and Christian Ungermann. 2021. 'Who's in Control? Principles of Rab GTPase Activation in Endolysosomal Membrane Trafficking and Beyond'. *Journal of Cell Biology* 220 (9): e202105120. <https://doi.org/10.1083/jcb.202105120>.
- Brady, Melanie V., and Flora M. Vaccarino. 2021. 'Role of SHH in Patterning Human Pluripotent Cells towards Ventral Forebrain Fates'. *Cells* 10 (4). <https://doi.org/10.3390/cells10040914>.
- 670 Briscoe, James, and Pascal P. Thérond. 2013. 'The Mechanisms of Hedgehog Signalling and Its Roles in Development and Disease'. *Nature Reviews Molecular Cell Biology* 14 (7): 418–31. <https://doi.org/10.1038/nrm3598>.
- Buglino, John A., and Marilyn D. Resh. 2008. 'Hhat Is a Palmitoyltransferase with Specificity for N-Palmitoylation of Sonic Hedgehog'. *Journal of Biological Chemistry* 283 (32): 22076–88. <https://doi.org/10.1074/jbc.M803901200>.
- 675 Burke, Richard, Denise Nellen, Manolo Bellotto, Ernst Hafen, Kirsten-André Senti, Barry J. Dickson, and Konrad Basler. 1999. 'Dispatched, a Novel Sterol-Sensing Domain Protein Dedicated to the Release of Cholesterol-Modified Hedgehog from Signaling Cells'. *Cell* 99 (7): 803–15. [https://doi.org/10.1016/S0092-8674\(00\)81677-3](https://doi.org/10.1016/S0092-8674(00)81677-3).
- 680 Chamberlain, Chester E., Juhee Jeong, Chaoshe Guo, Benjamin L. Allen, and Andrew P. McMahon. 2008. 'Notochord-Derived Shh Concentrates in Close Association with the Apically Positioned Basal Body in Neural Target Cells and Forms a Dynamic Gradient during Neural Patterning'. *Development* 135 (6): 1097–1106. <https://doi.org/10.1242/dev.013086>.
- 685 Chamoun, Zeina, Randall K. Mann, Denise Nellen, Doris P. von Kessler, Manolo Bellotto, Philip A. Beachy, and Konrad Basler. 2001. 'Skinny Hedgehog, an Acyltransferase Required for Palmitoylation and Activity of the Hedgehog Signal'. *Science* 293 (5537): 2080–84. <https://doi.org/10.1126/science.1064437>.
- Corrales, Jo Michelle D., Sandra Blaess, Eamonn M. Mahoney, and Alexandra L. Joyner. 2006. 'The Level of Sonic Hedgehog Signaling Regulates the Complexity of Cerebellar Foliation'. *Development* 133 (9): 1811–21. <https://doi.org/10.1242/dev.02351>.

- 690 Coulter, Michael E., Cristina M. Dorobantu, Gerrald A. Lodewijk, François Delalande, Sarah Cianferani, Vijay S. Ganesh, Richard S. Smith, et al. 2018. 'The ESCRT-III Protein CHMP1A Mediates Secretion of Sonic Hedgehog on a Distinctive Subtype of Extracellular Vesicles'. *Cell Reports* 24 (4): 973-986.e8. <https://doi.org/10.1016/j.celrep.2018.06.100>.
- 695 Dahmane, Nadia, and A Ruiz i Altaba. 1999. 'Sonic Hedgehog Regulates the Growth and Patterning of the Cerebellum.' *Development (Cambridge, England)* 126 (14): 3089–3100.
- D'Angelo, Gisela, Tamás Matussek, Sandrine Pizette, Pascal P. Théron, Gisela D'Angelo, Tamás Matussek, Sandrine Pizette, et al. 2015. 'Endocytosis of Hedgehog through Dispatched Regulates Long-Range Signaling'. *Developmental Cell* 32 (3): 290–303. <https://doi.org/10.1016/j.devcel.2014.12.004>.
- 700 D'Arcangelo, Jennifer G., Jonathan Crissman, Silvere Pagant, Alenka Čopič, Catherine F. Latham, Erik L. Snapp, and Elizabeth A. Miller. 2015. 'Traffic of P24 Proteins and COPII Coat Composition Mutually Influence Membrane Scaffolding'. *Current Biology* 25 (10): 1296–1305. <https://doi.org/10.1016/j.cub.2015.03.029>.
- 705 Deffieu, Maika S., Ieva Cesonyte, François Delalande, Gaëlle Boncompain, Cristina Dorobantu, Eli Song, Vincent Lucansky, et al. 2021. 'Rab7-Harboring Vesicles Are Carriers of the Transferrin Receptor through the Biosynthetic Secretory Pathway'. *Science Advances* 7 (2): eaba7803. <https://doi.org/10.1126/sciadv.aba7803>.
- 710 Di Minin, Giulio, Markus Holzner, Alice Grison, Charles E. Dumeau, Wesley Chan, Asun Monfort, Loydie A. Jerome-Majewska, Henk Roelink, and Anton Wutz. 2022. 'TMED2 Binding Restricts SMO to the ER and Golgi Compartments'. *PLoS Biology* 20 (3). <https://doi.org/10.1371/journal.pbio.3001596>.
- Dickson, Lindsey James, Shijie Liu, and Brian Storrie. 2020. 'Rab6 Is Required for Rapid, Cisternal-Specific, Intra-Golgi Cargo Transport'. *Scientific Reports* 10 (1): 16604. <https://doi.org/10.1038/s41598-020-73276-w>.
- 715 Fourriere, Lou, Amal Kasri, Nelly Gareil, Sabine Bardin, Hugo Bousquet, David Pereira, Franck Perez, Bruno Goud, Gaëlle Boncompain, and Stéphanie Miserey-Lenkei. 2019. 'RAB6 and Microtubules Restrict Protein Secretion to Focal Adhesions'. *Journal of Cell Biology* 218 (7): 2215–31. <https://doi.org/10.1083/jcb.201805002>.
- 720 Glashauser, Jade, Carolina Camelo, Manuel Hollmann, Wilko Backer, Thea Jacobs, Jone Isasti Sanchez, Raphael Schleutker, et al. 2023. 'Acute Manipulation and Real-Time Visualization of Membrane Trafficking and Exocytosis in *Drosophila*'. *bioRxiv*. <https://doi.org/10.1101/2022.03.25.483021>.
- 725 Gore, Tanvi, Tamás Matussek, Gisela D'Angelo, Cécile Giordano, Thomas Tognacci, Laurence Lavenant-Staccini, Catherine Rabouille, and Pascal P. Théron. 2021. 'The GTPase Rab8 Differentially Controls the Long- and Short-Range Activity of the Hedgehog Morphogen Gradient by Regulating Hedgehog Apico-Basal Distribution'. *Development* 148 (5): dev191791. <https://doi.org/10.1242/dev.191791>.
- 730 Goud, Bruno, Shijie Liu, and Brian Storrie. 2018. 'Rab Proteins as Major Determinants of the Golgi Complex Structure'. *Small GTPases* 9 (1–2): 66–75. <https://doi.org/10.1080/21541248.2017.1384087>.
- Groves, Ian, Marysia Placzek, and Alexander G. Fletcher. 2020. 'Of Mitogens and Morphogens: Modelling Sonic Hedgehog Mechanisms in Vertebrate Development: Modelling Shh Mechanisms in Development'. *Philosophical Transactions of the Royal Society B: Biological Sciences* 375 (1809). <https://doi.org/10.1098/rstb.2019.0660>.
- 735 Hatori, Ryo, and Thomas B. Kornberg. 2020. 'Hedgehog Produced by the *Drosophila* Wing Imaginal Disc Induces Distinct Responses in Three Target Tissues'. *Development (Cambridge, England)* 147 (22): dev195974. <https://doi.org/10.1242/dev.195974>.
- 740 Hill, Steven A, Andrew S Blaeser, Austin A Coley, Yajun Xie, Katherine A Shepard, Corey C Harwell, Wen-Jun Gao, and A Denise R Garcia. 2019. 'Sonic Hedgehog Signaling in Astrocytes Mediates Cell Type-Specific Synaptic Organization'. *ELife* 8: 1–23. <https://doi.org/10.7554/elife.45545>.

- Huber, Megan E., Regina Kurapova, Chelsea M. Heisler, Elham Karamooz, Fikadu G. Tafesse, and Melanie J. Harriff. 2020. 'Rab6 Regulates Recycling and Retrograde Trafficking of MR1 Molecules'. *Scientific Reports* 10 (1): 20778. <https://doi.org/10.1038/s41598-020-77563-4>.
- 745 Hutchings, Joshua, and Giulia Zanetti. 2019. 'Coat Flexibility in the Secretory Pathway: A Role in Transport of Bulky Cargoes'. *Current Opinion in Cell Biology* 59 (August): 104–11. <https://doi.org/10.1016/j.ceb.2019.04.002>.
- Kim, Pitna, Madeline R. Scott, and James H. Meador-Woodruff. 2019. 'Abnormal ER Quality Control of Neural GPI-Anchored Proteins via Dysfunction in ER Export Processing in the Frontal Cortex of Elderly Subjects with Schizophrenia'. *Translational Psychiatry* 9 (1): 1–9. <https://doi.org/10.1038/s41398-018-0359-4>.
- 750 Li, Xue, Yihui Wu, Chenghao Shen, Tatyana Y. Belenkaya, Lorraine Ray, and Xinhua Lin. 2015. 'Drosophila P24 and Sec22 Regulate Wingless Trafficking in the Early Secretory Pathway'. *Biochemical and Biophysical Research Communications* 463 (4): 483–89. <https://doi.org/10.1016/j.bbrc.2015.04.151>.
- 755 Lopez, Sergio, Ana Maria Perez-Linero, Javier Manzano-Lopez, Susana Sabido-Bozo, Alejandro Cortes-Gomez, Sofia Rodriguez-Gallardo, Auxiliadora Aguilera-Romero, Veit Goder, and Manuel Muñiz. 2020. 'Dual Independent Roles of the P24 Complex in Selectivity of Secretory Cargo Export from the Endoplasmic Reticulum'. *Cells* 9 (5): 1295. <https://doi.org/10.3390/cells9051295>.
- 760 Maity, Tapan, Naoyuki Fuse, and Philip A. Beachy. 2005. 'Molecular Mechanisms of Sonic Hedgehog Mutant Effects in Holoprosencephaly'. *Proceedings of the National Academy of Sciences* 102 (47): 17026–31. <https://doi.org/10.1073/pnas.0507848102>.
- 765 Major, E O, A E Miller, P Mourrain, R G Traub, E de Widt, and J Sever. 1985. 'Establishment of a Line of Human Fetal Glial Cells That Supports JC Virus Multiplication.' *Proceedings of the National Academy of Sciences* 82 (4): 1257–61. <https://doi.org/10.1073/pnas.82.4.1257>.
- Mañuel, Muniz, and Riezman Howard. 2016. 'Trafficking of Glycosylphosphatidylinositol Anchored Proteins from the Endoplasmic Reticulum to the Cell Surface'. *Journal of Lipid Research* 57 (3): 352–60. <https://doi.org/10.1194/jlr.R062760>.
- 770 Matusek, Tamás, Franz Wendler, Sophie Polès, Sandrine Pizette, Gisela D'Angelo, Maximilian Fürthauer, and Pascal P. Therond. 2014. 'The ESCRT Machinery Regulates the Secretion and Long-Range Activity of Hedgehog'. *Nature* 516 (729): 99–103. <https://doi.org/10.1038/nature13847>.
- McClelland, Kathryn, Weili Li, and Norman D. Rosenblum. 2022. 'Pallister-Hall Syndrome, GLI3, and Kidney Malformation'. *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* 190 (3): 264–78. <https://doi.org/10.1002/ajmg.c.31999>.
- 775 Melamed, Jilian R., Joshua T. Morgan, Stephen A. Ioele, Jason P. Gleghorn, Jennifer Sims-Mourtada, and Emily S. Day. 2018. 'Investigating the Role of Hedgehog/GLI1 Signaling in Glioblastoma Cell Response to Temozolomide'. *Oncotarget* 9 (43): 27000–15. <https://doi.org/10.18632/oncotarget.25467>.
- 780 Mitrovic, Sandra, Houchaima Ben-Tekaya, Eva Koegler, Jean Gruenberg, and Hans-Peter Hauri. 2008. 'The Cargo Receptors Surf4, Endoplasmic Reticulum-Golgi Intermediate Compartment (ERGIC)-53, and P25 Are Required to Maintain the Architecture of ERGIC and Golgi'. Edited by Benjamin Glick. *Molecular Biology of the Cell* 19 (5): 1976–90. <https://doi.org/10.1091/mbc.e07-10-0989>.
- 785 Palma, Verónica, Daniel A. Lim, Nadia Dahmane, Pilar Sánchez, Thomas C. Brionne, Claudia D. Herzberg, Yorick Gitton, Alan Carleton, Arturo Álvarez-Buylla, and Ariel Ruiz i Altaba. 2005. 'Sonic Hedgehog Controls Stem Cell Behavior in the Postnatal and Adult Brain'. *Development* 132 (2): 335–44. <https://doi.org/10.1242/dev.01567>.
- 790 Pastor-Cantizano, Noelia, Juan Carlos Montesinos, César Bernat-Silvestre, María Jesús Marcote, and Fernando Aniento. 2016. 'P24 Family Proteins: Key Players in the Regulation of Trafficking Along the Secretory Pathway'. *Protoplasma* 253 (4): 967–85. <https://doi.org/10.1007/s00709-015-0858-6>.

- 795 Patwardhan, Anand, Sabine Bardin, Stéphanie Miserey-Lenkei, Lionel Larue, Bruno Goud, Gracá Raposo, and Cédric Delevoye. 2017. 'Routing of the RAB6 Secretory Pathway towards the Lysosome Related Organelle of Melanocytes'. *Nature Communications* 8. <https://doi.org/10.1038/ncomms15835>.
- 800 Peotter, Jennifer, William Kasberg, Iryna Pustova, and Anjon Audhya. 2019. 'COPII-Mediated Trafficking at the ER/ERGIC Interface'. *Traffic* 20 (7): 491–503. <https://doi.org/10.1111/tra.12654>.
- Petrov, Kostadin, Bradley M. Wierbowski, Jingjing Liu, and Adrian Salic. 2020. 'Distinct Cation Gradients Power Cholesterol Transport at Different Key Points in the Hedgehog Signaling Pathway'. *Developmental Cell* 55 (3): 314–327.e7. <https://doi.org/10.1016/j.devcel.2020.08.002>.
- 805 Pizette, Sandrine, Tamás Matussek, Bram Herpers, Pascal P. Théron, and Catherine Rabouille. 2021. 'Hherisomes, Hedgehog Specialized Recycling Endosomes, Are Required for High Level Hedgehog Signaling and Tissue Growth'. *Journal of Cell Science* 134 (10). <https://doi.org/10.1242/jcs.258603>.
- 810 Port, Phillip, George Hausmann, and Konrad Basler. 2011. 'A Genome-Wide RNA Interference Screen Uncovers Two P24 Proteins as Regulators of Wingless Secretion'. *EMBO Reports* 12 (11): 1144–52. <https://doi.org/10.1038/embor.2011.165>.
- Porter, Jeffery A., Keith E. Young, and Philip A. Beachy. 1996. 'Cholesterol Modification of Hedgehog Signaling Proteins in Animal Development'. *Science* 274 (5285): 255–59. <https://doi.org/10.1126/science.274.5285.255>.
- 815 Qiu, Kaixin, Xiaojie Zhang, Shuai Wang, Chunyan Li, Xin Wang, Xuezhi Li, and Yili Wu. 2019. 'TMP21 in Alzheimer's Disease: Molecular Mechanisms and a Potential Target'. *Frontiers in Cellular Neuroscience* 13. <https://doi.org/10.3389/fncel.2019.00328>.
- Resh, Marilyn D. 2021. 'Palmitoylation of Hedgehog Proteins by Hedgehog Acyltransferase: Roles in Signalling and Disease'. *Open Biology* 11 (3): rsob.200414. <https://doi.org/10.1098/rsob.200414>.
- 820 Rubin, Lee L., and Frederic J. de Sauvage. 2006. 'Targeting the Hedgehog Pathway in Cancer'. *Nature Reviews Drug Discovery* 5 (12): 1026–33. <https://doi.org/10.1038/nrd2086>.
- Sandoval, Lisette, Mariana Labarca, Claudio Retamal, Paula Sánchez, Juan Larraín, and Alfonso González. 2022. 'Sonic Hedgehog Is Basolaterally Sorted from the TGN and Transcytosed to the Apical Domain Involving Dispatched-1 at Rab11-ARE'. *Frontiers in Cell and Developmental Biology* 10. <https://doi.org/10.3389/fcell.2022.833175>.
- 825 Saraste, Jaakko, and Michaël Marie. 2018. 'Intermediate Compartment (IC): From Pre-Golgi Vacuoles to a Semi-Autonomous Membrane System'. *Histochemistry and Cell Biology* 150 (5): 407–30. <https://doi.org/10.1007/s00418-018-1717-2>.
- 830 Schuiki, Irmgard, and Allen Volchuk. 2012. 'Diverse Roles for the P24 Family of Proteins in Eukaryotic Cells'. *Biomolecular Concepts* 3 (6): 561–70. <https://doi.org/10.1515/bmc-2012-0028>.
- Shin, Ji Hyun, So Jung Park, Doo Sin Jo, Na Yeon Park, Joon Bum Kim, Ji Eun Bae, Yoon Kyung Jo, et al. 2019. 'Down-Regulated TMED10 in Alzheimer Disease Induces Autophagy via ATG4B Activation'. *Autophagy* 15 (9): 1495–1505. <https://doi.org/10.1080/15548627.2019.1586249>.
- 835 Straface, Giuseppe, Tamar Aprahamian, Andrea Flex, Eleonora Gaetani, Federico Biscetti, Roy C. Smith, Giovanni Pecorini, et al. 2009. 'Sonic Hedgehog Regulates Angiogenesis and Myogenesis during Post-Natal Skeletal Muscle Regeneration'. *Journal of Cellular and Molecular Medicine* 13 (8b): 2424–35. <https://doi.org/10.1111/j.1582-4934.2008.00440.x>.
- 840 Strating, Jeroen R.P.M., and Gerard J.M. Martens. 2009. 'The P24 Family and Selective Transport Processes at the ER-Golgi Interface'. *Biology of the Cell* 101 (9): 495–509. <https://doi.org/10.1042/bc20080233>.
- Tabata, T., and T. B. Kornberg. 1994. 'Hedgehog Is a Signaling Protein with a Key Role in Patterning Drosophila Imaginal Discs'. *Cell* 76 (1): 89–102. [https://doi.org/10.1016/0092-8674\(94\)90175-9](https://doi.org/10.1016/0092-8674(94)90175-9).

- 845 Tang, Xiao, Rong Chen, Vince St Dollente Mesias, Tingxuan Wang, Ying Wang, Kristina Poljak, Xinyu Fan, et al. 2022. 'A SURF4-to-Proteoglycan Relay Mechanism That Mediates the Sorting and Secretion of a Tagged Variant of Sonic Hedgehog.' *Proceedings of the National Academy of Sciences of the United States of America* 119 (11): e2113991119. <https://doi.org/10.1073/pnas.2113991119>.
- 850 Tanimoto, H., S. Itoh, P. ten Dijke, and T. Tabata. 2000. 'Hedgehog Creates a Gradient of DPP Activity in *Drosophila* Wing Imaginal Discs'. *Molecular Cell* 5 (1): 59–71. [https://doi.org/10.1016/s1097-2765\(00\)80403-7](https://doi.org/10.1016/s1097-2765(00)80403-7).
- Tashima, Yuko, Tetsuya Hirata, Yusuke Maeda, Yoshiko Murakami, and Taroh Kinoshita. 2021. 'Differential Use of P24 Family Members as Cargo Receptors for the Transport of Glycosylphosphatidylinositol-Anchored Proteins and Wnt1'. *The Journal of Biochemistry* 171(1): 75–83. <https://doi.org/10.1093/jb/mvab108>.
- 855 Thauvin, Marion, Irène Amblard, Christine Rampon, Aurélien Mourton, Isabelle Queguiner, Chenge Li, Arnaud Gautier, Alain Joliot, Michel Volovitch, and Sophie Vríz. 2022. 'Reciprocal Regulation of Shh Trafficking and H2O2 Levels via a Noncanonical BOC-Rac1 Pathway.' *Antioxidants (Basel, Switzerland)* 11 (4): 841–56. <https://doi.org/10.3390/antiox11040718>.
- 860 Torroja, Carlos, Nicole Gorfinkiel, and Isabel Guerrero. 2005. 'Mechanisms of Hedgehog Gradient Formation and Interpretation'. *Journal of Neurobiology* 64 (4): 334–56. <https://doi.org/10.1002/neu.20168>.
- Vetrivel, Kulandaivelu S., Anitha Kodam, Ping Gong, Ying Chen, Angèle T. Parent, Satyabrata Kar, and Gopal Thinakaran. 2008. 'Localization and Regional Distribution of P23/TMP21 in the Brain'. *Neurobiology of Disease* 32 (1): 37–49. <https://doi.org/10.1016/j.nbd.2008.06.012>.
- 865 Weigel, Aubrey V, Chi-lun Chang, Gleb Shtengel, C. Shan Xu, David P. Hoffman, Melanie Freeman, Nirmala Iyer, et al. 2021. 'ER-to-Golgi Protein Delivery through an Interwoven, Tubular Network Extending from ER'. *Cell* 184 (9): 2412–2429.e16. <https://doi.org/10.1016/j.cell.2021.03.035>.
- 870 Wolpert, Lewis. 2016. *Positional Information and Pattern Formation. Current Topics in Developmental Biology*. Vol. 117. Elsevier Inc. <https://doi.org/10.1016/bs.ctdb.2015.11.008>.
- Xie, Jian, Yuan Yang, Jianbo Li, Jing Hou, Kun Xia, Weihong Song, and Shengchun Liu. 2014. 'Expression of Tmp21 in Normal Adult Human Tissues'. *International Journal of Clinical and Experimental Medicine* 7 (9): 2976–83.
- 875 Yam, Patricia T, and Frédéric Charron. 2013. 'Signaling Mechanisms of Non-Conventional Axon Guidance Cues: The Shh, BMP and Wnt Morphogens'. *Current Opinion in Neurobiology, Development of neurons and glia*, 23 (6): 965–73. <https://doi.org/10.1016/j.conb.2013.09.002>.
- 880 Zanetti, Giulia, Kanika Bajaj Pahuja, Sean Studer, Soomin Shim, and Randy Schekman. 2012. 'COPII and the Regulation of Protein Sorting in Mammals'. *Nature Cell Biology* 14 (1): 20–28. <https://doi.org/10.1038/ncb2390>.

Figures and figure legends




Figure 1

Figure 1. Characterisation of SHH-RUSH system. (A) Schematic illustrating the SHH-RUSH reporter system. The Streptavidin-KDEL hook (grey) was encoded on the same plasmid as SHH-RUSH (green)

with an internal ribosome entry site (IRES, black) to enable expression of both genes. **(B)** Cartoon illustrating the SHH-RUSH retention and release upon biotin addition. **(C)** Representative western blot of SVG-A cell lysates either non-transfected (NT), transfected with untagged SHH (SHH) or with SHH-RUSH probed with antibodies against SHH. GAPDH was used as a loading control. **(D)** SHH-ELISA of supernatant taken from cells either non-transfected (NT) or transfected with SHH-RUSH and cultured in the absence or presence of biotin for 24 h. Data is represented as mean +/- SD from two individual experiments. Unpaired t-test p value < 0.05 (*). **(E)** RT-qPCR analysis of *Gli1* mRNA expression (using primer set a) in RNA extracted from 3T3 fibroblasts treated with supernatants taken from cells either non-transfected (NT), mock-transfected (Mock), or transfected with SHH-RUSH and cultured in the absence or presence of biotin for 24 h. Data is represented as mean +/- SD from two individual experiments. Unpaired t-test p value < 0.01 (**). **(F)** RT-qPCR analysis of *Gli1* mRNA expression (using primer set b) from NIH 3T3 fibroblasts treated with supernatants taken from cells either non-transfected (NT), or transfected with untagged SHH (SHH WT) or SHH-RUSH. Data is represented as mean +/- SD from two independent experiments. Unpaired t-test p value < 0.05 (*). ns = non-significant. **(G)** Representative confocal microscopy images of SVG-A cells expressing SHH-RUSH and fixed at indicated timepoints after biotin addition. Scale bar = 10 μ m. **(H)** Representative confocal images of SVG-A cells co-transfected with SHH-RUSH (purple) and a medial Golgi marker, Mannosidase-II-EGFP (Mann-II-EGFP; green). The left micrograph shows a snapshot at 5 min post biotin addition. Scale bar = 10 μ m. The right panel highlights zoomed insets of the Golgi area at indicated timepoints post biotin addition from the white dotted square on the left image. **(I)** Quantification of the mean fluorescence intensity of SHH-RUSH in the Golgi area (black line) and colocalisation (blue line) of the two channels over time. Data is represented as mean +/- SD from three individual experiments.




Figure 2

Figure 2. SHH-RUSH follows the classical biosynthetic pathway. (A) Representative confocal images from a single plane in a z-stack of images from cells transfected with SHH-RUSH and stained with antibodies against Rab6 or Rab7, and fixed at 45 min post biotin addition. (B) Quantification of colocalisation between SHH-RUSH and endogenous Rab5, Rab6 or Rab7 from whole cells in which the Golgi was masked, at different timepoints post biotin addition. Data is represented as mean +/- SD, n

915

= 30 cells. Unpaired t-test p value < 0.0001 (****) or < 0.001 (***) or < 0.01 (**). (C) Representative TIRF images of cells co-transfected with SHH-RUSH and GFP-Rab6a under the weak L30 promoter associated to Video S3. The left panel shows a still image taken at 1898 s post biotin addition. Scale bar = 10 μ m. The right panel highlights zoomed insets showing a double-positive SHH-RUSH/Rab6a vesicle arriving into the focal plane then fusing with the plasma membrane. (D) Schematic illustrating the SHH-RUSH cell surface staining assay. (E) Quantification by flow cytometry of the ratio of surface staining (anti-SHH + AF-647) over the total mCherry signal mean fluorescence intensities is represented at different timepoints post biotin addition in cells treated with siRNA against Rab6a (siRab6a), or Rab7a (siRab7a), or a non-targeting control (siCTRL). Data was normalised to t = 0 and plotted as mean +/- SD from four independent experiments. Unpaired t-test p value < 0.05 (*).




Figure 3

930 **Figure 3. TMED10 regulates SHH transport through the biosynthetic pathway.** (A) Representative western blot of lysates taken from SVG-A cells treated with siCTRL or siTMED10. Blots were then probed with antibodies against TMED10. GAPDH was used as a loading control. (B) Quantification of western blot in (A), data plotted as mean \pm SD, n = 3 independent experiments, * $p < 0.05$. (C) Quantification of SHH-NLuc released into cell culture media from cells treated with siCTRL or
935 siTMED10 over 24 hrs. Data plotted as mean \pm SD, n = 6 independent experiments, ** $p < 0.01$. (D) Quantification by flow cytometry of the ratio of surface staining (anti-SHH + AF-647) over the total mCherry signal mean fluorescence intensities is represented at different timepoints post biotin addition in cells treated with siRNA against TMED10 (siTMED10) or siCTRL. Data was normalised to t = 0 and plotted as mean \pm SD from 3 independent experiments. Unpaired t-test p value < 0.0001
940 (****). (E) Flow cytometry analysis of TfR surface staining at different timepoints post biotin addition from SVG-A cells transfected with TfR-RUSH and treated with siTMED10 or siCTRL. Data was normalised to t = 0 and plotted as mean \pm SD from 3 independent experiments. No significant differences were measured between the siTMED10 and siCTRL. (F) Colocalisation analysis of SHH-RUSH with cis-Golgi marker GM130. The left panel shows representative confocal images of cells
945 treated with siCTRL or siTMED10 and transfected with SHH-RUSH (magenta), fixed at 15 min post biotin addition, permeabilized and stained for GM130 (green). Scale bar = 10 μm . The right panel shows the quantification of colocalisation between SHH-RUSH and GM130 at 0 min and 15 min post-biotin addition. Data is plotted as mean \pm SD (n = 20 cells). Unpaired t-test p value < 0.0001 (****). (G) Colocalisation analysis of SHH-RUSH with trans-Golgi. The left panel shows representative confocal images of cells treated with siCTRL or siTMED10 and transfected with SHH-RUSH (purple),
950 fixed at 45 min post biotin addition, permeabilized and stained for TGN46 (green). Scale bar = 10 μm . The right panel shows the quantification of colocalisation between SHH-RUSH and TGN46 at 0 min and 45 min post-biotin addition. Data is plotted as mean \pm SD (n = 20 cells). Unpaired t-test p value < 0.05 (*). (H) TMED10 dynamics during SHH intracellular dynamics associated to Video S5. The left
955 panel shows a still image at 0 s post biotin addition of SVG-A cells co-transfected with GFP-TMED10 and SHH-RUSH. Scale bar = 10 μm . The right panel highlights zoomed images of a double-positive TMED10:SHH-RUSH vesicle trafficking together towards a perinuclear region at indicated timepoints. Scale bar = 2 μm .




Figure 4

Figure 4. TMED10 associates with SHH-RUSH at ERES and aids in its exit from ER. (A) Live confocal imaging of ER exit site marker SEC24D (green) and SHH-RUSH (purple) associated to Video S7. The left panel shows a still image of a cell co-transfected with SEC24D-EGFP and SHH-RUSH at 0 s post biotin addition. Scale bar = 10 μm . The right panel highlights zoomed images of a SHH-RUSH vesicle
965 leaving a SEC24D-positive ERES. Scale bar = 2 μm . **(B)** Quantification of SHH-RUSH at SEC24D-positive puncta over time post-biotin addition. Data is plotted as mean fluorescence intensity of SHH-RUSH (magenta) and SEC24D-EGFP (green) and is representative of 5 individual cells. **(C)** Representative confocal images from a single panel in a z-stack of cells treated with siCTRL or siTMED10 and transfected with SHH-RUSH, fixed at either 0 min or 15 min post biotin addition, permeabilized and
970 stained with antibodies against SEC31. Scale bar = 10 μm . **(D)** Quantification of SHH-RUSH colocalisation with SEC31 over time in siCTRL and siTMED10 treated cells. Data is plotted as mean \pm SD (n = 30 cells). Unpaired t-test p value < 0.01 (**) or < 0.0001 (****). **(E)** Quantification of the ratio of SHH-RUSH intensity at SEC31-positive puncta at 15 min versus 0 min post biotin addition in siCTRL or siTMED10 treated cells. Dotted line (red) represents relative SHH-RUSH intensity at 0 min post
975 biotin addition. Data is plotted as mean \pm SD (n = 30 cells). Unpaired t-test p value < 0.001 (***)




Figure 5

Figure 5: Loss of *Baiser* function affects Hh secretion and signalling *in vivo*. (A) Left panel: cartoon illustrating pupal lethality rescue assay and larval wing imaginal disc (WID) outgrowth assay. In the “tester line”, expression of Hh is increased in the WID posterior domain (in pink) leading to distal anterior tissue outgrowth. This can be rescued upon decreasing Hh level or Hh extracellular secretion. Middle panel: lethality rescue assay - percentage of escaper animals in “tester line” control, and overexpressing *bai* RNAi-s in the posterior domain of the “tester line”. Right panel: WID outgrowth assay - percentage of outgrown, moderately and fully rescued discs with respect to their genotypes. Sample numbers (n) are indicated above the graphs. (B-D) Hh target gene expression in wild-type and *bai* RNAi wing discs. (B): Engrailed, (C): Patched (Ptc) and (D): anti- β -gal (*dpp-lacZ*). The anterior/posterior (A/P) compartment boundaries are marked with yellow dotted lines. Placement of insets are indicated by white rectangles. White arrows point to the areas in the anterior compartment, where proximal Hh target gene expression is different between wild-type and *bai* RNAi conditions. Scale bar – 50 μ m. (E-G) Quantification of Engrailed (E, n=8), Patched (F, n=10) and *dpp-lacZ* (G, n=10) in the anterior compartment of wild-type control (black) and *bai* RNAi (magenta) discs. The A/P boundary is indicated by black arrows. (H) Left panel: Cartoon illustrating the dorsal depletion of *bai* using RNAi, with ventral area serving as an internal control. Middle and right panels: Extracellular staining of endogenous Hh (middle) and filamentous Actin (right) in discs with *bai* RNAi. Dotted yellow line labels boundary between dorsal and ventral compartments, scale bar – 50 μ m. Quantification of extracellular Hh and Actin intensity levels between the dorsal and ventral compartments are shown next to the corresponding immunostainings. Data plotted as mean \pm SD, n = 10. (I-J) Distal ectopic *dpp-lacZ* gene expression measured in discs overexpressing Hh in producing cells for 1 day (I) or 4 days (J). Cartoons next to the fluorescent panels illustrate *dpp-lacZ* expression pattern (green). The pink rectangles indicate the domain of expression of *UAS-Hh* together with GFP RNAi (control), *disp* RNAi (positive control) and *bai* RNAi. White arrows point to ectopic *dpp-lacZ* expression (or in case of *disp* and *bai* RNAi the lack of it) in cuboidal cells of the WID. Note, that 1-day induction of Hh overexpression does not lead to changes in WID morphology, and that *bai* RNAi in Hh producing cells is capable of suppressing ectopic distal *dpp-lacZ* expression. Additional examples can be found on Supplementary Figure S6. Scale bar – 100 μ m.