



HAL
open science

Quantification of antibiotic resistance genes (ARGs) in clouds at a mountain site (puy de Dôme, central France)

Florent Rossi, Raphaëlle Péguilhan, Nathalie Turgeon, Marc Veillette,
Jean-Luc Baray, Laurent Deguillaume, Pierre Amato, Caroline Duchaine

► To cite this version:

Florent Rossi, Raphaëlle Péguilhan, Nathalie Turgeon, Marc Veillette, Jean-Luc Baray, et al.. Quantification of antibiotic resistance genes (ARGs) in clouds at a mountain site (puy de Dôme, central France). *Science of the Total Environment*, 2023, 865, pp.161264. 10.1016/j.scitotenv.2022.161264 . hal-04300300

HAL Id: hal-04300300

<https://cnrs.hal.science/hal-04300300>

Submitted on 22 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Quantification of Antibiotic Resistance Genes (ARGs) in Clouds at a**
2 **Mountain Site (puy de Dôme, Central France)**

3
4 Florent Rossi¹, Raphaëlle Péguilhan², Nathalie Turgeon¹, Marc Veillette¹, Jean-Luc Baray^{3,4},
5 Laurent Deguillaume^{3,4}, Pierre Amato² and Caroline Duchaine¹

6
7
8 ¹ Centre de Recherche de l'Institut Universitaire de Cardiologie et de Pneumologie de
9 Québec, Université Laval, Québec City, Québec, Canada, Département de Biochimie, de
10 Microbiologie et de Bio-informatique, Faculté des Sciences et de Génie, Université Laval,
11 Québec City, Québec, Canada

12 ² Université Clermont Auvergne, CNRS, SIGMA Clermont, ICCF, F-63000 CLERMONT-
13 FERRAND, France.

14 ³ Université Clermont Auvergne, CNRS, Observatoire de Physique du Globe de Clermont-
15 Ferrand, UAR 833, F-63000 CLERMONT-FERRAND, France.

16 ⁴ Université Clermont Auvergne, CNRS, Laboratoire de Météorologie Physique, UMR 6016,
17 F-63000 CLERMONT-FERRAND, France.


18
19
20
21
22
23 **Corresponding authors:** Caroline.Duchaine@bcm.ulaval.ca

24 CRIUCPQ, Building A-21 2725 Chemin Ste-Foy, Québec, Canada G1V 4G5

25
26
27
28
29
30
31
32
33
34 **Keywords:** Clouds, Bioaerosols, Antimicrobial resistance genes (ARGs), Atmosphere

36 **Graphical abstract**

37



49

50

51

52

53 **Highlights**

- 54 • Significant concentrations of ARGs are present in clouds in France
- 55 • The origin of air masses influenced the type and quantities of detected ARGs
- 56 • Veterinary use could be the main contributors of ARGs in the free atmosphere
- 57 • In such stressful environment, ARGs may provide selective advantage to
- 58 microorganisms

59

60

61

62

63 **Abstract**

64 Antibiotic resistance in bacteria is becoming a major sanitary concern worldwide. The
65 extensive use of large quantities of antibiotics to sustain human activity has led to the rapid
66 acquisition and maintenance of antibiotic resistant genes (ARGs) in bacteria and to their
67 spread into the environment. Eventually, these can be disseminated over long distances by the
68 atmospheric transport. Here, we assessed the presence of ARGs in clouds as an indicator of
69 long-distance travel potential of antibiotic resistance in the atmosphere. We hypothesized that
70 a variety of ARGs can reach the altitude of clouds mainly located within the free troposphere.
71 Once incorporated in the atmosphere, they are efficiently transported and their respective
72 concentrations should differ depending on the sources and the geographical origin of the air
73 masses. We deployed high-flow rate impingers and collected twelve cloud events between
74 September 2019 and October 2021 at the meteorological station of the puy de Dôme summit
75 (1,465 m a.s.l, France). Total airborne bacteria concentration was assessed by flow cytometry,
76 and ARGs subtypes of the main families of antibiotic resistance (quinolone, sulfonamide,
77 tetracycline; glycopeptide, aminoglycoside, β -lactamase, macrolide) including one mobile
78 genetic element (transposase) were quantified by qPCR. Our results indicate the presence of
79 29 different ARGs' subtypes at concentrations ranging from 1.01×10^3 to 1.61×10^4 copies m^{-3}
80 of air. Clear distinctions could be observed between clouds in air masses transported over
81 marine areas (Atlantic Ocean) and clouds influenced by continental surfaces. Specifically,
82 quinolones (mostly *qepA*) resistance genes were prevalent in marine clouds (54% of the total
83 ARGs on average), whereas higher contributions of sulfonamide, tetracycline; glycopeptide,
84 β -lactamase and macrolide were found in continental clouds. This study constitutes the first
85 evidence for the presence of microbial ARGs in clouds at concentrations comparable to other
86 natural environments. This highlights the atmosphere as routes for the dissemination of ARGs
87 at a large scale.

89 **Introduction**

90

91 Antibiotic resistance has been classified as one of the 10 biggest threats for human societies in
92 the near future. In 2019, more than 4.95 million deaths were associated with antimicrobial
93 resistance, and about 1.27 million were directly attributed to antibiotic-resistant bacterial
94 infections ¹. If no action is taken, this number could rise to up to 10 million deaths per year by
95 2050, making it the first cause of death worldwide and engendering cumulative global cost to
96 the economy over 100 trillion USD ². The resistance to antimicrobial substances is a natural
97 phenomenon that includes natural microbial evolution dynamics, and the extensive use of
98 antibiotics to sustain growing human populations have greatly promoted it ³. Each year, tens
99 of thousands of tons of antibiotics are used for treating and preventing human, animal and
100 plant diseases ⁴, and at least a fraction can be released in the environment. Nowadays,
101 antibiotics can be detected in waste water ⁵ and in the natural environment, from rivers ⁶ and
102 sea water ^{7,8} to soils ⁹, and therefore carrying antibiotic resistance can now be considered
103 beneficial for bacteria in the environment. A recent report from the French national agency for
104 sanitary, environment and work safety highlighted median total antibiotics concentrations in
105 French rivers of 12 ng L⁻¹, dominated by fluoroquinolones and sulfamethoxazole ¹⁰. In
106 hospital raw effluent and waste water treatment plants effluent, these concentrations were
107 17.10³ ng L⁻¹ and 300 ng L⁻¹ respectively. In soils, concentrations reaching up to 62 to 3.10³
108 ng g of dw⁻¹ in sewage sludge were reported. Given such conditions, the natural environment
109 is becoming a major actor in the emergence of antibiotic resistance genes (ARGs) in microbes
110 naturally present in the environment.

111 So far, the main antibiotic resistance mechanisms described in bacteria include direct
112 antibiotic inactivation through enzymatic digestion, and modifications of the biological targets

113 such as by methylation of the binding sites ¹¹. Other mechanisms prevent antibiotics from
114 entering the cell (e.g. biofilm formation, reduced membrane permeability) or are responsible
115 for their active evacuation (e.g. efflux pumps). These mechanisms are encoded by so-called
116 antimicrobial resistance genes (ARGs), carried either by chromosomes or plasmids. In the
117 latter case, these can be easily transferred horizontally between bacteria cells and species
118 through genetic material exchanges (conjugation, etc).

119 ARGs dispersal is logically tightly linked with that of bacteria cells. ARGs can transit
120 via microbes between ecosystems, and the atmosphere could act as a major route of their
121 dispersal, in particular over long distances ^{12,13}. Although the conditions encountered in the
122 atmosphere may severely limit bacterial survival, some amount of genetic material is
123 undoubtedly dispersed by atmospheric circulations. Investigating the occurrence of ARGs in
124 the air at high-altitude and clouds can thus provide valuable information regarding the long-
125 distance spread of such functions. Main bacteria cell aerosolization mechanisms include
126 mechanical disturbances, such as wind, raindrop impaction on surfaces, bubble-bursting, etc
127 ¹⁴⁻¹⁶. Once aerosolized, depending on air turbulence, airborne micro-organisms can eventually
128 reach higher altitudes, travel over continents ^{13,17}, and can integrate the atmospheric water
129 cycle, clouds and precipitation, before deposition ^{18,19}.

130 With the increase awareness that human health is closely related with the wider
131 environment, studying ARGs dispersal in the atmosphere, its potential to spread over long
132 distances and its relation with human activities could provide critical and valuable
133 information. Hence, this study is a part of the wider “Discovery Frontiers” research program
134 dedicated to the study of airborne dissemination of ARGs ²⁰. In compliance with the “One
135 Health” interdisciplinary approach ^{21,22}, this research program incorporates assessment of
136 several different indoor environments and their outside emissions, novel ARG tracking and

137 surveillance methods, selective culture approaches, and animal models to be incorporated into
138 exposure and risk assessment models.

139 Here, we examined ARGs in clouds in order to investigate their capacity to be
140 dispersed over long distances through the multiphase atmosphere. Clouds were sampled
141 between September 2019 and October 2021 from a high altitude site in central France (puy de
142 Dôme meteorological station, 1,465 m a.s.l.). For each event, total bacteria and a panel of 33
143 ARGs subtypes were quantified by flow cytometry and qPCR, respectively, and interpreted in
144 regards to the geographical origin of the corresponding air masses. The ARGs assessed were
145 related to quinolone, sulfonamide, tetracycline, glycopeptide, aminoglycoside, β -lactamase
146 and macrolide resistance and included one mobile genetic element (transposase). Overall, this
147 study constitutes the first report of the presence of microbial ARGs in clouds and evidences of
148 the atmosphere, and especially clouds, as pathways for the dissemination of antimicrobial
149 resistance functions worldwide.

150

151 **Material and Methods**

152 **Study site, collection of clouds and Air Masses Backward Trajectories**

153 Cloud sampling was performed at the puy de Dôme (PUY) meteorological observatory
154 (1,465 m a.s.l., 45.772° N, 2.9655° E - France) from September 2019 to October 2021. At
155 each sampling occasion, two high flow rate impingers²³ (HFRi, DS6, Kärcher SAS,
156 Bonneuil sur Marne, France) sampling at an air flow rate of $\sim 2 \text{ m}^3 \text{ h}^{-1}$ were deployed from the
157 roof platform of the station to collect the cloudy air, that were then used for qPCR analyses. In
158 parallel, a cloud collector allowing to sample cloud droplets was deployed nearby^{24,25} for
159 total cell quantification using flow cytometry. Sampling were performed for 2 to 5
160 consecutive hours depending on cloud events' duration. The HFRi were filled with 850 mL of

161 sterile nucleic acid preservation (NAP) buffer solution as the impaction liquid ²⁶. The volumes
162 of cloud water recovered were estimated by weighting, assuming a cloud water density of 1.0.
163 The NAP buffer solution was composed by 0.019 M of ethylenediaminetetra-acetic acid
164 (EDTA) disodium salt dihydrate, 0.018 M of sodium citrate trisodium salt dihydrate, 3.8 M of
165 ammonium sulfate, and H₂SO₄ to pH 5.2. HFRi samplers were sterilized prior each sampling
166 event by thorough rinses with 70% ethanol followed by 10 minutes of exposure to UV light.
167 Field blanks were conducted before each sampling and consisted of 850 mL of sterile NAP
168 buffer solution left in contact with the HFRi's tanks for several minutes before recovery. The
169 cloud droplet impactor was rinsed with ultrapure water and sterilized by autoclave before each
170 sampling.

171 The CAT three-dimensional (3D) forward/backward kinematic trajectory model was
172 used to calculate the 72-hours backward trajectories of clouds' air masses. The model uses
173 dynamic fields extracted from the ECMWF ERA-5 meteorological data archive
174 (<https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5> ²⁷), allowing for the
175 determination of air masses geographical origins and altitudes over the last 72 hours before
176 sampling at the PDD meteorological observatory ^{28,29}. The model also provides the altitude
177 relative to the planetary boundary layer (PBL) to estimate the influence of the underlying
178 surfaces. In the present work, the ECMWF data used to calculate back-trajectories have a
179 spatial resolution of 0.5° in latitude and longitude and a temporal resolution of 3 hours, and
180 back-trajectory points have been calculated every 15 min.

181

182 **Sample Processing and DNA Extraction**

183 Cloud water samples collected with the HFRi in NAP preservation buffer were
184 concentrated by individual filtration through 47 mm diameter mixed cellulose ester

185 membranes (MCE - 0.2 μm porosity) within a laminar flow hood, rolled into 5 mL bead tubes
186 (Macherey-Nagel) and added with 1.2 mL of lysis buffer (MWA1 - Macherey-Nagel) before
187 being stored at -80°C until further analysis. DNA extraction was carried out individually from
188 each separate filter using the NucleoMag DNA/RNA Water kit for water and air samples
189 (Macherey-Nagel), based on the manufacturer's instructions, with some minor modifications
190 to match with the modifications made on the original protocol (use of 47 mm diameter filters
191 instead of 25 mm, and larger volumes of lysis buffer): briefly, after the recommended 10 min
192 of mechanical lysis and 30 s of centrifugation (30 s at 1,000 g), 450 μL of the lysates was
193 transferred into two different microtubes. DNA extraction was then performed as
194 recommended, with final elution volumes of 50 μL of RNase-free H_2O provided in the kit.

195

196 **Quantification of Total Bacteria by Flow Cytometry**

197 The concentration of total bacteria in cloud water was determined from samples
198 collected with cloud droplets impactors. Briefly, the cells contained in 40 mL of water
199 samples were pre-concentrated into ~ 200 μL by ultrafiltration using vivaspin 20 (Sartorius, 10
200 Kda molecular weight cut off), from two successive centrifugation steps of 20 mL at 4000 g
201 for 16 min, and the addition of 1 mL of tris-EDTA buffer (1X), then transfer into 2 mL sterile
202 microtubes. The exact concentration factor for each sample was obtained by weighting initial
203 and final liquid volumes, considering a density of 1.0. Volumes of 200 μL of the pre-
204 concentrated samples were then stained with SYBR Green (1X final concentration,
205 Invitrogen) for 15 minutes in the dark at room temperature, and total bacteria counts were
206 performed using an LSR Fortessa Cell Analyzer flow cytometer (Becton Dickinson, Franklin
207 Lakes, NJ). Cytograms were obtained by plotting SYBR green fluorescence intensity (488
208 nm, voltage 350 V) over forward scattering (voltage 500 V) after 60 seconds of run, and flow

209 rates were measured before analyzes of samples by weighing of ultrapure water after 10
210 minutes of run. All flow cytometry analyses were conducted within 4 to 12 hours of sampling.

211

212 **ARGs Quantification by qPCR**

213 A panel of 33 ARGs belonging to the main families of antibiotic resistance was
214 assessed from samples collected with HFRi in NAP buffer: β -lactamases, quinolones,
215 tetracyclines, macrolides, aminoglycosides, sulfonamides, vancomycins as well as one mobile
216 genetic element (transposases). More details about the ARGs targeted, including the
217 corresponding sets of primers, are described in supplementary material (Table 1SI^{30,31}). PCR
218 reactions and cycling conditions were identical for all target genes, and were as following: 4
219 μ L of template DNA (diluted to 1/10), 10 μ L of IQ Green supermix (BioRad), 5.88 μ L of
220 RNase-free H₂O and 0.12 μ L of equimolarly mixed primers (50 μ M each) in 20 μ l final
221 volume. Cycling conditions were set at 95°C for 3 min as initial denaturation step, followed
222 by 40 cycles of 95°C for 10 s and 60°C for 30 s. The concentrations of each target gene in the
223 samples were determined relatively to 10-fold dilution series of a 10⁶ copies x μ l⁻¹ mix of
224 control plasmids (pUCIDT-AMP) including the target sequence. All qPCR reactions were
225 performed in technical duplicates from the pool of DNA extracts from samples collected
226 simultaneously. Field blanks were also processed under the same conditions. A Ct value
227 threshold of 37 was used as the limit of quantification, and all amplification peaks were
228 systematically checked based on the melting curves of the corresponding plasmid standard.

229

230 **Statistical Analyses**

231 All statistics were performed using R (R Core Team (2021)). Statistical differences between
232 each family of ARGs considering all cloud events (n = 12) was assessed by the nonparametric

233 Kruskal-Wallis test followed by multiple pairwise comparison using Tukey contrast from the
234 “nparcomp” function of the R package of the same name³². The relationships between the 4
235 variables associated with the 72-hours backward trajectories (time spent within and above the
236 planetary boundary layer, above continental and marine surfaces) and the relative
237 concentrations of the different families of antibiotics quantified (quinolones, β -lactamases,
238 vancomycin, tetracycline, sulfonamide, macrolide, aminoglycosides) including transposase in
239 the samples were assessed using partial least square regression analysis (PLSR). The analysis
240 was computed using the “PLS” function from the mixOmic R package^{33,34} and all data were
241 centered and reduced before analysis. Prior to the PLSR analysis, a hierarchical clustering was
242 performed considering only the 72-hours backward trajectory parameters (Table 2 SI) to
243 identify classes of cloud samples. The clustering involved Euclidean distances and was
244 performed using the ward aggregation method. Statistical differences between clusters were
245 assessed by multiple bootstrap resampling (100 bootstraps, $P < 0.05$) using the “pvclust”
246 function of the corresponding R package³⁵. Based on these groups, 95% level confidence
247 interval ellipses were drawn.

248

249 **Results**

250 **ARGs diversity and concentration in clouds**

251 Of the 33 ARGs subtypes assessed in our study on 12 independent cloud events, 29 were
252 detected at least once, and 6 were observed in 75% (9) or more of the samples (*qnrB* – *qepA* -
253 *aac(3)* – *ermB* – *tetO* and *vanA*); 4 of the targets were not detected in any of the samples (*tetS*
254 – *blaGes* - *blaMOX* and *vanSA*, Table 3 SI). The total absolute concentration of the ARGs
255 assessed, considering all clouds, averaged $2.08 \times 10^4 \pm 6.40 \times 10^3$ copies mL⁻¹ of cloud water
256 (mean \pm standard error of the mean), with values ranging from 3.45×10^3 to 6.47×10^4 copies

257 mL⁻¹ depending on samples (Table 1 and Figure 1A). This corresponded to an average
258 concentration of $5.42 \times 10^3 \pm 1.51 \times 10^3$ ARGs copies m⁻³ of air (1.01×10^3 to 1.61×10^4 copies m⁻³).
259 ³). The highest values were observed on the 03-Nov-20 and 10-Nov-20 and the lowest on the
260 22-Oct-19 and 28 Apr-21. The concentration of every family of ARGs varied significantly
261 between clouds (Kruskal-Wallis, $P < 0.05$). However, quinolone resistance genes were by far
262 the most prominent family (Kruskal-Wallis, $P < 0.05$). Their absolute concentration ranged
263 from 2.69×10^2 to 4.90×10^4 copies mL⁻¹ (1.15×10^2 to 1.24×10^4 copies m⁻³ of air respectively),
264 with an average of $1.19 \times 10^4 \pm 3.93 \times 10^3$ copies mL⁻¹ ($3.18 \times 10^3 \pm 9.87 \times 10^2$ copies m⁻³). Based
265 on the respective relative contributions of the different families of genes to the total pool of
266 ARGs, two different patterns of samples could be observed (Figure 1B): half of clouds were
267 characterized by limited proportions of resistance to quinolones (50% or less, 34% on
268 average) and relatively high proportions of genes related with resistance to either macrolides
269 (17-jul-20 and 10-Nov-20), β -lactamases (16-Oct-20 and 26-Oct-21), sulfonamides (28-Apr-
270 21), or the three of them (22-Oct-19). The other half of samples displayed an overwhelming
271 majority of quinolones resistance genes, reaching up to 88% (74% on average), and low
272 proportions of all the other families. Interestingly, the vast majority of quinolone resistance
273 was contributed by *qepA* (~98%), whereas *qnrB* represented only a minor fraction (< 2% on
274 average), regardless of clouds. Concerning the other genes, aminoglycoside resistance and
275 transposases were the lowest of all (Kruskal-Wallis, $P < 0.05$) with average concentrations of
276 1.08 ± 0.45 and 2.51 ± 1.35 copies mL⁻¹ respectively (0.26 ± 0.11 and 1.14 ± 0.76 copies m⁻³).
277 Their cumulated contribution to the total ARGs barely reached 0.05%, and there were many
278 events where they could not be detected. The relative contributions of the other families of
279 ARGs were, by decreasing importance: β -lactamases (~12%), sulfonamides (~11%),
280 vancomycins (~9%), macrolides (~8%) and tetracyclines (~6%).

281 The total absolute bacteria number concentration averaged $8.0 \times 10^3 \pm 2.9 \times 10^3$ cells per mL^{-1} of
282 cloud water, with, as observed for ARGs, large variations between samples (Figure 2 and
283 Table 1). Based on these data, each bacteria cell carried on average 7.1 ± 2.5 ARGs.
284 Nevertheless, no correlation could be observed between these two variables (Spearman
285 correlation coefficient $\rho = -0.23$, $P = 0.5$, figure 2), indicating that the total ARGs copies does
286 not necessarily covariate with the total number of bacteria cells in cloud water. Specifically,
287 there were >10 times more ARGs copies than bacteria cells in the samples on the 10-Nov-20,
288 19-Nov-20 and 13-Oct-21, and ~ 5 times more in the samples collected on the 11-Mar-20, 03-
289 Nov-20, 06-May-20, and 26-Oct-20. In contrast, in the samples collected on 22-Oct-19, 16-
290 Oct-20, 28-Oct-20 and 28-Apr-21, bacteria cells were more abundant than ARGs copies by
291 factors of up to one order of magnitude, with values that ranged between 1 ARG every 10
292 bacteria to 1 ARG per bacteria at most.

293

294 **Influence of the geographical origin of the air masses on the ARG and bacteria** 295 **abundances in clouds**

296 The clouds assessed originated from various geographical areas (Figure 1SI). Based on the
297 time spent by air masses above continental or marine surfaces within the 72-hours preceding
298 sampling, as well as their altitudes, 3 groups of samples could be distinguished (Figure 2SI
299 and Table 2SI): “Continental” clouds (events 22-Oct-19, 17-Jul-20, 16-Oct-20, 10 Nov-20
300 and 28-Apr-21), for which the air masses spent on average 70% of the time over continental
301 surfaces, “Marine high” clouds (events 11-Mar-20, 03-Nov-20, 19-Nov-20 and 06-May-21),
302 for which the air masses spent 85% of time over the Atlantic Ocean, mostly at high altitude in
303 the free troposphere, and finally “Marine low” clouds (events 28-Oct-20, 13-Oct-21 and 26-
304 Oct-21) for which the air masses spent most time (75%) above the Atlantic Ocean, for a large
305 part within the planetary boundary layer (PBL).

306 Based on the relative abundance of the main families of ARGs (95% level confidence
307 interval; Figure 3A), “Continental” clouds differed significantly and were more variable than
308 “Marine” clouds (both “low” and “high”), whereas the two “Marine” types clouds did not
309 significantly differ from each other. More precisely, quinolones resistance genes were more
310 abundant in “Marine” clouds (Figure 3B), in particular “Marine high” clouds, whereas
311 sulfonamide and tetracycline resistance genes appeared mostly associated with “Continental”
312 surfaces, especially for air masses that travelled within the PBL. Macrolide resistance genes
313 were associated with continental surfaces as well, but these were mostly prevalent in air
314 masses that travelled at higher altitude. The abundances of β -lactamase, vancomycin and
315 aminoglycoside resistance genes as well as transposases could not be explained by the
316 geographical origin of the air masses using this classification.

317

318 **Discussion**

319 **ARGs reach high altitudes and clouds**

320 Detecting ARGs in clouds robustly evidences that they are transported over large
321 geographical areas over the planet. Here, we monitored 33 different genes belonging to the
322 main classes of antimicrobial resistance. This is standard in atmospheric studies^{36–38},
323 although it is still non-exhaustive of the ARGs known. For this reason, most estimations of
324 ARGs concentrations in the environment, including ours in clouds, likely underestimate the
325 current situation. The ARGs known at the time of writing includes more than 6,500 ontology
326 terms covering resistant mechanisms from approximately 2,923 antimicrobial resistance
327 determinants³⁹. Most have functions not directly associated with antibiotic resistance but with
328 more general roles in stress response, for instance the increase of outer-membrane
329 permeability to face osmotic variations, efflux pumps toward a large variety of toxic

330 substances⁴⁰. Some antibiotic-degrading β -lactamases (ampC-type) were also found to be
331 related with ancestral peptidoglycan-modifying activities. The atmosphere is a highly
332 selective and stressful environment for microbial cells²⁴; some of these genes and functions
333 could therefore be favorable for survival⁴¹, and reinforce the capacity of bacterial ARGs to
334 spread by atmospheric transport.

335 Our data evidence cloud's resistome. Numerous antibiotic resistance genes were found
336 recurrent or more sporadic, at concentrations ranging from $\sim 10^3$ to $\sim 10^5$ gene copies mL^{-1} .
337 Comparatively to other environments, this is 2 to 3 orders of magnitude lower than in rivers
338⁴²⁻⁴⁴ (6.2×10^6 to 1×10^8 copies mL^{-1}) and lakes⁴⁵ (4.4×10^6 to 2.4×10^7 copies mL^{-1}), and waste
339 water effluents⁴⁶ (1.0×10^3 to 1.0×10^7 copies mL^{-1}) and sea in coastal areas (1.2×10^4 –
340 3.9×10^5 copies mL^{-1}). Our observations are even higher than the concentrations reported in
341 sea water⁴⁷ (21.1 to 8×10^3 copies mL^{-1}). However, on a volumetric basis, it has to be kept in
342 mind that water in clouds is distributed among droplets with micrometric radius, and cloud
343 liquid water content (LWC) is typically on the order of ~ 0.1 to 0.8 mL of water per m^{-3} of air.
344 In this respect, clouds on the planet represent diluted environments since they are
345 polydisperse medium, also composed of air. Additionally, direct comparisons of ARG
346 quantification between independent studies are not necessarily relevant as the number and
347 diversity of the targets may vary.

348

349 **Spatial and temporal variability of ARGs in clouds and links with potential sources**

350 The abundance of antimicrobial resistance functions in the environment is likely variable in
351 space and times in relation with the natural (seasonal) or anthropogenically induced
352 fluctuations of biomass and biodiversity in ecosystems, meteorological conditions,
353 atmospheric and oceanic circulation, etc. In the atmosphere, the microbial content is affected

354 by sources such as vegetation, so marked seasonal variations of the biodiversity are often
355 observed, including the puy de Dôme sampling site ⁴⁸. Here, we did not observe such seasonal
356 variations on the ARGs in clouds and their abundance. Many concurrent environmental
357 factors influence the composition of cloud water (e.g. ionic concentrations), and although our
358 dataset is relatively consequent regarding the difficulties of sampling, it remains too limited to
359 clearly identify environmental trends such as seasonality, requiring long-term monitoring.

360 We could detect links between the geographical origin of the air masses and the relative
361 abundances of the different ARG families in clouds, which attests of the heterogeneous
362 distribution of ARGs in the environment. These can be related with differences of usage
363 between countries and continents, as previously reported in waste water treatment plants ⁴⁶.
364 Quinolones resistance genes were dominant in our study, in particular in air masses
365 influenced by marine areas. This is consistent with literature as they were found to be the
366 second most abundant in urban air samples from various cities around the Earth, although still
367 far behind β -lactamases resistance genes ³⁸. However, their presence in the indoor-air of
368 human facilities, including waste water treatment plants and many different animal farms,
369 appears much lower ⁴⁹. While this partially traduces different detection methods, this could
370 suggest that these systems may not be the main contributors of quinolones resistance in the
371 atmosphere in comparison with natural sources. This is supported by metagenomics data from
372 ocean samples collected by past TARA expeditions from 2009 to 2012 ⁵⁰, where an important
373 proportion of quinolone resistance genes, in particular multidrug efflux pumps was observed
374 and especially in coastal areas ⁴⁷. Similar trends have been observed in soils, where efflux
375 pump resistance mechanisms represented nearly 60% of the total detected ARGs, out of 242
376 tested ⁵¹. In our study, the vast majority of quinolones were related with *qepA*, an exclusively
377 plasmid-mediated efflux pump ⁵² from the superfamily of major facilitators, likely derived
378 from soil β -proteobacteria strains' transmembrane transporters ^{53,54} and hence explaining its

379 occurrence in the environment . Its regulation involves the soxRS gene, a positive two-
380 component regulator part of the superoxide stress response and triggered by various stresses,
381 including damages to DNA, exposure to toxicants and reactive oxygen species ^{52,54}. This
382 supports that the environment is not only an important reservoir of bacterial resistance but
383 could promote their maintenance through environmental stresses, and that the atmosphere
384 would act as vector at a large scale of these genes. Quinolone antibiotics, including
385 fluoroquinolones, are broad spectrum synthetic antibiotics introduced in the early 1960s and
386 widely used until the late 1980s ⁵⁵. Since the recommendation of their restriction in Europe
387 emitted by the Pharmacovigilance Risk Assessment Committee (PRAC) in 2018, their
388 utilization has plummeted down to ~6% in human health and ~2% in food production ⁵⁶.
389 Notable molecules in this family of antibiotics includes ciprofloxacin, levofloxacin and
390 norfloxacin, used against a large panel of Gram-negative (including *Pseudomonas*
391 *aeruginosa*) and Gram-positive bacteria (including *Staphylococcus aureus*); their activity as
392 antibacterial agents rely on their interaction with topoisomerase II (DNA gyrase), which
393 prevents DNA replication. This class of molecules rapidly caused the emergence of resistance
394 in bacteria ⁵⁷. It therefore makes sense to find quinolone resistance genes among the prevalent
395 ARGs in the environment.

396 The other main classes of antibiotic resistance, namely β -lactamases, sulfonamides,
397 vancomycin, macrolide and tetracycline, were relatively more abundant in “Continental”
398 clouds. While this partly results from a relative decrease of the amounts quinolones resistance
399 genes, it also suggests continental surfaces as their main sources. Table 2 summarizes
400 antibiotic consumption for 29 countries in Europe for human and veterinary usages ⁵⁶.
401 Veterinary usages contribute ~61.4% (6558 tons of active products) of the total antibiotics
402 market, versus ~ 38.6% for human health (4122 tons) ⁵⁶. Most antibiotics related to human
403 health are expected to be caught by waste water treatment systems, where removal efficiency

404 has been estimated to reach up to 76% of the inputs ⁵⁸, with variations depending on
405 compounds ⁵⁹. If we omit the gene *qepA*, which we related to oceanic inputs as mentioned
406 previously, we found a fairly good agreement between the relative abundances of
407 antimicrobial functions observed in clouds and the proportions of the different families of
408 molecules used in the various countries. For instance, the highest concentration of ARGs were
409 observed on 10-Nov-20 in an air mass that flew over Spain. This country is the biggest
410 purchaser of antibiotic products for veterinary purposes in Europe (~ 1,244 tons of active
411 substance ⁶⁰). More local sources should also largely contribute to our observations.
412 Accordingly, sulfonamides and macrolides, which represent 21.7 % and 12.3%, respectively,
413 of the veterinary antibiotics sold in France ^{61,62}; accordingly, represented on average 21.5%
414 and 13.9% of the total ARGs quantified in our samples. However, resistance to tetracycline
415 was found at lower proportions than could be expected from such extrapolations. This may
416 relate with seasonal variations, as observed in rivers with the lowest abundance of *tetA*, *tetC*
417 and *tetM* during winter than the rest of the year ⁶³.

418

419 **The selective advantage of ARGs in clouds could promote their maintenance in the** 420 **environment**

421 There are increasing reports of the presence of multi-resistant bacteria in the environment, and
422 this represents one of the most worrisome threats to human societies nowadays. In our work,
423 although multi-resistance was not specifically addressed, we found ARGs at higher
424 abundance than bacteria cell by a factor of 5 in ~60% (7/12) of the samples. As the samples
425 were filtered through 0.22 µm porosity before analysis, viruses and cell-free DNA fragments
426 (i.e., external DNA, bacteriophage-associated DNA, and membrane vesicle-associated DNA)
427 were excluded. However, they are known to have critical roles in horizontal gene transfer and
428 dispersal including ARGs ^{66,67}. Viruses for example, exhibit high proportions of ARGs in

429 their genome, with a strong prevalence of antibiotic efflux mechanism ⁶⁸ Overall, these
430 evidences the presence of multiple ARG copies in genomes and possibly multiple resistance
431 combined together within single cells, and this was particularly true for “Marine” clouds.
432 Such prevalence of antimicrobial resistance genes is most likely a combination of both
433 intrinsic and acquired resistance, i.e. of ARGs carried by both chromosomes and plasmids; the
434 methods used here do not allow distinguishing them.

435 Many ARGs such as the efflux pump *qepA* are involved in global stress regulation as
436 already mentioned. We propose that these functions could participate to the survival of
437 bacteria in clouds, which would in turn contribute to global spread. Clouds are indeed
438 depicted as particularly harsh environments, selective for the most efficient multiple-stress
439 response phenotypes ^{69–71}. Metatranscriptomics indicated the expression of biological
440 responses to stress, including defense mechanisms like elevated transmembrane transports,
441 binding activities and protein phosphorylation ²⁴. The selective advantage of carrying ARGs
442 such as the efflux pump *qepA* in clouds could therefore contribute to their maintenance and
443 spread in the environment, even in the absence of such antimicrobial molecules. A connection
444 between antibiotic resistance acquisition and bacterial aerosolization was even recently
445 established ⁴¹. More specifically, the osmotic and desiccative stresses caused by aerosolization
446 triggered an immediate response to antibiotic compounds targeting cell wall and protein
447 synthesis (Cephalothin and Gentamicin) in *Escherichia coli*.

448

449 **Global ARGs Dispersal**

450 Based on our current dataset, that compiles 12 clouds with various geographical origins, we
451 attempt here to provide an evaluation of the global contribution of clouds in antimicrobial
452 genes dispersal (Table 3). The ARG concentrations observed in clouds were expressed as

453 copies m^{-3} of cloudy air, based on sampling times and sampler's air flow rate corrected for the
454 altitude of puy de Dôme (1465 m). Considering an average volume of clouds within the first 6
455 km above the Earth of $4.66 \times 10^{17} \text{ m}^3$ ⁷² and assuming ARGs concentrations similar as in our
456 study (5.42×10^3 copies of ARGs m^{-3} of air), we estimate that globally clouds carry in
457 permanence $\sim 2.53 \times 10^{21}$ copies of ARGs. Annually, this would represent fluxes of $\sim 1.29 \times 10^{25}$
458 to $\sim 2.06 \times 10^{26}$ copies of ARGs transiting through clouds. Assuming a cloud precipitation rate
459 of $4.10 \times 10^{20} \text{ m}^3 \text{ year}^{-1}$ ⁷², approximately $\sim 2.2 \times 10^{24}$ copies would get precipitated to the Earth
460 surface by wet processes every year, while the remaining fraction can be expected to
461 evaporate and remain airborne. These estimations are based on a limited number of events
462 collected at a very specific location, but there can be strong spatial and temporal variation as
463 emphasized previously. Still, since we considered only clouds (i.e. $\sim 15\%$ of the atmospheric
464 volume) and did not account for below-cloud aerosol scavenging processes by precipitation,
465 and that we did not account for all known ARGs, these probably represent low estimates of
466 the total actual inputs of ARGs from the atmosphere.

467

468 **Conclusion**

469 In this study, we evidenced for the first time the presence of microbial ARGs in clouds at
470 concentrations comparable to other natural environments. Using a unique dataset composed of
471 12 events sampled over more than a year and originating from very different geographical
472 origins, our results highlights that clouds' resistome can be viewed as a mosaic composed of
473 antibiotic resistance from both distant origins and local landscapes. More specifically, a
474 strong Atlantic Ocean signature was observed in most of the clouds, as shown by the
475 predominance of the of *qepA* efflux pump, that decreased as the air masses flew over
476 continental surfaces, to the benefit of antibiotic resistance in relation with the consumption of
477 antibiotics in food-producing animals, such as β -lactamase, tetracycline and macrolide

478 resistance. Overall, our results point out clouds as a tremendous potential reservoir of ARGs
479 within the environment, and highlights the high atmosphere as routes for the dissemination of
480 ARGs at a large scale.

481

482 **Funding**

483 This project is part of the NSERC Discovery Frontiers-Antimicrobial resistance in the
484 Environment funded research program (539024-2019) (PI: Caroline Duchaine).

485

486 **Acknowledgements**

487 The puy de Dôme station which is a part of the CO-PDD national instrumented site, is
488 managed by the OPGC observatory and by the LaMP laboratory, and supported by the
489 Université Clermont Auvergne (UCA), by the Centre National de la Recherche Scientifique
490 (CNRS-INSU), and by the Centre National d'Etudes Spatiales (CNES). The authors would
491 like to thank Angelica Bianco for her help in sampling cloud water, Muriel Joly from the
492 ICCF microbiology department for her help in sample processing and DNA extractions, and
493 Sandra Banson-Valleix for her implication in the CAT 3D model calculation and the ECMWF
494 data extraction. The authors would also like to thank Hermine BILLARD and Jonathan
495 Colombet, from the Plateforme CYSTEM – UCA PARTNER (Clermont-Ferrand, FRANCE),
496 for their technical support and expertise.

497

498 **Conflict of Interest**

499 The authors declare no conflict of interest.

500

501 **Authors' contribution**

502 Caroline Duchaine, Nathalie Turgeon and Pierre Amato secured the funding. Caroline
503 Duchaine and Pierre Amato designed the experimental setup and defined the goal of the
504 study. Jean-Luc Baray and Laurent Deguillaume organized field sampling at the puy de Dôme
505 station. Florent Rossi, Raphaëlle Péguilhan and Laurent Deguillaume sampled the clouds.
506 Florent Rossi performed the ARGs monitoring, analyzed the results and produced the first
507 draft of the manuscript. Jean-Luc Baray performed the atmospheric back-trajectories
508 calculation and analysis. Marc Veillette and Nathalie Turgeon contributed significantly to the
509 design of the qPCR and provided key information in data treatment and in the understanding
510 of the results. All authors contributed significantly to the writing and the reviewing of the
511 manuscript.

512

513

514

515

516

517

518

519

520

521

522

523 **References**

- 524 1. Murray, C. J. *et al.* Global burden of bacterial antimicrobial resistance in 2019: a systematic
525 analysis. *The Lancet* (2022).
- 526 2. O’Neill, J. Tackling drug-resistant infections globally: final report and recommendations. (2016).
- 527 3. Larsen, J. *et al.* Emergence of methicillin resistance predates the clinical use of antibiotics. *Nature*
528 **602**, 135–141 (2022).
- 529 4. World Health Organization. *World health statistics 2018: monitoring health for the SDGs,*
530 *sustainable development goals.* (World Health Organization, 2018).
- 531 5. Lien, L. T. Q. *et al.* Antibiotics in wastewater of a rural and an urban hospital before and after
532 wastewater treatment, and the relationship with antibiotic use—a one year study from Vietnam.
533 *International Journal of Environmental Research and Public Health* **13**, 588 (2016).
- 534 6. Singh, R., Singh, A. P., Kumar, S., Giri, B. S. & Kim, K.-H. Antibiotic resistance in major rivers in the
535 world: a systematic review on occurrence, emergence, and management strategies. *Journal of*
536 *Cleaner Production* **234**, 1484–1505 (2019).
- 537 7. Hernandez, F. *et al.* Occurrence of antibiotics and bacterial resistance in wastewater and sea
538 water from the Antarctic. *Journal of hazardous materials* **363**, 447–456 (2019).
- 539 8. Tahrani, L., Van Loco, J., Ben Mansour, H. & Reyns, T. Occurrence of antibiotics in pharmaceutical
540 industrial wastewater, wastewater treatment plant and sea waters in Tunisia. *Journal of water*
541 *and Health* **14**, 208–213 (2016).
- 542 9. Thiele-Bruhn, S. Pharmaceutical antibiotic compounds in soils—a review. *Journal of plant nutrition*
543 *and soil science* **166**, 145–167 (2003).
- 544 10. ANSES. Antibiorésistance et environnement État et causes possibles de la contamination des
545 milieux en France. (2020).
- 546 11. Blair, J., Webber, M. A., Baylay, A. J., Ogbolu, D. O. & Piddock, L. J. Molecular mechanisms of
547 antibiotic resistance. *Nature reviews microbiology* **13**, 42–51 (2015).

- 548 12. Burrows, S. M. *et al.* Bacteria in the global atmosphere—Part 2: Modeling of emissions and
549 transport between different ecosystems. *Atmospheric Chemistry and Physics* **9**, 9281–9297
550 (2009).
- 551 13. Smith, D. J. *et al.* Intercontinental Dispersal of Bacteria and Archaea by Transpacific Winds. *Appl*
552 *Environ Microbiol* **79**, 1134–1139 (2013).
- 553 14. Blanchard, D. C. & Syzdek, L. D. Water-to-Air Transfer and Enrichment of Bacteria in Drops from
554 Bursting Bubbles. *Appl Environ Microbiol* **43**, 1001–1005 (1982).
- 555 15. Burrows, S. M., Elbert, W., Lawrence, M. G. & Pöschl, U. Bacteria in the global atmosphere—Part
556 1: Review and synthesis of literature data for different ecosystems. *Atmospheric Chemistry and*
557 *Physics* **9**, 9263–9280 (2009).
- 558 16. Joung, Y. S., Ge, Z. & Buie, C. R. Bioaerosol generation by raindrops on soil. *Nat Commun* **8**,
559 14668 (2017).
- 560 17. Smith, D. J. *et al.* Airborne bacteria in earth’s lower stratosphere resemble taxa detected in the
561 troposphere: Results from a new NASA Aircraft Bioaerosol Collector (ABC). *Frontiers in*
562 *Microbiology* 1752 (2018).
- 563 18. Péguilhan, R. *et al.* Rainfalls sprinkle cloud bacterial diversity while scavenging biomass. *FEMS*
564 *microbiology ecology* **97**, fiab144 (2021).
- 565 19. Amato, P. *et al.* Active microorganisms thrive among extremely diverse communities in cloud
566 water. *PLoS ONE* **12**, e0182869 (2017).
- 567 20. George, P. B. L. *et al.* Antimicrobial Resistance in the Environment: Towards Elucidating the Roles
568 of Bioaerosols in Transmission and Detection of Antibacterial Resistance Genes. *Antibiotics* **11**,
569 974 (2022).
- 570 21. Destoumieux-Garzón, D. *et al.* The One Health Concept: 10 Years Old and a Long Road Ahead.
571 *Front. Vet. Sci.* **5**, 14 (2018).
- 572 22. Robinson, T. P. *et al.* Antibiotic resistance is the quintessential One Health issue. *Trans R Soc Trop*
573 *Med Hyg* **110**, 377–380 (2016).

- 574 23. Šantl-Temkiv, T. *et al.* High-Flow-Rate Impinger for the Study of Concentration, Viability,
575 Metabolic Activity, and Ice-Nucleation Activity of Airborne Bacteria. *Environ. Sci. Technol.* **51**,
576 11224–11234 (2017).
- 577 24. Amato, P. *et al.* Metatranscriptomic exploration of microbial functioning in clouds. *Sci Rep* **9**,
578 4383 (2019).
- 579 25. Deguillaume, L. *et al.* Classification of clouds sampled at the puy de Dôme (France) based on 10
580 yr of monitoring of their physicochemical properties. *Atmos. Chem. Phys.* **14**, 1485–1506 (2014).
- 581 26. Gorokhova, E. Effects of preservation and storage of microcrustaceans in RNAlater on RNA and
582 DNA degradation. *Limnology and Oceanography: Methods* **3**, 143–148 (2005).
- 583 27. Hoffmann, L. *et al.* From ERA-Interim to ERA5: the considerable impact of ECMWF’s next-
584 generation reanalysis on Lagrangian transport simulations. *Atmos. Chem. Phys.* **19**, 3097–3124
585 (2019).
- 586 28. Baray, J.-L. *et al.* Cézeaux-Aulnat-Opme-Puy De Dôme: a multi-site for the long-term survey of
587 the tropospheric composition and climate change. *Atmos. Meas. Tech.* **13**, 3413–3445 (2020).
- 588 29. Renard, P. *et al.* Classification of Clouds Sampled at the Puy de Dôme Station (France) Based on
589 Chemical Measurements and Air Mass History Matrices. *Atmosphere* **11**, 732 (2020).
- 590 30. Stedtfeld, R. D. *et al.* Primer set 2.0 for highly parallel qPCR array targeting antibiotic resistance
591 genes and mobile genetic elements. *FEMS Microbiology Ecology* **94**, fiy130 (2018).
- 592 31. Nijhuis, R. H. T. *et al.* Detection of the plasmid-mediated colistin-resistance gene *mcr-1* in clinical
593 isolates and stool specimens obtained from hospitalized patients using a newly developed real-
594 time PCR assay: Table 1. *J. Antimicrob. Chemother.* **71**, 2344–2346 (2016).
- 595 32. Gunawardana, A. & Konietzschke, F. Nonparametric multiple contrast tests for general
596 multivariate factorial designs. *Journal of Multivariate Analysis* **173**, 165–180 (2019).
- 597 33. Abdi, H. Partial least squares regression and projection on latent structure regression (PLS
598 Regression). *WIREs Comp Stat* **2**, 97–106 (2010).

- 599 34. Lê Cao, K.-A., Martin, P. G., Robert-Granié, C. & Besse, P. Sparse canonical methods for biological
600 data integration: application to a cross-platform study. *BMC Bioinformatics* **10**, 34 (2009).
- 601 35. Suzuki, R. & Shimodaira, H. Pvcust: an R package for assessing the uncertainty in hierarchical
602 clustering. *Bioinformatics* **22**, 1540–1542 (2006).
- 603 36. Kormos, D., Lin, K., Pruden, A. & Marr, L. C. Critical review of antibiotic resistance genes in the
604 atmosphere. *Environ. Sci.: Processes Impacts* **24**, 870–883 (2022).
- 605 37. Zhang, T. *et al.* Time-resolved spread of antibiotic resistance genes in highly polluted air.
606 *Environment International* **127**, 333–339 (2019).
- 607 38. Li, J. *et al.* Global Survey of Antibiotic Resistance Genes in Air. *Environ. Sci. Technol.* **52**, 10975–
608 10984 (2018).
- 609 39. Alcock, B. P. *et al.* CARD 2020: antibiotic resistome surveillance with the comprehensive
610 antibiotic resistance database. *Nucleic Acids Research* gkz935 (2019) doi:10.1093/nar/gkz935.
- 611 40. Cox, G. & Wright, G. D. Intrinsic antibiotic resistance: Mechanisms, origins, challenges and
612 solutions. *International Journal of Medical Microbiology* **303**, 287–292 (2013).
- 613 41. Smith, B. L. & King, M. D. Aerosolization triggers immediate antibiotic resistance in bacteria.
614 *Journal of Aerosol Science* **164**, 106017 (2022).
- 615 42. Ouyang, W.-Y., Huang, F.-Y., Zhao, Y., Li, H. & Su, J.-Q. Increased levels of antibiotic resistance in
616 urban stream of Jiulongjiang River, China. *Appl Microbiol Biotechnol* **99**, 5697–5707 (2015).
- 617 43. Guo, X. *et al.* Biofilms as a sink for antibiotic resistance genes (ARGs) in the Yangtze Estuary.
618 *Water Research* **129**, 277–286 (2018).
- 619 44. Zou, H.-Y. *et al.* Antibiotic resistance genes in surface water and groundwater from mining
620 affected environments. *Science of The Total Environment* **772**, 145516 (2021).
- 621 45. Rajasekar, A., Qiu, M., Wang, B., Murava, R. T. & Norgbey, E. Relationship between water quality,
622 heavy metals and antibiotic resistance genes among three freshwater lakes. *Environ Monit*
623 *Assess* **194**, 64 (2022).

- 624 46. Wang, J., Chu, L., Wojnárovits, L. & Takács, E. Occurrence and fate of antibiotics, antibiotic
625 resistant genes (ARGs) and antibiotic resistant bacteria (ARB) in municipal wastewater treatment
626 plant: An overview. *Science of The Total Environment* **744**, 140997 (2020).
- 627 47. Lu, J. *et al.* Occurrence and spatial distribution of antibiotic resistance genes in the Bohai Sea and
628 Yellow Sea areas, China. *Environmental Pollution* **252**, 450–460 (2019).
- 629 48. Tignat-Perrier, R. *et al.* Seasonal shift in airborne microbial communities. *Science of The Total*
630 *Environment* **716**, 137129 (2020).
- 631 49. Gwenzi, W., Shamsizadeh, Z., Gholipour, S. & Nikaeen, M. The air-borne antibiotic resistome:
632 Occurrence, health risks, and future directions. *Science of The Total Environment* **804**, 150154
633 (2022).
- 634 50. Cuadrat, R. R. C., Sorokina, M., Andrade, B. G., Goris, T. & Dávila, A. M. R. Global ocean resistome
635 revealed: Exploring antibiotic resistance gene abundance and distribution in TARA Oceans
636 samples. *GigaScience* **9**, giaa046 (2020).
- 637 51. Qian, X. *et al.* Metagenomic analysis reveals the shared and distinct features of the soil resistome
638 across tundra, temperate prairie, and tropical ecosystems. *Microbiome* **9**, 108 (2021).
- 639 52. Rodríguez-Martínez, J. M. *et al.* Plasmid-mediated quinolone resistance: Two decades on. *Drug*
640 *Resistance Updates* **29**, 13–29 (2016).
- 641 53. Yamane, K. *et al.* New Plasmid-Mediated Fluoroquinolone Efflux Pump, QepA, Found in an
642 *Escherichia coli* Clinical Isolate. *Antimicrob Agents Chemother* **51**, 3354–3360 (2007).
- 643 54. Brandis, G., Gockel, J., Garoff, L., Guy, L. & Hughes, D. Expression of the *qepA1* gene is induced
644 under antibiotic exposure. *Journal of Antimicrobial Chemotherapy* **76**, 1433–1440 (2021).
- 645 55. Fàbrega, A., Madurga, S., Giralt, E. & Vila, J. Mechanism of action of and resistance to
646 quinolones: Mechanism of action of and resistance to quinolones. *Microbial Biotechnology* **2**,
647 40–61 (2009).
- 648 56. European Food Safety Authority, European Centre for Disease Prevention and Control, &
649 European Medicines Agency. *Antimicrobial consumption and resistance in bacteria from humans*

- 650 *and animals : third joint inter-agency report on integrated analysis of antimicrobial agent*
651 *consumption and occurrence of antimicrobial resistance in bacteria from humans and food-*
652 *producing animals in the EU/EEA : JIACRA III 2016–2018.* (Publications Office, 2021).
- 653 57. Cattoir, V., Poirel, L., Aubert, C., Soussy, C.-J. & Nordmann, P. Unexpected Occurrence of
654 Plasmid-Mediated Quinolone Resistance Determinants in Environmental *Aeromonas* spp. *Emerg*
655 *Infect Dis* **14**, 231–237 (2008).
- 656 58. Aydin, S., Aydin, M. E., Ulvi, A. & Kilic, H. Antibiotics in hospital effluents: occurrence,
657 contribution to urban wastewater, removal in a wastewater treatment plant, and environmental
658 risk assessment. *Environ Sci Pollut Res* **26**, 544–558 (2019).
- 659 59. Wang, J. & Wang, S. Removal of pharmaceuticals and personal care products (PPCPs) from
660 wastewater: A review. *Journal of Environmental Management* **182**, 620–640 (2016).
- 661 60. European Medicines Agency. *Sales of veterinary antimicrobial agents in 31 European countries in*
662 *2019 and 2020: trends from 2010 to 2020.* (Publications Office, 2021).
- 663 61. European Centre for Disease Prevention and Control - ESAC-Net Database. European Centre for
664 Disease Prevention and Control. *European Centre for Disease Prevention and Control*
665 [https://www.ecdc.europa.eu/en/antimicrobial-consumption/surveillance-and-disease-](https://www.ecdc.europa.eu/en/antimicrobial-consumption/surveillance-and-disease-data/database)
666 [data/database](https://www.ecdc.europa.eu/en/antimicrobial-consumption/surveillance-and-disease-data/database) (2020).
- 667 62. European Medicines Agency - ESVAC Database. European Surveillance of Veterinary
668 Antimicrobial Consumption (ESVAC). *European Medicines Agency*
669 [https://www.ema.europa.eu/en/veterinary-regulatory/overview/antimicrobial-](https://www.ema.europa.eu/en/veterinary-regulatory/overview/antimicrobial-resistance/european-surveillance-veterinary-antimicrobial-consumption-esvac)
670 [resistance/european-surveillance-veterinary-antimicrobial-consumption-esvac](https://www.ema.europa.eu/en/veterinary-regulatory/overview/antimicrobial-resistance/european-surveillance-veterinary-antimicrobial-consumption-esvac) (2018).
- 671 63. Lu, L. *et al.* Occurrence and Distribution of Tetracycline Antibiotics and Resistance Genes in
672 Longshore Sediments of the Three Gorges Reservoir, China. *Front. Microbiol.* **9**, 1911 (2018).
- 673 64. Woegerbauer, M., Bellanger, X. & Merlin, C. Cell-Free DNA: An Underestimated Source of
674 Antibiotic Resistance Gene Dissemination at the Interface Between Human Activities and

- 675 Downstream Environments in the Context of Wastewater Reuse. *Front. Microbiol.* **11**, 671
676 (2020).
- 677 65. Marti, E., Variatza, E. & Balcázar, J. L. Bacteriophages as a reservoir of extended-spectrum β -
678 lactamase and fluoroquinolone resistance genes in the environment. *Clinical Microbiology and*
679 *Infection* **20**, O456–O459 (2014).
- 680 66. Chen, M.-L. *et al.* Viral Community and Virus-Associated Antibiotic Resistance Genes in Soils
681 Amended with Organic Fertilizers. *Environ. Sci. Technol.* **55**, 13881–13890 (2021).
- 682 67. Subirats, J., Sánchez-Melsió, A., Borrego, C. M., Balcázar, J. L. & Simonet, P. Metagenomic
683 analysis reveals that bacteriophages are reservoirs of antibiotic resistance genes. *International*
684 *Journal of Antimicrobial Agents* **48**, 163–167 (2016).
- 685 68. Debroas, D. & Siguret, C. Viruses as key reservoirs of antibiotic resistance genes in the
686 environment. *ISME J* **13**, 2856–2867 (2019).
- 687 69. Joly, M. *et al.* Survival of microbial isolates from clouds toward simulated atmospheric stress
688 factors. *Atmospheric Environment* **117**, 92–98 (2015).
- 689 70. Wirgot, N. *et al.* Metabolic modulations of *Pseudomonas graminis* in response to H₂O₂ in cloud
690 water. *Sci Rep* **9**, 12799 (2019).
- 691 71. Wirgot, N., Vinatier, V., Deguillaume, L., Sancelme, M. & Delort, A.-M. H₂O₂ modulates the
692 energetic metabolism of the cloud microbiome. *Atmos. Chem. Phys.* **17**, 14841–14851 (2017).
- 693 72. Pruppacher, H. R. & Jaenicke, R. The processing of water vapor and aerosols by atmospheric
694 clouds, a global estimate. *Atmospheric Research* **38**, 283–295 (1995).

695

696

697

698

699

700

701

Quantification of Antibiotic Resistance Genes (ARGs) in Clouds at a Mountain Site (puy de Dôme, Central France)

Tables and Figures

702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718

Table 1: Concentrations of antimicrobial resistance genes (ARGs) and bacteria cells in cloud water (mL⁻¹ of water) in the 12 cloud events investigated. The concentrations of the main families of ARGs are displayed along with their means (\pm SEM; n=12) and median values. Statistical differences between families of ARGs are highlighted by the subscript (a > b > c > d, pairwise comparison, $P < 0.05$). *N.D.* = not detected (Ct > 37) and *N.A.* = not acquired (missing value).

Cloud sampling date	Quinolones	Aminoglycosides	Macrolides	Sulfonamide	Tetracycline	β -Lactamase	Vancomycin	Transposase	Total ARGs	Total Bacteria
Gene copies ml ⁻¹ of cloud water										
	Cell ml ⁻¹ of cloud water									
22-Oct-19	1.46 10 ³	2.69 10 ⁻¹	7.21 10 ¹	6.19 10 ²	4.96 10 ²	6.17 10 ²	3.40 10 ²	1.57 10 ¹	3.62 10 ³	3.32 10 ⁴
11-Mar-20	1.40 10 ⁴	4.78 10 ⁻¹	2.14 10 ²	9.40 10 ²	8.44 10 ²	2.92 10 ⁻¹	1.02 10 ³	<i>N.D.</i>	1.70 10 ⁴	3.18 10 ³
17-Jul-20	1.81 10 ⁴	2.6	1.21 10 ⁴	2.30 10 ³	1.19 10 ³	1.19 10 ¹	2.06 10 ³	<i>N.D.</i>	3.58 10 ⁴	<i>N.A.</i>
16-Oct-20	1.71 10 ³	<i>N.D.</i>	5.81 10 ¹	<i>N.D.</i>	2.43 10 ¹	3.50 10 ³	8.50 10 ²	<i>N.D.</i>	6.14 10 ³	7.70 10 ³
28-Oct-20	9.17 10 ³	9.94 10 ⁻¹	1.82 10 ²	9.39 10 ²	9.58 10 ²	6.79 10 ¹	1.76 10 ³	<i>N.D.</i>	1.31 10 ⁴	9.96 10 ³
03-Nov-20	4.90 10 ⁴	4.67	1.18 10 ²	1.67 10 ³	2.24 10 ³	4.80 10 ³	5.64 10 ³	3.65	6.35 10 ⁴	1.06 10 ⁴
10-Nov-20	2.18 10 ⁴	3.3	2.95 10 ⁴	1.29 10 ³	4.42 10 ²	4.07	1.16 10 ⁴	<i>N.D.</i>	6.47 10 ⁴	5.53 10 ³
19-Nov-20	5.74 10 ³	1.78 10 ⁻¹	8.00 10 ¹	7.55 10 ²	1.27 10 ²	1.19 10 ³	1.75 10 ³	4.43	9.65 10 ³	3.33 10 ²
28-Apr-21.	2.69 10 ²	6.14 10 ⁻²	<i>N.D.</i>	2.24 10 ³	9.39 10 ²	<i>N.D.</i>	<i>N.D.</i>	2.60 10 ⁻¹	3.45 10 ³	1.33 10 ⁴
06-May-21	5.20 10 ³	6.33 10 ⁻²	<i>N.D.</i>	<i>N.D.</i>	2.05 10 ²	<i>N.D.</i>	4.90 10 ²	<i>N.D.</i>	5.89 10 ³	1.19 10 ³
13-Oct-21	1.32 10 ⁴	3.36 10 ⁻¹	7.73 10 ²	3.97 10 ³	6.32 10 ²	<i>N.D.</i>	1.23 10 ³	6.12	1.98 10 ⁴	1.36 10 ³
26-Oct-21	3.00 10 ³	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>	8.23 10 ¹	3.58 10 ³	<i>N.D.</i>	<i>N.D.</i>	6.67 10 ³	1.63 10 ³
Median	7.45 10 ³ a	3.03 10 ⁻¹ cd	9.91 10 ¹ bc	9.39 10 ² bc	5.64 10 ² b	3.99 10 ¹ bc	1.12 10 ³ b	0 d	1.14 10 ⁴	5.53 10 ³
Mean \pm SEM	1.19 10 ⁴ \pm 3.93 10 ³	1.08 \pm 4.52 10 ⁻¹	3.60 10 ³ \pm 2.56 10 ³	1.23 10 ³ \pm 3.39 10 ²	6.82 10 ² \pm 1.80 10 ²	1.15 10 ³ \pm 5.08 10 ²	2.23 10 ³ \pm 9.54 10 ²	2.51 \pm 1.35	2.08 10 ⁴ \pm 6.40 10 ³	8.00 10 ³ \pm 2.85 10 ³

719
720
721
722

723 Table 2: Proportions of the main families of antibiotics used over Europe (29 countries,
724 human and veterinary usages; from JIACRA III (2021), and relative abundances observed in
725 clouds (this study).

726

Families of Antibiotics	Proportion of antibiotic usages in Europe		Relative abundance in clouds (this study)					
	Human Health	Food-producing animals	With <i>qepA</i>			Without <i>qepA</i>		
			Continental clouds	Marine clouds	All clouds	Continental clouds	Marine clouds	All clouds
β -lactamases	54.1%	24.3%	14.8%	10.6%	12.3%	21.8%	21.4%	21.6%
Quinolones	5.9%	2.6%	32.0%	69.8%	54.0%	0.7%	3.7%	2.4%
Sulfonamides	3.5%	9.1%	18.1%	6.2%	11.1%	22.9%	20.5%	21.5%
Glycopeptides (vancomycin)	< 7%	< 1%	9.4%	8.7%	8.9%	14.6%	35.8%	26.3%
macrolides	6.1%	7.4%	16.5%	1.0%	7.5%	28.2%	3.7%	13.9%
tetracyclines	2.4%	30.5%	9.0%	3.6%	5.9%	12.1%	15.5%	14.0%
aminoglycosides	0.2%	4.6%	< 0.1%	< 0.1%	< 0.1%	< 0.1%	< 0.1%	< 0.1%

727

728

729

730

731 Table 3: Global estimates of ARGs in clouds. Data for clouds (upper part of the Table) are
732 estimates from Pruppacher and Jaenicke (1995).


733

	Average	min	max	Unit
Volume fraction of clouds in the first 6 km	15.53	-	-	%
Volume of the first 6 km of the atmosphere	$3.00 \cdot 10^{18}$	-	-	m^3
Total volume of clouds in the first 6 km	$4.66 \cdot 10^{17}$	-	-	m^3
Cloud precipitation rate	$4.10 \cdot 10^{20}$	-	-	$\text{m}^3 \text{ year}^{-1}$
ARGs concentration in clouds	$5.42 \cdot 10^3$	$1.01 \cdot 10^3$	$1.61 \cdot 10^4$	copies m^{-3}
ARGs transiting in clouds	$2.53 \cdot 10^{21}$	$4.71 \cdot 10^{20}$	$7.51 \cdot 10^{21}$	copies
Total ARGs transiting in clouds every year	$6.94 \cdot 10^{25}$	$1.29 \cdot 10^{25}$	$2.06 \cdot 10^{26}$	copies year^{-1}
Average precipitation rates of ARGs	$2.22 \cdot 10^{24}$	$4.14 \cdot 10^{24}$	$6.61 \cdot 10^{24}$	copies year^{-1}

734

735 Figure 1: Concentrations of the different families of ARGs in the 12 clouds (A) and their
 736 relative contributions to the pool of total ARGs (B).


737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 760
 761
 762
 763
 764
 765



766 Figure 2: Total concentration of ARGs and bacteria in the 12 clouds (left axis; dots / squares
 767 and lines) and the estimated numbers of ARGs normalized to that of bacterial cells (right axis;
 768 grey bars). An ARGs per bacterial cells ratio of 1 is indicated by the dotted line. *N.A.* = not
 769 available (missing value).

770

771



772

773

774

775

776

777

778

779

780

781

782

783 Figure 3: Partial least square regression (pls-r) plot of the 12 cloud events based on
 784 environmental variables. **(A)** Representation of individual clouds, where groups of samples
 785 classified as “Continental” (red), “Marine high” (blue) and “Marine low” (grey) based on
 786 backward trajectories are highlighted by ellipses at 95% confidence level. **(B)** Correlation plot
 787 of the pls-r: predictors (origin and altitude of the air mass in the 72-hours preceding sampling)
 788 are represented in black, and responses (families of ARGs) in blue.

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

