



**HAL**  
open science

## Chiral acoustoplasmonics

Edson Cardozo de Oliveira, Beatriz Castillo López de Larrinzar, Chushuang Xiang,  
N. D. Lanzillotti-Kimura, Antonio García-Martín

► **To cite this version:**

Edson Cardozo de Oliveira, Beatriz Castillo López de Larrinzar, Chushuang Xiang, N. D. Lanzillotti-Kimura, Antonio García-Martín. Chiral acoustoplasmonics. 17th International Conference on Phonon Scattering in Condensed Matter, Phonons 2023, Jul 2023, Paris, France. <hal-04350333>

**HAL Id: hal-04350333**

**<https://cnrs.hal.science/hal-04350333v1>**

Submitted on 18 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# Chiral acoustoplasmonics

Edson Rafael Cardozo de Oliveira<sup>a</sup>

B. Castillo López de Larrinzar<sup>b</sup>, C. Xiang<sup>a</sup>, N. D. Lanzillotti-Kimura<sup>a</sup>, A. García-Martín<sup>b</sup>

*a. Université Paris-Saclay, CNRS, Centre de Nanosciences et de Nanotechnologies, 10 Boulevard Thomas Gobert, 91120 Palaiseau, France*

*b. Instituto de Micro y Nanotecnología IMN-CNM, CSIC, CEI UAM + CSIC, Isaac Newton 8, Tres Cantos, Madrid 28760, Spain*

[edson.cardozo@c2n.upsaclay.fr](mailto:edson.cardozo@c2n.upsaclay.fr)

The possibility of creating and manipulating nanostructured materials encouraged the exploration of new strategies to control electromagnetic properties. Among the most intriguing nanostructures are those that respond differently to helical polarization, i.e., exhibit chirality [1,2]. Here, we present a simple structure based on crossed elongated nanobars with the long axis parallel to the polarization plane forming a given angle between them and vertically aligned, as shown in Fig. 1(a). We theoretically demonstrate that, in such structure, light-handedness defines the dominating cross-section absorption or scattering, with a 200% difference from its counterpart (scattering or absorption). The proposed chiral system opens the way to enhanced coherent phonon excitation and detection. In general, the generation of acoustic phonons is optimized by maximizing the absorption, while the detection is enhanced by engineering the scattering properties [3,4]. In chiral structures, this could be achieved by a simple pump-probe experiment using circularly polarized light at the same wavelength. The use of structures sustaining chiral optical modes appears as a natural, yet unexplored strategy to address these conditions. These results constitute one of the first steps towards harvesting chirality effects in the design and optimization of efficient and versatile acoustoplasmonic transducers.

## References


[1] Rodríguez-Álvarez, J., García-Martín, A., Fraile Rodríguez, A., Batlle, X. & Labarta, A. Tunable circular dichroism through absorption in coupled optical modes of twisted triskelia nanostructures. *Sci. Rep.* 12, 26 (2022).

[2] Auguie, B., Alonso-Gómez, J. L., Guerrero-Martínez, A. & Liz-Marzán, L. M. Fingers Crossed: Optical Activity of a Chiral Dimer of Plasmonic Nanorods. *J. Phys. Chem. Lett.* 2, 846–851 (2011).

[3] Lanzillotti-Kimura, N. D. et al. Polarization-controlled coherent phonon generation in acoustoplasmonic metasurfaces. *Phys. Rev. B* 97, 235403 (2018).

[4] O'Brien, K. et al. Ultrafast acousto-plasmonic control and sensing in complex nanostructures. *Nat. Commun.* 5, 4042 (2014).

## Figures



**Figure 1:** (a) Sketch of the geometrical layout of the chiral structure. Two gold nanobars separated by a distance  $D = 80$  nm, with the long axis parallel to the polarization plane forming a given angle between them and vertically aligned. The bottom bar is embedded in SiO<sub>2</sub> while the top bar sits on the surface of SiO<sub>2</sub>. (b) Absorption and scattering cross-sections for right (RCP) and left (LCP) circularly polarized incident beam.