



HAL
open science

The Winding Road from Origin to Emergence (of Life)

Wolfgang Nitschke, Orion Farr, Nil Gaudu, Chloé Truong, François Guyot,
Michael J Russell, Simon Duval

► **To cite this version:**

Wolfgang Nitschke, Orion Farr, Nil Gaudu, Chloé Truong, François Guyot, et al.. The Winding Road from Origin to Emergence (of Life). *Life*, 2024, 14 (5), pp.607. 10.3390/life14050607 . hal-04573011

HAL Id: hal-04573011

<https://cnrs.hal.science/hal-04573011>

Submitted on 13 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2 The Winding Road from Origin to Emergence (of Life)

3 Wolfgang Nitschke ¹, Orion Farr ^{1,2}, Nil Gaudu ¹, Chloé Truong ¹, François Guyot ³, Michael J. Russell ⁴
4 and Simon Duval ¹

5 ¹ BIP (UMR 7281), CNRS, Aix-Marseille-Univ., 13009 Marseille, France

6 ² CINaM, CNRS, Aix-Marseille-Univ., 13009 Marseille, France

7 ³ IMPMC (UMR 7590), Sorbonne Univ., CNRS, Paris, France

8 Dipartimento di Chimica, Università degli studi di Torino, Torino, Italy

9 * Correspondence: e-mail@e-mail.com; Tel.: (optional; include country code; if there are multiple
10 corresponding authors, add author initials)

11 **Abstract:** Humanity's strive to understand why and how life appeared on planet Earth dates back
12 to prehistoric times. At the beginning of the 19th century, empirical biology started to tackle this
13 question yielding both Charles Darwin's Theory of Evolution and the paradigm that the crucial
14 trigger putting life on its tracks was the appearance of organic molecules. In parallel to these
15 developments in the biological sciences, physics and physical chemistry saw the fundamental laws
16 of thermodynamics being unraveled. Towards the end of the 19th century and during the first half
17 of the 20th century, the tensions between thermodynamics and the "organic-molecules-paradigm"
18 became increasingly difficult to ignore, culminating in Erwin Schrödinger's 1944 formulation of a
19 thermodynamics-compliant vision of life and, consequently, the prerequisites for its appearance.
20 We will first review the major milestones over the last 200 years in the biological and the physical
21 sciences, relevant to making sense of life and its origins and then discuss the more recent
22 reappraisal of the relative importance of metal ions *vs.* organic molecules in performing the
23 essential processes of a living cell. Based on this reassessment and the modern understanding of
24 biological free energy conversion (aka bioenergetics), we consider that scenarios wherein life
25 emerges from an abiotic chemiosmotic process are both thermodynamics-compliant and the most
26 parsimonious proposed so far.

27 **Keywords:** keyword 1; keyword 2; keyword 3 (List three to ten pertinent keywords specific to the
28 article yet reasonably common within the subject discipline.)
29

30 1. The Prevailing Paradigm: It's the Organics, Stupid!

31 Judging from the communications of major research agencies all over the world
32 which directly or indirectly tackle the question of life's origins, the most fundamental
33 prerequisite for life to appear has been identified: the presence of organic molecules such
34 as amino acids, nucleobases, lipids and sugars. The actual open questions pertain to the
35 precise synthesis pathways for these organics and the processes which then have
36 allowed them to evolve into living cells. The introduction to NASA's Astrobiology
37 Strategy 2015 roadmap for example reads: "A major goal of research on this topic in
38 astrobiology is to understand how the abiotic (non-biological) production of small
39 molecules led to the production of large and more complex molecules, prebiotic
40 chemistry, and the origin of life on Earth" [1]. Obedient to this instruction, all recent
41 space missions had an eye on detecting organics, the "seeds of life". They intend to
42 "understand the role asteroids may have played in delivering life-forming compounds
43 to Earth" [2], "to investigate questions ... where the organic matter that forms life
44 originally came from" [3] and thus to "grant us direct access to some of the ingredients
45 that likely ended up in the prebiotic soup that eventually resulted in the origin of life on
46 Earth" [4]. This focus on the detection of organic molecules in the solar system appears

47 natural if “we begin with the premise that life emerged on Earth from chemistry that led
48 to the synthesis of molecular building blocks, which, in turn, self-assembled to form
49 cells” [5].

50 The above listed citations of course resonate with the conviction expressed more
51 than 150 years ago by Charles Darwin in his oft-cited correspondence with Joseph
52 Dalton Hooker [6]: “But if (and oh what a big if) we could conceive in some warm little
53 pond ..., that a protein compound was chemically formed, ready to undergo still more complex
54 changes, ...” (emphasis added by the authors). As we will see below, Darwin’s assurance
55 that it is the chemically formed “protein compounds” that would “readily” (that is,
56 spontaneously) transform into more complex structures and eventually bring forth life,
57 is deeply rooted in and informed by the scientific/intellectual background of his century.
58 As evidenced by the citations above, Darwin’s conviction is as unswerving today as it
59 was in his days, despite many scientific paradigms having substantially changed at the
60 turn from the 19th to the 20th century and even more so during the 2nd half of the 20th
61 century, as we will discuss in this article. While, as we will argue, Darwin has merely
62 expressed the scientific paradigm of his days rather than having put profound
63 intellectual efforts into coming up with this clause in a letter to a colleague, the search
64 for the “chemically formed protein compounds” nevertheless turned into the
65 overarching roadmap (up to the present day) for understanding the appearance of life
66 on our planet. The decisive big leap forward following the direction suggested by this
67 roadmap obviously was Stanley Miller’s demonstration that organic molecules (mainly
68 amino acids) are generated if suitable mixtures of inorganic gases (H₂, CO₂, CH₄) and
69 solutes (mainly ammonia in water) were exposed to electrical discharges [7, 8]. Several
70 years later, the realization that the initial conditions underlying the Miller/Urey
71 experiments were inappropriate for the early Earth (assumed by geologists a century
72 before) sparked the search for other pre-biotic synthesis pathways for organics
73 (preferably amino acids, nucleobases and lipids, the predominant ingredients of extant
74 life on Earth). 70 years of post-Miller/Urey efforts to uncover such pathways have led to
75 a veritable zoo of chemical reaction schemes applicable to a large number of putative
76 initial environmental conditions on the primordial planet and beyond (Fig. 1). Most of
77 these generally follow the tracks of the Miller/Urey experiment in that they involve
78 standard chemical reactions under Earth-relevant conditions [5, 9, 10] while a few appeal
79 to more exotic schemes such as involving high-energy protons from Coronal Mass
80 Ejection (CME) events [11] or ionizing radiation from natural nuclear reactors [12].
81 However, organic molecules can obviously also be formed under even more
82 “outlandish” conditions such as on comets, asteroids or in the interstellar medium as
83 evidenced by the space missions mentioned above. For example, certain amino acids
84 have been detected on comets [13], the nucleobases used by extant life have been found
85 in meteorites [14] and in samples returned to Earth from asteroid Ryugu during JAXA’s
86 Hayabusa-2 mission [15, 16] and polycyclic aromatic hydrocarbons found on Ryugu
87 were identified as having been formed in the interstellar medium and further processed
88 on the asteroid [17].

89 There therefore is no shortage of potential pre-biotic synthesis pathways for organic
90 molecules and the predominant argument in **origin-of-life** research presently focuses on
91 which of these pathways is the most pertinent one to the actual birth of life on our planet
92 and whether these organics were produced *in-situ* (*i.e.* on Earth) in the atmosphere, the
93 hydrosphere or the lithosphere, or whether they were delivered from space (Fig. 1).




Figure 1. Diverse derivations of the prevailing paradigm for how life came into being on planet Earth. As discussed in the text, this paradigm holds that the presence of an ensemble of organic molecules is the necessary and sufficient condition for life to appear. When the first abiotic synthesis pathway for organics was discovered by Stanley Miller in the 1950s (left side of the figure), the question of life’s origin was considered fundamentally solved. In the meantime, a plethora of different ways to abiotically generate organics were found (only a few representatives of which are shown in this figure) and the detection of organic molecules in space even opened up the possibility that these “building blocks for life” may have been delivered from extraterrestrial sources. On the background of the mentioned paradigm, the debate presently mainly turns around the question of sorting out which specific pre-biotic synthesis pathway did indeed lead to life on Earth.


This tacitly eschews explaining why organic molecules would spontaneously “evolve” into ever-complexifying systems, finally reaching a state which can be recognized as “living”, while no other type of matter seemed able to pull off this feat. The narrative of “organic molecules engendering life” has become so deeply ingrained in the mindset of both scholars and the general public during the second half of the 20th century that no questioning seemed appropriate. Incidentally, the same appears to be the case for the topic “life” in general, where immense progress in microbiology, biochemistry and genetics since the 1950s uncovered the “hows” of life’s mechanisms obscuring the question of why life does what it does. In striking contrast, from the early 1800s into the early 20th century, scholars were painfully aware of life’s apparent antagonism to the behaviour of the physical world.

2. Physics vs. Biology; from Sadi Carnot to Erwin Schrödinger and from Jöns Jacob Berzelius to Stanley Miller

These were the times when thermodynamics was elaborated, morphing from Sadi Carnot’s considerations of the efficiency limits of heat engines in 1824 through Rudolf Clausius’ and William Thomson’s empirical formulations of the 2nd law of thermodynamics in the middle of the 19th century into the statistical rationalization of the 2nd law initiated by J.W. Gibbs and definitively conceptualized by Ludwig Boltzmann during the 2nd half of the 19th century (for a timeline, see Fig. 2). One of the major take-home messages first formulated by Thomson and later further developed by von Helmholtz and William Rankine was the notion of the ultimate “heat death” of the universe. The 2nd law of thermodynamics indeed predicts an inevitable degradation over time of the free energy available to do work, a conclusion later rationalized by trajectories of physical systems from less to more probable states in Boltzmann’s statistical mechanics. 19th century scholars therefore were fully aware that physical systems tend towards increasing disorder while, in striking contrast, life seemed to do exactly the opposite. The antithetical behaviours of the physical world on one side and

133
134

life on the other obliged 19th century biologists to come up with novel hypotheses attempting to resolve this cognitive dissonance.



135

136

137

138

139

140

141

142

143

144

Figure 2. Timeline (from the beginning of the 19th century to the present day) of the development of the major concepts in the fields of thermodynamics (top part of the figure above the time-arrow) and in the biological/chemical sciences dealing with life and its birth on our planet (bottom part below the time-arrow). Up until the middle of the 20th century, these developments occurred with little or no cross-talk between the physical and the biological disciplines, reflecting the perceived incompatibilities of observed behaviours. Since Schrödinger's venture into the life sciences [26] (although this topic was already touched upon by Boltzmann), the barrier between thermodynamics and biology was understood to be artificial later culminating in the rationalization of life as a far-from-equilibrium dissipative structure [28, 29, 54, 55].

145

146

147

148

149

150

151

152

153

154

155

2.1. The Century of Vitalism (~1800 to ~1900)

It seems counterintuitive from our present perspective that the year 1859 saw both the publication of Louis Pasteur's refutation of spontaneous generation and Charles Darwin's "On the origin of species". Pasteur's brilliant series of experiments **severely challenged** the millenia-old notion, famously expressed by Aristotle as early as the 4th century BCE (footnote 1)¹, that life can spontaneously emerge from inanimate matter. Thus Pasteur held that life could only derive from life while Darwin's theory of evolution saw the extant diversity and complexity of ecosystems emerge from ever simpler organisms finally arriving at a point in history when life rose out of an inanimate world. In our days, Pasteur's and Darwin's views appear to represent irreconcilable opposites. Not so in the 19th century! Nascent organic chemistry in the late

¹ So with animals, some spring from parent animals according to their kind, whilst others grow spontaneously and not from kindred stock; and of these instances of spontaneous generation some come from putrefying earth or vegetable matter, as is the case with a number of insects, while others are spontaneously generated in the inside of animals out of the secretions of their several organs. [18]

1800s and early 1900s was strongly influenced by (but also intensely struggling with) the concept of “vitalism”. Chemists in those days seemingly observed an insurmountable barrier between inorganic and organic molecules, the latter considered to only being produced by living organisms. This led to the notion that organic molecules must be endowed with some ill-defined “vital force” which enables them to bring forth life and, *vice versa*, require organisms for their synthesis. Jöns Jacob Berzelius, often cited as the father of organic chemistry at the beginning of the 19th century, pleaded the cause of vitalism for the major part of his scientific life, and continued to do so even after Friedrich Wöhler, one of his former students, had shown in 1828 that the organic molecule urea could be synthesized from inorganic precursors (Fig. 2). Pasteur himself considered that the experiments disproving spontaneous generation provided strong support for vitalism. Although many scientists (such as Justus von Liebig and Berzelius himself) felt intense unease concerning the “mystical” aspects of the vital force, they nevertheless stuck to vitalism for lack of a better rationalization of life’s perceived anti-thermodynamic behaviour. Darwin’s idea that “protein compounds” might be synthesized from inorganic sources, just like Wöhler’s urea, and then spontaneously further evolve into living matter thanks to the vital force inherent to organics therefore was all but outrageous within the mindset of vitalism which strongly influenced chemistry’s and biology’s thinking during the major part of the 19th century.

The development of an atomistic understanding of molecules, initiated by the concept of chemical structure proposed independently by Friedrich August Kekulé and Archibald Scott Couper in 1859 and culminating in the discovery of the electron and its role in chemical bonding at the turn from the 19th to the 20th century, did away with the fundamental dichotomy between inorganic and organic chemistry. Organic molecules were henceforth “just molecules” removing any scientific basis for assigning special life-inducing properties to hydrocarbons, as complex as they may come. Vitalism became increasingly disreputable, although a handful of biologists subscribed to vitalist principles well into the first decades of the 20th century [19-22]. Ernst Mayr finally buried vitalism as a biological hypothesis by declaring that “no biologist alive today would want to be classified as a vitalist” [23].

2.2. Organicism and Life’s Origin as a One-Off Event

The vital force of organics having become obsolete, a few scientists trying to rationalize life’s whereabouts turned to “Organicism” and the idea that the “organic whole” of specific assemblies of hydrocarbons must possess properties which cannot be explained by reductionist analysis of their constituents [24]. J.B.S. Haldane, together with Aleksandr Oparin considered as the fathers of the “prebiotic-soup-hypothesis” for the origin of life, *i.e.* the conceptual framework leading to the Miller/Urey experiments, strongly built his views of life and its origin on organicism. Shifting the life-enabling properties from the building blocks, the organic molecules, towards the “organismal whole” seemingly did away with the requirement for vitalist concepts since abiotically produced organics now “only” needed to accidentally assemble into first organisms which then allowed the “organismal” properties to kick in. Such accidental order-increasing (anti-entropic) trajectories were indeed conceivable in the framework of Boltzmann’s statistical interpretation of thermodynamics. Just as the probability-driven mixing of two distinguishable, previously compartmentalized, gases has a non-zero probability to, over time, transiently go through the unmixed state again, random mixtures of organic molecules might well accidentally form more ordered assemblies. Still, two comments impose themselves here:

(i) According to statistical thermodynamics, the probability of fluctuation-induced increase in order decreases steeply with complexity of these systems and the degree of complexity required to arrive at even the simplest imaginable cellular structures is doubtlessly enormous. A number of biologists around the middle of the 20th century were perfectly aware of the vanishingly small probability for life to arise this way by

209 mere chance prompting for example Jacques Monod to proclaim that "... man at last
210 knows that he is alone in the unfeeling immensity of the universe, out of which he has
211 emerged only by chance" [25]. It is in the framework of Monod's thinking that the term
212 "origin", as describing life's coming into being out of a soup of organics, unveils its
213 profound sense: a freak event which obviously did happen (since we know for sure that
214 there is life on Earth) but which was so unlikely that there is no point expecting it to
215 have happened elsewhere in the cosmos. Needless to say that many citations listed in
216 Section 1., while deeply rooted in the prebiotic soup paradigm, nonchalantly ignore the
217 inevitability of Monod's logic!

218 (ii) To the post-19th-century biologists, the notion of anti-entropic properties
219 inherent in organismal assemblies of organics may have been intuitively more
220 acceptable than the idea of a vitalist force in hydrocarbons. We do observe this anti-
221 entropic tendency in all living things, don't we? Still, the order-generating properties of
222 organismal assemblies as envisaged by the organicists are no less mystical than the vital
223 force and relying on such properties takes the phenomenon "life" out of the explanatory
224 framework of the physical sciences. In 1944, Erwin Schrödinger's book "What is Life?"
225 finally put an end to speculations about life's specialness by rigorously treating life as
226 part of the physical world [26].

227 **3. The Demystification of Life and Its Integration into the Physical Sciences**

228 *3.1. Schrödinger and the Crux of the 2nd Law*

229 As Schrödinger pointed out, the main misconception dooming all efforts to
230 understand life and its beginnings since the days of the early vitalists lay in considering
231 living things as independent entities, that is, as thermodynamically closed systems. That
232 this notion is fatally misguided will be brought home to anybody trying to escape
233 Earth's atmosphere without taking the proper supply of molecular oxygen. We "burn"
234 the food that we eat with the oxygen we breathe and it is the free energy available in this
235 reaction which allows us to live. This is true for all life on Earth and only the sources of
236 reductants and oxidants involved in this "life-enabling" reaction vary between species
237 [27]. To Schrödinger-the-physicist, this fact was a trivial but also inescapable
238 consequence of the 2nd law of thermodynamics. Entropy decrease in living beings
239 necessarily implies (a) that they are part of a larger system which itself is in a state of
240 thermodynamic disequilibrium and (b) that order-generation within the living system is
241 only allowed by the 2nd law if it takes the larger system closer to equilibrium. In
242 Schrödinger's words: Life has to "eat neg-entropy" [26]. Unsurprisingly, biologists were
243 unable to find even a single exception to the rule that life requires the input of free
244 energy from the environment in order to exist. Schrödinger's considerations therefore
245 have made the existence of living beings compliant with the laws of the physical world.
246 However, while we now know that environmental free energy rather than a mystical
247 vital force or organismal hat-tricks allows the generation of order by living systems, why
248 does life behave this way while abiotic systems don't? Or do they?

249 *3.2. Non-Living Systems Can Display Entropy-Decreasing Features; the End of the Life/Non- 250 Life-Dichotomy*

251 The world of physical phenomena is indeed replete with locally entropy-decreasing
252 processes. Such processes are observed from fluid dynamics (*e.g.* turbulence above a
253 critical Reynolds number, Taylor-Couette flows or Benard cells) through atmospheric
254 systems (from Kelvin-Helmholtz instabilities to large scale cyclones) to chemical reaction
255 schemes (for example the iconic Belousov-Zhabotinsky reaction). The common feature
256 of all such phenomena is that they arise within larger systems far from thermodynamic
257 equilibrium and that their local entropy decrease is constantly fed by the flux of free
258 energy taking the larger system closer to equilibrium.

259 This conversion of free energy (*i.e.* low entropy) of the larger system into entropy
260 decrease (ordering) in the subsystem ensures nominal compliance with the 2nd law but
261 *per se* doesn't explain how this conversion is accomplished. Progress in the theoretical
262 description of dynamic trajectories of systems far from thermodynamic equilibrium over
263 the last 50 years [28] together with significantly improved capabilities of numerical
264 simulations now provide an in-depth understanding of how the respective systems
265 under specific circumstances occasionally decrease entropy, *i.e.* generate spatially and/or
266 temporally ordered states [29].

267 The term “dissipative structures”, coined by Ilya Prigogine in the 1970s (Fig. 2) to
268 describe such states, emphasizes both the structuring of the subsystems and the free
269 energy dissipation in the larger, embedding ones. The appearance of a dissipative
270 structure therefore corresponds to an “emergence”, that is, a phenomenon not inherent
271 in the reductionist components of a system but a result of the thermodynamic forces
272 driving the system's trajectory [28]. The emergence of a dissipative structure therefore is
273 contingent on the free energy environment driving the system's dynamics and not on
274 what the system is made of!

275 Erwin Schrödinger showed that life, to comply with the 2nd law, must necessarily be
276 part of a larger system (the living entities plus their environment) which itself is in
277 thermodynamic disequilibrium. It is therefore inescapable to recognize life as an
278 emergent dissipative structure following the same general principles as the
279 physical/chemical systems described above. In this framework of thinking, the fact that
280 life appeared on planet Earth is first and foremost due to the thermodynamic
281 circumstances at the time and place of its birth and not to the reductionist properties of
282 its constituents. The somewhat mystical vital forces of organic molecules or the
283 supposed order-generating features of organismal assemblies finally find their
284 rationalization in the tendency of systems which are far from thermodynamic
285 equilibrium to form dissipative structures. Rather than life having “originated” as a
286 thermodynamically extremely unlikely event, modern understanding of far-from-
287 equilibrium thermodynamics suggests that we better refer to this phenomenon as a
288 thermodynamically-driven “emergence of life” as proposed previously [30, 31].

289 However, does this change of paradigm really deprive organic molecules of their
290 central importance in this process? Surely the environmental thermodynamic
291 disequilibria driving the emergence of life must have been exploited by organic-
292 molecule-based mechanisms, just as they are generally considered to be in extant life?

293 **4. Reassessing the Organo-Centric View of Living Entities**


294 *4.1. CHONPS vs. FeNiMoWCo+; Giant vs. Dwarf?*

295 The conviction that life is all about organic molecules is related to the even more
296 fundamental notion that the prerequisites for life on Earth come down to 6 relatively
297 light elements, affectionately called CHONPS (some authors prefer the easier to
298 pronounce term SPONCH). “Life as we know it is based on six elements, carbon (C),
299 hydrogen (H), oxygen (O), nitrogen (N), phosphorous (P), and sulphur (S) – typically
300 abbreviated as CHONPS” [32], or “A living cell consists of two basic kinds of polymers
301 ... composed of six elements abbreviated CHONPS” [33]. Not only are the CHONPSes
302 considered the compositional bricks of life, they also seem to be responsible for life's
303 defining processes, since “Biochemistry on Earth makes use of the key elements ... (or
304 CHONPS)” [34]. Indeed, per dry weight of living matter, 96% are organic molecules and
305 3 % are light-atomic-weight ions such as sodium, potassium, chloride etc. It therefore is
306 unsurprising that the early organic chemists in the times of Berzelius found only carbon
307 (~50%) with small admixtures of oxygen (20-30%), nitrogen (~15%) and even lower
308 amounts of hydrogen, phosphorous and sulphur [35].

309 We now know that in addition to the CHONPSes, further elements, mainly
310 consisting of transition metals, make up roughly 1% of the dry weight of prokaryotic

311 cells (Fig. 3a). This marginal fraction of mass is widely taken to indicate a
312 correspondingly marginal role in life in general and in particular in its appearance on
313 the early Earth. However, vital roles of some of these elements, later to become the
314 “essential trace elements”, have been recognized almost two centuries ago. Such
315 essential trace elements were first observed in the middle of the 19th century when Jules
316 Raulin, a student of Pasteur’s, described that growth of *Aspergillus niger* fully depended
317 on the presence of zinc. By the beginning of the 20th century, microbiologists had worked
318 out a catalog already comprising the majority of the essential trace elements identified
319 today. We therefore know since more than a hundred years that life is not only based on
320 the CHONPSes but also on a number of further elements such as Mg, Mn, Fe, Cu, Zn,
321 Ni, Mo, W, Co, V and so on (summarily referred to in this work as FeNiMoWCo+²). But
322 surely these elements must be of lesser importance to life’s processes since they are so
323 rare as compared to the CHONPSes, earning them the attribute “trace”? This reasoning
324 commits a serious mistake: the biological units actually capable of “doing something
325 useful”, such as for example converting environmental free energy into the internal
326 (spatial and temporal) order of a living cell, are of course not individual CHONPS-atoms
327 but the enzymes. A single enzyme, however, is made up from an enormous number of
328 CHONPS-atoms! Middle-of-the-road prokaryotic cells (such as *E. coli*) contain slightly
329 more than 2 million proteins (not all of which are enzymes, *i.e.* catalyze reactions or
330 convert disequilibria) [39]. This same cell contains about 0.2 % (by dry weight) of iron,
331 certainly a ridiculously low quantity if compared to the close to 50 % carbon or 20-30 %
332 oxygen. However, recalculating the weight percentage into actual number of Fe-ions per
333 cell yields the staggering value of roughly 4 million. *E. coli* therefore contains more iron
334 atoms than proteins! Even considering (i) that part of these irons are probably stored in
335 ferritins for later use, (ii) that many enzymes actually harbour multi-iron metal clusters
336 and (iii) that not all enzymes are metalloproteins, still yields a situation where a
337 substantial fraction of all enzymes are indeed iron-containing metalloenzymes. Among
338 the trace elements, iron might be considered borderline since it is required in much
339 higher amounts than other transition metals (Fe alone accounts for one fifth of the total
340 dry weight of all non-CHONPS-elements). However, even for the very low-abundancy
341 trace metal molybdenum [40], numbers of atoms per cell reach about 5000, which is
342 roughly commensurate with the number of individual bioenergetic electron transfer
343 chains when *E. coli* grows by (anaerobic) respiration of nitrate. The comparison in Fig. 3
344 illustrates that the picture of metals as marginal components in a living cell is fatally
345 flawed.

2 A host of these heavier atoms play crucial roles in metabolic processes of extant life. We have chosen to call them “FeNiMoWCo+” since iron, nickel, molybdenum/tungsten and cobalt have specifically been put forward as essential agents in putatively ancient reaction schemes, such as H₂-oxidation, CO₂-reduction, CH₄-oxidation, NO₃-reduction etc. [27,36,37]. The “+”-sign allows for further transition metals, the importance of which may not yet be fully appreciated. Copper does not enter the list since it almost certainly was not bioavailable under the anoxic conditions of the primordial ocean [38]. Manganese admittedly plays a paramount role in oxygenic photosynthesis. However, as argued in the past, we do not consider photosynthesis (oxygenic or anoxygenic) to have been operating at the time of life’s emergence [27]. Incidentally, the proposed list makes for a “pronounceable” word to compete with the CHONPSes.



346

347

348

349

Figure 3. Comparison of the respective share of organic molecules and metal ions in the makeup of an *E. coli* cell when measured (a) by weight or (b) by numbers. The surface of the depicted ellipsoids is proportional to the relative proportions of the treated items.

350

4.2. CHONPS vs. FeNiMoWCo+; Which One Does the Work?

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

To the bioenergetics community, *i.e.* the field studying the molecular mechanisms performing free energy conversion, all this comes as no surprise since more than 50 years of research have firmly established that it is the metals (Fe, Cu, Ni, Mo, ...) that do the job and that the protein scaffolds holding the metal centres mainly serve to fine-tune the physico-chemical properties of these inorganic clusters. Rather than being marginal phenomena in life's processes, metal ions are the dominant players! This evidence is further brought home by the observations that roughly half of all enzymes are metalloenzymes [41, 42] and that there does not seem to be a single recognized metabolic pathway that doesn't contain at least one metalloenzyme [43].

There likely is one subgroup of the Origin-of-Life community which will remain unfazed by the above described considerations, that is, RNA-world proponents. In their framework of thinking, (metallo-)protein-based processes, certainly the major mechanisms in extant life, did not operate at life's beginnings. Rather, it were catalytic RNA-based molecules (so-called ribozymes) which in the beginning performed these tasks, later taken over by proteinaceous enzymes. While we will here refrain from discussing the pros and cons of the RNA-world hypothesis, we cannot help noticing that all naturally occurring ribozymes (whereof there aren't a great many anyway) are in fact metallo-ribozymes depending on metal ions for catalysis and/or 3D-structuring [44].

Our present understanding of both the quantities and the functional roles of metal ions in living cells therefore fully justifies the opening statement in a recent review

371 article on the young field of metallomics: “Without metal ions life is not possible and
372 would not have evolved in the first place” [45]. This statement obviously is a far cry
373 from Darwin’s idea that protein compounds would readily evolve towards living
374 entities.

375 This paramount role of metal ions in enabling life should not come as a surprise.
376 Metal-free peptides (*i.e.* chains of amino acids) have no *a priori* catalytic activities, largely
377 due to the fact that these activities require specific structural foldings - foldings which
378 moreover often depend, for both form and function, on binding specific metal ion
379 complexes (thereby conferring a more-or-less stringent constraint on the amino acid
380 sequence forming the fold). In this general picture, strikingly, one class of metabolic
381 processes, whose mediation almost invariably requires these metal-ion enabled protein
382 fold structures, stands above all others in the affairs of life. Indeed, as we will discuss
383 below, the most important processes in metabolism, that is, the mechanism
384 underpinning the possibility for life to exist according to the 2nd law, are redox reactions.
385 In the overwhelming majority of cases, redox-active enzymes are metalloenzymes and it
386 is the metal ion (or clusters thereof) that does the job. Only in very rare exceptions (*e.g.*
387 cytochrome oxidase or photosystem II), protein-integral amino acids participate in redox
388 reactions and in all these cases, the parent enzymes are multi-metal-ion systems with the
389 respective amino acids performing isolated electron transfer steps within overall metal-
390 based chains.

391 Ribozymes, strictly speaking also metalloenzymes as pointed out above, have only
392 extremely specific catalytic properties, mainly directed towards themselves, such as
393 auto-splicing, or are involved in the replicatory functions of the ribosome [46]. None of
394 these catalytic functions is even only remotely related to the life-enabling process of
395 metabolism.

396 In most biochemistry textbooks, the existence of metalloenzymes is treated as
397 Nature having cleverly realized that catalytic performance of an enzyme can be
398 improved by recruiting metal ions in specific binding sites. This way of thinking is both
399 true to the “organics-first” paradigm and likely influenced by the way technology (in
400 particular in the chemical industry) operates. Kinetically sluggish reaction schemes
401 (often requiring cost-intensive high temperatures and high pressures) are frequently
402 substantially accelerated by the additions of certain metals. A number of arguments,
403 however, speak against the “from metal-free enzymes to metalloproteins”-notion. (i) All
404 apo-metalloenzymes, *i.e.* deprived of their metal centres, involved in energy converting
405 metabolism aren’t less active than their holo-counterparts but simply dead! (ii) By
406 contrast, their protein-free metal-clusters, frequently encountered in certain minerals
407 [36], often perform the same reactions as when integrated in polypeptides [47]. (iii) In
408 the case of environmental scarcity of certain metals, life has found ways to use other
409 available metals as replacement. For example, in iron-depleted habitats, the canonical
410 tasks of Fe-containing enzymes are routinely taken over by structurally (and sequence-
411 wise) fully unrelated copper-enzymes as exemplified by the *cd1*-Nir to Cu-Nir (both
412 enzymes reducing nitrite to nitric oxide) transition. If pre-existing metal-free life would
413 have recruited the metal-ions, one would certainly have to assume that we should still
414 be able to find such organisms in environments highly depleted in all metal ions. This
415 does not seem to be the case. To the best of our knowledge, no species has been observed
416 so far which can live without metals. This is even true for species only deduced from
417 metagenome analyses, where routinely a full host of metalloenzymes are identified *via*
418 their tell-tale specific cofactor-binding sequences. Both in the environment and under
419 laboratory conditions, truly metal-free media are simply sterile! We therefore consider it
420 an inescapable fact that FeNiMoWCo+ vastly surpasses the importance of CHONPS in
421 rationalizing why life on Earth was able to emerge.

422 5. A 2nd Law-Compliant Emergence of Life; from Generalities to Specifics

423 5.1. The 2nd Law Meets Prebiotic Chemistry

424 Almost 80 years after the publication of Schrödinger's book, the thermodynamic
425 argument begins to make occasional inroads into the organic soup fortress which
426 previously had managed to remain hermetic to criticism over the 2nd half of the 20th
427 century (and criticism there was ... [48-52]. NASA's website describing research on
428 putative life on Mars for example reads: "... the origin of life requires the presence of
429 carbon-based molecules, liquid water *and an energy source*" (emphasis by the authors)
430 [53]. Tellingly, the nature of this "energy" remains unspecified. Even more worrisome is
431 the fact that the energies put forward in many articles are of the sort of photons, ionizing
432 radiation, heat or electrical discharge (a combination of heat and reducing power). Such
433 "energies" can indeed assist in overcoming activation barriers of specific reactions
434 taking the reactant/product couples more rapidly to their thermodynamic equilibria.
435 This is what happens when small amounts of organics are detected after exposing high
436 amounts of precursor molecules to UV-irradiation, bombardment with all sorts of
437 ionizing radiations or electrical discharge. However, this is clearly not what life does!
438 Life takes the precursors and produces extremely out-of-equilibrium concentrations of
439 the products. To be able to do this, these processes must be driven by disequilibria in
440 life's environment, *i.e.* by "free energy" in the sense of Gibbs' formulation of the 2nd law,
441 as discussed above [54-58]. The crucial distinction between energy (a strictly conserved
442 entity as enshrined in the 1st law of thermodynamics) and non-conserved "free energy"
443 (in fact not an energy at all but a measure of how far from equilibrium a system is and
444 thus of its ability to do work) has been pointed out repeatedly in the past [58-60]. For
445 example, it isn't heat (a 1st law type energy) but a temperature gradient (a 2nd law type
446 free energy, that is, a thermodynamic disequilibrium) that produces hurricanes. Of
447 course, energies can be transformed into Gibbs free energy, such as when for example
448 radioactive decay in Earth's mantle generates heat and hence convection which
449 eventually will *via* the process of serpentinization bring about redox disequilibria [61].
450 However, the Gibbs free energy potentially being converted into other disequilibria, *e.g.*
451 life's local decrease in entropy, is this ultimate redox disequilibrium and not the (1st law-
452 type) energy of radioactive decay.

453 Leaving this (major) detail aside, a growing number of articles admittedly now
454 acknowledge Schrödinger's stricture that the emergence of life must obey the 2nd law
455 [62-65]. The necessity of pushing the subsystem life away from thermodynamic
456 equilibrium is now discussed even in articles from the prebiotic soup community. For
457 example, a consecutive sequence of disequilibrium-maintaining inputs of energy (some
458 of which are 1st law-type energies rather than Gibbs free energies, see above) are
459 proposed to take prebiotic organics step-by-step up to ever more extreme disequilibria
460 and eventually to cellular life which, at supposedly "higher stages of evolution" [63]
461 converts free energy through a process the biologists call "chemiosmosis" [63].

462 5.2. Extant Life's Universal Steam Engine: Chemiosmosis

463 The mechanism of chemiosmosis (we will explain the process in more detail below)
464 is common to all life on planet Earth and is therefore widely considered to be the way by
465 which the so-called Last Universal Common Ancestor (LUCA) harvested environmental
466 free energy [27, 67-71]. It took the biological sciences more than half a century to unravel
467 the major mechanisms of the chemiosmotic system and several key processes are still not
468 well understood. Chemiosmotic enzymes are among the most complex molecular
469 machineries involved in cellular processes [72, 73] and it is therefore only natural that
470 evolutionary biologists generally regard chemiosmosis as a highly evolved mechanism,
471 despite its ubiquity in extant life and likely deep ancestry going back to the common
472 ancestor of Bacteria and Archaea. Chemiosmosis' deep evolutionary roots are further
473 emphasized by the observation that its common principle of action (see below) has been
474 retained since the times of the LUCA while adapting to a bewildering variety of sources
475 of environmental free energies (all of which ultimately come down to electrochemical
476 free energy, see [74]) allowing life to spread to virtually every habitat on the planet

477 providing redox disequilibria of some sorts. At later stages of evolution, photosynthetic
478 organisms managed to generate such redox disequilibria *via* charge separation induced
479 by visible light photons while still fully relying on the chemiosmotic principle [27,74].
480 Even the plethora of different enzyme systems converting these redox disequilibria into
481 life's spatial and temporal ordering have now been shown to be built up from a very
482 small number of basic metalloprotein domains, many of which can be phylogenetically
483 traced back to the LUCA [75-80]. The chemiosmotic principle for the conversion of
484 environmental free energy into the entropy decrease characterizing life from the LUCA
485 to extant life is therefore now firmly established.

486 So what does this ubiquitous process actually do? The ultimate result of the
487 chemiosmotic mechanism is the conversion of redox disequilibria, *i.e.* of the
488 simultaneous presence of sufficiently reducing and sufficiently oxidizing redox
489 compounds into a disequilibrium ratio of polyphosphates to monomeric phosphates
490 (where the "sufficiently" indicates that the flow of electrons from reductant to oxidant is
491 exergonic even taking into account the free energy required to generate the
492 phosphate/polyphosphate disequilibrium). In most metabolic processes, these
493 polyphosphates are linked to an adenosine moiety, yielding the bioenergetically
494 prominent molecules adenosine mono-, di- and triphosphate (AMP, ADP and ATP). In
495 thermodynamic equilibrium at standard conditions (1 atmosphere, room temperature,
496 pH 7, 10 mM Mg^{2+} , 10 mM P_i), a mixture of ATP and ADP will feature a ratio of 1 ATP to
497 10 million ADPs [59, 81]. The chemiosmotic free-energy-converting engine drives the
498 ATP/ADP ratio to between 10 and 1000 ATPs per 1 ADP, that is, roughly 9 orders of
499 magnitude away from where it wants to be [59]. It is the Gibbs free energy available in
500 this disequilibrium which drives virtually all, by themselves strongly endergonic,
501 cellular reactions "uphill", that is, towards the disequilibria (*i.e.* entropy decrease)
502 characterizing life. The fact that it is the ATP/ADP couple (or, more rarely, the ADP/AMP
503 or even the ATP/AMP couples) which makes metabolic reactions proceed towards the
504 right (for life, but wrong for thermodynamics) direction led to the somewhat
505 unfortunate, since misleading, notion of ATP as the cellular energy "currency". For
506 example, a one molar solution of ATP provides zero kilojoules if it is in the presence of a
507 10^7 -fold excess of ADP while its energy content is -57 kJ/mol^{-1} if ADP is present at 10^{-3}
508 molar concentration [59,60]. As a matter of example, one cannot pay for a single
509 biosynthetic conversion of pyruvate into tyrosine by inserting an ATP-coin into the
510 money slot of the involved enzymes but it is the compounded "weight" of the ATP
511 mountain wanting to reverse into an ADP mountain that pushes the bulk of the
512 precursor molecules all the way up to their thermodynamically highly unlikely product,
513 tyrosine [54, 58].

514 Noteworthy, the free energy that can be extracted from out-of-equilibrium
515 ATP/ADP (or ADP/AMP) couples has next to nothing to do with the adenosine-moiety
516 since equilibrium constants determined for the adenosine-devoid, *i.e.* purely
517 "inorganic", triphosphate/diphosphate (also called pyrophosphate) or
518 pyrophosphate/phosphate couples lie in the same range as those of their adenosine-
519 linked counterparts [81]. Consequently, certain ATP-driven enzymes also perform their
520 tasks in the presence of non-equilibrium polyphosphates [82, 83].

521 5.3. Chemiosmosis, a Physical Phenomenon in Biological Disguise

522 Chemiosmosis is fundamentally based on the presence of a topological
523 compartment (containing roughly 1 to a few femtoliters) sealed from the environment
524 by a dielectric barrier enabling the whole system to work like a capacitor (Fig. 4). This
525 barrier is erroneously considered (and even described as such in most biochemistry
526 textbooks) as "the lipid bilayer membrane". CryoEM images obtained on native
527 biological membranes have shown that the surfaces of these membranes are in fact by a
528 large margin dominated by proteins and not by lipids [84-87], a fact already known to

529
530
531
532
533
534
535

quantitative biochemists half a century ago [88]. The low dielectric constant of the interior space of proteins ($\epsilon \approx 4$) indeed makes them perfect constituents of the type of dielectric barrier encountered in and required by chemiosmotic systems (high dielectric-constant environments, such as water with its ϵ of about 80, strongly reduce electric fields via polarization-induced shielding of charges; the internal dielectric constant of proteins is commensurate with that of a pure lipid membrane allowing strong electric fields to be generated already by small charge asymmetries).


536
537
538
539
540

Figure 4. Schematic representation of life’s universal mechanism for converting environmental free energy into cellular entropy decrease, chemiosmosis. The correspondence between individual items in this process and terms of the 2nd law of thermodynamics (expressed in plain language on the left of the figure) is emphasized by colour-coding.

541
542
543
544
545
546
547
548
549
550
551
552
553
554
555

We therefore consider the commonly held view that first cellular entities were necessarily bounded by lipid-based membranes as conjecture and generally at odds with what we see in extant life. The paradigm of the lipid-bilayer membrane as the confinement separating the cellular interior (the “cytoplasm”) from the environment naturally led to numerous speculations in the origin-of-life field on how first lipids may have been synthesized prebiotically and how first micellar and then vesicular structures may have been formed [89-91]. We contend (based on the biological facts mentioned above) that the presence of lipids is not a strict prerequisite for the formation of (chemiosmotically functional) compartments but that the respective dielectric barriers may also have consisted purely of proteinaceous matter very much as is the case in extant gas vesicles [92] or bacterial microcompartments (BMCs) [93].

This dielectric barrier harbours electron transfer chains made up primarily from a variable number (according to the specific redox substrates used) of metalloproteins assisted in most but not all cases by a few small redox active organics (quinones, flavins, NAD etc). While electrons flow through these “bioenergetic chains”, an electrostatic and

cation-concentration gradient³ is built up over the dielectric barrier, partly as a direct result of opposite barrier-sidedness of oxidation/reduction reactions and partly due to active vectorial transport of small cations such as protons or, more rarely, sodium ions described in more detail previously [27, 74]. These gradients generate a cation-motive force over the roughly 5nm-wide dielectric barrier which pushes the mentioned cations back through dedicated free-energy-converting machineries within the dielectric barrier towards the inside of the compartment. Within these machineries, the movement of the cations is coupled to phosphorylation reactions, displacing the ratios of in general ATP/ADP, but also PP_i/P_i [94] far away from their equilibrium values (see previous paragraph).

The mechanism of chemiosmosis as summarized in Fig. 4 looks eerily non-biological and predominantly involves physical/electrochemical principles (capacitor-like boundaries, redox gradients, electric fields, inorganic metal-clusters, inorganic phosphates and polyphosphates). Rather than being a highly evolved biological process, the core of chemiosmosis resembles a basic electrochemical device which is conceivable in the absence of any life-like features. What is more, it is a direct manifestation of how the 2nd law authorizes localized entropy decrease by taking its encompassing, larger, system closer to equilibrium (Fig. 4, colour-coded correspondence between elements of chemiosmosis and individual terms of the 2nd law).

6. Free energy Conversion Prior to Chemiosmosis?

In the above paragraphs, we have tried to impress on the reader the extraordinarily “abiotic” features of the chemiosmotic principle, the free-energy-converting mechanism of extant life. It therefore is far from obvious that chemiosmosis is, as commonly assumed, the high end mechanism having required extended evolutionary timescales to appear and prior to which other, likely simpler but unknown, types of energy conversion must have existed. Rather, we contend that the chemiosmotic principle may have operated right at the beginning of living things and may in fact have been the very fundamental reason why life was able to appear in the first place.

Here we have to admit that “contend” isn’t “know”. There is, of course, no stringent argument excluding the frequently proposed series of pre-chemiosmosis types of free energy conversion. However, neither are there arguments necessarily requiring such processes! The empirical approach to elucidating the evolutionary pathway of free energy converting mechanisms is based on comparative biology and phylogenetic analyses. This approach allows to reach back to the last universal common ancestor (LUCA) of all living beings on Earth but no further! And all available evidence presently indicates that LUCA used chemiosmosis. This implies that putative pre-chemiosmotic processes, as intellectually appealing and intuitive as they may come, can never be proven to have indeed existed. Applying Occam’s razor, we therefore consider that the only truly empirical approach to elucidating the emergence of life-enabling free energy conversion is to try and draft scenarios which can rationalize a prebiotic appearance of the chemiosmotic mechanism. If this approach succeeds, we will have a plausible, thermodynamically viable and empirically supported, narrative for life’s emergence. If it doesn’t, a nearly infinite number of alternative scenarios can be thought of but we will never be able to tell whether they have anything to do with the actual evolutionary history of life on Earth.

³ The term chemiosmosis is essentially a misnomer, since, contrary to the definition of osmosis, it doesn’t imply movement of water molecules but rather of cations (protons or sodium ions) to restore thermodynamic equilibrium conditions. However, the term is so firmly established in the biological literature that we will stick to it throughout this article. Nevertheless, true to Confucius’ insight that “The beginning of wisdom is to call things by their right name”, we would urge the bioenergetics community to try and come up with a more fitting term.

7. Towards Scenarios Featuring Prebiotic Chemiosmotic Processes

The string of arguments developed so far in this article calls for emergence of life hypotheses based on the prebiotic presence of a chemiosmotic system of free energy conversion resembling the principle shown in Fig. 4. While there are potentially a number of possible settings fulfilling this requirement, historically only the alkaline hydrothermal vent theory (AVT) so far took this necessity seriously. In fact, the original version of the AVT scenario [30, 95] was strongly boosted by the realization that several aspects of chemiosmosis come as a natural consequence of the topology of the porous hydrothermal mounds and of the chemical and electrochemical properties of both the alkaline vent fluid and the surrounding ocean waters [68-71, 96, 97].

Several details of this newer version of the alkaline hydrothermal vent hypothesis [98] drew fierce criticism from both the bioenergetics [99] and the thermodynamics [55] communities. More recent versions of this hypothesis strive to address these well-taken points of criticism (a more detailed account of the “evolutionary” trajectory of the AVT from its emergence to the present day will be presented in Russell [100]).

While the authors of the present article attempt to design and experimentally test scenarios generating close-to-lifelike manifestations of chemiosmotic processes in the framework of the AVT scenarios, we can and will not exclude that chemiosmosis may have arisen in other settings. However, the argumentation detailed throughout this article to our mind leaves no alternative to searching for a chemiosmotic emergence of life if both Schrödinger’s strictures and the experimental evidence for extant life’s universal way to comply with the 2nd law are taken seriously.

8. The Other “What Is Life” Question

Schrödinger’s insight that the 2nd law of thermodynamics necessarily implies that the generation of order in living matter must somehow be driven by environmental disequilibria, put the emergence of life on par with that of other dissipative-structure-forming phenomena of the physical world. However, extant life undeniably differs from such physical dissipative structures by the property of (error-prone) coded reproduction exposing it to Darwinian natural selection and thereby enabling evolutionary adaptation and innovation. It is this property which allowed life to persist as a dissipative structure through 4 billion years of evolution, to explore (and continue to do so) the available space of redox-disequilibrium-offering “habitats” on our planet and to eventually manage to reproduce the required redox tensions by using photon energy from our parent star [101]. Mutability together with natural selection thus accomplished the extraordinary feat of adding to Earth-bound, geochemical, sources of free energy a truly extraterrestrial one through hooking chemiosmosis up to light energy from the sun [102]. Such mutability may already have been present in replication processes based on peptides, that is, before the RNA-mediated coding systems had appeared, as worked out during the last two decades [103-106] with (again) metal-ions (*e.g.* Zn) playing crucial roles in recognition between template- and daughter-chains of peptides [105].

These considerations shine light on the fact that we are still sorely lacking a commonly accepted definition for the meaning of (the term) life. Since it is coded reproduction that sets life apart from other dissipative structures, considering life to only commence when coded reproduction set in appears perfectly defensible to us. Would that imply that all the considerations laid out in this article actually have no bearing on the emergence of life? We firmly argue to the contrary and in doing so we have Schrödinger on our side. All processes involved in putatively leading to the emergence of coded replication ineluctably entail a reduction in the entropy of the system and therefore have to be driven by a mechanism (in our minds likely chemiosmosis) capable of paying for this entropy decrease with free energy from the environment. The abiotic existence of such a free energy converting system therefore is the sine-qua-non condition for life to emerge.

Funding: This work was supported by the French Agence Nationale pour la Recherche (grant no. ANR-22-CE30-0035-01).

Acknowledgments: W.N. is grateful to Vincent Breton and Alain Falvard (LPC, Univ. Clermont-Auvergne) for stimulating discussions and for providing the impetus for writing this article. Thanks from all authors go to the B10NUMB3R5-team at the Weizmann Institute of Science (Israel). The input of low entropy from Elbert Branscomb's thinking (fluctuatingly) drove an essential part of the structure formation eventually resulting in this contribution.

References

- Horneck, G.; Walter, N.; Westall, F.; Grenfell, J.L.; Martin, W.F.; Gomez, F., et al. AstRoMap European astrobiology roadmap. *Astrobiology* **2016**, *16*(3), 201-243.
- <https://solarsystem.nasa.gov/news/517/why-bennu-10-reasons/>
- <https://www.isas.jaxa.jp/en/missions/spacecraft/current/hayabusa2.html>
- https://www.esa.int/Science_Exploration/Space_Science/Rosetta/Rosetta_s_comet_contains_ingredients_for_life
- Sasselov, D.D.; Grotzinger, J.P.; Sutherland, J.D. The origin of life as a planetary phenomenon. *Science Adv.* **2022**, *6*: eaax3419
- Darwin, C. *Letter to Joseph Dalton Hooker on February 1, 1871*, cited in: Darwin F (ed) **1887** *The life and letters of Charles Darwin, including an autobiographical chapter*, 3 vols. John Murray, London
- Miller, S.L. A production of amino acids under possible primitive earth conditions. *Science* **1953**, *117*, 528-529.
- Miller, S.L.; Urey, H.C. Organic compound synthesis on the primitive Earth: Several questions about the origin of life have been answered, but much remains to be studied. *Science* **1959**, *130*, 245-251.
- Ferus, M.; Rimmer, P.; Cassone, G.; Knížek, A.; Civiš, S.; Šponer, J.E. et al. One-pot hydrogen cyanide-based prebiotic synthesis of canonical nucleobases and glycine initiated by high-velocity impacts on early Earth. *Astrobiology* **2020**, *20*, 1476-1488.
- Root-Bernstein, R.; Brown, A.W. Novel apparatuses for incorporating natural selection processes into origins-of-life experiments to produce adaptively evolving chemical ecosystems. *Life* **2022**, *12*, 1508.
- Kobayashi, K.; Ise, J.-i.; Aoki, R.; Kinoshita, M.; Naito, K.; Udo, T.; Kunwar, B.; Takahashi, J.-i.; Shibata, H.; Mita, H.; et al. Formation of Amino Acids and Carboxylic Acids in Weakly Reducing Planetary Atmospheres by Solar Energetic Particles from the Young Sun. *Life* **2023**, *13*, 1103.
- Ebisuzaki, T.; Maruyama, S. Nuclear geyser model of the origin of life: Driving force to promote the synthesis of building blocks of life. *Geoscience Frontiers* **2017**, *8*, 275-298.
- Altwegg, K.; Balsiger, H.; Bar-Nun, A.; Berthelier, J.J.; Bieler, A.; Bochslers, P.; Briois, C.; Calmonte, U.; Combi, M.R.; Cottin, H.; et al. Prebiotic chemicals - Amino acid and phosphorus in the coma of comet 67P/Churyumov-Gerasimenko. *Sci. Adv.* **2016**, *2*, e1600285
- Oba, Y.; Takano, Y.; Furukawa, Y.; Koga, T.; Glavin, D.P.; Dworkin, J.P.; Naraoka, H. Identifying the wide diversity of extraterrestrial purine and pyrimidine nucleobases in carbonaceous meteorites. *Nature Communications* **2022**, *13*, 2008.
- Oba, Y.; Koga, T.; Takano, Y.; Ogawa, N.O.; Ohkouchi, N.; Sasaki, K.; Sato, H.; Glavin, D.P.; Dworkin, J.P.; Naraoka, H.; et al. Uracil in the carbonaceous asteroid (162173) Ryugu. *Nature Communications* **2023**, *14*, 1292.
- Potiszil, C.; Yamanaka, M.; Sakaguchi, C.; Ota, T.; Kitagawa, H.; Kunihiro, T.; et al. Organic Matter in the Asteroid Ryugu: What We Know So Far. *Life* **2023**, *13*, 1448.
- Zeichner, S.S.; Aponte, J.C.; Bhattacharjee, S.; Dong, G.; Hofmann, A.E.; Dworkin, J.P. et al. Polycyclic aromatic hydrocarbons in samples of Ryugu formed in the interstellar medium. *Science* **2023**, *382*, 1411-1416.
- Aristotle, *On the History of Animals*, Book V, Part 1
- Driesch, H. Philosophy of Vitalism. *Nature* **1913**, *2301*, 400.
- Driesch, H. *The History and Theory of Vitalism*. **1914**. *Trans. C. K. Ogden*. London: MacMillan.
- Reinke, J. *Die schaffende Natur: mit Bezugnahme auf Schopenhauer und Bergson*. **1919**. Quelle & Meyer.
- Reinke, J. *Kritik der Abstammungslehre*. **1920**. Barth. Leipzig.
- Mayr, E. *Toward a new philosophy of biology: Observations of an evolutionist*. **1988**, (No. 211). Harvard University Press.
- Needham, J. Organicism in Biology. *J. Philosophical Studies* **1928**, *3*, 29-40.
- Monod, J. *Le Hasard et la Necessite : Essai sur la philosophie de la biologie moderne*. **1970**.
- Schrödinger, E. *What is life? The physical aspect of the living cell and mind*. **1944**. Cambridge: Cambridge University Press.
- Schoepp-Cothenet, B.; van Lis, R.; Atteia, A.; Baymann, F.; Capowiez, L.; Ducluzeau, A.-L.; Duval, S.; ten Brink, F.; Russell, M.J.; Nitschke, W. On the universal core of bioenergetics. *Biochim. Biophys. Acta, Bioenergetics* **2013**, *1827*, 79-93.
- Kondepudi D; Prigogine I. *Modern Thermodynamics; From Heat Engines to Dissipative Structures*. **2006** Wiley & Sons, New York.
- Nicolis, G.; Prigogine, I. *Self-Organization in Non-equilibrium Systems*. **1977**. John Wiley and Sons, New York.
- Russell, M.J.; Hall, A.J. The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and pH front. *J. Geol. Soc. Lond.* **1997**, *154*, 377-402.
- Barge, L.M.; Branscomb, E.; Brucato, J.R.; Cardoso, S.S.S.; Cartwright, J.H.E.; Danielache, S.O.; et al. Thermodynamics, disequilibrium, evolution: Far-from-equilibrium geological and chemical considerations for origin-of-life research. *Origins of Life and Evolution of Biospheres* **2017**, *47*, 39-56.

- 711 32. Chatterjee, S.; Yadav, S. The Coevolution of Biomolecules and Prebiotic Information Systems in the Origin of Life: A
712 Visualization Model for Assembling the First Gene. *Life* **2022**, *12*, 834.
- 713 33. Deamer D.W. Origin of Life. What everyone needs to know. **2020**, Oxford University Press, ISBN 9780190098995.
- 714 34. Todd, Z.R. Sources of Nitrogen-, Sulfur-, and Phosphorus-Containing Feedstocks for Prebiotic Chemistry in the Planetary
715 Environment. *Life* **2022**, *12*, 1268.
- 716 35. Neidhardt, F.C.; Ingraham, J.L.; Schaechter, M. **1990**. Physiology of the bacterial cell: a molecular approach. Sinauer
717 Associates, Sunderland, Mass./USA.
- 718 36. Nitschke, W.; McGlynn, S.E.; Milner-White, E.J.; Russell, M.J. On the antiquity of metalloenzymes and their substrates in
719 bioenergetics. *Biochim. Biophys. Acta, Bioenergetics* **2013**, *1827*, 871-881.
- 720 37. Schoepp-Cothenet, B.; van Lis, R.; Philippot, P.; Magalon, A.; Russell, J.J.; Nitschke, W. The ineluctable requirement for the
721 trans-iron elements molybdenum and/or tungsten in the origin of life. *Sci. Rep.* **2012**, *2*, 263.
- 722 38. Anbar. A.D. Elements and Evolution. *Science* **2008**, *322*, 1481-1483.
- 723 39. Brock, T.D.; Madigan, M.T.; Martinko, J.M.; Parker, J. Brock biology of microorganisms. **2003**. Upper Saddle River (NJ):
724 Prentice-Hall, 2003, chapter 2, table 2.2.
- 725 40. Scott, D.; Amy, N.K. Molybdenum accumulation in chlD mutants of Escherichia coli. *Journal of bacteriology* **1989**, *171*, 1284-
726 1287.
- 727 41. Thomson, A.J.; Gray, H.B. Bioinorganic Chemistry. Current Opinion in Chemical Biology **1998**, *2*, 155-158.
- 728 42. Martin, R.B. Bioinorganic chemistry. In: Encyclopedia of molecular cell biology and molecular medicine **2006**. Wiley
- 729 43. Monosson, E. Evolution in a toxic world. **2012** Island Press Washington DC/USA.
- 730 44. Pyle, A.M. Ribozymes: a distinct class of metalloenzymes. *Science* **1993**, *261*, 709-714.
- 731 45. Maret, W. The quintessence of metallomics: a harbinger of a different life science based on the periodic table of the
732 bioelements. *Metallomics* **2022**, *14*, mfac051.
- 733 46. Talini G.; Gallori, E.; Maurel, M.-C. Natural and unnatural ribozymes: Back to the primordial RNA world. Research in
734 Microbiology **2009**, *160*, 457-465.
- 735 47. Farr, O.; Gaudu, N.; Danger, G.; Russell, M.J.; Ferry, D.; Nitschke, W.; Duval, S. Methanol on the rocks: Green rust
736 transformation promotes the oxidation of methane. *J. R. Soc. Interface* **2023**, *20*, 20230386.
- 737 48. Rutten, M.G. Origin of life on Earth, its evolution and actualism. *Evolution* **1957**, *11*, 56-59.
- 738 49. Sillen, L.G. Regulation of O₂, N₂ and CO₂ in the atmosphere; thoughts of a laboratory chemist. *Tellus* **1966**, *18*, 198-206.
- 739 50. Maden, B.E.H. No soup for starters? Autotrophy and the origins of metabolism. *Trends in Biochemical Sciences* **1995**, *20*, 337-341.
- 740 51. Yockey, H.P. Comments on "Let there be life"; Thermodynamic reflections on biogenesis and evolution" by Avshalom C.
741 Elitzur. *J. Theor. Biol.* **1995**, *176*, 349-355.
- 742 52. Branscomb, E.; Russell M.J. Frankenstein or a submarine alkaline vent: Who is responsible for abiogenesis? Part 1: What is life
743 – that it might create itself? *BioEssays* **2018**, *40*, 1700179 <https://doi.org/10.1002/bies.201700179>
744 https://cneos.jpl.nasa.gov/about/life_on_earth.html
- 745 54. Branscomb, E.; Russell, M.J. Turnstiles and bifurcators: the disequilibrium converting engines that put metabolism on the road.
746 *Biochim. Biophys. Acta, Bioenergetics* **2013**, *1827*, 62-78.
- 747 55. Branscomb, E.; Biancalani, T.; Goldenfeld, N. Russell, M. Escapement mechanisms and the conversion of disequilibria; the
748 engines of creation. *Physics Reports* **2017**, *677*, 1-60.
- 749 56. Branscomb, E.; Russell M.J. Why the Submarine Alkaline Vent is the Most Reasonable Explanation for the Emergence of Life.
750 *BioEssays* **2018**, *41*, 1800208.
- 751 57. Branscomb, E.; Russell, M.J. On the beneficent thickness of water. *Interface focus* **2019**, *9*, 20190061.
- 752 58. Branscomb, E. Boltzmann's casino and the unbridgeable chasm in emergence of life research. **2023**. *arXiv preprint*
753 *arXiv:2312.00932*.
- 754 59. Nicholls D.G.; Ferguson, S.J. Bioenergetics 4. **2013**. Academic Press, Amsterdam/Netherlands
- 755 60. Guérin, B. Bioénergétique. **2004**. EDP Sciences. Les Ulis/France
- 756 61. Russell, M.J. A Self-sustaining Serpentinization Mega-engine Feeds the Fougérite Nanoengines Implicated in the Emergence
757 of Guided Metabolism. *Front. Microbiol.* **2023**, *14*, 1083.
- 758 62. Schulze-Makuch, D.; Irwin, L.N. Life in the Universe. **2004**, Springer-Verlag.
- 759 63. Pascal, R.; Pross, A.; Sutherland, J.D. Towards an evolutionary theory of the origin of life based on kinetics and
760 thermodynamics. *Open biology* **2013**, *3*, 130156.
- 761 64. Davies, P.C.; Walker, S.I. The hidden simplicity of biology. *Reports on Progress in Physics* **2016**, *79*, 102601.
- 762 65. Walker, S.I. Origins of life: a problem for physics, a key issues review. *Reports on Progress in Physics* **2017**, *80*, 092601.
- 763 66. Kompanichenko, V. Thermodynamic jump from prebiotic microsystems to primary living cells. *Sci* **2020**, *2*, 14.
- 764 67. Mitchell, P. Coupling of phosphorylation to electron and hydrogen transfer by a chemi-osmotic type of mechanism. *Nature*
765 **1961**, *191*, 144-8.
- 766 68. Martin, W.; Russell, M.J. On the origin of cells: An hypothesis for the evolutionary transitions from abiotic geochemistry to
767 chemoautotrophic prokaryotes, and from prokaryotes to nucleated cells. *Philosophical Transactions of the Royal Society of London*
768 **2003**, *358B*, 27-85.
- 769 69. Martin, W.; Russell, M.J. On the origin of biochemistry at an alkaline hydrothermal vent, *Philos. Trans. R. Soc. Lond. B* **2007**, *367*,
770 1887-1925.

- 771 70. Lane, N.; Martin, W. F. The origin of membrane bioenergetics. *Cell* **2012**, *151*, 1406-1416.
- 772 71. Lane, N.; Allen, J.F.; Martin, W.F. How did LUCA make a living? Chemiosmosis in the origin of life. *BioEssays* **2010**, *32*: 271–
- 773 280. <http://dx.doi.org/10.1002/bies.200900131>
- 774 72. Ohnishi, T. NADH-quinone oxidoreductase, the most complex complex. *Journal of bioenergetics and biomembranes* **1995**, *25*, 325-
- 775 329.
- 776 73. Grimaldi, S.; Schoepp-Cothenet, B.; Ceccaldi, P.; Guigliarelli, B.; Magalon, A. The prokaryotic Mo/W-bisPGD enzymes family:
- 777 a catalytic workhorse in bioenergetics. *Biochimica et Biophysica Acta-Bioenergetics* **2013**, *1827*, 1048-1085.
- 778 74. Nitschke, W.; Schoepp-Cothenet, B.; Duval, S.; Zuchan, K.; Farr, O.; Baymann, F.; Panico, F.; Minguzzi, A.; Branscomb, E.;
- 779 Russell, M.J. Aqueous electrochemistry; the toolbox for life's emergence from redox disequilibria. *Electrochemical Science*
- 780 *Advances* **2022**, *p.e2100192*.
- 781 75. Ducluzeau, A.L.; van Lis, R.; Duval, S.; Schoepp-Cothenet, B.; Russell, M.J.; Nitschke, W. Was nitric oxide the first deep
- 782 electron sink? *Trends in biochemical sciences* **2009**, *34*, 9-15.
- 783 76. Baymann, F.; Lebrun, E.; Brugna, M.; Schoepp-Cothenet, B.; Giudici-Ortoni, M.T.; Nitschke, W. The redox protein
- 784 construction kit: pre-last universal common ancestor evolution of energy-conserving enzymes. *Philosophical Transactions of the*
- 785 *Royal Society of London. Series B: Biological Sciences* **2003**, *358*, 267-274.
- 786 77. Grein, F.; Ramos, A.R.; Venceslau, S.S.; & Pereira, I.A. Unifying concepts in anaerobic respiration: insights from dissimilatory
- 787 sulfur metabolism. *Biochimica et Biophysica Acta-Bioenergetics* **2013**, *1827*, 145-160.
- 788 78. Refojo, P.N.; Sousa, F.L.; Teixeira, M.; Pereira, M.M. The alternative complex III: a different architecture using known building
- 789 modules. *Biochimica et Biophysica Acta-Bioenergetics* **2010**, *1797*, 1869-1876.
- 790 79. Karavaeva, V.; Sousa, F.L. Modular structure of complex II: An evolutionary perspective. *Biochimica et Biophysica Acta-*
- 791 *Bioenergetics* **2023**, *1864*, 148916.
- 792 80. Zuchan, K.; Baymann, F.; Baffert, C.; Brugna, M.; Nitschke, W. The dyad of the Y-junction- and a flavin module unites diverse
- 793 redox enzymes. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **2021**, *1862*, 148401.
- 794 81. Thauer, R.K.; Jungermann, K.; Decker, K. Energy conservation in chemotrophic anaerobic bacteria. *Bacteriological reviews* **1977**,
- 795 *41*, 100-180.
- 796 82. de Gomez-Puyou, M.T.; de Jesus Garcia, J.; Gomez-Puyou, A. Synthesis of pyrophosphate and ATP by soluble mitochondrial
- 797 F1. *Biochemistry* **1993**, *32*, 2213-2218.
- 798 83. Chi, A.; Kemp, R.G. The primordial high energy compound: ATP or inorganic pyrophosphate? *Journal of Biological Chemistry*
- 799 **2000**, *275*, 35677-35679.
- 800 84. Kirchoff, H. Molecular crowding and order in photosynthetic membranes. *Trends in plant science* **2008**, *13*, 201-207.
- 801 85. Kumar, S.; Cartron, M.L.; Mullin, N.; Qian, P.; Leggett, G.J.; Hunter, C.N.; Hobbs, J.K. Direct imaging of protein organization
- 802 in an intact bacterial organelle using high-resolution atomic force microscopy. *ACS nano* **2017**, *11*, 126-133.
- 803 86. Casella, S.; Huang, F.; Mason, D.; Zhao, G.Y.; Johnson, G.N.; Mullineaux, C.W.; Liu, L.N. Dissecting the native architecture and
- 804 dynamics of cyanobacterial photosynthetic machinery. *Molecular plant* **2017**, *10*, 1434-1448.
- 805 87. Mullineaux, C.W.; Liu, L.N. Membrane dynamics in phototrophic bacteria. *Annual Review of Microbiology* **2020**, *74*, 633-654.
- 806 88. Salton, M.R.J.; Freer, J.H. Composition of the membranes isolated from several Gram-positive bacteria. *Biochimica et Biophysica*
- 807 *Acta (BBA)-General Subjects* **1965**, *107*, 531-538.
- 808 89. Luisi, P.L.; Walde, P.; Oberholzer, T. Lipid vesicles as possible intermediates in the origin of life. *Current opinion in colloid &*
- 809 *interface science* **1999**, *4*, 33-39.
- 810 90. Deamer, D. The role of lipid membranes in life's origin. *Life* **2017**, *7*, 5.
- 811 91. Purvis, G.; Šiller, L.; Crosskey, A.; Vincent, J.; Wills, C.; Sheriff, J.; Xavier, C.; Telling, J. Generation of long-chain fatty acids by
- 812 hydrogen-driven bicarbonate reduction in ancient alkaline hydrothermal vents. *Communications Earth & Environment* **2024**, *5*,
- 813 30.
- 814 92. Bayro, M.J.; Daviso, E.; Belenky, M.; Griffin, R.G.; Herzfeld, J. An amyloid organelle, solid-state NMR evidence for cross-β
- 815 assembly of gas vesicles. *Journal of Biological Chemistry* **2012**, *287*, 3479-3484.
- 816 93. Kerfeld, C.A.; Aussignargues, C.; Zarzycki, J.; Cai, F.; Sutter, M. Bacterial microcompartments. *Nature Reviews Microbiology*
- 817 **2018**, *16*, 277-290.
- 818 94. Holmes, A.O.; Kalli, A.C.; Goldman, A. The function of membrane integral pyrophosphatases from whole organism to single
- 819 molecule. *Frontiers in molecular biosciences* **2019**, *6*, 132.
- 820 95. Russell, M.J.; Daniel, R.M.; Hall, A.J.; Sherringham, J.A hydrothermally precipitated catalytic iron sulphide membrane as a
- 821 first step toward life. *J. Mol. Evol.* **1994**, *39*, 231-243.
- 822 96. Nitschke, W.; Russell, M.J. Hydrothermal focusing of chemical and chemiosmotic energy, supported by delivery of catalytic
- 823 Fe, Ni, Mo/W, Co, S and Se, forced life to emerge. *Journal Molecular Evolution* **2009**, *69*, 481-496.
- 824 97. Lane, N. Why are cells powered by proton gradients. *Nat. Educ* **2010**, *3*, 2.
- 825 98. Russell, M.J. On Irish bacteriometallogenesis and its wider connotations. **2023**, in: Andrew, C.J., Hitzman, M.W.; Stanley, G.
- 826 *Irish-type Deposits around the world*, Irish Assoc. Econ. Geol., Dublin. 45-94.
- 827 99. Jackson, J. Natural pH Gradients in Hydrothermal Alkali Vents Were Unlikely to Have Played a Role in the Origin of Life.
- 828 *Journal of Molecular Evolution* **2016**, *83*, 1-11. <https://doi.org/10.1007/s00239-016-9756-6>
- 829 100. Russell, M.J.; Nitschke, W. ms in preparation

- 830 101. Nitschke, W.; Mühlhoff, U.; Liebl U. Evolution, in: *Photosynthesis: A Comprehensive Treatise*, Raghavendra, A.S. 2000,
831 Cambridge University Press, chapter 22.
- 832 102. Blankenship, R.E. Molecular mechanisms of photosynthesis. 2021, John Wiley & Sons.
- 833 103. Frenkel-Pinter, M.; Samanta, M.; Ashkenazy, G.; Leman, L.J. Prebiotic peptides: Molecular hubs in the origin of life. *Chemical*
834 *Reviews* 2020, 120, 4707-4765.
- 835 104. Adamski, P.; Eleveld, M.; Sood, A.; Kun, Á.; Szilágyi, A.; Czárán, T.; Szathmáry, E.; Otto, S. From self-replication to replicator
836 systems en route to de novo life. *Nature Reviews Chemistry* 2020, 4, 386-403.
- 837 105. Kauffman, S.A. Approaches to the origin of life on Earth. *Life* 11, 1, 34-48.
- 838 106. Ashkenasy, G.; Jagasia, R.; Yadav, M.; Ghadiri, M. R. Design of a directed molecular network. *Proc. Natl. Acad. Sci.*
839 *USA* 2004, 101, 10872-10877.
- 840

841 **Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual
842 author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury
843 to people or property resulting from any ideas, methods, instructions or products referred to in the content.