



HAL
open science

Investigation of UV Light-Promoted Synthesis of α -Sulfonyl Amides from N -Sulfonyl Ynamides

Aurélien Galibert-Guijarro, Jérémy Tronc, Dominique Mouysset, Didier Siri, Stéphane Gastaldi, Michèle Bertrand, Laurence Feray

► **To cite this version:**

Aurélien Galibert-Guijarro, Jérémy Tronc, Dominique Mouysset, Didier Siri, Stéphane Gastaldi, et al.. Investigation of UV Light-Promoted Synthesis of α -Sulfonyl Amides from N -Sulfonyl Ynamides. Journal of Organic Chemistry, 2024, 89 (13), pp.9695-9699. 10.1021/acs.joc.4c01013 . hal-04643937

HAL Id: hal-04643937

<https://cnrs.hal.science/hal-04643937v1>

Submitted on 10 Jul 2024


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Investigation of UV light-promoted synthesis of α -sulfonyl amides from *N*-sulfonyl ynamides.

Aurélien Galibert-Guijarro, Jérémy Tronc, Dominique Mouysset, Didier Siri, Stéphane Gastaldi, Michèle P. Bertrand* and Laurence Feray*

Aix Marseille Univ, CNRS, ICR, Marseille, 13013, France




Abstract: UV Light-promoted synthesis of α -sulfonyl amides from *N*-sulfonyl ynamides without any additives is reported. The reaction proceeds through a radical chain mechanism involving the photo-induced cleavage of the nitrogen-sulfur bond, addition of electrophilic sulfonyl radical to the triple bond of the ynamide followed by beta-fragmentation of the sulfonyl group leading to a ketenimine hydrated upon work-up. This highly efficient rearrangement leads, after acidic treatment, to a wide range of α -sulfonyl amides in high yields.

Introduction. α -Sulfonyl amides are building blocks for organic synthesis and key structural motifs present in biologically active compounds.¹ Therefore, the development of efficient and rapid syntheses of these compounds is of great interest to the community of organic chemists. Current methods for the construction of the C—SO₂ bond of α -sulfonyl amides are based on a two-step strategy involving the formation of a C—S bond followed by oxidation of the sulfur atom in the presence of *m*-CPBA or H₂O₂.² Direct introduction of the SO₂R moiety can be achieved through sulfonylation of α -amide-carbanion intermediates^{1d} or by reacting amides with sulfinates in the presence of oxidants.³ However, these methods suffer from the use of strong bases or oxidants, and for some of them, i.e. sulfonylation of α -amide-carbanion intermediates, low to moderate yields are obtained.

Due to the presence of a nitrogen atom—bearing an electron-withdrawing substituent— directly linked to their carbon-carbon triple bond, ynamides behave as activated polarized alkynes that show unique reactivity and stability. These compounds are currently known as powerful synthetic intermediates for elegant constructions of versatile *N*-containing

molecules.^{4,5} Regarding the direct synthesis of α -sulfonyl amides, only few reports involve the use of ynamides as substrates⁶ (Scheme 1). Wudl⁷ described in 2006 that ynamide **A** can undergo upon heating the N-to-C thermal migration of the tosyl group leading to a ketenimine intermediate that can be hydrated to give the α -tosylated amide **A'** (Scheme 1(a)). More recently, Yuan and Zhu⁸ have reported a radical chain isomerization of *N*-sulfonyl ynamides in the presence of catalytic tosylchloride and an iridium-based photocatalyst to give stable ketenimines. They have shown that this process can be combined with an aza-Claisen rearrangement. During their work, closely related to our ongoing study, as an application of their study, they showed that the ketenimine resulting from the use of tosyl ynamide **B** can be transformed in α -tosyl amide **B'** via hydration (Scheme 1(b)). Finally, Yue and Feng⁹ described a visible-light-promoted oxo-sulfonylation of ynamides **C** to give α -sulfonyl imides **C'** in the presence of sulfonic acids through the in-situ formation of a vinylsulfonate **D**. They proposed that this latter, activated through energy transfer process could undergoes homolytic cleavage of the O—S bond (Scheme 1(c)), that would imply subsequent sulfonyl radical addition/ β -fragmentation sequence.




Scheme 1. Direct synthesis of α -sulfonylated amides from ynamides. (PC : Photocatalyst)

For some years now, our group has been interested in the involvement of ynamides in free-radical transformations.¹⁰ In the course of our investigations, we found out that upon activation under irradiation at 365 nm at room temperature, in the absence of any additives, *N*-sulfonyl ynamides **3** could be converted into α -sulfonyl amides **4** in high yield through a N to C migration of the sulfonyl group¹¹ (Scheme 1(d)). As compared with the other methodologies, our strategy offers several advantages such as easy implementation, use of readily available starting materials, neutral conditions, diversity with respect to the sulfonyl groups that can be used, and neither expensive metals nor photocatalysts are needed. The scope


and limits of this light-promoted synthesis of a wide range of α -sulfonyl amides from *N*-sulfonyl ynamides are discussed in the following section.

Results and discussion. To find the appropriate reaction conditions, ynamide **3aa** prepared from bromohexyne **1a** and *N*-benzyl-4-methylbenzenesulfonamide **2a** according to Hsung methodology¹² was selected as model substrate and submitted to irradiation using 18 W LED lamps (Evoluchem® lamps, λ nm, light intensity: 16.7 mW/cm² at 10 cm distance) at different wavelengths in degassed dichloromethane by varying the reaction time (Table 1). For a reaction time fixed at 24 h, irradiation at 365 nm gives excellent results compared with the other wavelengths studied, i.e. 400 nm and 450 nm, for which ynamide **3aa** is not fully consumed (entry 3 / entries 1,2). The reaction time can be reduced to 18 h or even 4 h¹³ without loss of efficiency, leading to the desired compound in 98% yield after acidic treatment (entries 4,5). Optimized reaction conditions, i.e. irradiation at 365 nm in degassed dichloromethane followed by work-up with a 2 M aqueous solution of HCl, were used to explore the scope of the transformation .



entry	λ (nm)	Time (h)	4aa Isolated yield (%)
1	450	24	40
2	400	24	Not detected
3	365	24	97
4	365	18	98
5	365	4	98


Table 1. Optimization of synthesis of α -sulfonyl amide **4aa** from *N*-sulfonyl ynamide **3aa**. ^a18 W LEDs (Evoluchem® lamps, λ nm, light intensity: 16.7 mW/cm² at 10 cm distance)



Scheme 2. Synthesis of α -sulfonyl amides. ³¹⁸W LEDs (Evoluchem® lamps, 365 nm, light intensity: 16.7 mW/cm² at 10 cm distance)

The results of irradiation experiments conducted with *N*-sulfonyl ynamides **3xy** are reported in Scheme 2. All ynamides were submitted to irradiation at 365 nm for 18 h at room temperature,¹⁴ and all reactions were conducted on 0.16 mmol of substrates. Most of the tested ynamides behave as expected leading to the desired products in high yields, especially those bearing aryl- and heteroaryl sulfonyl groups. It is to be noted that *N*-mesyl ynamides were also reactive giving α -mesyl amides **4al**, **4bl**, **4am** and **4bo** in moderate to good yields. Surprisingly ynamide **3ah** derived from *N*-tosylaniline did not lead to any transformation and was totally recovered after 18 h of irradiation. This point will be discussed later on.

Two plausible mechanisms are proposed in scheme 3. A radical chain mechanism involving the photo-induced homolytic cleavage of the nitrogen-sulfur bond, addition of electrophilic sulfonyl radical **E** to the ynamide triple bond followed by beta-fragmentation of the sulfonyl group of **G** leading to ketenimine **5** can explain the formation of α -sulfonyl amide **4** after hydration¹⁵ (Scheme 3(a)). Calculations of the first singlet excited state (vertical π - π^* transition) at the TD/ ω B97X-D/6-31+G(d,p) level of theory show that the latter spontaneously evolves via homolytic cleavage of the N-S bond when optimizing energy (see SI). Concerted 1,3-migration or in cage recombination of radicals **E** and **F** formed through the above initiation step could also be suggested at this stage (Scheme 3(b)).




Scheme 3. Mechanistic Studies. ^aRadical chain process. ^bIn cage radical recombination. ^cBlank experiments: i) DCM (0.2M), 18 h. ii) DCM (0.2M), 18 h, light protected. iii) hv 365 nm, DCM (0.2M), 18 h, TEMPO (1 equiv). ^dCross experiments, ¹H NMR molar ratio of **4aa**:**4al**:**4bd**:**4bo** = 0.4:0.5:0.35:0.5.

Several experiments were carried out to prove the involvement of radicals and to determine which of the two mechanisms envisaged was involved in the process. Reactions were conducted in the presence of sun light (Scheme 3(c) conditions i)), in the dark (Scheme 3(c) conditions ii)) or in the presence of one equivalent of TEMPO under irradiation at 365 nm (Scheme 3(c) conditions iii)) using ynamide **3aa** as substrate. Throughout any of these experiments, no transformation occurred after 18 h. This confirms both the necessity to use UV light and the radical nature of the process (inhibi-

tion by TEMPO). To confirm that the chain radical mechanism was responsible for the rearrangement, a cross-over experiment involving an equimolar mixture of ynamides **3al** and **3bd** was also performed (Scheme 3(d)). If cage recombination was involved, then only products **4al** and **4bd** should be obtained. In the case of a chain process, products **4al** and **4bd** in admixture with **4aa** and **4bo** should be expected. After 18h, four α -sulfonyl amides were formed, thus supporting the radical chain process. The 0.4:0.5:0.35:0.5 molar ratio of compounds **4aa**:**4al**:**4bd**:**4bo** was determined from the ^1H NMR spectrum of the crude reaction mixture.

Since the proposed mechanism involves the homolytic cleavage of the N—S bond, theoretical investigations were conducted in order to estimate N—S bond dissociation energy in ynamides **H**, **I**, and **J** chosen as models for *N*-aryl- and *N*-alkyl-sulfonyl ynamides (Scheme 4). Additional calculation of the C—S BDE in ketenimines showed the exothermicity of the process. It is to be noted that the presence of the triple bond lowers the N—S BDE by approximately 18 kcal/mol as compared to *N,N*-dimethyl-*N*-phenylsulfonyl amine **K**.




Scheme 4. Molecular Modelling: N—S and C—S bond dissociation enthalpies calculated in kcal/mol at the Mo6-2X/6-311++G(3df,3pd)//Mo6-2X/6-31+G(d,p) level for model ynamides and ketenimines.

The behaviour of **3ah** recovered unchanged after 18 h of irradiation is surprising.¹⁶ Both the similarity of UV absorption spectra and the lowering of the N—S bond BDE (Scheme 4, ynamide **J**) with respect to the others ynamides would lead to the conclusion that rearrangement should be very easy in that particular case. Reaction pathways, i.e., addition followed by β -fragmentation, calculated at the Mo6-2X/6-311++G(3df,3pd)//Mo6-2X/6-31+G(d,p) level for model ynamides **H_{Ph}** and **J_{Ph}**, confirmed the expected similar reactivity. Even the possibility of a competitive photo-Fries rearrangement¹⁷ was simulated for **J_{Ph}**; it might be a plausible alternative path (see SI). Finally, we tried to carry out N to C migration of sulfonyl group of **3ah** under thermic activation. While the reaction conducted in decaline at 120 °C on ynamide **3aa** led after acidic treatment to α -sulfonyl amide **4aa** in 96 %yield, a complex mixture was obtained when reacting **3ah** under the same ex-

perimental conditions. The desired compound **4ah** was detected as traces in the crude mixture on the ¹H NMR spectrum.

Thus the presence of a phenyl group at nitrogen remains the unexplained limitation of the scope of the reaction.



Scheme 5. Scale up

Finally, this protocol was scaled up to 1.6 mmol of ynamides **3aa** and **3al**. (Scheme 5). α -Sulfonyl amides **4aa** and **4al** were isolated in quantitative yields, showing the synthetic utility of the process.

Conclusion. In summary, a light-promoted synthesis of α -sulfonylated amides from *N*-sulfonyl ynamides has been developed. A wide range of α -sulfonyl amides were prepared through a radical chain process initiated by homolytic cleavage of the N—S bond under irradiation at 365 nm. This photochemical rearrangement features high atom economy, large substrate scope, mild / neutral reaction conditions, and high efficiency.

ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are available in the published article and its supporting information.

Supporting Information

Experimental procedures, Computational details, and NMR spectra (PDF file). The Supporting Information is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Author

* Aix Marseille Univ, CNRS, ICR, Marseille, 13013, France. laurence.feray@univ-amu.fr, ORCID : 0000-0001-8931-9621

* Aix Marseille Univ, CNRS, ICR, Marseille, 13013, France. michele.bertrand@univ-amu.fr

Notes

The manuscript was written through contributions of all authors. / All authors have given approval to the final version of the manuscript. The authors declare no competing financial interest.

REFERENCE

(1) (a) Chang, M.-Y.; Chang, B.-R.; Tai, H.-M.; Chang, N.-C. An Efficient Synthesis of N-Benzyl-3-Sulfonyl Glutarimides. Formal Synthesis of the Aromatase Inhibitor AG-1. *Tetrahedron Letters* **2000**, *41*, 10273–10276. (b) Chang, M.-Y.; Chen, S.-T.; Chang, N.-C. Reaction of Different A-Sulfonyl Acetamides with Methyl Acrylate. *Tetrahedron* **2002**, *58*, 5075–5080. (c) Ghosh, S.; Tiwari, P.; Pandey, S.; Misra, A. K.; Chaturvedi, V.; Gaikwad, A.; Bhatnagar, S.; Sinha, S. Synthesis and Evaluation of Antitubercular Activity of Glycosyl Thio- and Sulfonyl Acetamide Derivatives. *Bioorganic & Medicinal Chemistry Letters* **2008**, *18*, 4002–4005. (d) Kim, S.-H.; Johnson, J. A.; Jiang, J.; Parkhurst, B.; Phillips, M.; Pi, Z.; Qiao, J. X.; Tora, G.; Ye Chen, A.; Liu, E.; Yin, X.; Yang, R.; Zhao, L.; Taylor, D. S.; Basso, M.; Behnia, K.; Onorato, J.; Chen, X.-Q.; Abell, L. M.; Lu, H.; Locke, G.; Caporuscio, C.; Adam, L. P.; Gordon, D.; Wexler, R. R.; Finlay, H. J. Identification of Substituted Benzothiazole Sulfones as Potent and Selective Inhibitors of Endothelial Lipase. *Bioorg. Med. Chem. Lett.* **2019**, *29*, 1918–1921. (e) Khazir, J.; Mir, B. A.; Pandita, M.; Pilcher, L.; Riley, D.; Chashoo, G. Design and Synthesis of Sulphonyl Acetamide Analogues of Quinazoline as Anticancer Agents. *Med Chem Res* **2020**, *29*, 916–925.

(2) Liu, C.; Li, Z.; Weng, Z.; Fang, X.; Zhao, F.; Tang, K.; Chen, J.; Ma, W. Transition-Metal-Free Selective C(Sp³)-H Thiolation of Arylacetamides with Substituted Benzenethiols, Aryl Sulfonylchlorides and Diaryl Disulfides. *Asian J. Org. Chem.* **2020**, *9*, 668–672, and refs cited there in.

(3) (a) Prasad, Ch. D.; Sattar, Moh.; Kumar, S. Transition-Metal-Free Selective Oxidative C(Sp³)-S/Se Coupling of Oxindoles, Tetralone, and Arylacetamides: Synthesis of Unsymmetrical Organochalcogenides. *Org. Lett.* **2017**, *19* (4), 774–777. (b) Wang, X.; Bai, X.; Wu, C.; Dong, Y.; Zhang, M.; Fan, L.; Tang, L.; Yang, Y.; Zhang, J. Direct C(Sp³)-H Sulfonylation and Sulfuration Reactions of Isoquinoline-1,3(2*H*,4*H*)-diones under Metal-free Conditions. *Asian J Org Chem* **2021**, *10* (2), 386–391.

(4) For reviews reminding of historical background see: (a) Zifcsak, C. A.; Mulder, J. A.; Hsung, R. P.; Rameshkumar, C.; Wei, L.-L. Recent advances in the chemistry of ynamines and ynamides. *Tetrahedron* **2001**, *57*, 7575–7606. (b) DeKorver, K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. Ynamides: A Modern Functional Group for the New Millennium. *Chem. Rev.* **2010**, *110*, 5064–5106. (c) Evano, G.; Coste, A.; Jouvin, K. Ynamides: Versatile Tools in Organic Synthesis. *Angew. Chem. Int. Ed.* **2010**, *49*, 2840–2859. (d) Wang, X.-N.; Yeom, H.-S.; Fang, L.-C.; He, S.; Ma, Z.-X.; Kedrowski, B. L.; Hsung, R. P. Ynamides in Ring Forming Transformations. *Acc. Chem. Res.* **2014**, *47*, 560–578. (e) Evano, G.; Theunissen, C.; Leconte, M. Ynamides: Powerful and Versatile Reagents for Chemical Synthesis. *Aldrichimica Acta* **2015**, *48*, 59–70. (f) Evano, G.; Michelet, B.; Zhang, C. The anionic chemistry of ynamides: A review. *C. R. Chimie* **2017**, *20*, 648–664.

(5) For reviews describing recent developments see: (a) Zhou, B.; Tan, T.-D.; Zhu, X.-Q.; Shang, M.; Ye, L.-W. Reversal of Regioselectivity in Ynamide Chemistry. *ACS Catal.* **2019**, *9*, 6393–6406. (b) Hong, F.-L.; Ye, L.-W. Transition Metal-Catalyzed Tandem Reactions of Ynamides for Divergent N-Heterocycle Synthesis. *Acc. Chem. Res.* **2020**, *53*, 2003–2019. (c) Luo, J.; Chen, G.-S.; Chen, S.-J.; Yu, J.-S.; Li, Z.-D.; Liu, Y.-L. Exploiting Remarkable Reactivities of Ynamides: Opportunities in Designing Catalytic Enantioselective

Reactions. *ACS Catal.* **2020**, *10*, 13978–13992. (d) Lynch, C. C.; Sripada, A.; Wolf, C. Asymmetric Synthesis with Ynamides: Unique Reaction Control, Chemical Diversity and Applications. *Chem. Soc. Rev.* **2020**, *49*, 8543–8583. (e) Hu, Y.-C.; Zhao, Y.; Wan, B.; Chen, Q.-A. Reactivity of Ynamides in Catalytic Intermolecular Annulations. *Chem. Soc. Rev.* **2021**, *50*, 2582–2625. (f) Shandilya, S.; Protim Gogoi, M.; Dutta, S.; Sahoo, A. K. Gold-Catalyzed Transformation of Ynamides. *The Chemical Record* **2021**, *21*, 4123–4149. (g) Dutta, S.; Mallick, R. K.; Sahoo, A. K. Regioselective Difunctionalization and Annulation of Ynamide. *Angew Chem Int Ed* **2023**, *62*, e202300816. (h) Iftikhar, R.; Mazhar, A.; Iqbal, M. S.; Khan, F. Z.; Askary, S. H.; Sibtain, H. Ring Forming Transformations of Ynamides via Cycloaddition. *RSC Adv.* **2023**, *13*, 10715–10756. (i) Yang, L.; Hou, A.; Jiang, Q.; Cheng, M.; Liu, Y. Methodological Development and Applications of Tryptamine-Ynamide Cyclizations in Synthesizing Core Skeletons of Indole Alkaloids. *J. Org. Chem.* **2023**, *88*, 11377–11391.

(6) For the two-step synthesis of α -sulfonyl amides from terminal ynamides see: Shen, Y.; Li, Q.; Xu, G.; Cui, S. Coupling of Carboxylic Acids with Ynamides and Subsequent Rearrangement for the Synthesis of Imides/Amides. *Org. Lett.* **2018**, *20*, 5194–5197.

(7) Bendikov, M.; Duong, H. M.; Bolanos, E.; Wudl, F. An Unexpected Two-Group Migration Involving a Sulfonynamide to Nitrile Rearrangement. Mechanistic Studies of a Thermal N–C Tosyl Rearrangement. *Org. Lett.* **2005**, *7*, 783–786.

(8) Shen, J.; Wu, Z.; Liu, Y.; Bai, Y.; Qiu, J.; Zhang, Z.; Yuan, Z.; Zhu, G. Radical Chain Isomerization of N-Sulfonyl Ynamides to Ketenimines and Its Application to Furan Dearomatization. *Org. Lett.* **2021**, *23*, 9321–9326.

(9) Wang, L.; Lu, C.; Yue, Y.; Feng, C. Visible-Light-Promoted Oxo-Sulfonylation of Ynamides with Sulfonic Acids. *Org. Lett.* **2019**, *21*, 3514–3517.

(10) (a) Dwadnia, N.; Lingua, H.; Mouysset, D.; Mimoun, L.; Siri, D.; Feray, L. Intermolecular Addition of Carbon-Centered Radicals to Ynamides. A Regio- and Stereoselective Route to Persubstituted α -Iodo-Enamides. *J. Org. Chem.* **2020**, *85*, 4114–4121. (b) Galibert-Guijarro, A.; Mouysset, D.; Mimoun, L.; Bertrand, M. P.; Feray, L. Ynamides in Radical Reactions: A Route to Original Persubstituted 2-Aminofurans. *J. Org. Chem.* **2023**, *88*, 2464–2473.

(11) The fragmentation of beta-sulfonyl radicals is well documented in the literature. (a) For a selected example involving N-S bond cleavage see: Zhang, H.; Hay, E. B.; Geib, S. J.; Curran, D. P. Radical Cyclizations of Cyclic Ene Sulfonamides Occur with β -Elimination of Sulfonyl Radicals to Form Polycyclic Imines. *J. Am. Chem. Soc.* **2013**, *135*, 16610–16612. (b) For examples of vinyl radical fragmentation, see: García Ruano, J. L.; Alemán, J.; Parra, A.; Marzo, L. Sulfonyl Acetylenes as Alkynylating Reagents Under Radical or Anionic Conditions. *Eur. J. Org. Chem.* **2014**, 1577–1588. For recent general reviews see: (c) Chu, X.-Q.; Ge, D.; Cui, Y.-Y.; Shen, Z.-L.; Li, C.-J. Desulfonylation via Radical Process: Recent Developments in Organic Synthesis. *Chem. Rev.* **2021**, *121*, 12548–12680. (d) Flynn, A. J.; Ford, A.; Maguire, A.R. Synthetic and mechanistic aspects of sulfonyl migrations. *Org. Biomol. Chem.* **2020**, *18*, 2549–2610.

(12) Zhang, X.; Zhang, Y.; Huang, J.; Hsung, R. P.; Kurtz, K. C. M.; Oppenheimer, J.; Petersen, M. E.; Sagamanova, I. K.; Shen, L.; Tracey, M. R. Copper(II)-Catalyzed Amidations of Alkynyl Bromides as a General Synthesis of Ynamides and Z-Enamides. An Intramolecular Amidation for the Synthesis of Macrocyclic Ynamides. *J. Org. Chem.* **2006**, *71*, 4170–4177.

(13) 4 h reaction time is needed to consume all the substrate.

(14) Tosylated derivatives react much faster than mesylated substrates that necessitate 18 h for reaction completion. All experiments were conducted over 18 h reaction time for the sake of homogeneity.

(15) Ketenimine derived from ynamide **3aa** has been clearly identified through ¹H NMR proton realized on the crude reaction mixture before work-up with HCl (2 M).

(16) A longer irradiation time (48 h) and a change in wavelength (400 or 450 nm) do not change the result.

(17) Park, K. K.; Lee, J. J.; Ryu, J. Photo-Fries Rearrangement of N-Arylsulfonamides to Aminoaryl Sulfone Derivatives. *Tetrahedron* **2003**, *59*, 7651–7659.