



HAL
open science

On the kinetics of oxidation of ammonia and interactions with nitric oxide and sulfur dioxide

Philippe Dagaut

► **To cite this version:**

Philippe Dagaut. On the kinetics of oxidation of ammonia and interactions with nitric oxide and sulfur dioxide. International workshop on science and technology supporting energy conversion towards carbon neutrality, Tohoku University, Université d'Orléans, Jul 2024, ORLEANS, France. hal-04645045

HAL Id: hal-04645045

<https://cnrs.hal.science/hal-04645045v1>

Submitted on 11 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

On the kinetics of oxidation of ammonia and interactions with nitric oxide and sulfur dioxide.

Philippe Dagaut

CNRS, Orléans, France.

Ammonia has been used extensively for the selective non-catalytic reduction (SNCR) of nitric oxide (NO). This process efficiently reduces NO over a limited temperature range centered around 1200 to 1250 K. Experimental and kinetic modelling studies of SNCR are numerous. The presence of SO₂ may alter the SNCR process.

With increasing interest for ammonia as energy carrier or fuel, it is interesting to consider interactions between ammonia, nitric oxide and sulfur dioxide since they may play a role. under practical conditions where turbulence and recirculation favor the interaction between ammonia and nitrogen or sulfur.

We report results obtained in a jet-stirred reactor (JSR) at 1 bar for several equivalence ratios in the range 0.1 to 2, for several initial mole fractions of ammonia, sulfur dioxide and nitric oxide. Kinetic modelling of JSR experiments and burning velocities was performed and reaction pathway analyses were performed to interpret the results. The effect of SO₂ on the extent of NO reduction was controversial in the literature. We confirm that SO₂ does not lower the efficiency of the thermal de-NO_x process but shifts the optimal temperature to higher values through chain terminating processes ($H+H+M=H_2+M$) and radical pool reduction ($H+O_2+M=HO_2+M$) via the sequence of reactions $SO_2+H+M=>HOSO+M$, $HOSO+O_2=>HO_2+SO_2$, $HOSO+H=>SO_2+H_2$.

Acknowledgements: Support from the CAPRYSES project (ANR- 11-LABX-006–01) funded by ANR through the PIA (Programme d'Investissement d'Avenir) is gratefully acknowledged.