



HAL
open science

Orienting gaze toward a visual target: Neurophysiological synthesis with epistemological considerations

Laurent Goffart

► **To cite this version:**

Laurent Goffart. Orienting gaze toward a visual target: Neurophysiological synthesis with epistemological considerations. 2024. hal-04698698v2

HAL Id: hal-04698698

<https://cnrs.hal.science/hal-04698698v2>

Preprint submitted on 26 Nov 2024 (v2), last revised 14 Dec 2024 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Orienting gaze toward a visual target:

Neurophysiological synthesis with epistemological considerations

Laurent Goffart ^{1*}

1 : Aix Marseille Université, CNRS, Centre Gilles Gaston Granger, Aix-en-Provence, France

* : Correspondence: laurent.goffart@cnr.fr

Abstract: The appearance of an object triggers an orienting gaze movement toward its location. The movement consists of a rapid rotation of the eyes, the saccade, which is accompanied by a head rotation if the target eccentricity exceeds the oculomotor range, by a slow eye movement if the target moves. Completing a previous report, we explain the numerous points that lead to questioning the validity of a one-to-one correspondence relation between measured physical values of gaze or head orientation and neuronal activity. Comparing the sole kinematic (or dynamic) numerical values with neurophysiological recordings carries the risk of believing that central neuron activity directly encodes gaze or head physical orientation rather than mediating changes in extraocular and neck muscle contraction, not to mention possible changes happening elsewhere (in posture, in the autonomous nervous system and more centrally). Rather than reducing mismatches between extrinsic physical parameters (such as position or velocity errors), eye and head movements are behavioral expressions of intrinsic processes that restore a poly-equilibrium, i.e., balances of activities opposing antagonistic visuomotor channels. Past results obtained in the cat and monkey left a treasure of data allowing a synthesis, which illustrates the tremendous complexity underlying the mere changes in the orientations of the eyes and head. The aim of this synthesis is to serve as a guide for further investigations or for comparison with other species.

Keywords: poly-equilibrium; saccade; pursuit; fixation; neurophysiology; monkey; cat; kinematics; dynamics; model



1. Introduction


Among the numerous relations that an entire animal establishes with its environment, the orienting movement was once proposed to underlie this interrogative process which "*in Man has strongly developed in this highest form which is curiosity - the parent of that scientific method by which we hope one day to achieve a correct orientation in the knowledge of the world around us*" [1]. This view remains popular in the domain of contemporary cognitive neuroscience wherein orienting is a module underlying exploratory behavior, gathering sensory signals, transforming them into information and storing them in some sort of internal copies of the external world.

In this article, we shall take a different point of view, avoiding teleological and cognitivist arguments as much as possible. We shall focus on the orienting movements of the eyes and head toward the location of an object. We shall explain how these two movements are behavioral outcomes of intrinsic brain processes, which are blind to the physical and informational relations occurring at the macroscopic level, that is, between the animal organism and objects in the external world. We shall start from the standpoint that the inner milieu consists of multiple equilibria maintaining some kind of stability. Whenever asymmetrical changes disrupt it, intrinsic processes restore the homeostasis. This restoration involves a coordinated set of reactions measurable at the musculoskeletal, vascular and vegetative levels. Orienting movements of gaze, head or body toward a target are the overt, externally visible consequences of activities traveling through the sensorimotor neural networks during such restorations. Other changes are less visible, such as transient pupillary dilation or cardio-vascular changes (bradycardia associated with peripheral vasoconstriction and cephalic vasodilation) [2-3], but also postural adjustments [4-6]. Synaptic changes and the persistent traces left by repetition are internal consequences that are indirectly visible by longitudinal studies. In the same way as the sandy soil conforms to the current of a river while determining at the same time its flow, the sensorimotor neuronal channels assimilate to some extent the regularity of activity flows while constituting at the same time the set of constraints for the on-going flows.

Before detailing the neurophysiological substrate of orienting eye and head movements, it is useful to recall that they do not constitute the only type of reaction to an external stimulus. Other types include distancing, avoidance and escape. Orienting differs from them by the radically opposite way in which it manifests spatially in relation to the stimulus. Contrary to them, it consists of directing the body segment that carries the teleceptive organs towards the location of the stimulus, be it detected or coveted. In this article, we shall focus on two physical characteristics of animal reactions toward a visual stimulus: its velocity and its accuracy. Indeed, the orienting response consists neither of a sliding movement nor of a sequence of small movements directed toward the location of a targeted object. For the vast majority of animals, it manifests itself rather by a sudden and extremely rapid rotation on the mobile segment (body, head or eye) that carries the photosensitive epithelium. The amplitude and direction of the rotation are confined within a range of those sufficient to symmetrize back the functional neuronal image of the stimulus within the sensorimotor networks.

At the end of the orienting movement, measuring the amplitude of the angular displacement performed by the mobile segment gives a set of numerical values that are compared to the eccentricity of the target. Figure 1 illustrates their relationship in a goldfish [7]. The amplitude of the body rotation is not constant; it increases with the angular eccentricity of the stimulus relative to the body. This relation has been documented in several other species: in frogs [8], salamanders [9], barn owls [10],

cats [11], macaque monkeys [12-13] and human subjects [14-16]. Also observed in invertebrate animals such as insects (mantis: [17-18]; dragonfly: [19]) and crustaceans [20], it suggests that, beyond the diversity of photosensitive surfaces (retina or compound eyes), of muscle fibers, of their attachments - indirect or direct - to the mobile segment, the extension of physical space has somehow been embedded within the neural networks.



In cats and monkeys, the relation is not a fixed mapping between the topology of the photosensitive surface and a set of specific temporal commands changing the orientation of eyes and head. Experiments in these animals showed that this behavioral invariant is robust to changes in saccade trajectory induced by a brief electrical stimulation in some territories in the brainstem (cat: [21-22]; monkey: [23-27]) or in the cerebral cortex (monkey: [28-29]). However, when the perturbation is applied in some sites of the pontine reticular formation [24] or in the medio-posterior cerebellum [30-31], the tested monkey orients its gaze toward a location that is shifted with respect to the target location. The saccade landing position is shifted by an offset equal to the stimulation-induced change in eye position. This targeting error confirms that extraocular muscle proprioception does not participate in guiding saccades toward their target [32-33].

2. Fundamental and epistemological issues

In response to the same initial physical conditions, the amplitude of the response is not constant but variable within some limits. Beyond the multiplicity of possible channels by which retinal signals can reach the motor neurons and beyond the time course of the persistence of signals, this variability, which is visible in Figure 1, is also due to the facts that objects in the external world are not points and that visual fixation does not involve the projection of their image onto one single light sensing cell, but several ones. In some species, the photosensitive cells are grouped within the foveal pit. We shall use the notion of “gaze” instead of the so-called “line of sight” to denote the

imaginary “tube” through which photons travel, from the fixated object in the external world to the foveal photoreceptors.

Despite quasi-inevitable numerical differences, some degree of congruence must nevertheless be considered between the spatial location of a visual stimulus and the zone toward which gaze is directed. As mentioned above, this overlap is preserved even when the functioning of some key sensorimotor regions is perturbed. Before reporting these findings and discussing their implications, we shall address a few epistemological issues related to the risk of importing in the neurophysiological study of visual orienting notions whose biological plausibility remains uncertain or dubious [see also 34].

2.1. Distinguishing the medium of cerebral activity from physical space

The “binocular fixation point” is the intersection between two imaginary lines that characterize the direction of each visual axis while the “vergence angle” is the angle made by these two lines [35]. When that angle decreases, the eyes are said to converge and one speaks of convergence. When it increases, they are said to diverge (divergence). In the subject with no strabismus, when a saccade shifts gaze between two targets located along an iso-vergence curve, the spatial trajectory of the binocular fixation point follows neither the shortest physical path between the targets (the straight line) nor a path along the iso-vergence curve. Relative to the frontal plane, the binocular fixation point takes a path away from the iso-vergence curve followed by a return to it: a transient divergence happens during the saccade. The path traveled by gaze lies in a “geodesics” that does not belong to the zero-curvature space of Euclidean geometry. There is no force field in the external world that constrains gaze to take such a path. The path is intrinsically determined, caused by the neuronal networks upstream of the extraocular muscles.

As pointed out by Pellionisz and Llinas, *“since within the brain there is no ‘instantaneous’ simultaneity agent comparable to the light, the classical usage of separate space and time coordinates is inapplicable in the case of describing the inner workings of the central nervous system”* [36]. From a naive point of view, the timeless entity that geometers and physicists designate as “space”, the receptacle occupied by objects, might correspond to the medium within which flows of neuronal activity propagate, from the excitation of sensory cells by external objects to the time when the orienting reaction ends. However, the cerebral networks radically differ from the three-dimensional framework that assigns numerical values to positions and displacements of objects. They constitute a medium that is neither isotropic nor homogeneous, but composed of multiple and diverse channels. Each physical place captured by gaze is the outcome of multiple flows of activity through this medium. The divergence characterizing the afferent pathways first broadens the neural image of a small spot of light, an image then compressed by the convergence of activities towards the motor nerves. Transforming the recruitment of excitatory and inhibitory synapses and the firing rate of neurons to algebraic time functions engenders a continuum that is not in accordance with the remote and distributed conditions of action potential triggering [34].

The multiple sensorimotor channels cannot correspond to a neural representation (or correlate) of physical “space” for other reasons. The numerical values assigned to different physical target positions or movement endpoints are independent of each other. They are ordered and aggregate to each other within another continuum, which corresponds to an infinite number of

simultaneous points (mathematical space). In the brain, there is no corresponding entity. The elements are neither compact nor rigid, and their number is much smaller than the number of objects present in the external environment of a subject. Activities in the brain corresponding to different physical locations are neither points nor lines. Those leading to the generation of movements towards nearby but different locations can involve common neurons. Indeed, neurons have extended and overlapping response fields [37]. The sharing of neuronal elements is such that two simultaneous visual targets can lead to a saccade (called averaging saccade), which directs gaze toward a location situated midway between the two targets or along the line connecting them [38-39].

Moreover, measuring a sequence of motor responses to targets located at different places in the visual field is not equivalent to measuring the entire field. The measurement values obtain their status of simultaneity after their plot on two- or three-dimensional graphs. Whatever corresponds to physical space or to objects in the brain, it is not permanent, but short-lived. Its persistence requires either the presence of objects in the external world or in working memory, which adds numerous contextual non-spatial values, past and present [40-41]. When we consider the accuracy and precision with which saccades indicate the location of a target, they depend considerably on its presence. When it is produced shortly after target appearance, the saccade is sufficiently accurate to confirm the presence of the object by a set of sensations, which are relatively always the same and characterize the foveation. Regarding the object location, the kind of "measurement" made by saccades is, so to speak, reliable. By contrast, when the saccade is made after a delay, even after a relatively short one, the "measurement" is no longer reliable [42-46]. Visual sensations can no longer confirm the presence of the object since it has disappeared while the change in saccade accuracy and precision make the movement-related sensations uncertain. Saccades are also strongly inaccurate (hypermetric) when they are made with the eyes closed [6, 47]. In other words, it is as if the persistence of visual signals in the brain required the presence of objects to create a "space", in which case its internal representation (or neuronal copy) is either weakly reliable or unnecessary since it is already out there [48].

Finally, neural plasticity, the irreversibility of activities and their canalization argue for a fundamental difference between the cerebral medium and the Newtonian receptacle. Yet, the adaptation of movements to environmental constraints led some authors to propose that brain activity emulates internal models of the physical laws of motion. Unfortunately, they missed to explain how the diversity of animals and nervous systems is compatible with the three Newton's laws of motion. Leaving aside the principle according to which rest corresponds to a balance of forces that cancel each other out, they did not explain either how brain activity could sustain at constant speed the motion of a body segment with muscles of finite length. If the laws of mechanics are conventions [49-50] used to quantify the movement of objects in space, then it draws into question whether it is reasonable to assume their applicability to the brain sciences. The consistency and viscosity of the cerebral medium in which activities propagate are not as homogeneous as those of the environments (water, ground and air) in which the animal body can evolve. As for the eye movements, their study in untrained monkeys shows a tracking that is not as smooth as the continuous and uniform motion of a visual target. The visual tracking is discontinuous and saltatory, interrupted by saccades [51-53] indicating that the streams of visuo-motor activity do not spontaneously mimic the constant motion of the target. The fact that repeated practice is required to perform spatially and temporally accurate movements does not imply cerebralization of physical laws. It merely illustrates the flexibility of the brain's functioning to perform accurate movements, perhaps by eliminating irrelevant or distracting stimuli.

Refusing to import notions pertaining to the kinematic or kinetic description of movements onto the activity of neurons does not lead the neurophysiology of eye-head movements to a dead end. We shall see that it actually offers the opportunity to make a fresh start and investigate how the brain works differently from the way that has been promoted during the last decades.

2.2. The spatiotemporal transformation

A fundamental challenge in the brain sciences is to identify and extract processes from the complex networks of cells that are often polyvalent. Between the retinal ganglion cells and the extraocular and neck muscle fibers, some neurons are involved in multiple tasks, making the delineation of processes far from being a trivial task. The mere excitation of a small group of ganglion cells in the retina by a stimulus as simple as a brief spot of light is transmitted to a multitude of post-synaptic neurons in several deep cerebral nuclei (e.g., lateral geniculate nucleus, superior colliculus, nucleus of the optic tract). The action potentials emitted by these neurons are then transmitted to more numerous post-synaptic cells distributed in multiple areas of the cerebral cortex. The spot of light elicits a neural activation that spreads out within the brain's networks. Its expansion is associated with a temporal persistence insofar as the duration of elicited activities outlasts that of the stimulus. This maintained activity is necessary for changing the firing rate of specific groups of motor neurons and for driving the contraction of extraocular and neck muscles until gaze is oriented toward the location of the spot of light, sometimes a long time after its disappearance.


The process by which different loci and more distant stimuli in the visual field (i.e., more eccentric loci of retinal activity relative to the fovea) lead to different and longer motoneuronal discharges has been called "spatio-temporal transformation". Such a transformation is not limited to orienting movements toward a visual target. It characterizes all movements directed toward events that are spatially delimited such as an object in the visual field or a point stimulation on the surface of the skin, as well as toward events whose physical location is more diffuse such as sounds or screams. In all cases, a premature arrest of motoneuronal activities will make the movement hypometric, i.e., falling short of the target location.

Initially, the dysmetria of movements made by patients suffering from cerebellar dysfunction led to propose that the cerebellum performed this spatiotemporal transformation [54-55]. The afferent signals from retinotopically organized maps such as the one found in the superior colliculus were thought to be converted into a specific temporal pattern of motor activity (see Fig. 10 in [56]). However, the observation of accurate saccades in patients suffering from spinocerebellar degeneration, despite dramatic slowing and increased duration, led to abandon this hypothesis [57]. The conjecture was then proposed that saccade duration be determined by the work of a negative feedback loop [57-59]. The premotor neurons would be driven by a motor error command that results from the continuous comparison between two signals: a reference signal specifying the goal or desired state (desired eye orientation or desired change in eye orientation), and a feedback signal estimating the current state (current eye orientation or actual change in eye orientation). As long as a mismatch persists between these two signals, the motor error command drives the premotor neurons until the estimated current state matches the desired state. In this theoretical framework, there is no signal specifying the duration of saccades; it merely corresponds to the time taken by a feedback signal to zero out the motor error command. Saccade inaccuracy would result either from an altered reference signal or from an impaired feedback signal. Transient changes in the trajectory of saccades should not

affect their accuracy as long as it does not alter the signal that specifies the saccade goal, or the amplitude and direction required to foveate the target [20-28].

For several years, the negative feedback loop remained the simplest and widely shared hypothesis to account for the spatiotemporal transformation. The reference signal was shown to be conveyed by the population of active neurons in the intermediate and deep layers of the superior colliculus [60-62] while the comparison with feedback signals took place downstream in the reticular formation, under the influence of neurons in the medio-posterior cerebellum [29].

Figure 2 illustrates how the direction (A) and amplitude (B) of saccades toward a visual target are preserved in spite of their dramatic slowing when the activity of some neurons in the deep superior colliculus is suppressed by local injection of a blocker of sodium channels [63]. After each injection of a small amount of lidocaine, the peak velocity of saccades is greatly reduced (C) while their duration is increased (D) by an amount that appears proportional to the amount of velocity reduction. In this experiment, the injection did not silence all collicular neurons, but only cells that fired maximally for saccades having a particular direction and amplitude. Saccades with amplitudes and directions that were close to this optimal vector were inaccurate [61]. In addition to demonstrating that saccade metrics was specified by the population of active neurons in the superior colliculus [60], this study brought evidence for a causal link between collicular activity and saccade velocity and for a feedback control of saccade amplitude and direction (see also [64]).



Around the same time, works in head unrestrained cats led to propose that the comparison between the reference and feedback signals was performed within the superior colliculus itself, by a caudo-rostral migration of activity [65-66]. The change in the locus of collicular activity would switch from the recruitment of saccade-related neurons to the recruitment of fixation-related cells. The former neurons would be distributed across the entire extent of the SC whereas the latter would be concentrated in its rostral part. The saccade would stop as the “fixation” cells resume their sustained presaccadic firing rate, which in turn, would excite cells in the nucleus raphe interpositus (RIP) of the medial pontine reticular formation. Therein, neurons called omnidirectional pause neurons (OPNs) exhibit a sustained spontaneous firing rate and pause during each saccade, regardless of its amplitude and direction. Their sustained activity was proposed to inhibit premotor burst neurons and pause to trigger a saccade.

However, subsequent studies failed to observe any migration of activity across the SC during saccades [67-70, but see 71]. This lack of reproducibility led to abandon the conjecture (called “moving hill” hypothesis) of a caudo-rostral change in the locus of collicular activity (exception in [72]). Later, confirming observations reported in an earlier study [73], neurons in the rostral SC were shown to emit a burst of action potentials during miniature saccades [74-75], extending the saccade-related function to this part of the SC, as a microstimulation study suggested [25]. Indeed, contrary to electrical microstimulation of RIP, which only decelerates and interrupts visual saccades, microstimulation of the rostral SC deviates their trajectory toward the contralateral side. The absence of hypermetria after pharmacological inactivation of the rostral SC [76-77] further called into question its role in terminating saccades whereas the reduced rate of microsaccades during visual fixation [74, 77] confirmed its involvement in the generation of miniature saccades [78]. Finally, the lack of impairment in either the latency or the accuracy of saccades after RIP lesion [79-80] questioned the involvement of OPN in fixation and saccade termination. It also casted doubt on the idea that silencing the activity of OPNs disinhibit premotor burst neurons and allow their excitation by descending SC commands [81]. Indeed, the suggestions were made i) that OPNs’ sustained activity prevents the visual response of SC neurons from driving premotor burst neurons and ii) that saccades with very short latencies (also called express saccades) were driven by SC visual responses [82-83]. The demonstration of a loose coupling between the onset of saccades and the visual (rather than motor) responses of SC neurons [84-85] contradicted these ideas. As for saccades with regular latencies [86], the onset of express saccades is tightly coupled with the onset of motor burst of SC neurons [84]. The variability of saccade onsets likely depends upon the synchronies of action potentials impinging upon the premotor burst neurons: weaker synchrony yielding scattered and longer reaction times. As explained elsewhere [52], simultaneous excitatory action potentials from an assembly of neurons are indeed more efficient in exciting their post-synaptic targets than temporally scattered action potentials.

Altogether, these numerous studies and others led to realize that the spatiotemporal transformation is not explicitly performed by distinct groups of neurons, which neurophysiologists ought to search and identify within the brain. Instead, the transformation would be supported by activities of populations of neurons that are massively interconnected and distributed over several neural territories [87]. We shall see how this distributed character also holds for the cerebral control of gaze direction during visual fixation.

2.3. Gaze direction as a poly-equilibrium

Several lines of evidence led to propose an alternative to the “fixation-saccade” dichotomy [88-91]. First, the absence of fixation instability during lesion or during pharmacological inactivation of RIP [70-71] questions the necessary involvement of OPNs in maintaining stable gaze. Second, several studies made in awake animals report cases in which the animal does not direct accurately its gaze toward a visual target during fixation, or does not correctly orient its head toward a food target. During unilateral inactivation of the caudal fastigial nucleus (cFN), cats direct their gaze and mouth toward a location that is offset with respect to the physical target location [10]. A comparable eccentric fixation happens in head restrained monkeys during unilateral pharmacological inactivation of cFN. Therein, the local and unilateral injection of an inhibitory agent (muscimol) leads to an ipsilesional fixation offset [92-95] whereas the injection of a disinhibitory agent (bicuculline) yields a contralesional offset [96]. The muscimol-induced fixation offset is not a defect in positioning the eyes in the orbit, but a gaze-related disorder because its size is quasi-identical when comparing cases in which visual fixation is made with the head restrained versus unrestrained [97]. Moreover, when the head is free to move, a large ipsilesional deviation affects its orientation and the deviation of the eyes in their orbit is contralesional [97]. Finally, a weakening of stability accompanies the fixation offset insofar as monkeys have difficulty to maintain their gaze near the central target when another behaviorally relevant target appears in the peripheral visual field. Indeed, the animals make irrepressible saccades toward the peripheral target location during delayed oculomotor tasks (unpublished observations; see also the increased rate of no-go errors in [98]).

Anatomical studies report that cFN neurons project neither to the nucleus prepositus hypoglossi nor to the medial vestibular nucleus [99], nuclei in which tonic eye position neurons are located [100]. However, cFN neurons project to the rostral part of both superior colliculi [101]. Thus, during unilateral pharmacological perturbation (inactivation or disinhibition), asymmetrical and sustained activities from neurons in the left and right cFNs would affect the balance of bilateral activity in rostral SC whenever gaze is directed toward a target in the central visual field. This conjecture is confirmed by the consequence of unilaterally inactivating small sectors in the rostral SC. After local injection of small amounts of muscimol, monkeys exhibit an offset when they fixate a target, regardless of whether it is static [77] or moving [102]. They also make irrepressible saccades toward the peripheral target during a memory-guided saccade task [76]. Asymmetrical and sustained activities between the left and right cFNs and rostral SCs are transmitted to pre-cerebellar pontine nuclei, explaining eccentric fixation during pursuit eye movements [102-103]. Concerning the ipsilesional deviation of the head during unilateral cFN inactivation [10, 97], it would also result from asymmetrical sustained input from fastigial nuclei to the group of reticulospinal neurons innervating the neck motor neurons [104-106].

Offsets while directing the eyes toward the location of a static visual target have also been reported in monkeys after unilateral lesion [107] of frontal eye fields or during unilateral inactivation of a small part of them [108]. By contrast, unilateral lesion of area LIP, another major source of cortical input to SC [109-110], do not report fixation offsets [111-113]. However, a spontaneous nystagmus in the dark is observed during the first week after unilateral ablation of area 7 of parietal cortex [114]. Its slow eye movements are directed toward the contralesional side. Clinical studies of patients suffering from an acute hemisphere stroke report a conjugate deviation of the eyes [115-116] and head [117] toward the lesioned side, even when subjects explore a visual scene. However,

to our knowledge, their fixational eye movements have not been studied. By contrast, an asymmetric distribution of fixational saccade amplitudes toward the blind visual field has been shown in patients suffering from homonymous hemianopia. Some subjects also exhibit eccentric fixation [118].

Fixation offsets are not restricted to lesional cases. An upward offset is observed when macaque monkeys look at a small visual target in the dark [119-121]. Asymmetrical representations of the upper and lower visual fields in these primates [122-123] can account for the offset as well as for the upward bias of horizontal saccades [121]. While the population of active neurons is immune of this asymmetry in darkness, it may be different when a visual target is presented in an illuminated background. More cells sensitive to the lower visual field may indeed be recruited and counteract the upward bias caused by the asymmetrical representation of the upper and lower visual fields. In darkness, the absence of their recruitment would then account for the upward bias of horizontal saccades. Further investigations are still required to test whether bilateral activity from neurons in the cerebellar interpositus nuclei adjusts this asymmetry [121].

Altogether, these results led to revise the view of fixation as a process inhibiting the generation of saccades [76, 88-90] and to propose the conjecture of fixation as an equilibrium [77, 124-125], later revised a “poly-equilibrium” [126]. According to this theory (Figure 3), a saccade or a slow eye movement is not initiated as long as the visuo-oculomotor system is in a mode where opposing commands counterbalance each other. For generating saccadic and pursuit eye movements, symmetry breaking involves different groups of neurons in the left and right parts of the brainstem. Saccades involve the recruitment of burst neurons located in the pontomedullary reticular formation whereas slow eye movements involve the recruitment of neurons in both nuclei of the optic tract, pontine nuclei, cerebellum (medio-posterior territories and floccular complex) and thalamus. Prior to generating a saccadic or slow eye movement, the target-related signals do not travel toward the motor neurons across a vacuous medium. They travel through a large network of cells that already evince a spontaneous sustained firing rate, which maintains the stationarity of the eyes and head. This set of tonic premotor activities, once called “neural integrator”, altogether forms an equilibrium of commands that counterbalance each other. Under normal conditions, gaze direction is relatively stable during visual fixation, occasionally interrupted by saccadic intrusions or by ocular drifts.

The notion of poly-equilibrium is more appropriate for naming this balanced activity because multiple equilibria participate (Figure 3). It involves action potentials conveyed through the channels that lead to the generation of saccades, the channels that lead to the generation of slow eye movements, those yielding the near response (accommodation and vergence) and those holding the head upright [126]. The change in head orientation during unilateral lesion of the cFN [11, 97] or the SC [127] indicates that bilateral activity also contributes to the muscle tone that specifies the static orientation of the head in the earth horizontal (yaw) plane. Sustained activities are distributed within different groups of neurons in the left and right parts of the reticular formation (including pontine long-lead burst neurons and pursuit-related neurons), the cerebellum, the deep superior colliculi and the cerebral cortex. Concerning the deep superior colliculi, the activity involves their rostral pole but it is not restricted to them. It extends more caudally, even during steady fixation. Indeed, during the gap interval, saccade-related neurons in both SC fire, even those located at sites that are sensitive to target eccentricities of 8 deg (see Fig. 4 in [84]). Any suppression of activities within the set of commands participating to this poly-equilibrium alters the direction of gaze while fixating a target if it

is not counterbalanced. As mentioned earlier, gaze is directed toward a location that is offset relative to the target, regardless of whether it is static [63, 70-75, 83] or moving [80-81].




Figure 3. Considering static gaze direction as a poly-equilibrium leads to view unbalanced activities as the cause triggering eye movements, saccadic and slow. Unbalanced activities between the left and right superior colliculi promote the generation of saccades toward the side opposite to the most active SC. By contrast, unbalanced activities between the left and right nuclei of the optic tract promotes a slow eye movement toward the side to the most active nucleus (Figure 3). This contrast is indicated by the contralateral direction of saccades evoked by electrical stimulation in the SC [128-131] and the ipsilateral direction of slow eye movements evoked by stimulation of the nucleus of the optic tract [132-133]. It is also indicated by unit recording studies. Neurons in the nucleus of the optic tract increase their firing rate during ipsilateral pursuit and decrease during contralateral movements [133-135] whereas SC neurons burst during contralateral saccades [60, 128, 136].

2.4. Movement onset as a kind of symmetry breaking

Considering static gaze direction as a poly-equilibrium leads to view unbalanced activities as the cause triggering eye movements, saccadic and slow. Unbalanced activities between the left and right superior colliculi promote the generation of saccades toward the side opposite to the most active SC. By contrast, unbalanced activities between the left and right nuclei of the optic tract promotes a slow eye movement toward the side to the most active nucleus (Figure 3). This contrast is indicated by the contralateral direction of saccades evoked by electrical stimulation in the SC [128-131] and the ipsilateral direction of slow eye movements evoked by stimulation of the nucleus of the optic tract [132-133]. It is also indicated by unit recording studies. Neurons in the nucleus of the optic tract increase their firing rate during ipsilateral pursuit and decrease during contralateral movements [133-135] whereas SC neurons burst during contralateral saccades [60, 128, 136].

2.5. The wanderings of the “brain machine”

At the time when bioengineering approaches started to guide the neurophysiological inquiry of eye movements with systems analysis techniques, David A. Robinson already warned that “block diagrams of oculomotor organization serve as a compact description of system behavior but seldom have much bearing on the way in which the real system, composed of nerve and muscle, actually operates. The models thus do not contribute much to the neurophysiology (or neurology) of eye movements and incur the danger of suggesting that there actually are segregated portions of the nervous system which perform the differentiation, integration and other operations indicated in the boxes of the diagrams” [150]. Neuroanatomical and neurophysiological studies reveal indeed functional properties and connectivity that are not visible in cybernetic diagrams. Establishing correspondences between cybernetic diagrams and the brain is analog to comparing an automaton to a living animal. It led some authors to search for various sources of signal “noise” within the brain activity rather than for differences between brains and machines. Contrary to a machine, recruiting the same group of photosensitive cells does not lead to the same response.

In spite of evidence for reliable saccade motor commands [151], some authors posited their variability [152] and even imagined various kinds of signal noise [153] to explain cerebellar control of saccade precision. Between the retinal activity and the ultimate muscle fiber twitch, the distribution of action potentials within the brain is inevitably variable between two movements toward the same target. However, the search for noisy signals should ignore neither the importance of starting eye position in determining the locus of activity on the retina nor the different brain connectivity between different subjects [153]. Moreover, even though two brief and spatially distinct spots of light excite different retinal regions, the resulting activity will likely involve common neuronal elements in the brain because of the divergent pattern of connectivity. Response fields of central neurons that overlap are the main evidence for the sharing of neuronal elements during visual stimulations that are spatially distinct. This situation does not lead to unreliable visuomotor transformation if “the precision or accuracy of a saccade results from the summation of the movement tendencies produced by the population of neurons rather than the discharge of a small number of finely tuned neurons” [60]. The effects of variable discharges of neurons is reduced by averaging the activity over a population of numerous neurons. Thus, multiple combinations of active neurons can lead to kinematically (or dynamically) identical eye movements, reducing further the strength of the correlation between the firing rate of single neuron and movement kinematics (or dynamics) [34]. While point values are assigned to the target location and the place to which gaze is directed, a geometric segment (called

error) to their difference and a curve to the movement trajectory, their brain correlates, streams of activity, are not comparable.

3. Orienting gaze toward a moving target

Many objects in the world are moving and orienting movements are also made toward them. In primates, their appearance in the peripheral visual field elicits a saccade that brings the target image onto the foveae. Then, the foveation is maintained by slow pursuit eye movements interrupted by catch-up saccades. Movement accuracy consists of directing gaze toward the location where the target is here and now, the target location at the saccade landing time. The temporal interval between the onset of sensory signals and the landing time led some authors to defend a view according to which the functioning brain would be endowed with so-called “predictive” capabilities. In this section, we shall explain that this inference is the consequence of conflating physical events and numerical values attributed to them with their physiological correspondence.

3.1. The interceptive saccade

Benefiting from high-resolution recording of eye movements, studies showed that human and non-human primates manage to direct their gaze toward the actual location of a small target moving at constant speed [154-157] with an accuracy that is quasi-similar to that of saccades toward a static target [158]. This ability to intercept with gaze a small moving target seems to depend neither upon prior experience [159] nor upon the occurrence frequency of path taken by the target [160]. It occurs even when prior to its onset the saccade is perturbed by an unexpected change in eye position [161]. However, saccades toward a moving target are not always accurate. Accuracy is lost when the target does not move at constant speed: saccades to an accelerating target fall short of it whereas those to a decelerating target overshoot it [162].

The ability to bring the image of a moving target within the central visual field led some authors to suggest an ability to foresee (or “predict”) the location that the target will occupy at the time of saccade landing. Indeed, because of the polysynaptic path between the retinal ganglion cells and the extraocular muscles, the signals that drive the firing rate of motoneurons cannot originate from the retinal site that the target is crossing presently. They originate from a site that was crossed several tens of milliseconds earlier. The delay of visuomotor transmission has no fixed value; its value likely depends upon the number of active neurons, upon their firing rate and the simultaneity of presynaptic action potentials at each relay [52]. As for saccades toward a static target [163-164], the latency of interceptive saccades may also depend upon the target contrast.

A study in non-human primates regarding saccades toward a transient target moving with a constant speed reports several saccades landing beyond the location where the target disappeared, but along the path that it would have taken if it had remained visible [162]. Because the monkeys had no prior experience with such targets (small and moving), the saccades could not be guided by memory-related signals, but by signals that persisted after the target had disappeared. Moreover, the kinetic of the retinal streak (the time course of the target motion in the visual field) was taken into account. For identical paths, the pattern of landing positions depended upon the rate of change in target speed. As mentioned earlier, saccades are accurate if the target moved at constant speed; they overshoot its current location if it decelerated and fell short if the target accelerated. It is not known

whether accelerating and decelerating targets recruit different neuronal networks, or whether the different landing positions result from different spatiotemporal patterns of activity within the same plexus of neurons. As for curved saccades made in response to a target that steps from one location to another (double step saccades [168]), the recruitment of the same plexus for accelerating and decelerating targets would be another evidence for the importance of considering the distribution of activity within the neuronal networks for the causal determination of behavioral responses. Although important, the knowledge of anatomical connectivity is not sufficient to understand the neural control of movements. Lesion studies teach us that static and moving targets involve different sets of neurons. Indeed, impairment of some brain regions can affect saccades toward a moving target without altering saccades toward a static target [165-167].

3.2. The post-saccadic slow eye movement (pursuit)

A slow eye movement (called pursuit eye movement) follows the interceptive saccade toward a moving target. Studies in monkeys revealed that its speed does not spontaneously match with the target speed and that it increases with the animal's training and prior experience [159, 169]. Comparable observations were made in children [170-171]. In response to a continuous motion, naive monkeys and children track a target with a succession of catchup saccades interspersed by intervals during which gaze direction lags behind the target, with a gaze-target distance increasing until the next catchup saccade.

Clinical studies in human subjects [172-175] and lesion studies in macaques [165, 176-179] indicate that saccadic and pursuit eye movements recruit different sets of neurons. Indeed, some lesions impair slow pursuit eye movements without altering saccades, and vice-versa. The segregation of saccade-related and pursuit-related networks is confirmed by: i) developmental studies reporting different maturation speeds of saccadic and pursuit performances [170, 180-183], ii) separate adaptive adjustments of saccadic and pursuit eye movements [184-186], iii) different consequences on postural stability [187], iv) imaging studies in healthy human subjects [188-189], and v) the electrophysiological identification of saccade-related and pursuit-related neurons in distinct regions of the pontine tegmentum [190-192] and cerebral cortex [193-199]. In the cerebral cortex, the saccade-related subregions are preferentially interconnected with each other, and likewise for the pursuit-related subregions [199]. In summary, from the cerebral cortex to the reticular formation, the networks involved in the generation of pursuit and saccadic eye movements are relatively independent and parallel.

This parallelism contrasts with the claim that a "single sensorimotor process" drives the generation of both movement categories [200-201], a conjecture that is mostly based on behavioral observations and on a mistaken interpretation of neurophysiological observations. First, the fact that neurons in the rostral SC modulate their firing rate during both miniature saccades and pursuit [202-205] is the neural substrate of looking straight ahead, i.e. moving gaze neither rightward nor leftward, neither upward nor downward, but promoting movements "in all directions at the same time" (Figure 3). As explained above, bilateral rostral SC activity maintains the equilibrium necessary to maintain gaze direction toward the target, regardless of whether it is static or moving, actual or virtual. Second, it is premature to conclude an involvement of OPNs in pursuit eye movements because their electrical microstimulation slows them [206-207]. The slowing could result from inhibiting motor neurons by the retrograde [208-209] excitation of synapses from inhibitory burst neurons. Third, the observation that

subthreshold electrical microstimulation in lobules VIc-VII of the cerebellar vermis perturbs both saccadic and pursuit eye movement [210] remains compatible with a segregation of saccade-related and pursuit-related neurons. Therein, Purkinje cells firing during both saccades and pursuit are not numerous (approximately 10% [211]). Based on a large sample of neurons, another study reported that only 4% of saccade-related Purkinje cells modulated their firing rate during pursuit [212]. More recently, the direction tuning of Purkinje cells was found to differ between saccadic and pursuit eye movements, suggesting an independent processing of their action potentials in the fastigial nuclei [213]. In the latter, approximately one third of the neurons emit a burst of action potentials during saccades and modulate their firing rate during pursuit eye movements [214]. However, emitting a burst of action potentials does not necessarily imply that a neuron and its post-synaptic target belong to the saccade premotor network. The burst of “saccade-pursuit” neurons may be transmitted to precerebellar pursuit-related pontine nuclei and contribute to the post-saccadic enhancement of slow eye movements [215]. Finally, during unilateral inactivation of cFN, disorders in saccadic and pursuit eye movements are not correlated [103], whereas during bilateral lesions lead to bilateral hypermetria of saccades [92] while the gain of pursuit eye movements is either reduced [216] or unchanged [217].

In conclusion, behavioral observations provide helpful information for guiding neurophysiological investigations when they take into account anatomo-physiological constraints. Before giving several examples illustrating this benefit, we shall discuss the capacity to foresee the invisible and to “predict the future”.

3.3. About the “predictive” power of the brain

Because saccades direct gaze toward locations that have not yet been crossed by the target, interceptive saccades have been called “predictive” and scenarios were proposed to explain anticipation. For instance, the visuo-oculomotor system would be endowed with either a temporal countdown estimating “the time remaining until the collision of the target with the line of sight” [218] or a predictive clock announcing “the time at which the eye trajectory will cross the target” [219]. This conjecture led to imagine unrealistic options such as “a spatial lead of the gaze at the saccade end, instead of attempting a precise capture of the target” [218, 220]. Indeed, the overshoot of saccades is incompatible with most published data. The majority of studies report that interceptive saccades either fall short, or land approximately at the location where the target is at their landing time [154-160, 221], even when their trajectory is perturbed by a change in eye position elicited by a brief electrical microstimulation in the SC [161]. Recent studies in monkeys showed that knowing the target path in advance does not lead to an overshoot of saccades either. Neither the accuracy nor the precision of saccades differ between blocks of trials in which the target path is always the same for one trial to the next and blocks in which the target path varies across the trials [160]. Saccades landing beyond the target location may occur in human subjects, but their occurrence is rare and conditioned by specific task instructions. They occasionally happen when the target moves at constant velocity on a periodic path. However, they are almost eliminated when the subject must accurately look at the target in order to detect a change in shape [222].

At the neuronal level, when a saccade is made toward a target moving at constant speed, saccade-related neurons in the deep SC emit a burst of action potentials as during saccades toward a static target. However, the population of active neurons does not contain commands related to future locations of a moving target. No action potentials are emitted by cells whose response field correspond

to upcoming saccadic vectors [223]. The active population consists of a continuum of cells ranging from neurons issuing commands related to past locations of the target to neurons issuing commands related to the current target location. The topography of active neurons does not change as fast as the target in the visual field because residual activities related to recently traveled locations persist.

The same conclusion holds for the results obtained in a preliminary study [224]. When comparing the response fields during saccades toward a static versus a moving target, a shift was found in the amplitude associated with the maximum firing rate. Cells exhibited a maximum discharge during saccades of larger amplitude when they were made toward a centrifugal target [see also 223]. However, in that study, the target moved so fast ($60^\circ/\text{s}$) that it was possible that gaze did not capture it, unless the saccade was anticipatory. Indeed, a target that moves away from the central visual field with a speed of $60^\circ/\text{s}$ after an initial jump of $2-6^\circ$ reaches an eccentricity of $14-18^\circ$ two hundred milliseconds later (approximate duration of response time). Since most measurements illustrated in this study's representative figure [224] correspond to saccades smaller than 14 degrees, these could have been anticipatory, or their amplitude was strongly hypometric. If they were hypometric, then we do not understand why the putative additional command failed to compensate for the saccade undershoot. If the saccades were neither hypometric nor anticipatory, then the recorded bursts ought to correspond to that elicited by a past position of the target. Thus, the size of the shift in maximum firing rate between saccades to static and moving targets should be larger as the activity corresponds to positions of the moving target further in the past. Current knowledge does not allow determining whether a horizontal diffusion of neuronal activity happens at the level of the SC or at the level of collicular afferent input from structures of the cerebral cortex, such as area LIP [225-226] or FEF [227-228].

4. Neurophysiological explanations

Conflating kinematic (or dynamic) numerical values with neurophysiological recordings carries the risk of believing that central neuron activity directly encodes gaze or head orientation rather than mediating changes in extraocular and neck muscle contraction. Some studies examined how the firing rate of some central neurons could be accounted for by a combination of kinematic parameters such as position, velocity and acceleration of the eyes [229-232], sometimes without separating the cells whose firing rate precedes the combination from those whose firing rate follows it [233-234]. Given i) the spatial and temporal transformations that a brief spot of retinal excitation undergoes on its way to the motor neurons, ii) the temporal overlap of activities corresponding to successive target positions, iii) the multiple possible patterns of muscle contraction specifying any static orientation of gaze, and iv) the multiplicity of possible neuronal drives, identifying a neurophysiological correspondence of notions such as dynamic position error, velocity error or even velocity becomes challenging, otherwise vain (see also [34]).

Physical parameters are certainly convenient for interpreting and comparing the activity of neurons, sorting them in different groups and promoting scientific debates. However, the reservations that David A. Robinson made about the block diagrams of oculomotor organization should not be restricted to the operations (differentiation, integration, gain ...) performed by the blocks [150]. They also concern the signals proposed to be "encoded" in the firing rate of neurons such as instantaneous motor error, velocity error or even velocity. The motor error command that feeds the premotor neurons during saccades toward a static target [202, 235-236] might be less problematic because it is

an oculocentric command [237] that can result from the preserved retinotopic arrangement of fibers connecting different neuronal assemblies.

To sum up, questioning the validity of a one-to-one correspondence relation between the sequence of values measured at the macroscopic level and the activity of microscopic neuronal elements does not lead the neurophysiology of gaze orientation to an epistemological dead end where the advancement of knowledge becomes vain or impossible. An alternative approach remains possible. Rather than reducing mismatches between extrinsic physical parameters (such as position error or velocity error), eye and head movements can be viewed as the behavioral expression of intrinsic processes that restore balances of activities opposing antagonistic visuomotor channels.

Interpreting the firing rate of neurons with notions belonging to the physical description of the movement of a body segment should be completed by the identification of muscle groups and neuronal networks involved. Comparative studies of the consequences of their functional perturbation, clinical and neuroanatomical studies should remain a privileged guide for interpreting neuronal activities and understanding their contribution to behavioral performances [238-239]. Building upon the knowledge gathered during the last decades [240-249], the following sections explain how the poly-equilibrium is restored during eye saccades and during combined eye-head movements, while demonstrating the possibility of accounting for the neuronal basis of gaze visual orientation, without resorting to kinematic (or dynamic) notions.


4.1. Extraocular muscles

Each eye movement involves synergies between anatomic-physiological elements that are invisible to investigations limited to behavioral analyses. However, the latter are helpful for discovering constraints in the contraction changes of the six muscles attached to the surface of each ocular globe (Figure 4). Combined with the relaxation of the medial rectus (MR) muscle fibers, the contraction of the lateral rectus (LR) muscle fibers of the same eye turns the eyeball toward the temporal side. Conversely, the contraction of MR combined with the relaxation of LR rotates the opposite eye toward the nasal side. In both cases, the eye rotates within the plane passing through the two muscles (toward the right in Fig. 4A). These muscles are antagonist to each other because their insertions are symmetrically distributed on opposite sides of the globe [250].

When the head is upright, the eye movement is horizontal provided that no change happens in the balances made by the contraction level of the other four extraocular muscles, between, on the one hand, the pair composed of superior rectus (SR) and inferior oblique (IO) muscles, and on the other hand, the pair composed of the inferior rectus (IR) and superior oblique (SO) muscles (Fig. 5A). Any change in the balance of forces exerted by these two pairs will add a vertical component to the horizontal eye movement. An upward deflection of horizontal saccades if the infraductive actions of IR and SO muscle fibers does not counterbalance changes in the contraction of SR and IO muscle fibers. Reciprocally, horizontal saccades will exhibit a downward deflection if the supraductive action of SR and IO does not counterbalance changes in the contraction of IR and SO muscle fibers.

An upward eye movement results from the contraction of SR and IO muscle fibers of both eyes combined with the relaxation of IR and SO of both eyes (Fig. 4B) whereas a downward eye movement results from the contraction of latter combined with the relaxation of the former (Fig. 4C). Their trajectory is straight vertical if the balance between the adduction and abduction does not change.

An upward eye movement results from the contraction of SR and IO muscle fibers of both eyes combined with the relaxation of IR and SO of both eyes (Fig. 4B) whereas a downward eye movement results from the contraction of latter combined with the relaxation of the former (Fig. 4C). We shall see below that their trajectory is straight vertical because of bilateral activity from the medio-posterior cerebellum.



Anatomo-physiological studies indicated that for each eye, distinct motor and premotor cells innervate these different muscles while behavioral studies report horizontal and vertical movements of both eyes with trajectories that are quasi-rectilinear (not curved) and torsionless. Visible to the external observer only, such trajectories imply the existence of synergies in the recruitment of extraocular muscles, synergies that the sole anatomical survey cannot deduce easily.

4.2. Binocular synergy

Orienting movements of gaze toward a visual target involve both eyeballs. The rotation of the left eye results from changes in the contraction of the fibers of its extra-ocular muscles, which are innervated by motor neurons that are distinct from the cells innervating the muscles of the right eye. A synergy is thus required between the groups of premotor neurons for moving both eyes quasi-simultaneously. We shall see below the neural substrate of Hering's law of equal innervation, whose failure would lead to diplopia.

4.2.1. Horizontal eye movements

Figure 5 illustrates the network responsible for leftward rotations of both eyeballs, i.e., for the contraction of the left eye's LR muscle and the right eye's MR, combined with the relaxation of antagonist muscles (left eye's MR muscle and right eye's LR). As explained above, the contraction of the other four extraocular muscles (SO and IR against IO and SR) is maintained if the movement is straight horizontal (Fig. 5A).

The degree of contraction of an extraocular muscle depends on the frequency of action potentials emitted by motor neurons innervating its fibers. For a horizontal eye movement, the motor cells innervating the LR muscle are grouped in the ipsilateral abducens nucleus whereas those innervating the MR muscle are in the ipsilateral oculomotor nucleus. The abducens nuclei are located in the pontine reticular formation and the oculomotor nuclei in the midbrain (Fig. 5B).

The contraction of agonist muscles is caused by an increase in the firing rate of motor neurons (MNs) in the abducens nucleus for the LR muscle and in the oculomotor nucleus for the MR muscle (Fig. 5C). During saccades, the muscle contraction is brisk not only because of the contractile properties of extraocular muscles [188, 190, 196], but also because the incoming input comes in a burst of action potentials with short interspike intervals [251-256]. In the abducens nucleus, internuclear neurons (AINs) relay to the motor neurons that innervate the MR muscle (synapse b) the excitation they receive at the same time as MN from excitatory burst neurons (EBNs, synapses a) located in the paramedian pontine reticular formation (ppRF) [146, 257].

In monkeys, a ppRF lesion eliminates the possibility of producing ipsilesional saccades (ipsilateral gaze palsy) whereas the generation of slow eye movements (during the vestibular ocular reflex or pursuit) remains possible [172, 259]. Electrical microstimulation applied to the ppRF evokes an ipsilateral movement of both eyes, which does not resemble a saccade. Its speed is most often constant and depends on the stimulation frequency [261-262]. The electrically-evoked movement is not saccade-like for two reasons at least. First, the stimulation parameters (constant current and frequency of pulses) do not replicate the complete population burst that usually precedes each saccade. Indeed, a limitation of the electrical stimulation technique is that its strength cannot be enhanced without exciting unrelated fibers of passage and eliciting reactions other than oculomotor. Secondly, the antagonist muscles were not relaxed as they usually are during saccades. Their relaxation results from a pause in the firing rate of motor neurons, which is caused by action potentials emitted by inhibitory burst neurons (IBNs) located in the contralateral dorsal paragigantocellularis reticular formation (dPGRF) [263-266].




Figure 5. Neuronal network involved in the generation of leftward saccades. *A:* The thickness of arrows attached to the eyeballs schematizes the strength of muscle contraction. LR: lateral rectus, MR: medial rectus, SR: superior rectus, IR: inferior rectus, SO: superior oblique, IO: inferior oblique. *B:* parasagittal section of the brainstem and cerebellum showing the approximate locations of the oculomotor nucleus (OMN), paramedian pontine reticular formation (ppRF), abducens nucleus (ABD), dorsal paragigantocellularis reticular formation (dPGRF), caudal Fastigial Nucleus (cFN) and superior colliculus (SC). *C:* Connecting lines ended by an arrow indicate excitatory connections; those ended by a circle indicate inhibitory synaptic connections. The thickness of connecting lines schematizes the strength with which the neurons fire. SRBNs: saccade-related burst neurons, OPNs: omnipause neurons, EBNs: excitatory burst neurons, IBNs: inhibitory burst neurons, ABDs: abducens nucleus, MNs: motoneurons, AINs: abducens internuclear neurons. Read text for explanations.

The consequences of dPGRF lesion have not yet been studied. However, because of the inhibition they exert on the motoneurons innervating the antagonist muscles and neurons (synapses d), the suppression of their burst is expected to hinder saccades and reduce their speed and amplitude. By disinhibiting IBNs located in the opposite side (synapses f), the unilateral suppression of IBNs' activity is also expected to increase the agonist drive during contralesional saccades and render them hypermetric. Indeed, recordings in monkeys reveal that approximately half of IBNs in the side opposite to the recruited EBN emit action potentials during contralateral saccades. These action potentials combine with those emitted by EBNs (synapses a and f) to compose the agonist oculomotor command [94, 257, 266-268].

Excitatory and inhibitory burst neurons receive descending inputs from saccade related neurons in the contralateral SC (synapses g and h) and cFN (synapses i and j). The collicular input is monosynaptic in cats [269] but disynaptic in monkeys [270]. This difference between feline and

primate species also concerns the fastigio-collicular projections. In monkeys, they are concentrated in the rostral part of both SCs [101] whereas they target more caudal regions in cats [271].

Finally, although premotor burst neurons receive lateralized input from the contralateral SC and cFN, the premotor control of saccades should be considered as bilateral [94, 257, 266-268] because of the omnidirectional bursts of saccade-related fastigial neurons. Indeed, for each horizontal saccade, a burst is emitted by neurons not only in the contralateral, but also in the ipsilateral cFN [272-275]. The contralateral burst was proposed to accelerate the saccade by exciting the agonist drive from EBNs (synapses i and j) whereas the ipsilateral burst was supposed to decelerate or brake the saccade by recruiting IBNs in the opposite side (synapse k).

Several facts led to revise this sequential activation and “biphasic” view of the fastigial control of saccade acceleration and deceleration [276] and to propose bilateral fastigial control of saccades [94, 266-268]. First, the biphasic hypothesis is not applicable to the generation of vertical saccades [94, 268]. Second, all IBNs do not fire during the late part of contralateral saccades; some of them burst at saccade onset, as EBNs do [264-265]. The same remark holds for the burst of saccade related fastigial neurons during ipsiversive saccades [275]. Moreover, the idea that the contralateral burst contributes to the acceleration of saccades is refuted by its later occurrence with larger amplitude of contralateral gaze shifts [277]. Yet, the peak velocity of contralateral gaze shifts is significantly reduced during unilateral cFN inactivation [278]. Finally, still after unilateral cFN inactivation, ipsilateral saccades exhibit increased peak velocity and enhanced displacement during the accelerating epoch, regardless of whether they are associated with a head movement [278]. Altogether, these observations are not compatible with the biphasic hypothesis, but remain compatible with the bilateral hypothesis.


4.2.2. Vertical eye movements

Contrary to the unilateral origin of the excitatory drive of binocular horizontal saccades, the excitatory phasic drive is bilateral for saccades directed upward or downward. An upward saccade involves the contraction of SR and IO muscle fibers of both eyes combined with the relaxation of IR and SO muscle fibers, while the contraction of adductive and abductive muscles (LR and MR) does not change (Fig. 6A).


The contraction of agonist muscles is driven by a burst of action potentials emitted by motor neurons located in the ipsilateral and contralateral oculomotor nuclei (for the IO and SR muscle fibers, respectively) [279-281]. The relaxation of antagonist muscles results from a pause in the firing rate of motor neurons located in the ipsilateral oculomotor nucleus (for the IR muscles) and the contralateral trochlear nucleus (for the SO muscles) (Fig. 6B). The agonist motor neurons receive a bilateral burst command from uEBNs located in both (left and right) rostral interstitial nuclei of the medial longitudinal fasciculus (riMLF; synapses a and b in Fig. 6C) [281-284]. Likewise, the inhibition of antagonist motor neurons is bilateral and driven by uEBNs (synapses c). It originates from uIBNs located in the contralateral interstitial nuclei of Cajal (iNC; synapses d and e).

The generation of downward saccade involves the contraction of IR and SO muscle fibers of both eyes combined with the binocular relaxation of SR and IO muscle fibers (Fig. 7A). The contraction of agonist muscles is driven by a burst of action potentials emitted by motor neurons located in the ipsilateral oculomotor nucleus (for the IR muscles) and the contralateral trochlear nucleus (for the SO muscles) while the relaxation of antagonist muscles results from a pause in the firing rate of motor

neurons located in the ipsilateral oculomotor nucleus (for the IO muscles) and the contralateral one (for the SR muscles).



Contrary to the bilateral origin of the excitatory premotor drive of upward saccades, it seems to be unilateral for downward saccades [285], originating from dEBNs located in the ipsilateral riMLF (synapses a and b). The origin of the inhibition of antagonist motor neurons has not yet been identified [286]. The asymmetry of the excitatory input is consistent with the effects of unilateral lesion of riMLF since downward saccades are more impaired than upward ones [287].



It is worth noting that i) the contraction of SR and IO muscle fibers does not only supraduct the eye during upward saccades and that ii) the contraction of IR and SO muscle fibers does not only infraduct the eye during downward saccades. The contraction of SR and SO muscle fibers also causes incycloduction (also called intorsion) of the globe while the contraction of IO and IR muscle fibers causes excycloduction (also called extorsion). Thus, during upward saccades, the incycloduction caused by the motor commands sent to the SR combines with the excycloduction caused by the commands to the IO, whereas during downward saccades, the excycloduction caused by activation of the IR muscle combines with the incycloduction caused by activation of the SO muscle. Behavioral measurements made in healthy subjects reveal negligible torsional component during vertical saccades [288], suggesting that the incycloduction and the excycloduction cancel each other out.

During upward saccades, such a balance must result from bilateral activity in the midbrain because for each eye, the motor neurons innervating the SR muscle are located in the contralateral

OMN whereas the motor neurons innervating the IO muscle are located in the ipsilateral OMN (Fig. 6C). Likewise, during downward saccades, the balance must also result from bilateral activity in the midbrain. The motor neurons innervating the IR muscle are located in the ipsilateral OMN whereas the motor neurons innervating the SO muscle are located in the contralateral trochlear nucleus (Fig. 7C).

Thus, the generation of vertical saccades that are straight and torsionless rests upon adjusted bilateral motor activation, which must take into account both the structural (neurons and their connectivity) and functional (firing properties, secondary and tertiary consequences of muscle contractions) asymmetries existing between oculomotor territories distributed on either side of the brain stem. In monkeys, stimulation of the right riMLF produces a conjugate clockwise rotation of both eyes, whereas left riMLF stimulation produces counter clockwise rotation [289].

Since the generation of vertical saccades is bilateral, a torsional component should appear in vertical saccades after unilateral lesion of riMLF with a direction that depends upon the side of the lesion. When dEBNs in the right riMLF are lesioned, activity in the opposite (unimpaired) riMLF should provoke a binocular counter clockwise torsion, i.e., an incycloduction of the right eye (caused by SO contraction) associated with an excyclotorsion of the left eye (caused by IR contraction). Clinical observations made in a patient suffering from a unilateral midbrain lesion affecting the right riMLF confirm this oculo-motor disorder. A counter clockwise torsional eye rotation accompanies each voluntary vertical saccades, downward and upward [290]. The fact that the patient's upward saccades exhibited a counter clockwise torsion means that the saccade of the right eye (caused by SR contraction) was associated with an incycloduction while the saccade of the left eye (caused by IO contraction) was associated with an excycloduction.

These clinical observations are important because they suggest that the ipsilateral projection of uEBNs toward the oculomotor nucleus is stronger than the contralateral projection. Otherwise, during upward saccades, the counter clockwise torsion would have been counterbalanced by the clockwise torsion (resulting from the IO contraction in the right eye and from the SR contraction in the left eye) promoted by the crossed projections from uEBNs to the right OMN. More fundamentally, such inferences illustrate the mutual benefit of combining neurophysiological and neuroanatomical studies in non-human primates with clinical studies in patients for advancing our understanding of oculomotor disorders as well as normal physiology.

Finally, bilateral control of vertical saccades likely involves the two superior colliculi, with both medial halves driving upward saccades (Fig. 6C) and both lateral halves driving downward saccades (Fig. 7C). As mentioned earlier with the bilateral conjecture, saccade-related neurons in the midline cerebellum may be involved in compensating for asymmetries and regulating the bilateral balance of activity between the left and right parts of the brainstem. Indeed, most neurons in the caudal fastigial nuclei [272-273] and in the vermis [291-292] emit a burst of action potentials during vertical saccades. The paucity of fastigial projections to the midbrain [293] and the numerous studies showing that asymmetrical functional perturbation of the medio-posterior cerebellum primarily impairs the horizontal component of saccades [94] suggest a weak involvement in the generation of vertical and torsional components of saccades. Vertical saccades exhibit a deflection of their trajectory toward the side of caudal fastigial inactivation [94, 268, 294-295] or toward the opposite side when the vermis is asymmetrically lesioned [296-297].

However, further neuroanatomical studies should investigate the cerebellar connectivity with the midbrain because another clinical study reported a transient counter clockwise torsion during horizontal saccades in a patient suffering from a lesion involving the left deep cerebellar nuclei and the left lateral medulla oblongata [298]. Furthermore, future clinical studies should test whether bilateral lesions account for the rare changes in Listing's law in patients suffering from cerebellar ataxia [299].

4.2.3. Synergy of horizontal and vertical saccades

The burst neurons involved in the generation of horizontal and vertical saccades are located in the pontomedullary reticular formation and in the midbrain, respectively. Since the pontomedullary reticular formation is located more caudally than the midbrain, the distance "traveled" by a collicular action potential to post-synaptic premotor burst neurons differs between horizontal or vertical saccades. Yet, when we examine the velocity profiles of the horizontal and vertical components of oblique saccades, we see that the two components start at the same time. This synchrony may result from releasing the premotor burst neurons from the inhibition exerted by omnipause neurons [300]. Simultaneously released from their tonic inhibition (synapse e in Fig. 5), premotor burst neurons then can emit their burst of action potentials synchronously. Accordingly, lesion of RIP should desynchronize the onset of horizontal and vertical components during oblique saccades. No experiment has yet investigated this prediction.

Another coupling exists between the horizontal and vertical components of oblique saccades. Its neurophysiological substrate has not been identified either. When the speed of one component diminishes, the other component is also slowed down. The duration of both component is lengthened during oblique saccades. When one compares cardinal and oblique saccades having equal horizontal displacement amplitudes, the horizontal component is slower during oblique saccades. Thus, the two components are not executed independently of each other. Their interaction, called component stretching, does not seem to originate only from the density of post-synaptic inputs from different collicular sectors to the populations of burst neurons in the ppRF and the midbrain [301]. A component stretching even occurs in oblique saccades evoked by an electrical microstimulation of the same collicular site, when the amplitude of one component is artificially varied by triggering the electrically-evoked saccade at different times after a visual saccade [302].

The most spectacular demonstration of component stretching was brought after transient inactivation of neurons in the ppRF [303-304]. Therein, local injection of lidocaine reduced both horizontal and vertical components of oblique saccades. In some experiments, the inactivation was subtle enough to slow down the saccades without altering their accuracy. Consistent with the hypothesis of negative feedback control of saccade amplitude, the reported results also indicate that the elements that have been inactivated do not participate in the feedback that updates the residual motor error. They are also compatible with the hypothesis of poly-equilibrium insofar as saccade duration is prolonged if the activity of neurons participating in its restoration is suppressed.

4.3. The synergy of eye and head movements

Orienting gaze toward the location of a stimulus often involves combined movements of the eyes and head. A change in the orientation of the trunk may also accompany the head movement [13]. When the body is at rest, a head movement carrying the eyeballs, gaze movement amplitude corresponds to the sum of the amplitudes of the eyes and head movements. Most of the time, gaze

captures the target (i.e., its image is focused on the fovea) before the head stops moving because the saccadic eye movement is much faster than the head movement. Then, while the head continues to move, the eyes rotate in the opposite direction. The interval during which the eyes rotate in the direction opposite to the head movement is called vestibulo-ocular reflex. A slow counter-rotation of the eyes in their orbit can also occur before the saccade, the head rotation preceding that of the eyes as during orienting toward an expected target location [305]. Thus, depending upon the amount of head rotation, the eye saccade will be of greater or lesser amplitude. The amplitude also depends on the orbital deviation of the eyeballs. If the eyes are deviated in the direction opposite to the direction of the impending saccade, the saccade amplitude is greater [12, 306-308]; the eyes having a wider range of mobility before reaching the maximum deviation imposed by the oculomotor range. Consequently, the contribution of the head to the gaze displacement amplitude is smaller than during gaze movements initiated when the deviation of the eyes is closer to the limits of the oculomotor range. In the latter case, the head will start moving sooner. Thus, the contribution of the head to the gaze displacement amplitude depends also upon its onset relative to saccade onset. Regardless of whether the head moves or not, small gaze movements are equally accurate [308-309]. However, for eccentric targets requiring large gaze shifts, the accuracy is improved when a head movement accompanies the eye saccade [308].

In macaques, changing the orientation of the head involves at least twenty muscles [310]. Three groups can be distinguished: i) the groups of muscles whose contraction changes the orientation of the head with respect to the trunk, ii) the “orthogonal” groups of muscles whose contraction stabilizes the atlas while it rotates under the influence of the first group, and iii) the group of muscles that do not change their contraction level [311-312]. For instance, the co-contraction of the two recti capitis posterior minor (RCPm) muscles stabilizes the atlas while the contraction of each rectus capitis posterior major (RCPM) and obliquus capitis inferior (OCI) muscles produces ipsilateral horizontal rotation. The contralateral RCPM and OCI muscles, however, show no change in electromyographic (EMG) activity. This lack of change in the activity of antagonist muscles indicates that stopping the head rotation does not involve their contraction. For movements of amplitude greater than 20 degrees, the contraction of the splenius capitis muscle completes the contraction of RCPM and OCI muscles. The co-contraction of the RCPm, RCPM, OCI and obliquus capitis superior (OCS) muscles stabilizes the head at the end of its change in orientation.

In summary, when a head movement is involved, gaze orienting response mobilizes several muscles, the synergy of which results from an activity involving a complex set of motor and premotor neurons, in the reticular formation, the vestibular nuclei and the spinal cord. In the continuation of studies performed in cats, neurophysiological investigations of eye-head movements in monkeys boomed in the 1990s and the beginning of the 21st century. They were dominated by the assumption that gaze direction, i.e., the direction of the line of sight in space, was an intrinsic cerebral command involving saccade-related neurons in SC [88-89, 131, 313-316] and FEF [317]. This command would subsequently be fractionated into separate eye and head premotor commands, downstream of the SC (for an alternative view see [318-319]).

Three major obstacles hampered the development of experimental investigations on eye-head movements. The first one was the technical difficulty of carrying out experiments with monkeys whose head is free to move. Added to this was the challenge posed by the need to merge the new neurophysiological data collected in non-human primates with the considerable body of knowledge

that had been obtained during prior experiments with cats. When differences between feline and primate species were noticed, the relevance of data collected in non-primate animals to understand the physiology of the human brain was questioned. Third, the attractiveness of this developing research field in macaque monkeys was diminished by socio-political factors that promoted fast publishing research works as well as the development of brain imaging and computer modeling studies. Faced with a migration of young investigators towards scientific fields in which the work was less risky, the neurophysiological study of eye-head coupling in monkeys declined. Societal changes in European countries also complicated and rendered socially thankless the use of non-human primates for the advancement of scientific knowledge, discouraging the few researchers who remained engaged.

Fortunately, past results mostly obtained in cats left a treasure trove of data that allows a synthesis, which, although incomplete, illustrates the tremendous complexity that underlies the orienting movements of the eyes and head. The following text completes a recent report [126] with more detailed explanations and relevant references. It will serve as a guide for further investigations with marmosets or for comparison with other species.

4.3.1. Reticulo-vestibulo-reticular synergies

In cats, the recruitment of neck muscles during horizontal head movements exhibits some peculiarities. Although some muscles are systematically recruited, either phasically (splenius muscles, OCI, levator scapulae and complexus) or tonically (biventer cervicis), the recruitment of other muscles (semispinalis cervicis, longissimus, levator scapulae, scalenus anterior and OCS) depends on the orientation of the head relative to the trunk [320]. At the neurophysiological level, the motor network responsible for horizontal movements can be divided into three groups: an excitatory medial reticulospinal system and two medial vestibulospinal subsystems, a contralateral excitatory subsystem and an ipsilateral inhibitory subsystem. Contrary to macaques, almost every vestibulospinal neuron also projects to the ocular motor nuclei in cats. In monkeys, the relative absence of cells carrying the same signals to motor neurons innervating the extraocular and neck muscles accounts for the dominance of the vestibulo-ocular reflex over the vestibulo-collic reflex [321].

4.3.1.1. *The reticulospinal channel*

Motor neurons that innervate the agonist muscles receive signals from neurons located ipsilaterally in the rostradorsal portion of the nucleus reticularis gigantocellularis (NRGC), the nucleus reticularis pontis caudalis (NRPC), and the dorsal portion of the nucleus reticularis gigantocellularis [322]. The combined lesion of the NRPC and NRGC abolishes the generation of ipsilateral head movements almost completely [323]. Ipsilesional saccadic eye movements are then also absent. However, contralateral saccades and vertical movements of the eyes and head do not seem to be affected. The slow compensatory movements of the vestibulo-ocular and vestibulo-collic reflexes are also spared.

The firing properties and connectivity of reticulospinal neurons are compatible with their involvement in the generation of combined ipsilateral eye and head movements [324]. Indeed, eye-neck reticulospinal neurons (EN-RSNs) emit bursts of action potentials during ipsilateral eye saccades associated with ipsilateral EMG activity of neck muscles. When the eyes are deviated toward the contralateral side and the ipsilateral neck muscles relaxed, the neurons are silent. Action potentials

are emitted only when the saccade leads the eyes beyond the orbital sagittal plane. None is emitted when the saccade is made in the contralateral orbital hemifield. This sensitivity to the orbital eye deviation, which appears neither in tecto-reticulo-spinal neurons [325-326] nor in cortico-reticulo-spinal neurons [327], may result from afferents from neurons in the nucleus prepositus hypoglossi (NPH). Therein, neurons display a sustained firing rate that increases with ipsilateral deviation of the eyes in the orbit. When the eyes are deviated toward the contralateral side, they are silent [328]. Thus, NPH neurons may account for silencing the EN-RSNs because their projection are primarily contralateral and inhibitory.

A projection of OPNs onto the EN-RSNs has been interpreted as evidence for an involvement of OPNs not only in the generation of eye saccades but also in eye-head gaze saccades [329]. This idea is compatible with a correlation between the duration of their pause and gaze saccade duration stronger than between the pause duration and the duration of eye saccades [330]. It is also compatible with the slowing of head movement when a short electrical microstimulation is applied in the RIP nucleus [330]. However, the alternative interpretation that OPNs remain primarily involved in the generation of ocular saccades cannot be rejected. The fact that the eye saccade duration is shorter than the gaze saccade duration does not necessarily imply that the burst duration of premotor neurons is also shorter. The shorter duration of eye saccades may result from the action of the vestibulo-ocular reflex downstream, at the level of motor neurons. Thus, the correlation between the pause duration of OPNs and the duration of eye saccades would be less strong than the correlation with the duration of gaze saccades. The slowing of head movement by electrical microstimulation of RIP may result from a transient reactivation of the vestibulo-colic and vestibulo-ocular reflexes because of the inhibition that OPNs exert upon excitatory and inhibitory burst neurons [91]. Finally, it is worth signaling that in monkeys, i) a slowing of head movement is rarely observed when RIP is stimulated [27] and ii) the end of the OPN's pause is better correlated with the eye saccade end than with the gaze shift end [331].

4.3.1.2. The vestibulospinal channels

The medial vestibulospinal tract (MVST) is the major pathway by which afferent signals from the horizontal semicircular canals influence the axial musculature of a cat's body. A tri-synaptic pathway connects the labyrinth to the motor neurons innervating the neck muscles [332-333]. The neurons at the origin of the vestibulo-nucal pathway are secondary vestibular neurons located in the medial vestibular nucleus (MVN). Activated by ipsilateral vestibular afferents, the firing rate of these neurons increases during an ipsilateral rotation of the head and decreases during a contralateral rotation, a response pattern called of type I [334-335]. The post-synaptic influence of the MVST onto the motor neurons of the contralateral cervical cord is excitatory. An inhibitory monosynaptic projection from the MVN to ipsilateral neck motoneurons has also been shown [333-334, 336]. The fact that a stimulation of the horizontal canal evokes a head movement directed towards the contralateral side [337] is compatible with this organization. Many neurons of the MVST, which are mainly located in the rostral region of the MVN, emit a collateral towards the contralateral abducens nucleus [338-340]. This double connection with the extraocular and head motor neurons demonstrates their involvement in combined movements of the eyes and head.

Electrophysiological studies report relatively high sustained activity of Eye-Neck Vestibular neurons when the eyes are centered in the orbit [341]. This tonic firing rate results either from an intrinsic property of neurons or from their sustained excitation by primary vestibular neurons [342]. In

some vestibular neurons, a sensitivity of their firing rate to the deviation of the head relative to the trunk suggests an influence also from neck proprioceptive signals [343].

During the passive rotation of the head, the vestibular signals inhibit the motor neurons innervating the ipsilateral extraocular and neck muscles while facilitating the activity of those innervating the contralateral muscles. Thus, rotation of the head promotes the emission of vestibular signals that lead to the counter-rotation of the eyes (vestibulo-ocular reflex) and of the head (vestibulo-colic reflex) in the direction opposite to the head rotation. These reflex movements also oppose active rotations of the head. We shall now examine how saccade-related commands remove this opposition.


4.3.2. Inhibition of vestibulo-ocular and vestibulo-colic reflexes

In addition to excitatory input from EBNs and EN-RSNs (synapses b in Figure 8), the motor neurons innervating the agonist eye and neck muscles (E-MNs and N-MNs, respectively) also receive inhibitory afferents from the IVNs-1 located in the ipsilateral MVN (synapses c) and excitatory afferents from the EVNs-1 in the contralateral MVN (synapses d). The inhibition of agonist motor neurons (by IVNs-1, synapse c) is prevented by action potentials emitted by IVNs-2 (synapse e), which are recruited by EN-RSNs and EBNs (synapses a) and by action potentials emitted by EVNs-1 located in the contralateral MVN (synapses f). These contralateral EVNs-1 are indeed released from the inhibition of IVNs-2 (synapse g) by the burst of action potentials emitted by IBNs (synapse h), which are recruited by agonist premotor neurons (synapses i). Thus, a disinhibition facilitates the recruitment of agonist muscles while the relaxation of antagonist muscles is promoted by the inhibition that IBNs exert on IVNs-2 in the contralateral MVN (synapse h) and on antagonist ocular premotor and motor neurons (synapses j).

During a combined horizontal movement of the eyes and head, the eyes do not counter-rotate in their orbit (vestibulo-ocular reflex) because the motor and internuclear neurons (E-MNs) in the contralateral abducens nucleus are inhibited by IBNs (synapse j). By exciting the IVNs-2 (synapses a), the bursts emitted by RSNs and EBNs prevent the ipsilateral IVNs-1 from thwarting the recruitment of agonist motor neurons (synapses c). The counter-rotation of the eyes restarts as the firing rate of ipsilateral EBNs and IBNs diminishes. The declining firing rate of EBNs removes the inhibition that ipsilateral IVNs-2 exert upon IVNs-1 and EVNs-1 (synapses e and k) while a declining firing rate of IBNs disinhibits IVNs-2 in the contralateral side (synapse h), which start inhibiting the EVNs-1 and IVNs-2 (synapses g and l). The ipsilateral MNs are inhibited by the IVNs-1 (synapse c) while the contralateral MNs can emit again action potentials in proportion to the excitation they receive from the contralateral EVNs-1 (synapses m) while the head movement terminates. Since this excitation recruits the antagonist neck muscles, it contributes to stopping the head movement [344-345]. However, it is worth reminding that in monkeys, this co-contraction has been questioned by EMG studies [310-311].

In summary, the firing rate reduction of IVNs-2 contributes to end the orienting gaze shift by two parallel processes. The first process is the consequence of the gradual decline of excitation from EBNs and EN-RSNs, while the second process results from an inhibition by IBNs in the contralateral side. In monkeys, the decline of EBNs and EN-RSNs firing rate might not result from a reduced drive from collicular neurons because the latter often continue to fire after saccade end [67, 315, 346-350]. Instead of a reduced drive, a desynchronization of presynaptic input to premotor neurons may happen. As explained elsewhere [52], temporally scattered pre-synaptic input are less efficient than

simultaneous ones for exciting post-synaptic neurons. The decline of EBNs and EN-RSNs activity may result from the inhibitory input from IBNs in the contralateral side. Thus, action potentials emitted by IBNs during contralateral saccades would participate in the inhibition of contralateral IVNs-2 and promote the counter-rotation of the eyes.



5. Conclusions

Orienting gaze movements enable an animal to localize and capture an object in its physical environment. They are convenient means for investigating how networks of neurons govern oculomotor behavior [351] and underlie behavioral performances, notably when neurological damage alters their normal functioning [238-239]. During the last decades, following the steps of predecessors who conducted their research in feline species, neurophysiologists and neuroanatomists gathered considerable knowledge in monkeys that contributed to identify the core networks involved in the generation of eye and head movements and to understand oculomotor disorders in human patients.

Technological advancements brought the possibility to measure precisely the time course of eye and head movements and to study correlations between the firing rate of neurons and various kinematic and dynamic parameters (amplitude, velocity and acceleration) or the angular distance between gaze and target directions. Analytical habits, which became conventions, resulted and gradually led some neuroscientists to assume a one-to-one correspondence relation between measured physical values of gaze or head orientation and neuronal activity. From there, the search for neurophysiological evidence supporting their theoretical diagram was promoted.

However, comparing kinematic or dynamic numerical values with neurophysiological recordings carries the risk of believing that central neuron activity directly encodes gaze or head orientation rather than mediating changes in extraocular and neck muscle contraction, in posture, in the autonomous nervous system and more centrally. Neuroanatomical and neurophysiological studies reveal functional properties and connectivity that are not visible in cybernetic diagrams. The comparison between a diagram and the brain is analog to that between an automaton and a living animal.

Complementing a previous synthesis [126], this review article is enriched by epistemological comments on the notion of “space” in the brain and by numerous additional precisions and references. It also provides the solid ground of evidence for a new departure in the neurophysiological study of orienting movements by considering static gaze orientation as poly-equilibrium, i.e., multiple balances of neuronal activities opposing mutually antagonistic channels. A saccade or a slow eye movement is not initiated as long as the visuo-oculomotor system is within in a mode where opposing commands counterbalance each other. For generating saccadic and pursuit eye movements, the symmetry breaking involves different groups of neurons in the left and right parts of the brainstem. Thus, rather than reducing position or error signals - which are numerical values belonging to the physical domain of kinematics -, orienting movements of the eyes and head are the behavioral expressions of intrinsic neuronal processes restoring the poly-equilibrium. Further investigations will connect them with processes involved in postural adjustments as well as with those that do not belong to classical visuomotor functions, but to autonomous functions [352-353] and alertness [354].

Acknowledgments:

This research was funded by Centre National de la Recherche Scientifique (CNRS). The author thanks Dr Blas Torres and Dr David L. Sparks for accepting the inclusion of one of their figure in this synthesis article. He also thanks the reviewers for their constructive comments.

Abbreviations:

ABD	abducens nucleus
AIN	abducens internuclear neuron
cFN	caudal fastigial nucleus
dPGRF	dorsal paragigantocellularis reticular formation
EBN	excitatory burst neuron
E-MN	eye-motoneuron
EVN-1	type 1 excitatory vestibular neuron
EVN-2	type 2 excitatory vestibular neuron
FEF	frontal eye field
IBN	inhibitory burst neuron
iNC	interstitial nucleus of Cajal
IO	inferior oblique
IR	inferior rectus
IVN-1	type 1 inhibitory vestibular neuron
IVN-2	type 2 inhibitory vestibular neuron
LR	lateral rectus
MLF	medial longitudinal fasciculus
MN	motoneuron
MR	medial rectus
MVN	medial vestibular nucleus
MVST	medial vestibulospinal tract
N-MN	neck motoneuron
NOT	nucleus of the optic tract
NPH	nucleus prepositus hypoglossi

NRGC	nucleus reticularis gigantocellularis
NRPC	nucleus reticularis pontis caudalis
OMN	oculomotor nucleus
ppRF	paramedian pontine reticular formation
RIP	raphe interpositus
riMLF	rostral intersitial nucleus of MLF
EN-RSN	eye-neck reticulospinal neuron
SC	superior colliculus
SO	superior oblique
TRO	trochlear nucleus

References

1. Pavlov, I.P. Conditioned reflexes. An investigation of the physiological activity of the cerebral cortex. Dover Publ., New York, 1960.
2. Sokolov, E.N. Higher nervous functions: The orienting reflex. *Ann. Rev. Physiol.* 1963, 25, 545-580.
3. Lynn, R. Attention, arousal and the orientation reaction. 1966, Pergamon Press.
4. Bertenthal, B.; von Hofsten, C. Eye, head and trunk control: The foundation for manual development. *Neurosci. Biobehav. Rev.* 1998, 22, 515-520.
5. Koceja, D.M.; Allway, D.; Earles, D.R. Age differences in postural sway during volitional head movement. *Arch. Phys. Med. Rehab.* 1999, 80, 1537-1541.
6. Stoffregen, T.A.; Bardy, B.G.; Bonnet, C.T.; Pagulayan, R.J. Postural stabilization of visually guided eye movements. *Ecol. Psychol.* 2006, 18, 191-222.
7. Torres, B.; Luque, M.A.; Pérez-Pérez, M.P.; Herrero, L. Visual orienting response in goldfish: a multidisciplinary study. *Brain Res. Bull.* 2005, 66, 376-380.
8. Kostyk, S. K.; Grobstein, P. Neuronal organization underlying visually elicited prey orienting in the frog—I. Effects of various unilateral lesions. *Neurosci.* 1987, 21, 41-55.
9. Werner, C.; Himstedt, W. Mechanism of head orientation during prey capture in Salamander. *Zool. Jb. Physiol.*, 1985, 89, 359-368.
10. Knudsen, E.I.; Blasdel, G.G.; Konishi, M. Sound localization by the barn owl (*Tyto alba*) measured with the search coil technique. *J. Comp. Physiol.* 1979, 133, 1-11.
11. Goffart, L.; Pélisson, D. Orienting gaze shifts during muscimol inactivation of caudal fastigial nucleus in the cat. I. Gaze dysmetria. *J. Neurophysiol.* 1998, 79, 1942-1958.
12. Freedman, E.G.; Sparks, D.L. Eye-head coordination during head-unrestrained gaze shifts in rhesus monkeys. *J. Neurophysiol.* 1997, 77, 2328-2348.
13. McCluskey, M.K.; Cullen, K.E. Eye, head, and body coordination during large gaze shifts in rhesus monkeys: movement kinematics and the influence of posture. *J. Neurophysiol.* 2007, 97, 2976-2991.
14. Collewijn, H.; Erkelens, C.J.; Steinman, R.M. Binocular co-ordination of human horizontal saccadic eye movements. *J. Physiol.* 1988, 404, 157-182.
15. Collewijn, H.; Erkelens, C.J.; Steinman, R.M. Binocular co-ordination of human vertical saccadic eye movements. *J. Physiol.* 1988, 404, 183-197.

16. Guitton, D.; Volle, M. Gaze control in humans: eye-head coordination during orienting movements to targets within and beyond the oculomotor range. *J. Neurophysiol.* 1987, 58, 427-459.
17. Lea, J.Y.; Mueller, C.G. Saccadic head movements in mantids. *J. Comp. Physiol.* 1977, 114, 115-128.
18. Rossel, S. Foveal fixation and tracking in the praying mantis. *J. Comp. Physiol.* 1980, 139, 307-331.
19. Olberg, R. M.; Seaman, R.C.; Coats, M. I.; Henry, A.F. Eye movements and target fixation during dragonfly prey-interception flights. *J. Comp. Physiol. A* 2007, 193, 685-693.
20. Consi, T. R.; Passani, M.B.; Macagno, E.R. Eye movements in *Daphnia magna*: Regions of the eye are specialized for different behaviors. *J. Comp. Physiol. A*, 1990, 166, 411-420.
21. Pélisson, D.; Guitton, D.; Munoz, D.P. Compensatory eye and head movements generated by the cat following stimulation-induced perturbations in gaze position. *Exp. Brain Res.* 1989, 78, 654-658.
22. Pélisson, D.; Goffart, L.; Guitton, D. On-line compensation of gaze shifts perturbed by micro-stimulation of the superior colliculus in the cat with unrestrained head. *Exp. Brain Res.* 1995, 106, 196-204.
23. Sparks, D.L.; Mays, L.E. Spatial localization of saccade targets. I. Compensation for stimulation-induced perturbations in eye position. *J. Neurophysiol.* 1983, 49, 45–63.
24. Sparks, D.L.; Mays, L.E.; Porter, J.D. Eye movements induced by pontine stimulation: interaction with visually triggered saccades. *J. Neurophysiol.* 1987, 58, 300–318.
25. Keller, E.L.; Gandhi, N.J.; Shieh, J.M. Endpoint accuracy in saccades interrupted by stimulation in the omnipause region in monkey. *Visual Neurosci.* 1996, 13, 1059-1067.
26. Gandhi, N. J.; Keller, E.L. Comparison of saccades perturbed by stimulation of the rostral superior colliculus, the caudal superior colliculus, and the omnipause neuron region. *J. Neurophysiol.* 1999, 82(6), 3236-3253.
27. Gandhi, N. J.; Sparks, D.L. Dissociation of eye and head components of gaze shifts by stimulation of the omnipause neuron region. *J. Neurophysiol.* 2007, 98, 360-373.
28. Schiller, P.H.; Sandell, J.H. Interactions between visually and electrically elicited saccades before and after superior colliculus and frontal eye field ablations in the rhesus monkey. *Exp. Brain Res.* 1983, 49, 381–392.
29. Kurylo, D.D. Interaction of visually guided saccades with saccades induced by electrical stimulation of the posterior parietal cortex in the monkey. *Vision Res.* 1991, 31, 2065-2073.
30. Keller, E. L., Slakey, D. P., & Crandall, W. F. (1983). Microstimulation of the primate cerebellar vermis during saccadic eye movements. *Brain research*, 288(1-2), 131-143.

31. Noda, H.; Murakami, S.; Warabi, T. Effects of fastigial stimulation upon visually-directed saccades in macaque monkeys. *Neurosci. Res.* 1991, 10: 188–199.
32. Guthrie, B.L.; Porter, J.D.; Sparks, D.L. Corollary discharge provides accurate eye position information to the oculomotor system. *Science* 1983, 221, 1193-1195.
33. Lewis, R.; Zee, D.; Hayman, M.; Tamargo, R. Oculomotor function in the rhesus monkey after deafferentation of the extraocular muscles. *Exp. Brain Res.* 2001, 141, 349-358.
34. Goffart, L. Cerebralization of mathematical quantities and physical features in neural science: a critical evaluation. *Eur. Phys. J. Web Conf.* 2024, 300, 01007.
35. Collewijn, H.; Erkelens, C.J.; Steinman, R.M. Trajectories of the human binocular fixation point during conjugate and non-conjugate gaze-shifts. *Vision Res.* 1997, 37, 1049-1069.
36. Pellionisz, A.; Llinás, R. Space-time representation in the brain. The cerebellum as a predictive space-time metric tensor. *Neurosci.* 1982, 7, 2949-2970.
37. McIlwain, J.T. Large receptive fields and spatial transformations in the visual system. *Int. Rev. Physiol.* 1976, 10, 223–248.
38. Ottes, F.P.; Van Gisbergen, J.A.; Eggermont, J.J. Metrics of saccade responses to visual double stimuli: two different modes. *Vision Res.* 1984, 24, 1169-1179.
39. Glimcher, P.W.; Sparks, D.L. Representation of averaging saccades in the superior colliculus of the monkey. *Exp. Brain Res.* 1993, 95, 429-435.
40. Leavitt, M.L.; Mendoza-Halliday, D.; Martinez-Trujillo, J.C. Sustained activity encoding working memories: not fully distributed. *TINS* 2017, 40, 328-346.
41. Jaffe, R.J.; Constantinidis, C. Working memory: from neural activity to the sentient mind. *Compr. Physiol.* 2021, 11, 2547.
42. Funahashi, S.; Bruce, C.J.; Goldman-Rakic, P.S. Mnemonic coding of visual space in the monkey's dorsolateral prefrontal cortex. *J. Neurophysiol.* 1989, 61, 331-349.
43. Gnadt, J.W.; Bracewell, R.M.; Andersen, R.A. Sensorimotor transformation during eye movements to remembered visual targets. *Vision Res.* 1991, 31, 693-715.
44. Funahashi, S.; Bruce, C.J.; Goldman-Rakic, P.S. Dorsolateral prefrontal lesions and oculomotor delayed-response performance: evidence for mnemonic "scotomas". *J. Neurosci.* 1993, 13, 1479-1497.
45. White J.M.; Sparks D.L.; Stanford T.R. Saccades to remembered target locations: an analysis of systematic and variable errors. *Vision Res.* 1994, 34, 79-92.
46. Barton E.J.; Sparks D.L. Saccades to remembered targets exhibit enhanced orbital position effects in monkeys. *Vision Res.* 2001, 41: 2393-2406.

47. Kirchner, J.; Watson, T.; Bauer, J.; Lappe, M. Eyeball translations affect saccadic eye movements beyond brainstem control. *J. Neurophysiol.* 2023, 130, 1334-1343.
48. O'Regan, J.K. Solving the "real" mysteries of visual perception: the world as an outside memory. *Can J. Psychol.* 1992, 46, 461-488.
49. Pulte, H. From axioms to conventions and hypotheses: The foundations of mechanics and the roots of Carl Neumann's "Principles of the Galilean-Newtonian theory". In: *The Significance of the hypothetical in the natural sciences.* Heidelberger, M. and Schiemann, G. (eds) Gruyter, Berlin, pp. 77-98, 2009.
50. De Paz, M. From jurisprudence to mechanics: Jacobi, Reech, and Poincaré on convention. *Sci. Context* 2018, 31, 223–250.
51. Burrelly, C.; Quinet, J.; Cavanagh, P.; Goffart, L. Learning the trajectory of a moving visual target and evolution of its tracking in the monkey. *J. Neurophysiol.* 2016, 116, 2739-2751.
52. Goffart, L.; Burrelly, C.; Quinet, J. Synchronizing the tracking eye movements with the motion of a visual target: basic neural processes. *Prog. Brain Res.* , 2017, 236, 243-268.
53. Botschko, Y.; Yarkoni, M.; Joshua, M. Smooth pursuit eye movement of monkeys naive to laboratory setups with pictures and artificial stimuli. *Front. Syst. Neurosci.* 2018, 12, 15.
54. Kornhuber, H.H. Motor functions of cerebellum and basal ganglia: the cerebellocortical saccadic (ballistic) clock, the cerebellonuclear hold regulator, and the basal ganglia ramp (voluntary speed smooth movement) generator. *Kybernetik* 1971, 8, 157-162.
55. Robinson, D.A. Models of the saccadic eye movement control system. *Kybernetik* 1973, 14, 71-83.
56. Ohtsuka, K.; Noda, H. Burst discharges of mossy fibers in the oculomotor vermis of macaque monkeys during saccadic eye movements. *Neurosci. Res.* 1992, 15, 102-114.
57. Zee, D.S.; Optican, L.M.; Cook, J.D.; Robinson, D.A.; Engel, W.K. Slow saccades in spinocerebellar degeneration. *Arch. Neurol.* 1976, 33, 243-251.
58. Robinson, D.A. Oculomotor control signals. In: *Basic Mechanisms of Ocular Motility and Their Clinical Implications*, edited by Lennerstrand G, Bachy-Rita P. Oxford, UK: Pergamon, 1975, p. 337–374.
59. Jürgens, R.; Becker, W.; Kornhuber, H.H. Natural and drug-induced variations of velocity and duration of human saccadic eye movements: evidence for a control of the neural pulse generator by local feedback. *Biol. Cybern.* 1981, 39, 87-96.
60. Sparks, D.L.; Holland, R.; Guthrie, B.L. Size and distribution of movement fields in the monkey superior colliculus. *Brain Res.* 1976, 113, 21-34.
61. Lee, C.; Rohrer, W.H.; Sparks, D.L. Population coding of saccadic eye movements by neurons in the superior colliculus. *Nature* 1988, 332, 357-360.

62. Sparks, D.L. Commands for coordinated eye and head movements in the primate superior colliculus. In: *The oculomotor system: new approaches for studying sensorimotor integration*. 2004, Hall, W.C. and Moschovakis, A.K. (Eds), CRC Press, pp. 303-318.
63. Sparks, D.L.; Lee, C.; Rohrer, W.H. Population coding of the direction, amplitude, and velocity of saccadic eye movements by neurons in the superior colliculus. *Cold Spring Harbor Symp. Quant. Biol.* 1990, 55, 805-811.
64. Kato, R.; Grantyn, A.; Dalezios, Y.; Moschovakis, A.K. The local loop of the saccadic system closes downstream of the superior colliculus. *Neuroscience* 2006, 143, 319-337.
65. Munoz, D.P., Guitton, D.; Pélisson, D. Control of orienting gaze shifts by the tectoreticulospinal system in the head-free cat. III. Spatiotemporal characteristics of phasic motor discharges. *J. Neurophysiol.* 1991, 66, 1642-1666.
66. Munoz, D.P.; Wurtz, R.H. Saccade-related activity in monkey superior colliculus. II. Spread of activity during saccades. *J. Neurophysiol.* 1995, 73, 2334-2348.
67. Anderson, S.R.; Porrill, J.; Sklavos, S.; Gandhi, N. J.; Sparks, D.L.; Dean, P. Dynamics of primate oculomotor plant revealed by effects of abducens microstimulation. *J. Neurophysiol.* 2009, 101, 2907-2923.
68. Kang, I.; Lee, C. Properties of saccade-related neurons in the cat superior colliculus: patterns of movement fields and discharge timing. *Exp. Brain Res.* 2000, 131, 149-164.
69. Moschovakis, A.K.; Gregoriou, G.G.; Savaki, H.E. Functional imaging of the primate superior colliculus during saccades to visual targets. *Nature Neurosci.* 2001, 4, 1026-1031.
70. Soetedjo, R.; Kaneko, C.R.; Fuchs, A.F. Evidence against a moving hill in the superior colliculus during saccadic eye movements in the monkey. *J. Neurophysiol.* 2002, 87, 2778-2789.
71. Port, N.L.; Sommer, M.A.; Wurtz, R.H. Multielectrode evidence for spreading activity across the superior colliculus movement map. *J. Neurophysiol.* 2000, 84, 344-357.
72. Parr, T.; Friston, K.J. Active inference and the anatomy of oculomotion. *Neuropsychologia.* 2018, 111, 334-343.
73. Sparks, D.L.; Mays, L.E. Movement fields of saccade-related burst neurons in the monkey superior colliculus. *Brain Res* 1980, 190, 39-50.
74. Hafed, Z.M.; Goffart, L.; Krauzlis, R.J. A neural mechanism for microsaccade generation in the primate superior colliculus. *Science* 2009, 323, 940-943.
75. Hafed, Z.M.; Krauzlis, R.J. Similarity of superior colliculus involvement in microsaccade and saccade generation. *J. Neurophysiol.* 2012, 107, 1904-1916.
76. Munoz, D.P.; Wurtz, R.H. Fixation cells in monkey superior colliculus. II. Reversible activation and deactivation. *J. Neurophysiol.* 1993b, 70, 576-589.

77. Goffart, L.; Hafed, Z.M.; Krauzlis, R.J. Visual fixation as equilibrium: evidence from superior colliculus inactivation. *J. Neurosci.* 2012, 32, 10627-10636.
78. Krauzlis, R.J., Goffart, L.; Hafed, Z.M. Neuronal control of fixation and fixational eye movements. *Philos. Trans. Roy. Soc. B: Biol. Sci.* 2017, 372, 20160205.
79. Kaneko, C.R. Effect of ibotenic acid lesions of the omnipause neurons on saccadic eye movements in rhesus macaques. *J. Neurophysiol.* 1996, 75, 2229-2242.
80. Soetedjo, R.; Kaneko, C.R.; Fuchs, A.F. Evidence that the superior colliculus participates in the feedback control of saccadic eye movements. *J. Neurophysiol.* 2002, 87, 679-695.
81. Corneil, B.; Munoz, D. Overt responses during covert orienting. *Neuron* 2014, 82, 1230-1243.
82. Edelman, J.A.; Keller, E.L. Activity of visuomotor burst neurons in the superior colliculus accompanying express saccades. *J. Neurophysiol.* 1996, 76, 908-926.
83. Dorris, M.C.; Pare, M.; Munoz, D.P. Neuronal activity in monkey superior colliculus related to the initiation of saccadic eye movements. *J. Neurosci.* 1997, 17, 8566-8579.
84. Sparks, D.L.; Rohrer, W.H.; Zhang, Y. The role of the superior colliculus in saccade initiation: a study of express saccades and the gap effect. *Vision Res.* 2000, 40, 2763-2777.
85. Marino, R.A.; Levy, R.; Boehnke, S.; White, B.J.; Itti, L.; Munoz, D.P. Linking visual response properties in the superior colliculus to saccade behavior. *Eur. J. Neurosci.* 2012, 35, 1738-1752.
86. Sparks, D.L. Functional properties of neurons in the monkey superior colliculus: coupling of neuronal activity and saccade onset. *Brain Res.* 1978, 156, 1-16.
87. Robinson, D.A. Implications of neural networks for how we think about brain function. *Behav. Brain Sci.* 1992, 15: 644-655.
88. Munoz, D.P.; Guitton, D. Fixation and orientation control by the tecto-reticulo-spinal system in the cat whose head is unrestrained. *Rev. Neurol.* 1989, 145, 567-579.
89. Guitton, D.; Munoz, D.P. Control of orienting gaze shifts by the tectoreticulospinal system in the head-free cat. I. Identification, localization, and effects of behavior on sensory responses. *J. Neurophysiol.* 1991, 66, 1605-1623.
90. Munoz, D.P.; Guitton, D. Control of orienting gaze shifts by the tectoreticulospinal system in the head-free cat. II. Sustained discharges during motor preparation and fixation. *J. Neurophysiol.* 1991, 66, 1624-1641.
91. Takahashi, M.; Sugiuchi, Y.; Na, J.; Shinoda, Y. Brainstem circuits triggering saccades and fixation. *J. Neurosci.* 2022, 42, 789-803.
92. Robinson, F.R.; Straube, A.; Fuchs, A.F. Role of the caudal fastigial nucleus in saccade generation. II. Effects of muscimol inactivation. *J. Neurophysiol.* 1993, 70, 1741-1758.

93. Ohtsuka, K.; Sato, H.; Noda, H. Saccadic burst neurons in the fastigial nucleus are not involved in compensating for orbital nonlinearities. *J. Neurophysiol.* 1994, 71, 1976-1980.
94. Goffart, L.; Chen, L.L.; Sparks, D. L. Deficits in saccades and fixation during muscimol inactivation of the caudal fastigial nucleus in the rhesus monkey. *J. Neurophysiol.* 2004, 92, 3351-3367.
95. Guerrasio, L.; Quinet, J.; Büttner, U.; Goffart, L. Fastigial oculomotor region and the control of foveation during fixation. *J. Neurophysiol.* 2010, 103, 1988-2001.
96. Sato, H.; Noda, H. Saccadic dysmetria induced by transient functional decortication of the cerebellar vermis. *Exp. Brain Res.* 1992, 88, 455-458.
97. Quinet, J.; Goffart, L. Saccade dysmetria in head-unrestrained gaze shifts after muscimol inactivation of the caudal fastigial nucleus in the monkey. *J. Neurophysiol.* 2005, 93, 2343-2349.
98. Kurkin, S.; Akao, T.; Fukushima, J.; Shichinohe, N.; Kaneko, C. R.; Belton, T.; Fukushima, K. No-go neurons in the cerebellar oculomotor vermis and caudal fastigial nuclei: planning tracking eye movements. *Exp. Brain Res.* 2014, 232, 191-210.
99. Noda, H.; Sugita, S.; Ikeda, Y. Afferent and efferent connections of the oculomotor region of the fastigial nucleus in the macaque monkey. *J. Comp. Neurol.* 1990, 302, 330-348.
100. McFarland, J.L.; Fuchs, A.F. Discharge patterns in nucleus prepositus hypoglossi and adjacent medial vestibular nucleus during horizontal eye movement in behaving macaques. *J. Neurophysiol.* 1992, 68, 319-332.
101. May, P.J.; Hartwich-Young, R.; Nelson, J.D.; Sparks, D.L.; Porter, J.D. Cerebellotectal pathways in the macaque: implications for collicular generation of saccades. *Neurosci.* 1990, 36, 305-324.
102. Hafed, Z.M.; Goffart, L.; Krauzlis, R.J. Superior colliculus inactivation causes stable offsets in eye position during tracking. *J. Neurosci.* 2008, 28, 8124-8137.
103. Burrely, C.; Quinet, J.; Goffart, L. Pursuit disorder and saccade dysmetria after caudal fastigial inactivation in the monkey. *J. Neurophysiol.* 2018, 120, 1640-1654.
104. Fukushima, K.; Peterson, B.W.; Uchino, Y.; Coulter, J.D.; Wilson, V.J. Direct fastigiospinal fibers in the cat. *Brain Res.* 1977, 126, 538-542.
105. Wilson, V.J.; Uchino, Y.; Susswein, A.; Fukushima, K. Properties of direct fastigiospinal fibers in the cat. *Brain Res.* 1977, 126, 543-546.
106. Takahashi, M.; Sugiuchi, Y.; Shinoda, Y. Convergent synaptic inputs from the caudal fastigial nucleus and the superior colliculus onto pontine and pontomedullary reticulospinal neurons. *J. Neurophysiol.* 2014, 111, 849-867.
107. Latto, R.; Cowey, A. Fixation changes after frontal eye-field lesions in monkeys. *Brain Res.* 1971, 30, 25-36.

108. Dias, E.C.; Segraves, M.A. Muscimol-induced inactivation of monkey frontal eye field: effects on visually and memory-guided saccades. *J. Neurophysiol.* 1999, 81, 2191-2214.
109. Paré, M.; Wurtz, R.H. Progression in neuronal processing for saccadic eye movements from parietal cortex area lip to superior colliculus. *J. Neurophysiol.* 2001, 85, 2545-2562.
110. Wurtz, R.H., Sommer, M.A., Paré, M.; Ferraina, S. Signal transformations from cerebral cortex to superior colliculus for the generation of saccades. *Vision Res.* 2001, 41, 3399-3412.
111. Lynch, J.C.; McLaren, J.W. Deficits of visual attention and saccadic eye movements after lesions of parietooccipital cortex in monkeys. *J. Neurophysiol.* 1989, 61, 74-90.
112. Li, C.S.R., Mazzoni, P.; Andersen, R.A. Effect of reversible inactivation of macaque lateral intraparietal area on visual and memory saccades. *J. Neurophysiol.* 1999, 81, 1827-1838.
113. Wardak, C.; Olivier, E.; Duhamel, J.R. Saccadic target selection deficits after lateral intraparietal area inactivation in monkeys. *J. Neurosci.* 2002, 22, 9877-9884.
114. Ventre, J.; Faugier-Grimaud, S. Effects of posterior parietal lesions (area 7) on VOR in monkeys. *Exp. Brain Res.* 1986, 62, 654-658.
115. Simon, J.E.; Morgan, S.C.; Pexman, J.H.W.; Hill, M.D.; Buchan, A.M. CT assessment of conjugate eye deviation in acute stroke. *Neurol.* 2003, 60, 135-137.
116. Becker, E.; Karnath, H.O. Neuroimaging of eye position reveals spatial neglect. *Brain* 2010, 133, 909-914.
117. Fruhmann Berger, M.; Johannsen, L.; Karnath, H.O. Time course of eye and head deviation in spatial neglect. *Neuropsychol.* 2008, 22, 697.
118. Reinhard, J.I.; Damm, I.; Ivanov, I.V.; Trauzettel-Klosinski, S. Eye movements during saccadic and fixation tasks in patients with homonymous hemianopia. *J. Neuro-ophthalm.* 2014, 34(4), 354-361.
119. Snodderly, D.M. Effects of light and dark environments on macaque and human fixational eye movements. *Vision Res.* 1987, 27, 401-415.
120. Barash, S.; Melikyan, A.; Sivakov, A.; Tauber, M. Shift of visual fixation dependent on background illumination. *J. Neurophysiol.* 1998, 79, 2766-2781.
121. Goffart, L.; Quinet, J.; Chavane, F.; Masson, G. Influence of background illumination on fixation and visually guided saccades in the rhesus monkey. *Vision Res.* 2006, 46, 149-162.
122. Van Essen, D.C.; Newsome, W.T.; Maunsell, J.H. The visual field representation in striate cortex of the macaque monkey: asymmetries, anisotropies, and individual variability. *Vision Res.* 1984, 24, 429-448.
123. Hafed, Z.M.; Chen, C.Y. Sharper, stronger, faster upper visual field representation in primate superior colliculus. *Curr. Biol.* 2016, 26, 1647-1658.

124. Goffart, L.; Bourrelly, C.; Quinton, J.C. Neurophysiology of visually guided eye movements: critical review and alternative viewpoint. *J. Neurophysiol.* 2018, 120, 3234-3245.
125. Goffart, L. Kinematics and the neurophysiological study of visually-guided eye movements. *Prog. Brain Res.* 2019, 249, 375-384.
126. Goffart, L.; Quinet, J.; Bourrelly, C. Neurophysiology of gaze orientation: Core neuronal networks. In: Grafman, J.H. (Ed.), *Encyclopedia of the Human Brain, Second Edition 2005*, 1, 681–699. USA: Elsevier. 10.1016/B978-0-12-820480-1.00062-0.
127. Isa T., Itouji T., Sasaki S. Control of head movements in the cat: two separate pathways from the superior colliculus to neck motoneurons and their roles in orienting movements. In: *Vestibular and Brain Stem Control of Eye, Head and Body Movements*, edited by Shimazu H. and Shinoda Y. Tokyo: Japan Scientific Societies Press, 1992, p. 275–284. doi:10.1159/000421388.
128. Schiller, P.H.; Stryker, M. Single-unit recording and stimulation in superior colliculus of the alert rhesus monkey. *J. Neurophysiol.* 1972, 35, 915-924.
129. Robinson, D.A. Eye movements evoked by collicular stimulation in the alert monkey. *Vision Res.* 1972, 12, 1795-1808.
130. Stanford, T.R.; Freedman, E.G.; Sparks, D.L. Site and parameters of microstimulation: evidence for independent effects on the properties of saccades evoked from the primate superior colliculus. *J. Neurophysiol.* 1996, 76, 3360-3381.
131. Freedman, E.G.; Stanford, T.R.; Sparks, D.L. Combined eye-head gaze shifts produced by electrical stimulation of the superior colliculus in rhesus monkeys. *J. Neurophysiol.* 1996, 76, 927-952.
132. Schiff, D.; Cohen, B.; Raphan, T. Nystagmus induced by stimulation of the nucleus of the optic tract in the monkey. *Exp. Brain Res.* 1988, 70, 1-14.
133. Mustari, M.J.; Fuchs, A.F. Discharge patterns of neurons in the pretectal nucleus of the optic tract (NOT) in the behaving primate. *J. Neurophysiol.* 1990, 64, 77-90.
134. Ilg, U.J.; Hoffmann, K.P. Responses of monkey nucleus of the optic tract neurons during pursuit and fixation. *Neurosci. Res.* 1991, 12, 101-110.
135. Ilg, U.J.; Hoffmann, K.P. Responses of neurons of the nucleus of the optic tract and the dorsal terminal nucleus of the accessory optic tract in the awake monkey. *Eur. J. Neurosci.* 1996, 8, 92-105.
136. Wurtz, R.H.; Goldberg, M.E. Activity of superior colliculus in behaving monkey. 3. Cells discharging before eye movements. *J. Neurophysiol.* 1972, 35, 575-586.
137. Chubb, M.C.; Fuchs, A.F. Contribution of γ group of vestibular nuclei and dentate nucleus of cerebellum to generation of vertical smooth eye movements. *J. Neurophysiol.* 1982, 48, 75-99.
138. Tomlinson, R.D.; Robinson, D.A. Signals in vestibular nucleus mediating vertical eye movements in the monkey. *J. Neurophysiol.* 1984, 51, 1121-1136.

139. Chubb, M.C.; Fuchs, A.F.; Scudder, C.A. Neuron activity in monkey vestibular nuclei during vertical vestibular stimulation and eye movements. *J. Neurophysiol.* 1984, 52, 724-742.
140. Scudder, C.A.; Fuchs, A.F. Physiological and behavioral identification of vestibular nucleus neurons mediating the horizontal vestibuloocular reflex in trained rhesus monkeys. *J. Neurophysiol.* 1992, 68, 244-264.
141. McFarland, J.L.; Fuchs, A.F. Discharge patterns in nucleus prepositus hypoglossi and adjacent medial vestibular nucleus during horizontal eye movement in behaving macaques. *J. Neurophysiol.* 1992, 68, 319-332.
142. Böhmer, A.; Straumann, D. Pathomechanism of mammalian downbeat nystagmus due to cerebellar lesion: a simple hypothesis. *Neurosci. Lett.* 1998, 250, 127-130.
143. Marti, S.; Straumann, D.; Büttner, U.; Glasauer, S. A model-based theory on the origin of downbeat nystagmus. *Exp. Brain Res.* 2008, 188, 613-631.
144. Yakushin, S.B.; Gizzi, M.; Reisine, H.; Raphan, T.; Büttner-Ennever, J. Cohen, B. Functions of the nucleus of the optic tract (NOT). II. Control of ocular pursuit. *Exp. Brain Res.* 2000, 131, 433-447.
145. Inoue, Y.; Takemura, A.; Kawano, K.; Mustari, M.J. Role of the pretectal nucleus of the optic tract in short-latency ocular following responses in monkeys. *Exp. Brain Res.* 2000, 131, 269-281.
146. Strassman, A.; Highstein, S.M.; McCrea, R.A. Anatomy and physiology of saccadic burst neurons in the alert squirrel monkey. I. Excitatory burst neurons. *J. Comp. Neurol.* 1986, 249, 337-357.
147. Mustari, M.J.; Fuchs, A.F.; Kaneko, C.R.; Robinson, F.R. Anatomical connections of the primate pretectal nucleus of the optic tract. *J. Comp. Neurol.* 1994, 349, 111-128.
148. Büttner-Ennever J.A.; Cohen, B.; Horn, A.K.E.; Reisine, H. Efferent pathways of the nucleus of the optic tract in monkey and their role in eye movements. *J. Comp. Neurol.* 1996, 373, 90-107.
149. Appell, P.P.; Behan, M. Sources of subcortical GABAergic projections to the superior colliculus in the cat. *J. Comp. Neurol.* 1990, 302, 143-158.
150. Robinson D.A. Models of oculomotor neural organization. In: Bach-Y-Rita P. and Collins C.C. (Eds), *The control of eye movements* 1971, 519-538, Academic Press, New York.
151. Hu, X.; Jiang, H.; Gu, C.; Li, C.; Sparks, D.L. Reliability of oculomotor command signals carried by individual neurons. *Proc. Natl. Acad. Sci. U. S. A.* 2007, 104, 8137-8142.
152. Xu-Wilson, M.; Chen-Harris, H.; Zee, D. S. and Shadmehr R. Cerebellar contributions to adaptive control of saccades in humans. *J. Neurosci.* 2009, 29, 12930-12939.
153. Eggert, T.; Robinson, F.R.; Straube, A. Modeling inter-trial variability of saccade trajectories: effects of lesions of the oculomotor part of the fastigial nucleus. *PLoS Comput. Biol.* 2016, 12, e1004866.

154. Barmack, N.H. Modification of eye movements by instantaneous changes in the velocity of visual targets. *Vision Res.* 1970, 10, 1431-1441.
155. Cassanello, C.R.; Nihalani, A.T.; Ferrera, V.P. Neuronal responses to moving targets in monkey frontal eye fields. *J. Neurophysiol.* 2008, 100, 1544-1556.
156. Gellman, R.S.; Carl, J.R. Motion processing for saccadic eye movements in humans. *Exp. Brain Res.* 1991, 84, 660-667.
157. Keller, E.L.; Johnsen, S.S. Velocity prediction in corrective saccades during smooth-pursuit eye movements in monkey. *Exp. Brain Res.* 1990, 80, 525-531.
158. Fleuriot, J.; Hugues, S.; Perrinet, L.; Goffart, L. Saccadic foveation of a moving visual target in the rhesus monkey. *J. Neurophysiol.* 2011, 105, 883-895.
159. Bourrelly, C.; Quinet, J.; Cavanagh, P.; Goffart, L. Learning the trajectory of a moving visual target and evolution of its tracking in the monkey. *J. Neurophysiol.* 2016, 116, 2739-2751.
160. Orlando-Dessaints, N.; Goffart, L. Tracking a moving visual target in the rhesus monkey: Influence of the occurrence frequency of the target path. *J. Neurophysiol.* 2023, 130, 1425-1443.
161. Fleuriot, J.; Goffart, L. Saccadic interception of a moving visual target after a spatiotemporal perturbation. *J. Neurosci.* 2012, 32, 452-461.
162. Quinet, J.; Goffart, L. Does the brain extrapolate the position of a transient moving target? *J. Neurosci.* 2015a, 35, 11780-11790.
163. Barnes, C.S. Human eye saccades to targets of low energy. Doctoral thesis, University of Toronto, 1995.
164. Bell, A.H.; Meredith, M.A.; Van Opstal, A.J.; Munoz, D. Stimulus intensity modifies saccadic reaction time and visual response latency in the superior colliculus. *Exp. Brain Res.* 2006, 174, 53-59.
165. Newsome, W.T.; Wurtz, R.H.; Dursteler, M.R.; Mikami, A. Deficits in visual motion processing following ibotenic acid lesions of the middle temporal visual area of the macaque monkey. *J. Neurosci.* 1985, 5, 825-840.
166. May, J.G.; Keller, E.L.; Suzuki, D.A. Smooth-pursuit eye movement deficits with chemical lesions in the dorsolateral pontine nucleus of the monkey. *J. Neurophysiol.* 1988, 59, 952-977.
167. Schiller, P.H.; Lee, K. The effects of lateral geniculate nucleus, area V4, and middle temporal (MT) lesions on visually guided eye movements. *Visual Neurosci.* 1994, 11, 229-241.
168. Becker, W.; Jürgens, R. An analysis of the saccadic system by means of double step stimuli. *Vision Res.* 1979, 19, 967-983.
169. Botschko, Y.; Yarkoni, M.; Joshua, M. Smooth pursuit eye movement of monkeys naive to laboratory setups with pictures and artificial stimuli. *Front. Syst. Neurosci.* 2018, 12, 15.

170. Salman, M.S.; Sharpe, J.A.; Lillakas, L.; Dennis, M.; Steinbach, M.J. Smooth pursuit eye movements in children. *Exp. Brain Res.* 2006, 169, 139-143.
171. Ingster-Moati, I.; Vaivre-Douret, L.; Quoc, E.B.; Albuissou, E.; Dufier, J.L.; Golse, B. Vertical and horizontal smooth pursuit eye movements in children: a neuro-developmental study. *Eur. J. Paediatr. Neurol.* 2009, 13, 362-366.
172. Hanson, M.R.; Hamid, M.A.; Tomsak, R.L.; Chou, S.S.; Leigh, R.J. Selective saccadic palsy caused by pontine lesions: clinical, physiological, and pathological correlations. *Ann. Neurol.* 1986, 20, 209-217.
173. Pierrot-Deseilligny, C.; Amarenco, P.; Rouillet, E.; Marteau, R. Vermal infarct with pursuit eye movement disorders. *J. Neurol. Neurosurg. Psychiatr.* 1990, 53, 519-521.
174. Ohtsuka, K.; Igarashi, Y.; Chiba, S. Cerebellar peduncle lesion without saccadic abnormalities. *Ophthalmol.* 1992, 204, 44-48.
175. Büttner U, Straube A, Spuler A. Saccadic dysmetria and "intact" smooth pursuit eye movements after bilateral deep cerebellar nuclei lesions. *J. Neurol. Neurosurg. Psychiatr.* 1994, 57, 832–834.
176. Segraves, M.A.; Goldberg, M.E.; Deng, S.Y.; Bruce, C.J.; Ungerleider, L.G.; Mishkin, M. The role of striate cortex in the guidance of eye movements in the monkey. *J. Neurosci.* 1987, 7, 3040-3058.
177. MacAvoy, M.G.; Gottlieb, J.P.; Bruce, C.J. Smooth-pursuit eye movement representation in the primate frontal eye field. *Cereb. Cortex* 1991, 1, 95-102.
178. Shi, D.; Friedman, H.R.; Bruce, C.J. Deficits in smooth-pursuit eye movements after muscimol inactivation within the primate's frontal eye field. *J. Neurophysiol.* 1998, 80, 458-464.
179. Ono, S.; Mustari, M.J. Horizontal smooth pursuit adaptation in macaques after muscimol inactivation of the dorsolateral pontine nucleus (DLPN). *J. Neurophysiol.* 2007, 98, 2918-2932.
180. Daniel, B.M.; Lee, D.N. Development of looking with head and eyes. *J. Exp. Child Psychol.* 1990, 50, 200-216.
181. Salman, M.S.; Sharpe, J.A.; Eizenman, M.; Lillakas, L.; Westall, C.; To, T.; Deniis M.; Steinbach, M.J. Saccades in children. *Vision Res.* 2006, 46, 1432-1439.
182. Ego, C.; Orban de Xivry, J.J.; Nassogne, M.C.; Yüksel, D.; Lefèvre, P. The saccadic system does not compensate for the immaturity of the smooth pursuit system during visual tracking in children. *J. Neurophysiol.* 2013, 110, 358-367.
183. Sinno, S.; Najem, F.; Abouchacra, K.S.; Perrin, P.; Dumas, G. Normative values of saccades and smooth pursuit in children aged 5 to 17 years. *J. Am. Acad. Audiol.* 2020, 31, 384-392.
184. Ogawa, T.; Fujita, M. Adaptive modifications of human postsaccadic pursuit eye movements induced by a step-ramp-ramp paradigm. *Exp. Brain Res.* 1997, 116, 83-96.

185. Chou, I.H.; Lisberger, S.G. Spatial generalization of learning in smooth pursuit eye movements: implications for the coordinate frame and sites of learning. *J. Neurosci.* 2002, 22, 4728-4739.
186. Sun, Z.; Smilgin, A.; Junker, M.; Dicke, P.W.; Thier, P. Short-term adaptation of saccades does not affect smooth pursuit eye movement initiation. *J. Vision* 2017, 17, 19-19.
187. 187. Bucci, M.P.; Ajrezo, L.; Wiener-Vacher, S. Oculomotor tasks affect differently postural control in healthy children. *Int. J. Dev. Neurosci.* 2015, 46, 1-6.
188. Petit, L.; Haxby, J.V. Functional anatomy of pursuit eye movements in humans as revealed by fMRI. *J. Neurophysiol.* 1999, 82, 463-471.
189. Rosano, C.; Krisky, C.M.; Welling, J.S.; Eddy, W.F.; Luna, B.; Thulborn, K.R.; Sweeney, J.A. Pursuit and saccadic eye movement subregions in human frontal eye field: a high-resolution fMRI investigation. *Cerebral Cortex* 2002, 12, 107-115.
190. Yamada T., Suzuki D.A.; Yee R.D. Smooth pursuit-like eye movements evoked by microstimulation in macaque nucleus reticularis tegmenti pontis. *J. Neurophysiol.* 1996, 76, 3313–3324.
191. Suzuki D.A.; Yamada T.; Yee R.D. Smooth-pursuit eye-movement-related neuronal activity in macaque nucleus reticularis tegmenti pontis. *J. Neurophysiol.* 2003, 89, 2146–2158.
192. Ono, S.; Das, V.E.; Mustari, M.J. Gaze-related response properties of DLPN and NRTP neurons in the rhesus macaque. *J. Neurophysiol.* 2004, 91, 2484-2500.
193. Bruce, C.J.; Goldberg, M.E.; Bushnell, M.C.; Stanton, G.B. Primate frontal eye fields. II. Physiological and anatomical correlates of electrically evoked eye movements. *J. Neurophysiol.* 1985, 54, 714-734.
194. Gottlieb, J.P.; Bruce, C.J.; MacAvoy, M.G. Smooth eye movements elicited by microstimulation in the primate frontal eye field. *J. Neurophysiol.* 1993, 69, 786-799.
195. Gottlieb, J.P.; MacAvoy, M.G.; Bruce, C.J. Neural responses related to smooth-pursuit eye movements and their correspondence with electrically elicited smooth eye movements in the primate frontal eye field. *J. Neurophysiol.* 1994, 72, 1634-1653.
196. Stanton, G.B.; Friedman, H.R.; Dias, E.C.; Bruce, C.J. Cortical afferents to the smooth-pursuit region of the macaque monkey's frontal eye field. *Exp. Brain Res.* 2005, 165, 179-192.
197. Tian, J.R.; Lynch, J.C. Corticocortical input to the smooth and saccadic eye movement subregions of the frontal eye field in Cebus monkeys. *J. Neurophysiol.* 1996, 76, 2754-2771.
198. Mahaffy, S.; Krauzlis, R.J. Neural activity in the frontal pursuit area does not underlie pursuit target selection. *Vision Res.* 2011, 51, 853-866.
199. Lynch, J.C.; Tian, J.R. Cortico-cortical networks and cortico-subcortical loops for the higher control of eye movements. *Prog. Brain Res.* 2006, 151, 461-501.

200. Orban de Xivry, J.J.; Lefevre, P. Saccades and pursuit: two outcomes of a single sensorimotor process. *J. Physiol.* 2007, 584, 11-23.
201. Goettker, A.; Gegenfurtner, K.R. A change in perspective: The interaction of saccadic and pursuit eye movements in oculomotor control and perception. *Vision Res.* 2021, 188, 283-296.
202. Krauzlis, R.J.; Basso, M.A.; Wurtz R.H. Discharge properties of neurons in the rostral superior colliculus of the monkey during smooth-pursuit eye movements. *J. Neurophysiol.* 2000, 84, 876–91.
203. Krauzlis, R.J. Neuronal activity in the rostral superior colliculus related to the initiation of pursuit and saccadic eye movements. *J. Neurosci.* 2003, 23, 4333–44.
204. Hafed, Z.M.; Krauzlis, R.J. Goal representations dominate superior colliculus activity during extrafoveal tracking. *J. Neurosci.* 2008, 28, 9426-9439.
205. Krauzlis, R.J., Goffart, L.; Hafed, Z.M. Neuronal control of fixation and fixational eye movements. *Philos. Trans. Roy. Soc. B: Biol. Sci.* 2017, 372, 20160205.
206. Missal, M.; Keller, E.L. Common inhibitory mechanism for saccades and smooth-pursuit eye movements. *J. Neurophysiol.* 2002, 88, 1880-1892.
207. Keller, E.L.; Missal, M. Shared brainstem pathways for saccades and smooth-pursuit eye movements. *Ann. N. Y. Acad. Sci.* 2003, 1004, 29-39.
208. Strassman, A.; Evinger, C.; McCrea, R.A.; Baker, R.G.; Highstein, S.M. Anatomy and physiology of intracellularly labelled omnipause neurons in the cat and squirrel monkey. *Exp. Brain Res.* 1987, 67, 436-440.
209. Takahashi, M.; Sugiuchi, Y.; Na, J.; Shinoda, Y. Brainstem circuits triggering saccades and fixation. *J. Neurosci.* 2022, 42, 789-803.
210. Krauzlis, R.J.; Miles, F.A. Role of the oculomotor vermis in generating pursuit and saccades: effects of microstimulation. *J. Neurophysiol.* 1998, 80, 2046-2062.
211. Suzuki, D.A.; Keller, E.L. The role of the posterior vermis of monkey cerebellum in smooth-pursuit eye movement control. II. Target velocity-related Purkinje cell activity. *J. Neurophysiol.* 1988, 59, 19-40.
212. Sato, H.; Noda, H. Posterior vermal Purkinje cells in macaques responding during saccades, smooth pursuit, chair rotation and/or optokinetic stimulation. *Neurosci. Res.* 1992, 12, 583-595.
213. Raghavan, R.T.; Lisberger, S.G. Responses of Purkinje cells in the oculomotor vermis of monkeys during smooth pursuit eye movements and saccades: comparison with floccular complex. *J. Neurophysiol.* 2017, 118, 986-1001.
214. Fuchs, A. F., Robinson, F. R.; Straube, A. Participation of the caudal fastigial nucleus in smooth-pursuit eye movements. I. Neuronal activity. *J. Neurophysiol.* 1994, 72, 2714-2728.

215. Lisberger S.G. Postsaccadic enhancement of initiation of smooth pursuit eye movements in monkeys. *J. Neurophysiol.* 1998, 79: 1918–1930.
216. Robinson, F.R.; Straube, A.; Fuchs, A.F. Role of the caudal fastigial nucleus in pursuit eye movements. II. Effects of muscimol inactivation. *J. Neurophysiol.* 1997, 78, 848–859.
217. Helmchen, C.; Machner, B.; Schwenke, H.; Sprenger, A. Bilateral lesion of the cerebellar fastigial nucleus: Effects on smooth pursuit acceleration and non-reflexive visually-guided saccades. *Front. Neurol.* 2022, 13, 883213.
218. Klam, F.; Petit, J.; Grantyn, A.; Berthoz, A. Predictive elements in ocular interception and tracking of a moving target by untrained cats. *Exp. Brain Res.* 2001, 139, 233–247.
219. de Brouwer, S.; Yuksel, D.; Blohm, G.; Missal, M.; Lefèvre, P. What triggers catch-up saccades during visual tracking? *J. Neurophysiol.* 2002, 87, 1646-1650.
220. Berthoz A. *Simplexity: Simplifying Principles for a Complex World* (translated by Weiss G). New Haven, CT: Yale Univ. Press, 2012.
221. Fuchs, A.F. Saccadic and smooth pursuit eye movements in the monkey. *J. Physiol.* 1967a, 191, 609–631.
222. Van Gelder, P.; Lebedev, S.; Liu, P.M.; Tsui, W.H. Anticipatory saccades in smooth pursuit: task effects and pursuit vector after saccades. *Vision Res.* 1995, 35, 667-678.
223. Goffart, L.; Cecala, A. L.; Gandhi, N. J. The superior colliculus and the steering of saccades toward a moving visual target. *J. Neurophysiol.* 2017, 118, 2890-2901.
224. Keller, E.L.; Gandhi, N.J.; Weir, P.T. Discharge of superior collicular neurons during saccades made to moving targets. *J. Neurophysiol.* 1996, 76, 3573–3577.
225. Paré, M.; Wurtz, R. H. Monkey posterior parietal cortex neurons antidromically activated from superior colliculus. *J. Neurophysiol.* 1997, 78, 3493-3497.
226. Churan, J.; Kaminiarz, A.; Schwenk, J.C.; Bremmer, F. Coding of interceptive saccades in parietal cortex of macaque monkeys. *Brain Struct. Funct.* 2021, 226, 2707-2723.
227. Sommer, M.A., & Wurtz, R.H. Frontal eye field neurons orthodromically activated from the superior colliculus. *J. Neurophysiol.* 1998, 80, 3331–3335.
228. Cassanello, C.R.; Nihalani, A.T.; Ferrera, V.P. Neuronal responses to moving targets in monkey frontal eye fields. *J. Neurophysiol.* 2008, 100, 1544-1556.
229. Shidara, M.; Kawano, K; Gomi, H.; Kawato, M. Inverse-dynamics model eye movement control by Purkinje cells in the cerebellum. *Nature* 1993, 365, 50–52.
230. Das, V.E.; Economides, J.R.; Ono, S.; Mustari, M.J. Information processing by parafoveal cells in the primate nucleus of the optic tract. *Exp. Brain Res.* 2001, 140, 301–310.

231. Inoue, Y.; Takemura, A.; Kawano, K.; Mustari, M.J. Role of the pretectal nucleus of the optic tract in short-latency ocular following responses in monkeys. *Exp. Brain Res.* 2000, 131, 269–281.
232. Ono, S.; Das, V.E.; Economides, J.R.; Mustari, M.J. Modeling of smooth pursuit-related neuronal responses in the DLPN and NRTP of the rhesus macaque. *J. Neurophysiol.* 2005, 93, 108-116.
233. Sun, Z.; Dicke, P.W.; Thier, P. Differential kinematic encoding of saccades and smooth-pursuit eye movements by fastigial neurons. *Neurosci. Bull.* 2022, 38, 927-932.
234. Sun, Z.; Smilgin, A.; Junker, M.; Dicke, P.W.; Thier, P. The same oculomotor vermal Purkinje cells encode the different kinematics of saccades and of smooth pursuit eye movements. *Scient. Rep.* 2017, 7, 40613.
235. Mays, L.E.; Sparks, D.L. Dissociation of visual and saccade-related responses in superior colliculus neurons. *J. Neurophysiol.* 1980, 43, 207-232.
236. Sparks, D.L.; Porter, J.D. Spatial localization of saccade targets. II. Activity of superior colliculus neurons preceding compensatory saccades. *J. Neurophysiol.* 1983, 49, 64-74.
237. Sparks, D.L. Conceptual issues related to the role of the superior colliculus in the control of gaze. *Cur. Opin. Neurobiol.* 1999, 9, 698-707.
238. Leigh, R.J.; Zee, D.S. *The neurology of eye movements.* Oxford University Press, USA, 2015.
239. Rucker, J.C.; Lavin, P.J. *Neuro-ophthalmology: ocular motor system.* Bradley's Neurology in Clinical Practice E-Book 2021, 18, 1-43.
240. Büttner-Ennever, J.A. *Neuroanatomy of the oculomotor system.* Elsevier, USA, 2006.
241. Raphan, T.; Cohen, B. Brainstem mechanisms for rapid and slow eye movements. *Annu. Rev. Physiol.* 1978, 40, 527-552.
242. Fuchs, A.F.; Kaneko, C.R.; Scudder, C.A. Brainstem control of saccadic eye movements. *Annu. Rev. Neurosci.* 1985, 8, 307-337.
243. Henn, V.; Henn, K.; Vilis, T.; Cohen, B. Brainstem regions related to saccade generation. *Rev. Oculom. Res.* 1989, 3, 105-212.
244. Sparks, D.L.; Mays, L.E. Signal transformations required for the generation of saccadic eye movements. *Annu. Rev. Neurosci.* 1990, 13, 309-336.
245. Keller, E.L.; Heinen, S.J. Generation of smooth-pursuit eye movements: neuronal mechanisms and pathways. *Neurosci. Res.* 1991, 11, 79-107.
246. Moschovakis, A.K.; Scudder, C.A.; Highstein, S.M. The microscopic anatomy and physiology of the mammalian saccadic system. *Prog. Neurobiol.* 1996, 50, 133-254.
247. Scudder, C.A.; Kaneko, C.R.S.; Fuchs, A.F. The brainstem burst generator for saccadic eye movements: a modern synthesis. *Exp. Brain Res.* 2002, 142, 439-462.

248. Horn, A.K.; Leigh, R.J. The anatomy and physiology of the ocular motor system. *Handb. Clin. Neurol.* 2011, 102, 21-69.
249. Mustari, M.J.; Ono, S.; Das, V.E. Signal processing and distribution in cortical-brainstem pathways for smooth pursuit eye movements. *Ann. N. Y. Acad. Sci.* 2009, 1164, 147-154.
250. Spencer, R.F.; McNeer, K.W. The periphery: extraocular muscles and motor neurones. In *Eye movements*; Carpenter, R.H.S.; MacMillan Press, London, UK, Volume 8, pp.175-199.
251. Fuchs, A.F.; Luschei, E.S. Firing patterns of abducens neurons of alert monkeys in relationship to horizontal eye movement. *J. Neurophysiol.* 1970, 33, 382-392.
252. Schiller, P.H. The discharge characteristics of single units in the oculomotor and abducens nuclei of the unanesthetized monkey. *Exp. Brain Res.* 1970, 10, 347-362.
253. Fuchs, A.F.; Scudder, C.A.; Kaneko, C.R., 1988. Discharge patterns and recruitment order of identified motoneurons and internuclear neurons in the monkey abducens nucleus. *J. Neurophysiol.* 60, 1874-1895.
254. Sylvestre, P.A.; Cullen, K.E., 1999. Quantitative analysis of abducens neuron discharge dynamics during saccadic and slow eye movements. *J. Neurophysiol.* 82, 2612-2632.
255. Luschei, E.S.; Fuchs, A.F. Activity of brain stem neurons during eye movements of alert monkeys. *J. Neurophysiol.* 1972, 35, 445-461.
256. Keller, E.L. Participation of medial pontine reticular formation in eye movement generation in monkey. *J. Neurophysiol.* 1974, 37, 316-332.
257. Van Gisbergen, J.A.M.; Robinson, D.A.; Gielen, S.A. Quantitative analysis of generation of saccadic eye movements by burst neurons. *J. Neurophysiol.* 1981, 45, 417-442.
258. Horn, A.K.; Büttner-Ennever, J.A.; Suzuki, Y.; Henn, V. Histological identification of premotor neurons for horizontal saccades in monkey and man by parvalbumin immunostaining. *J. Comp. Neurol.* 1995, 359, 350-363.
259. Henn, V.; Lang, W.; Hepp, K.; Reisine, H. Experimental gaze palsies in monkeys and their relation to human pathology. *Brain* 1984, 107, 619-636.
260. Cohen, B.; Komatsuzaki, A. Eye movements induced by stimulation of the pontine reticular formation: evidence for integration in oculomotor pathways. *Exp. Neurol.* 1972, 36, 101-117.
261. Anderson, S.R.; Porrill, J.; Sklavos, S.; Gandhi, N. J.; Sparks, D.L.; Dean, P. Dynamics of primate oculomotor plant revealed by effects of abducens microstimulation. *J. Neurophysiol.* 2009, 101, 2907-2923.
262. Strassman, A.; Highstein, S.M.; McCrea, R.A. Anatomy and physiology of saccadic burst neurons in the alert squirrel monkey. II. Inhibitory burst neurons. *J. Comp. Neurol.* 1986, 249, 358-380.

263. Scudder, C.A.; Fuchs, A.F.; Langer, T.P. Characteristics and functional identification of saccadic inhibitory burst neurons in the alert monkey. *J. Neurophysiol.* 1988, 59, 1430-1454.
264. Cullen, K.E.; Guitton, D. Analysis of primate IBN spike trains using system identification techniques. I. Relationship to eye movement dynamics during head-fixed saccades. *J. Neurophysiol.* 1997, 78, 3259-3282.
265. Kojima, Y.; Iwamoto, Y.; Robinson, F.R.; Noto, C.T.; Yoshida, K. Premotor inhibitory neurons carry signals related to saccade adaptation in the monkey. *J. Neurophysiol.* 2008, 99, 220-230.
266. Sparks, D.L.; Barton, E.J. Neural control of saccadic eye movements. *Cur. Opin. Neurobiol.* 1993, 3, 966-972.
267. Goffart, L. Saccadic eye movements: basic neural processes. *Ref. Mod. Neurosci. Biobehav. Psychol.* 2017.
268. Burrelly, C.; Quinet, J.; Goffart, L. Bilateral control of interceptive saccades: evidence from the ipsipulsion of vertical saccades after caudal fastigial inactivation. *J. Neurophysiol.* 2021, 125, 2068-2083.
269. Takahashi, M.; Shinoda, Y. Brain stem neural circuits of horizontal and vertical saccade systems and their frame of reference. *Neurosci.* 2018, 392, 281-328.
270. Keller, E.L., McPeck, R.M.; Salz, T. Evidence against direct connections to PPRF EBNs from SC in the monkey. *J. Neurophysiol.* 2000, 84, 1303-1313.
271. Roldan, M.; Reinoso-Suarez, F. Cerebellar projections to the superior colliculus in the cat. *J. Neurosci.* 1981, 1, 827-834.
272. Ohtsuka, K.; Noda, H. Saccadic burst neurons in the oculomotor region of the fastigial nucleus of macaque monkeys. *J. Neurophysiol.* 1991, 65, 1422-1434.
273. Fuchs, A.F.; Robinson, F.R.; Straube, A. Role of the caudal fastigial nucleus in saccade generation. I. Neuronal discharge pattern. *J. Neurophysiol.* 1993, 70, 1723-1740.
274. Helmchen, C.; Straube, A.; Büttner, U. Saccade-related activity in the fastigial oculomotor region of the macaque monkey during spontaneous eye movements in light and darkness. *Exp. Brain Res.* 1994, 98, 474-482.
275. Kleine, J.F.; Guan, Y.; Büttner, U. Saccade-related neurons in the primate fastigial nucleus: what do they encode? *J. Neurophysiol.* 2003, 90, 3137-3154.
276. Robinson, F.R.; Fuchs, A.F. The role of the cerebellum in voluntary eye movements. *Ann. Rev. Neurosci.* 2001, 24, 981-1004.
277. Fuchs, A.F.; Brettler, S.; Ling, L. Head-free gaze shifts provide further insights into the role of the medial cerebellum in the control of primate saccadic eye movements. *J. Neurophysiol.* 2010, 103, 2158-2173.

278. Quinet, J.; Goffart, L. Head-unrestrained gaze shifts after muscimol injection in the caudal fastigial nucleus of the monkey. *J. Neurophysiol.* 2007, 98, 3269-3283.
279. Robinson, D.A. Oculomotor unit behavior in the monkey. *J. Neurophysiol.* 1970, 33: 393–403.
280. Fuchs, A.F.; Luschei, E.S. The activity of single trochlear nerve fibers during eye movements in the alert monkey. *Exp. Brain Res.* 1971 13: 78-89.
281. Walton, M.M. Reduced activity of vertically acting motoneurons during convergence. *J. Neurophysiol.* 2022, 128, 671-680.
282. Büttner, U; Büttner-Ennever, J.A.; Henn V. Vertical eye movement related unit activity in the rostral mesencephalic reticular formation of the alert monkey. *Brain Res.* 1977, 130,239-252.
283. King, W.M.; Fuchs, A.F. Reticular control of vertical saccadic eye movements by mesencephalic burst neurons. *J. Neurophysiol.* 1979, 42, 861-876.
284. Moschovakis, A.K.; Scudder, C.A.; Highstein, S.M. The structure of the primate oculomotor burst generator. I. Medium-lead burst neurons with upward on-directions. *J. Neurophysiol.* 1991, 65, 203-217.
285. Moschovakis, A.K.; Scudder, C.A.; Highstein, S.M.; Warren, J.D.M. The structure of the primate oculomotor burst generator. II. Medium-lead burst neurons with downward on-directions. *J. Neurophysiol.* 1991, 65, 218-229.
286. Horn, A.K.; Helmchen, C.; Wahle, P. GABAergic neurons in the rostral mesencephalon of the macaque monkey that control vertical eye movements. *Ann. N. Y. Acad. Sci.* 2003, 1004, 19-28.
287. Suzuki, Y.; Büttner-Ennever, J.A.; Straumann, D.; Hepp, K.; Hess, B.J.M.; Henn, V. Deficits in torsional and vertical rapid eye movements and shift of Listing's plane after uni- and bilateral lesions of the rostral interstitial nucleus of the medial longitudinal fasciculus. *Exp. Brain Res.* 1995, 106, 215-232.
288. Straumann, D.; Zee, D.S.; Solomon, D.; Lasker, A.G.; Roberts, D.C. Transient torsion during and after saccades. *Vision Res.* 1995, 35, 3321-3334
289. Crawford, J.D.; Vilis, T. Symmetry of oculomotor burst neuron coordinates about Listing's plane. *J. Neurophysiol.* 1992, 68, 432-448.
290. Kremmyda, O., Büttner-Ennever, J.A., Büttner, U.; Glasauer, S. Torsional deviations with voluntary saccades caused by a unilateral midbrain lesion. *Case Reports* 2009, bcr0820080807.
291. Ohtsuka, K.; Noda, H. Discharge properties of Purkinje cells in the oculomotor vermis during visually guided saccades in the macaque monkey. *J. Neurophysiol.* 1995, 74, 1828-1840.
292. Soetedjo, R.; Fuchs, A.F. Complex spike activity of purkinje cells in the oculomotor vermis during behavioral adaptation of monkey saccades. *J. Neurosci.* 2006 26, 7741-7755.

293. Sato, H.; Noda, H. Divergent axon collaterals from fastigial oculomotor region to mesodiencephalic junction and paramedian pontine reticular formation in macaques. *Neurosci. Res.* 1991, 11, 41-54.
294. Iwamoto, Y.; Yoshida, K. Saccadic dysmetria following inactivation of the primate fastigial oculomotor region. *Neurosci. Lett.* 2002, 325, 211-215.
295. Goffart, L.; Chen, L.L.; Sparks, D. L. Saccade dysmetria during functional perturbation of the caudal fastigial nucleus in the monkey. *Ann. N. Y. Acad. Sci.* 2003, 1004, 220-228.
296. Takagi, M.; Zee, D.S.; Tamargo, R.J. Effects of lesions of the oculomotor vermis on eye movements in primate: saccades. *J. Neurophysiol.* 1998, 80, 1911-1931.
297. Nitta, T.; Akao, T.; Kurkin, S.; Fukushima, K. Involvement of the cerebellar dorsal vermis in vergence eye movements in monkeys. *Cereb. Cortex* 2008, 18, 1042-1057.
298. Helmchen, C.; Glasauer, S.; Büttner, U. Pathological torsional eye deviation during voluntary saccades: a violation of Listing's law. *J. Neurol. Neurosurg. Psychiat.* 1997, 62, 253-260.
299. Fetter, M.; Anastasopoulos, D.; Haslwanter, T. Three-dimensional properties of saccadic eye movements in patients with cerebellar ataxia. In *Current Oculomotor Research: Physiological and Psychological Aspects* 1999, pp. 391-396, Boston, MA: Springer US.
300. Ohgaki, T.; Markham, C.H.; Schneider, J.S.; Curthoys, I.S. Anatomical evidence of the projection of pontine omnipause neurons to midbrain regions controlling vertical eye movements. *J. Comp. Neurol.* 1989, 289, 610-625.
301. Grantyn, A.; Brandi, A.M.; Dubayle, D.; Graf, W.; Ugolini, G.; Hadjidimitrakis, K.; Moschovakis, A. Density gradients of trans-synaptically labeled collicular neurons after injections of rabies virus in the lateral rectus muscle of the rhesus monkey. *J. Comp. Neurol.* 2002, 451, 346-361.
302. Nichols, M.J.; Sparks, D.L. Component stretching during oblique stimulation-evoked saccades: the role of the superior colliculus. *J. Neurophysiol.* 1996a, 76, 582-600.
303. Sparks, D.L.; Barton, E.J.; Gandhi, N.J.; Nelson, J. Studies of the role of the paramedian pontine reticular formation in the control of head-restrained and head-unrestrained gaze shifts. *Ann. N. Y. Acad. Sci.* 2002, 956, 85-98.
304. Barton, E.J.; Nelson, J.S.; Gandhi, N.J.; Sparks, D.L. Effects of partial lidocaine inactivation of the paramedian pontine reticular formation on saccades of macaques. *J. Neurophysiol.* 2003, 90, 372-386.
305. Bizzi, E.; Kalil, R.E.; Morasso, P. Two modes of active eye-head coordination in monkeys. *Brain Res.* 1972, 40, 45-48.
306. Volle, M.; Guitton, D. Human gaze shifts in which head and eyes are not initially aligned. *Exp. Brain Res.* 1993, 94, 463-470.

307. Tomlinson, R.D.; Bahra, P.S. Combined eye-head gaze shifts in the primate. I. Metrics. *J. Neurophysiol.* 1986, 56, 1542-1557.
308. Becker, W.; Jürgens, R. Gaze saccades to visual targets: does head movement change the metrics. *The head-neck sensory motor system* 1992, 427-433.
309. Quinet, J.; Goffart, L. Influence of head restraint on visually triggered saccades in the Rhesus monkey. *Ann. N. Y. Acad. Sci.* 2003, 1004, 404-408.
310. Richmond, F.J.; Singh, K.; Corneil, B.D. Neck muscles in the rhesus monkey. I. Muscle morphometry and histochemistry. *J. Neurophysiol.* 2001, 86, 1717-1728.
311. Lestienne, F.G., Le Goff, B.; Liverneaux, P.A. Head movement trajectory in three-dimensional space during orienting behavior toward visual targets in rhesus monkeys. *Exp. Brain Res.* 1995, 102, 393-406.
312. Corneil, B.D.; Olivier, E.; Richmond, F.J., Loeb, G.E.; Munoz, D. P. Neck muscles in the rhesus monkey. II. Electromyographic patterns of activation underlying postures and movements. *J. Neurophysiol.* 2001, 86, 1729-1749.
313. Paré, M.; Crommelinck, M.; Guitton, D. Gaze shifts evoked by stimulation of the superior colliculus in the head-free cat conform to the motor map but also depend on stimulus strength and fixation activity. *Exp. Brain Res.* 1994, 101, 123-139.
314. Freedman, E.G.; Sparks, D.L. Activity of cells in the deeper layers of the superior colliculus of the rhesus monkey: evidence for a gaze displacement command. *J. Neurophysiol.* 1997, 78, 1669-1690.
315. Rodgers, C.K.; Munoz, D.P.; Scott, S.H.; Paré, M. Discharge properties of monkey tectoreticular neurons. *J. Neurophysiol.* 2006, 95, 3502-3511.
316. Choi, W.Y.; Guitton, D. Firing patterns in superior colliculus of head-unrestrained monkey during normal and perturbed gaze saccades reveal short-latency feedback and a sluggish rostral shift in activity. *J. Neurosci.* 2009, 29, 7166-7180.
317. Knight, T.A.; Fuchs, A.F. Contribution of the frontal eye field to gaze shifts in the head-unrestrained monkey: effects of microstimulation. *J. Neurophysiol.* 2007, 97, 618-634.
318. Chen, L.L.; Tehovnik, E.J. Cortical control of eye and head movements: integration of movements and percepts. *Eur. J. Neurosci.* 2007, 25, 1253-1264.
319. Knight, T.A. Contribution of the frontal eye field to gaze shifts in the head-unrestrained rhesus monkey: neuronal activity. *Neurosci.* 2012, 225, 213-236.
320. Thomson, D.B.; Loeb, G. E.; Richmond, F.J. Effect of neck posture on the activation of feline neck muscles during voluntary head turns. *J. Neurophysiol.* 1994, 72, 2004-2014.
321. Highstein, S.M.; Holstein, G.R. The anatomy of the vestibular nuclei. *Prog. Brain Res.* 2006, 151, 157-203.

322. Peterson, B.W.; Maunz, R.A.; Pitts, N.G.; Mackel, R. G. Patterns of projection and branching of reticulospinal neurons. *Exp. Brain Res.* 1975, 23, 333-351.
323. Isa, T.; Sasaki, S. Effects of lesion of paramedian pontomedullary reticular formation by kainic acid injection on the visually triggered horizontal orienting movements in the cat. *Neurosci. Lett.* 1988, 87, 233-239.
324. Grantyn, A.; Berthoz, A. Reticulo-spinal neurons participating in the control of synergic eye and head movements during orienting in the cat: I. Behavioral properties. *Exp. Brain Res.* 1987, 66, 339-354.
325. Grantyn, A.; Berthoz, A. Burst activity of identified tecto-reticulo-spinal neurons in the alert cat. *Exp. Brain Res.* 1985, 57, 417-421.
326. Grantyn, A. How visual inputs to the ponto-bulbar reticular formation are used in the synthesis of premotor signals during orienting. *Prog. Brain Res.* 1989, 80, 159-170.
327. Alstermark, B.; Pinter, M.J.; Sasaki, S. Pyramidal effects in dorsal neck motoneurons of the cat. *J. Physiol.* 1985, 363, 287-302.
328. Escudero, M; de la Cruz, R.R.; Delgado-Garcia, J.M. A physiological study of vestibular and prepositus hypoglossi neurons projecting to the abducens nucleus in the alert cat. *J. Physiol.* 1992, 458: 539-560.
329. Grantyn, A.; Kuze, B.; Brandi, A.M.; Thomas, M.A.; Quenech'du, N. Direct projections of omnipause neurons to reticulospinal neurons: A double-labeling light microscopic study in the cat. *J. Comp. Neurol.* 2010, 518, 4792-4812.
330. Paré, M.; Guitton, D. Brain stem omnipause neurons and the control of combined eye-head gaze saccades in the alert cat. *J. Neurophysiol.* 1998, 79(6), 3060-3076.
331. Phillips, J.O.; Ling, L.; Fuchs, A.F. Action of the brain stem saccade generator during horizontal gaze shifts. I. Discharge patterns of omnidirectional pause neurons. *J. Neurophysiol.* 1999, 81(3), 1284-1295.
332. Wilson, V.J.; Yoshida, M. Monosynaptic inhibition of neck motoneurons by the medial vestibular nucleus. *Exp. Brain Res.* 1969, 9, 365-380.
333. Wilson, V.J.; Yoshida, M. Comparison of effects of stimulation of Deiters' nucleus and medial longitudinal fasciculus on neck, forelimb, and hindlimb motoneurons. *J. Neurophysiol.* 1969, 32, 743-758.
334. Wilson, V.J.; Maeda, M. Connections between semicircular canals and neck motoneurons in the cat. *J. Neurophysiol.* 1974, 37, 346-357.
335. Isu, N.; Yokota, J. Morphophysiological study on the divergent projection of axon collaterals of medial vestibular nucleus neurons in the cat. *Exp. Brain Res.* 1983, 53, 151-162.

336. Rapoport, S.; Susswein, A.; Uchino, Y.; Wilson, V. J. Synaptic actions of individual vestibular neurones on cat neck motoneurons. *J. Physiol.* 1977, 272, 367-382.
337. Suzuki, J.I.; Cohen, B. Head, eye, body and limb movements from semicircular canal nerves. *Exp. Neurol.* 1964, 10, 393-405.
338. Baker, R.G.; Mano, N.; Shimazu, H. Postsynaptic potentials in abducens motoneurons induced by vestibular stimulation. *Brain Res.* 1969, 15, 577-580.
339. Hikosaka, O.; Igusa, Y.; Imai, H. Inhibitory connections of nystagmus-related reticular burst neurons with neurons in the abducens, prepositus hypoglossi and vestibular nuclei in the cat. *Exp. Brain Res.* 1980, 39, 301-311.
340. McCrea, R.A.; Yoshida, K.; Berthoz, A.; Baker, R. Eye movement related activity and morphology of second order vestibular neurons terminating in the cat abducens nucleus. *Exp. Brain Res.* 1980, 40, 468-473.
341. Berthoz, A.; Droulez, J.; Vidal, P.P.; Yoshida, K. Neural correlates of horizontal vestibulo-ocular reflex cancellation during rapid eye movements in the cat. *J. Physiol.* 1989, 419, 717-751.
342. Boyle, R.; Goldberg, J.M.; Highstein, S.M. Vestibular nerve inputs to vestibulospinal and vestibulo-ocular neurons of the squirrel monkey. In: *The Head-Neck Sensory Motor System*, A Berthoz, W Graf et PP Vidal (Eds), 1992, Oxford University Press, New York, p 255-258.
343. Anastasopoulos, D.; Mergner, T. Canal-neck interaction in vestibular nuclear neurons of the cat. *Exp. Brain Res.* 1982, 46, 269-280.
344. Hannaford, B.; Stark, L. Roles of the elements of the triphasic control signal. *Exp. Neurol.* 1985, 90, 619-634.
345. Roucoux, A.; Crommelinck, M.; Decostre, M.F. Neck muscle activity in eye—head coordinated movements. *Prog. Brain Res.* 1989, 80, 351-362.
346. Munoz D.P.; Wurtz R.H. Saccade-related activity in monkey superior colliculus. I. Characteristics of burst and buildup cells. *J. Neurophysiol.* 1995, 73, 2313-2333.
347. Munoz, D.P.; Waitzman D. M.; Wurtz, R.H. Activity of neurons in monkey superior colliculus during interrupted saccades. *J. Neurophysiol.* 1996, 75, 2562-2580.
348. Everling, S., Paré, M.; Dorris, M.C.; Munoz, D.P. Comparison of the discharge characteristics of brain stem omnipause neurons and superior colliculus fixation neurons in monkey: implications for control of fixation and saccade behavior. *J. Neurophysiol.* 1998, 79, 511-528.
349. Goossens H.H.L.M.; Van Opstal, A.J. Blink-perturbed saccades in monkey. II. Superior colliculus activity. *J. Neurophysiol.* 2000, 83, 3430-3452.
350. Keller E.L.; Gandhi N.J.; Vijay Sekaran, S. Activity in deep intermediate layer collicular neurons during interrupted saccades. *Exp. Brain Res.* 2000, 130, 227-237.

351. Moschovakis, A.K. Are laws that govern behavior embedded in the structure of the CNS? The case of Hering's law. *Vision Res.* 1995, 35, 3207-3216.
352. McDougal, D.H.; Gamlin, P.D. Autonomic Control of the Eye. *Comprehens. Physiol.* 2014, 5, 439-473.
353. Wu, F.; Zhao, Y.; Zhang, H. Ocular autonomic nervous system: an update from anatomy to physiological functions. *Vision* 2022, 6, 6.
354. Moruzzi G, Magoun HW. Brain stem reticular formation and activation of the EEG. *Electroencephalogr. Clin. Neurophysiol.* 1949, 1,455-473.
355. von Helmholtz H. Treatise on physiological optics. J. P. C. Southall (ed) 1924, Optical Soc. America. <http://psych.upenn.edu/backuslab/helmholtz>
356. Simpson, J.I.; Graf, W. Eye-muscle geometry and compensatory eye movements in lateral-eyed and frontal-eyed animals. *Ann. N. Y. Acad. Sci.* 1981, 374, 20-30.
357. Cox, P.G.; Jeffery, N. Morphology of the mammalian vestibulo-ocular reflex: The spatial arrangement of the human fetal semicircular canals and extraocular muscles. *J. Morphol.* 2007, 268, 878-890.
358. Cox, P.G.; Jeffery, N. Geometry of the semicircular canals and extraocular muscles in rodents, lagomorphs, felids and modern humans. *J. Anat.* 2008, 213, 583-596.