



HAL
open science

INVARIANTS OF PERSISTENCE MODULES DEFINED BY ORDER-EMBEDDINGS

Claire Amiot, Thomas Brüstle, Eric Hanson

► **To cite this version:**

Claire Amiot, Thomas Brüstle, Eric Hanson. INVARIANTS OF PERSISTENCE MODULES DEFINED BY ORDER-EMBEDDINGS. 2024. hal-04708065

HAL Id: hal-04708065

<https://cnrs.hal.science/hal-04708065v1>

Preprint submitted on 24 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INVARIANTS OF PERSISTENCE MODULES DEFINED BY ORDER-EMBEDDINGS

CLAIRE AMIOT, THOMAS BRÜSTLE, ERIC J. HANSON

ABSTRACT. One of the main objectives of topological data analysis is the study of discrete invariants for persistence modules, in particular when dealing with multiparameter persistence modules. In many cases, the invariants studied for these non-totally ordered posets \mathcal{P} can be obtained from restricting a given module to a subposet \mathcal{X} of \mathcal{P} that is totally ordered (or more generally, of finite representation type), and then computing the barcode (or the general direct sum decomposition) over \mathcal{X} .

We consider in this paper general order-preserving embeddings of representation-finite subposets \mathcal{X} into \mathcal{P} and study systematically the invariants obtained by decomposing the restriction of a given \mathcal{P} -module M to \mathcal{X} into its indecomposable summands. The restriction functor from $\text{mod } \mathcal{P}$ to $\text{mod } \mathcal{X}$ is well-studied, and it is known to be exact and admits both left and right adjoint functors, known as induction and co-induction functors. This allows us to obtain new homological insights, and also to re-interpret previous results. We use this approach also to determine bases, thus generalizing the concept of signed barcodes which is considered in the literature in relation to stability results.

It turns out that considering only order-embeddings of *one fixed* poset \mathcal{X} into the poset \mathcal{P} , and studying the set of all indecomposables obtained from \mathcal{X} introduces a lot of redundancy. We therefore also study iterated embeddings of several posets of increasing sizes, while limiting attention to only some indecomposables (that have not been obtained from embedding of smaller posets previously).

CONTENTS

1. Introduction	1
2. \mathbb{Z} -linear invariants	4
3. Invariants defined by order-embeddings	6
4. Invariants and exact structures	10
5. Interval bases via families of embeddings	15
References	21

1. INTRODUCTION

1.1. **Setting.** The study of discrete invariants for persistence modules is one of the main objectives of topological data analysis (TDA). When a pointwise finite-dimensional persistence module M is defined over a totally ordered poset \mathcal{P} , it decomposes into a finite sum of intervals. This data is called the *barcode* of M and completely describes the module up to isomorphism, see [BC-B20, Theorem 1.2] and [GR92, Section 3.6]. However, when studying multiparameter persistence modules, the underlying poset \mathcal{P} usually has wild representation type, and thus does not admit such a complete discrete invariant. We refer to the summary article [BL23] for an introduction to multiparameter persistence modules, and why it is relevant in practice to study more general posets than totally ordered ones.

To address the need of TDA for discrete invariants even in the multiparameter case, several invariants have been introduced and studied: the dimension vector $\mathbf{dim}_{\mathcal{P}}$ (aka Hilbert function), the rank invariant $\mathbf{rk}_{\mathcal{P}}$, and a number of generalizations thereof, see [BL23, BOOS] for an overview of recently studied invariants.

It turns out that, in many cases, these invariants studied for non-totally ordered posets \mathcal{P} can be obtained from restricting a given module to a subposet \mathcal{X} of \mathcal{P} that is totally ordered (or more generally, of finite representation type), and then computing the barcode (or the general direct sum decomposition) over \mathcal{X} . This approach has been described in e.g. [DKM24, AENY23b, HNOX] introducing certain subposets \mathcal{X} having the form of “zigzags” or “tours” and “courses”, all referring to a poset whose indecomposable representations can be described via intervals.

In this paper, we consider general embeddings of representation-finite subposets \mathcal{X} into \mathcal{P} and study invariants obtained by decomposing the restriction of a given \mathcal{P} -module M to \mathcal{X} into its indecomposable summands. The restriction functor $\text{mod } \mathcal{P} \rightarrow \text{mod } \mathcal{X}$ is well-studied, and it is known to be exact and admits both left and right adjoint functors. This allows one to obtain new homological insights, and also to re-interpret previous results. It also allows one to determine bases, thus generalizing the concept of signed barcodes from [BOO].

To fix notation, consider an *order-embedding* $f : \mathcal{X} \rightarrow \mathcal{P}$; that is, a poset morphism that is an isomorphism onto its image. The restriction of a \mathcal{P} -module M to $f(\mathcal{X})$ can be described completely by counting the multiplicities of indecomposable direct summands. We assume \mathcal{X} to be representation-finite, and thus obtain a discrete invariant $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^f$ of rank $|\text{Ind } \mathcal{X}|$, the number of indecomposable \mathcal{X} -modules. Repeating this process for all possible embeddings $f : \mathcal{X} \rightarrow \mathcal{P}$ and collecting the result into a vector indexed over all embeddings, we obtain the invariant $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$ describing all multiplicities of indecomposable \mathcal{X} -modules obtained from all possible ways to restrict the \mathcal{P} -module M to a subposet of form \mathcal{X} . One also defines an equivalent invariant $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}$ by calculating the Hom-dimensions from the set of indecomposable \mathcal{X} -modules to the restrictions of the persistence module along all order embeddings. See Definition 3.3 and Proposition 3.6 for details.

Consider for example the poset \mathcal{X}_1 with one element. Then \mathcal{X}_1 has only one indecomposable representation, and its multiplicity in a general module is given by the dimension. Thus listing all possible embeddings of \mathcal{X}_1 into \mathcal{P} and counting multiplicities amounts to describing the dimension vector, so $\mathbf{mult}_{\mathcal{X}_1, \mathcal{P}} = \mathbf{dim}_{\mathcal{P}}$. Consider now the poset $\mathcal{X}_2 = \{1 \rightarrow 2\}$. This poset is representation-finite, and the multiplicity of the indecomposable representation given by the interval $[1, 2]$ can be computed by calculating the rank of the map $M_1 \rightarrow M_2$. This allows one to show that the multiplicity invariant $\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}$ is equivalent to the well-studied rank invariant $\mathbf{rk}_{\mathcal{P}}$, see Corollary 3.8. More generally, as discussed in Example 3.5, the “generalized rank invariant” of [KM21] and the “compressed multiplicities” of [AENY23b] both record the multiplicity of the sincere interval modules $\mathbb{I}_{\mathcal{X}}$ over certain subsets $\mathcal{X} \subseteq \mathcal{P}$.

1.2. Results. For several subposets \mathcal{X} , we address the question of how the multiplicities in $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$ can be computed in terms of linear algebra. See Examples 3.7 and 3.14 and the calculations in Section 3. In each case, the multiplicities of indecomposable summands of M can be obtained computing ranks or intersections given by the structure maps of the module M . More generally, we show in Proposition 3.6 that the invariant $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$ can be obtained by calculating the Hom-dimensions from the set of indecomposable representations to the given persistence module, thus reducing the problem to solving a system of linear equations.

Besides computability, the second and equally important quality of an invariant for persistence modules is the stability. For the rank invariant, this question has been addressed in [BOO, BOOS]: they single out a class of special persistence modules, the “hook modules”, which are defined as indicator modules of certain shapes called hooks: For $a < b$ in \mathcal{P} , let $[a, b[= \{c \in \mathcal{P} \mid a \leq c \not\leq b\}$. The interval module $\mathbb{I}_{[a, b[}$ is called a *hook module*. Note that $\mathbb{I}_{[a, b[}$ is the cokernel of the natural map $\mathbb{I}_{[b, \infty)} \rightarrow \mathbb{I}_{[a, \infty)}$ between indecomposable projective \mathcal{P} -modules. It is shown in [BOO] that the hook modules play the role of projective objects in a relative homology theory given by some exact structure \mathcal{E} , thus every persistence module M admits a resolution by hook modules. One can then show that these hooks form a basis for the (image of the) rank invariant (see Definition 4.11), which in turn allows one to define a “signed barcode”, leading to a stability result for the considered invariant [BOOS].

The viewpoint of relative homological algebra has further been extended in many recent works (see e.g. [AENY23a, BBH24, BBH, BOO, BOOS, CGRST, OS]), but we are only now able to conceptually explain the appearance and importance of hook modules and their generalizations: Studying embeddings $f : \mathcal{X} \rightarrow \mathcal{P}$, the corresponding restriction functor $\text{mod } \mathcal{P} \rightarrow \text{mod } \mathcal{X}$ admits both a left adjoint $f_! : \text{mod } \mathcal{X} \rightarrow \text{mod } \mathcal{P}$ and a right adjoint $f^* : \text{mod } \mathcal{X} \rightarrow \text{mod } \mathcal{P}$. One can then use the adjunction formula to show that the set

$$\mathcal{H} = \{f_!U \mid U \text{ indecomposable } \mathcal{X}\text{-module and } f : \mathcal{X} \rightarrow \mathcal{P} \text{ an embedding}\}$$

forms exactly the indecomposable relative projective objects for the exact structure defined as the class \mathcal{E} of short exact sequences $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ in $\text{mod } \mathcal{P}$ which have the property that

$$0 \rightarrow f^*L \rightarrow f^*M \rightarrow f^*N \rightarrow 0$$


splits for all embeddings $f : \mathcal{X} \rightarrow \mathcal{P}$. See Proposition 4.8. Coming back to the rank invariant, we study embeddings of the poset \mathcal{X}_2 , and since every indecomposable \mathcal{X}_2 -module is the cokernel of a map between

indecomposable projectives, we see easily that the set \mathcal{H} above is formed exactly by the hook modules in this case.

Moving on to more general posets that detect other aspects of the persistence module M , we consider $\mathcal{X} = \mathcal{X}'_3$ or \mathcal{X}''_3 :

$$\mathcal{X}'_3 = \begin{array}{c} & & 2 \\ & \nearrow & \\ 1 & & \\ & \searrow & \\ & & 3 \end{array} \quad \mathcal{X}''_3 = \begin{array}{c} & & 3 \\ & \nearrow & \\ 1 & & \\ & \searrow & \\ & & 2 \end{array}$$

We notice that the corresponding generalized hook modules are not always given just by specifying a subposet of \mathcal{P} , but they are given by vector spaces of dimension two at some vertices, see Example 4.10. These can be illustrated as follows:



Already in [BOO], they specified a second basis for the rank invariant, given by rectangle modules $\mathbb{I}_{[a,b]}$ (with support $[a,b]$), which are somewhat easier to deal with than the hook modules. We aim to generalise this result here in terms of general embeddings: Given a poset embedding $f : \mathcal{X} \rightarrow \mathcal{P}$ and $U \in \text{mod } \mathcal{X}$, the adjunction formulas yield a natural map $\theta_f U : f_! U \rightarrow f_* U$. We denote $\Theta_f U = \text{Im}(\theta_f U)$. It is shown in [AET] that the association $U \mapsto \Theta_f U$ extends to a functor $\text{mod } \mathcal{X} \rightarrow \text{mod } \mathcal{P}$. One easily verifies that in case $\mathcal{X} = \mathcal{X}_2$, the set

$$\mathcal{R} = \{ \Theta_f U \mid U \text{ indecomposable } \mathcal{X}\text{-module and } f : \mathcal{X} \rightarrow \mathcal{P} \text{ an embedding} \}$$

coincides with the set of rectangle modules. However, before moving on to find bases in a more general setting, we aim to reduce redundancy. Considering sets like

$$\mathcal{H} = \{ f_! U \mid U \text{ indecomposable } \mathcal{X}\text{-module and } f : \mathcal{X} \rightarrow \mathcal{P} \text{ an embedding} \}$$

produces the same \mathcal{P} -modules several times from different embeddings f and different indecomposables U . It seems more efficient to first collect all simple \mathcal{P} -modules via embeddings of the one-point poset \mathcal{X}_1 , then consider all \mathcal{P} -modules obtained from the interval $[1, 2]$ by embedding the poset $\mathcal{X}_2 = \{1 \rightarrow 2\}$ into \mathcal{P} and so on. In other words, instead of studying the set of all indecomposables obtained from embeddings of one poset \mathcal{X} , we study now embeddings of several posets of increasing sizes, but limit attention to only some indecomposables (that have not been obtained from embedding of smaller posets previously). The setting we consider here is where we fix a family of posets $\mathcal{F} = \{\mathcal{X}_i\}$ together with certain embeddings into \mathcal{P} , and then we define $\mathbf{mult}_{\mathcal{F}}$ to count the multiplicity of the characteristic indecomposable module $\mathbb{I}_{\mathcal{X}_i}$ in the restriction of M to \mathcal{X}_i along the various embeddings. So only one indecomposable is considered for each poset embedding. As discussed in Example 5.8, both the generalized rank invariant of [KM21] and the various compressed multiplicities of [AENY23b] can be realized in the form $\mathbf{mult}_{\mathcal{F}}$.

As explained in Section 5, there are examples of the invariants $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$ which are equivalent to invariants of the form $\mathbf{mult}_{\mathcal{F}}$. In such cases, the perspective of iterated embeddings is useful for removing some of the redundant information stored in the invariant $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$. Aside from this decrease in redundancy, this change in perspective also allows us to describe explicit bases of some of the invariants $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$. See Corollary 5.11.

With $\mathcal{F} := \{\mathcal{X}_1, \mathcal{X}_2\}$ for \mathcal{X}_1 and \mathcal{X}_2 the one- and two-point poset as before, we can recover the result that the set \mathcal{R} (the rectangles in this case) forms a basis for $\mathbf{mult}_{\mathcal{F}}$.


Corollary 1.1 (Corollary 5.15). *Let \mathcal{P} be a finite connected poset, and let \mathcal{X} be one of the following posets*

$$\mathcal{X}'_3 = \begin{array}{c} & & 2 \\ & \nearrow & \\ 1 & & \\ & \searrow & \\ & & 3 \end{array} \quad \mathcal{X}''_3 = \begin{array}{c} & & 3 \\ & \nearrow & \\ 1 & & \\ & \searrow & \\ & & 2 \end{array}$$

Assume that for any $a < b$ in \mathcal{P} there exists $f \in \text{Emb}_{\mathcal{X}}^{\mathcal{P}}$ with $\{a, b\} \in \text{Im } f$, then the set

$$\mathcal{R} = \{\Theta_f(U) \mid (f, U) \in \text{Emb}_{\mathcal{X}}^{\mathcal{P}} \times \text{Ind } \mathcal{X}\}$$

forms a basis of $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$.



1.3. Acknowledgements. Claire Amiot is supported by the ANR project CHARMS and by the Institut Universitaire de France. Thomas Brüstle was supported by the CRM (Centre de recherches mathématiques, Montreal) and Eric Hanson by IUF to visit Institut Fourier, Grenoble. Most of this project was realised during these stays. Thomas Brüstle and Eric Hanson were partially supported by NSERC Discovery Grant RGPIN/04465-2019. Eric Hanson was also partially supported by the Canada Research Chairs program CRC-2021-00120 and NSERC Discovery Grant RGPIN-2022-03960. A portion of this work was completed while Eric Hanson was a postdoctoral student at l'Université de Sherbrooke. The authors also thank Benjamin Blanchette, Steffen Oppermann, Steve Oudot, and Luis Scoccola for stimulating discussions on the subject.

2. \mathbb{Z} -LINEAR INVARIANTS

For a finite poset \mathcal{P} , we study the category $\text{mod } \mathcal{P}$ as the category of functors $\mathcal{P} \rightarrow \text{mod } k$ for some fixed field k . It is a k -linear abelian category with finite dimensional Hom-spaces which satisfies the Krull-Schmidt property; that is, every object is isomorphic to a finite direct sum of indecomposable objects, and this decomposition is unique up to permutation.

An object $M \in \text{mod } \mathcal{P}$ is also called a \mathcal{P} -module. We denote by M_a the vector space $M(a)$, and by $M_{a \rightarrow b}$ the element in $\text{Hom}_k(M_a, M_b)$ given by the functor M applied to the pair $a \leq b \in \mathcal{P}$. We further denote $\mathbf{dim}_a M := \dim_k M_a$, and $\mathbf{rk}_{a,b} M := \text{rk } M_{a \rightarrow b}$.

The *split Grothendieck group* $K_0^{\text{sp}}(\mathcal{P})$ of $\text{mod } \mathcal{P}$ is the free abelian group generated by isoclasses $[M]$ of objects M in $\text{mod } \mathcal{P}$ modulo the relation $[M \oplus N] = [M] + [N]$ for any objects M, N in $\text{mod } \mathcal{P}$. Fixing a set $\text{Ind } \mathcal{P}$ of representatives of isoclasses of the indecomposable \mathcal{P} -modules, the set $\{[U] \in K_0^{\text{sp}}(\mathcal{P}) \mid U \in \text{Ind } \mathcal{P}\}$ forms a basis for $K_0^{\text{sp}}(\mathcal{P})$. In other words, decomposing M into indecomposable direct summands $M \simeq \bigoplus_{U \in \text{Ind } \mathcal{P}} U^{\alpha_U}$, the vector $(\alpha_U)_{U \in \text{Ind } \mathcal{P}}$ uniquely determines M up to isomorphism, by the Krull-Schmidt property.

We say that a poset \mathcal{P} is of *finite representation type* (or *representation-finite*) if $\text{Ind } \mathcal{P}$ is finite. Therefore the free abelian group $K_0^{\text{sp}}(\mathcal{P})$ is finitely generated if and only if \mathcal{P} is of finite representation type.

Some indecomposable \mathcal{P} -modules have been object of intense studies in the persistence theory literature. One example of such modules are the *rectangle modules* $\mathbb{I}_{[a,b]}$ (with support $[a, b]$), see e.g. [BLO22, KM21]: Every pair $a \leq b$ in \mathcal{P} yields an indecomposable \mathcal{P} -module $U = \mathbb{I}_{[a,b]}$ with $U_x = k$ for all $a \leq x \leq b$ in \mathcal{P} and $U_{x \rightarrow y}$ the identity map for all $a \leq x \leq y \leq b$ in \mathcal{P} , while all other vector spaces and linear maps are zero.

More generally, a subset $\mathcal{S} \subseteq \mathcal{P}$ is called an *interval* if \mathcal{S} is order convex (in the sense that any $a \leq b \leq c$ in \mathcal{P} with $a, c \in \mathcal{S}$ also has $b \in \mathcal{S}$) and connected (in the sense that any $a, b \in \mathcal{S}$ are related by an alternating chain $a = s_0 \leq s_1 \geq \dots \geq s_k = b$ with each $s_i \in \mathcal{S}$). Each interval determines an *interval module*¹ $\mathbb{I}_{\mathcal{S}}$ analogous to the rectangle module $\mathbb{I}_{[a,b]}$. Note that interval modules are always indecomposable, see e.g.

¹Note that interval modules are also called “spread modules” in some works, e.g. [BBH24, CGRST].

[BL18, Proposition 2.2]. These modules have featured prominently in many recent works, see e.g. Sections 9 and 10 of the survey article [BL23] and the references therein. We will consider the following special classes of interval modules in this paper.

- Example 2.1.** (1) For $a \in \mathcal{P}$, let $[a, \infty) = \{b \in \mathcal{P} \mid a \leq b\}$. The interval module $P_a := \mathbb{I}_{[a, \infty)}$ is an indecomposable projective, and all indecomposable projective \mathcal{P} -modules have this form.
- (2) For $a \in \mathcal{P}$, let $(-\infty, a] = \{b \in \mathcal{P} \mid a \geq b\}$. The interval module $I_a := \mathbb{I}_{(-\infty, a]}$ is an indecomposable injective. Note also that all indecomposable injective \mathcal{P} -modules have this form.
- (3) For $a \in \mathcal{P}$, the interval module $S_a := \mathbb{I}_{\{a\}}$ (which is also a rectangle module since $\{a\} = [a, a]$) is the simple top of P_a and the simple socle of I_a . Note that all simple \mathcal{P} -modules have this form.
- (4) For $a < b$, let $[a, b[= \{c \in \mathcal{P} \mid a \leq c \not\leq b\}$. The interval module $H_{a,b} := \mathbb{I}_{[a, b[}$ is called a *hook module*, see [BOO]. Note that $H_{a,b}$ is the cokernel of the natural injective map $P_b \rightarrow P_a$.
- (5) For $a < b$, let $]a, b] = \{c \in \mathcal{P} \mid b \geq c \not\geq a\}$. The interval module $C_{a,b} := \mathbb{I}_{]a, b]}$ is called a *cohook module*. Note that $C_{a,b}$ is the kernel of the natural surjective map $I_a \rightarrow I_b$.
- (6) If \mathcal{P} is connected, then it admits a unique sincere interval module $\mathbb{I}_{\mathcal{P}}$. (Recall that *sincere* means that $(\mathbb{I}_{\mathcal{P}})_a \neq 0$ for all $a \in \mathcal{P}$.)

2.1. Definitions and first examples.

Definition 2.2. A \mathbb{Z} -invariant (or *invariant* for short) on $\text{mod } \mathcal{P}$ is a \mathbb{Z} -linear map $\Phi : K_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}^I$ for some set I . In other words, Φ can be viewed as a map $\text{mod } \mathcal{P} \rightarrow \mathbb{Z}^I$ which is constant on isoclasses of modules and such that $\Phi(M \oplus N) = \Phi(M) + \Phi(N)$. An invariant Φ is *finite* if its image is finitely generated. We denote by $\text{rk } \Phi$ the rank of its image.

Example 2.3.

- (1) Counting the multiplicities of all indecomposable direct summands yields an invariant $\mathbf{mult}_{\mathcal{P}} : K_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}^{\text{Ind } \mathcal{P}}$ of rank $|\text{Ind } \mathcal{P}|$. It is defined by $\mathbf{mult}_{\mathcal{P}}(M)_U = \alpha_U$ where $M \simeq \bigoplus_{U \in \text{Ind } \mathcal{P}} U^{\alpha_U}$ is the decomposition of M into indecomposable objects. If \mathcal{P} is a poset of finite representation type, then $\mathbf{mult}_{\mathcal{P}}$ is a finite invariant. More generally, if $\mathcal{U} \subseteq \text{Ind } \mathcal{P}$ is a finite family of indecomposable modules, we define $\mathbf{mult}_{\mathcal{P}}^{\mathcal{U}} : K_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}^{\mathcal{U}}$ by composing $\mathbf{mult}_{\mathcal{P}}$ with the natural projection of $\mathbb{Z}^{\mathcal{U}} \rightarrow \mathbb{Z}^{\text{Ind } \mathcal{P}}$.

For instance, if $\mathcal{U} = \{\mathbb{I}_{[a,b]} \mid a \leq b\}$ is the set of all rectangle modules then a module M is called *rectangle-decomposable* if $\mathbf{mult}_{\mathcal{P}}(M)_U = 0$ for $U \notin \mathcal{U}$, or equivalently when $\mathbf{mult}_{\mathcal{P}}(M)$ is obtained from $\mathbf{mult}_{\mathcal{P}}^{\mathcal{U}}(M)$ by the natural inclusion of $\mathbb{Z}^{\mathcal{U}} \rightarrow \mathbb{Z}^{\text{Ind } \mathcal{P}}$.

- (2) For $a \in \mathcal{P}$, $\mathbf{dim}_a : K_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}$ is an invariant of rank 1 given by $\mathbf{dim}_a(M) = \dim_k M_a$. The map $\mathbf{dim}_{\mathcal{P}} := \bigoplus_{a \in \mathcal{P}} \mathbf{dim}_a$ is an invariant of rank $|\mathcal{P}|$. This invariant is often known as the *dimension vector* or *Hilbert function*.
- (3) Denote the set of ordered pairs in \mathcal{P} by

$$\{\leq_{\mathcal{P}}\} := \{(a, b) \in \mathcal{P}^2 \mid a \leq b\}.$$

For $(a, b) \in \{\leq_{\mathcal{P}}\}$, the map $\mathbf{rk}_{a,b} : K_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}$ given by $\mathbf{rk}_{a,b}(M) = \text{rk } M_{a \rightarrow b}$ is an invariant of rank 1. The map $\mathbf{rk}_{\mathcal{P}} := \bigoplus_{(a,b) \in \{\leq_{\mathcal{P}}\}} \mathbf{rk}_{a,b}$ is an invariant of rank $|\{\leq_{\mathcal{P}}\}|$ called the *rank invariant* (see [CZ09, BOO]).

- (4) For any finite family $\mathcal{U} \subseteq \text{Ind } \mathcal{P}$, the map $\mathbf{dimh}_{\mathcal{P}}^{\mathcal{U}} : K_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}^{\mathcal{U}}$ defined by $\left(\mathbf{dimh}_{\mathcal{P}}^{\mathcal{U}}(M)\right)_U := \dim_k \text{Hom}_{\mathcal{P}}(U, M)$ for $U \in \mathcal{U}$ is an invariant of rank $|\mathcal{U}|$. Following [BBH24], we call this kind of invariant a *dim-Hom invariant*.

2.2. Comparing invariants. Of course, a finite invariant can only be complete when the poset \mathcal{P} is representation-finite. One goal of persistence theory is to extract useful information from $M \in \text{mod } \mathcal{P}$ by some finite invariants even when the poset \mathcal{P} is representation-finite. The following definitions provide tools to compare different invariants.

Definition 2.4. Two invariants Φ_1, Φ_2 on $\text{mod } \mathcal{P}$ are *equivalent* (in symbols $\Phi_1 \simeq \Phi_2$) if there exists an isomorphism $\varphi : \text{Im } \Phi_1 \rightarrow \text{Im } \Phi_2$ such that $\varphi \circ \Phi_1 = \Phi_2$.

Definition 2.5. Let Φ_1, Φ_2 be two invariants on $\text{mod } \mathcal{P}$. We say that Φ_1 is *finer* than Φ_2 , and denote it by $\Phi_1 \geq \Phi_2$, if there exists a \mathbb{Z} -linear map $\varphi : \text{Im } \Phi_1 \rightarrow \text{Im } \Phi_2$ such that $\varphi \circ \Phi_1 = \Phi_2$ (in particular, φ is surjective). This defines a partial order on the equivalence classes of invariants on $\text{mod } \mathcal{P}$.

Example 2.6.

- (1) The diagonal inclusion $\mathcal{P} \rightarrow \{\leq_{\mathcal{P}}\}$ induces an injective map $\varphi : \mathbb{Z}^{\mathcal{P}} \rightarrow \mathbb{Z}^{\{\leq_{\mathcal{P}}\}}$ whose transpose $\varphi^T : \mathbb{Z}^{\{\leq_{\mathcal{P}}\}} \rightarrow \mathbb{Z}^{\mathcal{P}}$ is surjective and satisfies $\varphi^T \circ \mathbf{rk}_{\mathcal{P}} = \mathbf{dim}_{\mathcal{P}}$, hence $\mathbf{rk}_{\mathcal{P}} \geq \mathbf{dim}_{\mathcal{P}}$. In general, the rank invariant is strictly finer than the dimension vector, in fact for a connected poset \mathcal{P} the rank invariant is equivalent to the dimension vector if and only if $|\mathcal{P}| = 1$. It is a classical result in representation theory that the rank invariant $\mathbf{rk}_{\mathcal{P}}$ is complete for a totally ordered poset \mathcal{P} .
- (2) Let $\text{Proj } \mathcal{P} \subseteq \text{Ind } \mathcal{P}$ the subset of indecomposable projective modules (see Example 2.1(1)). It is classical that $\mathbf{dim}_{\mathcal{P}} = \mathbf{dimh}_{\mathcal{P}}^{\text{Proj } \mathcal{P}}$, so these invariants are equivalent. See [BBH24, Proposition 5.1].
- (3) Let $\mathcal{H} \subset \text{Ind } \mathcal{P}$ be the subset formed by the hook modules and the projective indecomposable modules (see Example 2.1). It is shown in [BOO] and [BBH24] that the invariants $\mathbf{rk}_{\mathcal{P}}$ and $\mathbf{dimh}_{\mathcal{P}}^{\mathcal{H}}$ are equivalent.

Proposition 2.7. *Let \mathcal{P} be a poset of finite representation type. Then the invariants $\mathbf{mult}_{\mathcal{P}}$ and $\mathbf{dimh}_{\mathcal{P}}^{\text{Ind } \mathcal{P}}$ are equivalent.*

Proof. Denote by $\{U_1, \dots, U_{\ell}\} = \text{Ind } \mathcal{P}$. For $1 \leq i, j \leq \ell$ we have $\mathbf{mult}_{U_i}(U_j) = \delta_{i,j}$ and $\mathbf{dimh}_{U_i}(U_j) = \dim_k \text{Hom}_{\mathcal{P}}(U_i, U_j)$. Hence if we denote by $\Phi \in \text{Mat}_{\ell}(\mathbb{Z})$ the matrix defined by $\Phi_{i,j} := \dim_k \text{Hom}_{\mathcal{P}}(U_i, U_j)$, we have that $\Phi \circ \mathbf{mult}_{\mathcal{P}} = \mathbf{dimh}_{\mathcal{P}}^{\text{Ind } \mathcal{P}}$ showing that $\mathbf{mult}_{\mathcal{P}} \geq \mathbf{dimh}_{\mathcal{P}}^{\text{Ind } \mathcal{P}}$.

Since \mathcal{P} is representation-finite, it is representation directed by [Si10, Theorem 1.4]. This means that the transitive closure of the relation $U_i \leq U_j$ whenever $\text{Hom}(U_i, U_j) \neq 0$ is a partial order on $\text{Ind } \mathcal{P}$. It also implies that every indecomposable module is a brick; that is, that $\text{End}_{\mathcal{P}}(U_i) \cong k$ for all $U_i \in \text{Ind } \mathcal{P}$. We conclude that Φ is an isomorphism using the following lemma with $Y = \text{Ind } \mathcal{P}$ and $\varphi_i(j) = \Phi_{i,j}$. \square

Lemma 2.8. *Let Y be a finite poset. For each $y \in Y$ assume that we have $\varphi_y \in \mathbb{Z}^Y$ satisfying*

- $\varphi_y(y) = 1$
- $\varphi_y(x) \neq 0 \Rightarrow y \geq x$

Then the family $(\varphi_y, y \in Y)$ is a basis of the abelian group \mathbb{Z}^Y .

Proof. The proof is classical, but since we use this result in several places in the paper, we give here a proof for the convenience of the reader. We first show that the family $(\varphi_y)_{y \in Y}$ generates the free abelian group \mathbb{Z}^Y . For $\Psi \in \mathbb{Z}^Y$, denote

$$\mathcal{S}_{\Psi} := \{y \in Y \mid \exists y' \geq y \text{ with } \Psi(y') \neq 0\}.$$

We show by induction on $|\mathcal{S}_{\Psi}|$ that Ψ is a linear combination of the φ_y with $y \in \mathcal{S}_{\Psi}$. If $\mathcal{S}_{\Psi} = \emptyset$ then $\Psi = 0$ and there is nothing to show. Assume now that $\mathcal{S}_{\Psi} \neq \emptyset$ and let $y \in \mathcal{S}_{\Psi}$ be a maximal element, hence $\Psi(y) \neq 0$. Define $\Psi' := \Psi - \Psi(y)\varphi_y$. We will show that $\mathcal{S}_{\Psi'} \subseteq \mathcal{S}_{\Psi}$. Let z be in $\mathcal{S}_{\Psi'}$. Then there exists $y' \geq z$ such that $\Psi'(y') \neq 0$. That is $\Psi(y') \neq \Psi(y)\varphi_y(y')$. If $\Psi(y') \neq 0$ then $z \in \mathcal{S}_{\Psi}$. If $\Psi(y') = 0$, then $\varphi_y(y') \neq 0$ which implies by hypothesis that $y \geq y'$. Hence we have $y \geq z$ with $\Psi(y) \neq 0$, that is $z \in \mathcal{S}_{\Psi}$. Hence we have $\mathcal{S}_{\Psi'} \subseteq \mathcal{S}_{\Psi}$. Moreover $y \in \mathcal{S}_{\Psi}$ but $y \notin \mathcal{S}_{\Psi'}$. Thus we know by induction that Ψ' is a linear combination of the φ_y with $y \in \mathcal{S}_{\Psi'}$, and so $\Psi = \Psi' + \Psi(y)\varphi_y$ is a linear combination of the φ_y with $y \in \mathcal{S}_{\Psi}$.

Let $\sum_{y \in Y} \alpha_y \varphi_y = 0$ be a linear combination. Assume that $\{y \in Y \mid \alpha_y \neq 0\}$ is non empty and let z be a maximal element in this set. Then we obtain $-\alpha_z = \sum_{y \neq z} \alpha_y \varphi_y(z) \neq 0$. Therefore there exists y with $\alpha_y \neq 0$ and $\varphi_y(z) \neq 0$ which implies $z \leq y$ and contradicts the maximality of z . Hence the family is linearly independent. \square

3. INVARIANTS DEFINED BY ORDER-EMBEDDINGS

Definition 3.1. Let \mathcal{X} and \mathcal{P} be finite posets. A map $f : \mathcal{X} \rightarrow \mathcal{P}$ is an *order-embedding* if $x \leq x'$ precisely when $f(x) \leq f(x')$, for any $x, x' \in \mathcal{X}$. Denote by $\text{Emb}_{\mathcal{X}}^{\mathcal{P}}$ the set of all order-embeddings of \mathcal{X} into \mathcal{P} up to automorphism of \mathcal{X} .

Remark 3.2. To avoid redundancy, we chose to consider the set $\text{Emb}_{\mathcal{X}}^{\mathcal{P}}$ of order-embeddings up to automorphism. For instance, if the posets \mathcal{X} and \mathcal{P} are given by

$$\mathcal{X} = \begin{array}{c} & & 2 \\ & \nearrow & \\ 1 & & \\ & \searrow & \\ & & 3 \end{array} \quad \mathcal{P} = \begin{array}{ccccc} & & & b & \\ & & & \nearrow & \\ & & a & & d \\ & & \searrow & & \\ & & c & & \end{array}$$

then $|\text{Emb}_{\mathcal{X}}^{\mathcal{P}}| = 1$, even if there exist two different order-embeddings of \mathcal{X} into \mathcal{P} .

Any $f \in \text{Emb}_{\mathcal{X}}^{\mathcal{P}}$ induces a restriction functor $f^* : \text{mod } \mathcal{P} \rightarrow \text{mod } \mathcal{X}$ which is exact. It is also dense since order-embeddings are injective. Therefore f^* induces a surjective \mathbb{Z} -linear map $K_0^{\text{SP}}(\mathcal{P}) \rightarrow K_0^{\text{SP}}(\mathcal{X})$.

Definition 3.3. Let \mathcal{X} be a poset of finite representation type, \mathcal{P} be a finite poset and, E be a subset of $\text{Emb}_{\mathcal{X}}^{\mathcal{P}}$. We denote by $I := \text{Ind } \mathcal{X}$.

We define $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^E : K_0^{\text{SP}}(\mathcal{P}) \rightarrow \mathbb{Z}^{E \times I}$ by

$$\left(\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^E(M) \right)_{(f, U)} := \text{mult}_{\mathcal{X}}(U, f^*M) \quad \text{for } (f, U) \in E \times I.$$

We define $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^E : K_0^{\text{SP}}(\mathcal{P}) \rightarrow \mathbb{Z}^{E \times I}$ by

$$\left(\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^E(M) \right)_{(f, U)} := \dim \text{Hom}_{\mathcal{X}}(U, f^*M) \quad \text{for } (f, U) \in E \times I.$$

When $E = \text{Emb}_{\mathcal{X}}^{\mathcal{P}}$ we write $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$ and $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}$ instead of $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^E$ and $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^E$.

Remark 3.4. See Remark 4.9 for a discussion of how to realize the invariant $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^E$ as a dim-Hom invariant (as defined in Example 2.3(4)).

We show in Proposition 3.6 that the invariants $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^E$ and $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^E$ are equivalent. The aim of this paper is to study these (equivalent) invariants. We first review a pair of examples which have appeared recently in the literature.

Example 3.5. (1) Consider the poset

$$\mathcal{X} = \begin{array}{ccccc} & & & b & \\ & & & \nearrow & \\ a & & & & d \\ & & & \searrow & \\ & & & c & \end{array}$$

This poset has 10 indecomposable modules, all of which are interval modules. Of these, there are two which are not rectangle modules, namely $\mathbb{I}_{\{a, b, c\}}$ and $\mathbb{I}_{\{b, c, d\}}$. For \mathcal{P} a product of two totally ordered sets, it is shown in [BLO22] that a \mathcal{P} -module M is rectangle decomposable if and only if it satisfies

$$\left(\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^{\text{Emb}_{\mathcal{X}}^{\mathcal{P}}}(M) \right)_{(f, U)} = 0 \quad \text{for all } (f, U) \in \text{Emb}_{\mathcal{X}}^{\mathcal{P}} \times \{\mathbb{I}_{\{a, b, c\}}, \mathbb{I}_{\{b, c, d\}}\}.$$

Note that the authors of [BLO22] do not assume \mathcal{P} to be finite.

- (2) The ‘‘generalized rank invariant’’ of [KM21] and the ‘‘compressed multiplicities’’ of [AENY23b] both record the multiplicity of the sincere interval modules $\mathbb{I}_{\mathcal{X}}$ over certain subsets $\mathcal{X} \subseteq \mathcal{P}$. (The authors of [AENY23b] assume that \mathcal{P} is a product of two finite totally ordered sets and the authors of [KM21] work over an arbitrary locally finite poset.) See [BBH24, Section 3.2] for a summary of these and related results. In Section 5, we will similarly examine situations where we allow the poset \mathcal{X} to vary and record only the multiplicity of the sincere interval modules.

Proposition 3.6. *Let \mathcal{X} be a representation-finite poset and \mathcal{P} be a finite poset. Then for any $E \subseteq \text{Emb}_{\mathcal{X}}^{\mathcal{P}}$ the invariants $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^E$ and $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^E$ are equivalent.*

Proof. By Proposition 2.7, there is an invertible map $\Phi : \mathbb{Z}^I \rightarrow \mathbb{Z}^I$ such that $\Phi \circ \mathbf{mult}_{\mathcal{X}} = \mathbf{dimh}_{\mathcal{X}}^I$. We then obtain the following commutative diagram

$$\begin{array}{ccc} K_0^{\text{SP}}(\mathcal{P}) & \xrightarrow{\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^E} & \mathbb{Z}^{E \times I} \\ \parallel & & \downarrow \Phi^E \\ K_0^{\text{SP}}(\mathcal{P}) & \xrightarrow{\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^E} & \mathbb{Z}^{E \times I} \end{array}$$

where Φ^E is an isomorphism. □

Example 3.7. (1) Let \mathcal{X}_1 be the poset with one element. Then $\text{Emb}_{\mathcal{X}_1}^{\mathcal{P}}$ is naturally in bijection with \mathcal{P} , and $I = \text{Ind } \mathcal{X}_1$ has one element. For $a \in \mathcal{P}$, one immediately checks that $\mathbf{mult}_{\mathcal{X}_1, \mathcal{P}} = \mathbf{dim}_{\mathcal{P}} = \mathbf{dimh}_{\mathcal{X}_1, \mathcal{P}}$.

- (2) Let \mathcal{X}_2 be the poset $\{1 \rightarrow 2\}$. It is well-known that the poset \mathcal{X}_2 is of representation-finite type, with $\mathbb{I} = \text{Ind } \mathcal{X}_2 = \{P_1 = \mathbb{I}_{\{1,2\}}, P_2 = \mathbb{I}_{\{2\}}, H_{1,2} = \mathbb{I}_{\{1\}}\}$.

Let $f : \mathcal{X}_2 \rightarrow \mathcal{P}$ be an order-embedding. Then one easily checks that for any $M \in \text{mod } \mathcal{P}$

$$\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}(M)_{(f,U)} = \begin{cases} \dim M_{f_1} - \text{rk } M_{f_1 \rightarrow f_2} & \text{if } U = \mathbb{I}_{\{1\}} \\ \text{rk } M_{f_1 \rightarrow f_2} & \text{if } U = \mathbb{I}_{\{1,2\}} \\ \dim M_{f_2} - \text{rk } M_{f_1 \rightarrow f_2} & \text{if } U = \mathbb{I}_{\{2\}} \end{cases}$$

And we have

$$\mathbf{dimh}_{\mathcal{X}_2, \mathcal{P}}(M)_{(f,U)} = \begin{cases} \dim M_{f_1} - \text{rk } M_{f_1 \rightarrow f_2} & \text{if } U = \mathbb{I}_{\{1\}} \\ \dim M_{f_1} & \text{if } U = \mathbb{I}_{\{1,2\}} \\ \dim M_{f_2} & \text{if } U = \mathbb{I}_{\{2\}} \end{cases}$$

The isomorphism Φ which yields the equivalence between $\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}^f$ and $\mathbf{dimh}_{\mathcal{X}_2, \mathcal{P}}^f$ is triangular with diagonal entries 1 as constructed in Lemma 2.8:

$$\Phi = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

From this example we deduce the following

Corollary 3.8. *Let \mathcal{P} be a finite connected poset with at least 2 elements. Then the invariants $\mathbf{rk}_{\mathcal{P}}$ and $\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}$ are equivalent.*

Proof. Let $\mathbb{E} = \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}}$ and consider the map $p : \mathbb{E} \times \mathbb{I} \rightarrow \{\leq_{\mathcal{P}}\}$ defined by

$$p(f, U) = \begin{cases} (f_1, f_2) & \text{if } U = \mathbb{I}_{\{1,2\}} \\ (f_1, f_1) & \text{if } U = \mathbb{I}_{\{1\}} \\ (f_2, f_2) & \text{if } U = \mathbb{I}_{\{2\}} \end{cases}$$

Since the poset \mathcal{P} is connected and has at least two elements, this map is surjective and induces an injective morphism $\mathbb{Z}^{\{\leq_{\mathcal{P}}\}} \rightarrow \mathbb{Z}^{\mathbb{E} \times \mathbb{I}}$. Let $\Phi : \mathbb{Z}^{\mathbb{E} \times \mathbb{I}} \rightarrow \mathbb{Z}^{\{\leq_{\mathcal{P}}\}}$ defined by

$$\Phi_f := \begin{pmatrix} 0 & 1 & -1 \\ 0 & 0 & 1 \\ 1 & 0 & -1 \end{pmatrix} \text{ for any } f \in \mathbb{E}.$$

Then we have $\Phi \circ p^* \circ \mathbf{rk}_{\mathcal{P}} = \mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}$. Moreover since $\mathbf{rk}_{\mathcal{P}}$ is surjective, and since $\Phi \circ p^*$ is injective, we obtain that $\Phi \circ p^*$ is an isomorphism from $\mathbb{Z}^{\{\leq_{\mathcal{P}}\}}$ to the image of $\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}$. \square

Remark 3.9. Note that the set $\mathbb{E} = \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}}$ has cardinality $3 \cdot |\{\leq_{\mathcal{P}}\}|$, where $\{\leq_{\mathcal{P}}\} := \{(a, b) \in \mathcal{P}^2 \mid a < b\}$. This is in general strictly greater than $|\{\leq_{\mathcal{P}}\}| = |\{\leq_{\mathcal{P}}\}| + |\mathcal{P}|$, so the invariant $\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}$ is not surjective. We will see in Section 5 a way to reduce this redundant information.

Proposition 3.10. *Let \mathcal{P} be a connected poset such that every element lies in a chain of length ≥ 3 . Let \mathcal{X}_3 be the poset $\{1 \rightarrow 2 \rightarrow 3\}$. Then the invariants $\mathbf{rk}_{\mathcal{P}}$ and $\mathbf{mult}_{\mathcal{X}_3, \mathcal{P}}$ are equivalent.*

Proof. There are 6 indecomposable objects in $\text{mod } \mathcal{X}_3$ which are

$$\mathbb{I} = \left\{ H_{1,2} = \mathbb{I}_{\{1\}}, H_{2,3} = \mathbb{I}_{\{2\}}, P_3 = \mathbb{I}_{\{3\}}, H_{1,3} = \mathbb{I}_{\{1,2\}}, P_2 = \mathbb{I}_{\{2,3\}}, P_3 = \mathbb{I}_{\{1,2,3\}} \right\}.$$

Then one computes

$$\left(\mathbf{mult}_{\mathcal{X}_3, \mathcal{P}}(M) \right)_{(f,U)} = \begin{cases} \dim M_{f_1} - \text{rk } M_{f_1 \rightarrow f_2} & \text{if } U = \mathbb{I}_{\{1\}} \\ \dim M_{f_2} - \text{rk } M_{f_1 \rightarrow f_2} - \text{rk } M_{f_2 \rightarrow f_3} + \text{rk } M_{f_1 \rightarrow f_3} & \text{if } U = \mathbb{I}_{\{2\}} \\ \dim M_{f_3} - \text{rk } M_{f_2 \rightarrow f_3} & \text{if } U = \mathbb{I}_{\{3\}} \\ \text{rk } M_{f_1 \rightarrow f_2} - \text{rk } M_{f_1 \rightarrow f_3} & \text{if } U = \mathbb{I}_{\{1,2\}} \\ \text{rk } M_{f_2 \rightarrow f_3} - \text{rk } M_{f_1 \rightarrow f_3} & \text{if } U = \mathbb{I}_{\{2,3\}} \\ \text{rk } M_{f_1 \rightarrow f_3} & \text{if } U = \mathbb{I}_{\{1,2,3\}} \end{cases}$$

Now for $a < b$ in \mathcal{P} , by hypothesis there exists $c \in \mathcal{P}$ with either $c < a$ or with $b < c$. Therefore, there exists $f \in \text{Emb}_{\mathcal{X}_3}^{\mathcal{P}}$ such that $f(1) = a$ and $f(2) = b$, or such that $f(2) = a$ and $f(3) = b$. Using this embedding f , one can easily check the equivalence. \square

Remark 3.11. Note that here again, in general, the cardinality of $\text{Emb}_{\mathcal{X}_3}^{\mathcal{P}} \times \mathbb{I}$ is strictly greater than the cardinality of $\{\leq_{\mathcal{P}}\}$. Therefore the invariant $\mathbf{mult}_{\mathcal{X}_3, \mathcal{P}}$ is not surjective. For example, if \mathcal{P} is as follows

$$\mathcal{P} = \begin{array}{c} & & c \\ & \nearrow & \\ a & \longrightarrow & b \\ & \searrow & \\ & & d \end{array}$$

one has $|\text{Emb}_{\mathcal{X}_3}^{\mathcal{P}}| = 2$, so $|\text{Emb}_{\mathcal{X}_3}^{\mathcal{P}} \times \mathbb{I}| = 12$. However the rank of the invariant is $9 = |\{\leq_{\mathcal{P}}\}|$. One can check the following 3 relations

$$\begin{aligned} \mathbf{mult}(\mathbb{I}_{\{b,c,d\}}) - \mathbf{mult}(\mathbb{I}_{\{b,c\}}) - \mathbf{mult}(\mathbb{I}_{\{b,d\}}) + \mathbf{mult}(\mathbb{I}_{\{b\}}) &= 0; \\ \mathbf{mult}(\mathbb{I}_{\{a,b,c,d\}}) - \mathbf{mult}(\mathbb{I}_{\{a,b,c\}}) - \mathbf{mult}(\mathbb{I}_{\{a,b,d\}}) + \mathbf{mult}(\mathbb{I}_{\{a,b\}}) &= 0; \\ \mathbf{mult}(M) - \mathbf{mult}(\mathbb{I}_{\{a,b\}}) - \mathbf{mult}(\mathbb{I}_{\{b,c\}}) - \mathbf{mult}(\mathbb{I}_{\{b,d\}}) + \mathbf{mult}(\mathbb{I}_{\{b\}}) &= 0, \end{aligned}$$

where M denotes the indecomposable \mathcal{P} -module with dimension vector $(1, 2, 1, 1)$.

Proposition 3.12. Let \mathcal{X}'_3 and \mathcal{X}''_3 be the following posets

$$\mathcal{X}'_3 = \begin{array}{c} & 2 \\ & \nearrow \\ 1 & \\ & \searrow \\ & 3 \end{array} \quad \mathcal{X}''_3 = \begin{array}{c} 1 & \searrow \\ & 3 \\ 2 & \nearrow \end{array}$$

Let \mathcal{P} be a connected poset such that any pair $a < b \in \mathcal{P}$ is in the image of an order-embedding $\mathcal{X}'_3 \rightarrow \mathcal{P}$ (resp. of an order embedding $\mathcal{X}''_3 \rightarrow \mathcal{P}$), then the invariant $\mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}$ (resp. $\mathbf{mult}_{\mathcal{X}''_3, \mathcal{P}}$) is strictly finer than the rank invariant $\mathbf{rk}_{\mathcal{P}}$.

Proof. The indecomposable \mathcal{X}'_3 -modules are

$$\text{Ind } \mathcal{X}'_3 = \left\{ S_1 = \mathbb{I}_{\{1\}}, P_2 = \mathbb{I}_{\{2\}}, P_3 = \mathbb{I}_{\{3\}}, H_{1,3} = \mathbb{I}_{\{1,2\}}, H_{1,2} = \mathbb{I}_{\{1,3\}}, P_1 = \mathbb{I}_{\{1,2,3\}} \right\}.$$

Now for $f \in \text{Emb}_{\mathcal{X}'_3}^{\mathcal{P}}$, denote by $\text{rk } M_{f_1 \rightarrow f_2 \oplus f_3}$ the rank of the map $M_{f_1} \rightarrow M_{f_2} \oplus M_{f_3}$ induced by $M_{f_1 \rightarrow f_2}$ and $M_{f_1 \rightarrow f_3}$.

Then one computes that

$$\left(\mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}(M) \right)_{(f,U)} = \begin{cases} \dim M_{f_1} - \text{rk } M_{f_1 \rightarrow f_2} - \text{rk } M_{f_1 \rightarrow f_3} + \text{rk } M_{f_1 \rightarrow f_2 \oplus f_3} & \text{if } U = \mathbb{I}_{\{1\}} \\ \dim M_{f_2} - \text{rk } M_{f_1 \rightarrow f_2} & \text{if } U = \mathbb{I}_{\{2\}} \\ \dim M_{f_3} - \text{rk } M_{f_1 \rightarrow f_3} & \text{if } U = \mathbb{I}_{\{3\}} \\ \text{rk } M_{f_1, f_2} - \text{rk } M_{f_1 \rightarrow f_2 \oplus f_3} & \text{if } U = \mathbb{I}_{\{1,2\}} \\ \text{rk } M_{f_1 \rightarrow f_3} - \text{rk } M_{f_1 \rightarrow f_2 \oplus f_3} & \text{if } U = \mathbb{I}_{\{1,3\}} \\ \text{rk } M_{f_1 \rightarrow f_2 \oplus f_3} & \text{if } U = \mathbb{I}_{\{1,2,3\}} \end{cases}$$

Let $a < b$ be in \mathcal{P} . Then by hypothesis, there exists an embedding $f : \mathcal{X}'_3 \rightarrow \mathcal{P}$ with $f(1) = a$ and $f(2) = b$. Then we have

$$\left(\mathbf{rk}_{\mathcal{P}}(M) \right)_{(a,b)} = \left(\mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}(M) \right)_{(f, \mathbb{I}_{\{1,2\}})} - \left(\mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}(M) \right)_{(f, \mathbb{I}_{\{1,2,3\}})}.$$

For $a \in \mathcal{P}$, by convexity, there exists $f \in \text{Emb}_{\mathcal{X}'_3}^{\mathcal{P}}$ with either $f(1) = a$ or with $f(2) = a$.

If $f(1) = a$, we check that $\left(\mathbf{rk}_{\mathcal{P}}(M) \right)_{(a,a)}$ is in the free abelian group generated by

$$\left\{ \delta_{(f, \mathbb{I}_{\{1\}})}, \delta_{(f, \mathbb{I}_{\{1,2\}})}, \delta_{(f, \mathbb{I}_{\{1,3\}})}, \delta_{(f, \mathbb{I}_{\{1,2,3\}})} \right\},$$

where $\left\{ \delta_{(f,U)} \mid (f,U) \in \mathbb{E} \times \mathbb{I} \right\}$ denotes the canonical dual basis in $\mathbb{Z}^{\mathbb{E} \times \mathbb{I}}$.

If $f(2) = a$, one can also check that $\left(\mathbf{rk}_{\mathcal{P}}(M) \right)_{(a,a)}$ is in the free abelian group generated by

$$\left\{ \delta_{(f, \mathbb{I}_{\{2\}})}, \delta_{(f, \mathbb{I}_{\{1,2\}})}, \delta_{(f, \mathbb{I}_{\{1,2,3\}})} \right\}.$$

Therefore we have $\mathbf{rk}_{\mathcal{P}} \leq \mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}$.

To prove that the invariant $\mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}$ is strictly finer than $\mathbf{rk}_{\mathcal{P}}$ consider the case $\mathcal{P} = \mathcal{X}'_3$. The rank of $\mathbf{rk}_{\mathcal{P}}$ is the cardinal of $\{\leq_{\mathcal{P}}\}$ which is 5, while the rank of $\mathbf{mult}_{\mathcal{X}'_3}$ is 6 since there are 6 indecomposable modules in $\text{mod } \mathcal{X}'_3$.

The proof is completely similar for \mathcal{X}''_3 . □

Remark 3.13. (1) Note that here again, the cardinality of $\text{Emb}_{\mathcal{X}'_3}^{\mathcal{P}}$ is strictly greater than the rank of the invariant $\mathbf{mult} := \mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}$. For instance if \mathcal{P} is the poset

$$\mathcal{P} = \begin{array}{c} & & c \\ & & \swarrow \\ a & \longrightarrow & b \\ & & \searrow \\ & & d \end{array}$$

then $|\text{Emb}_{\mathcal{X}'_3}^{\mathcal{P}}| = 2$ so $|\text{Emb}_{\mathcal{X}'_3}^{\mathcal{P}} \times \mathbb{I}| = 12$, but one can check that the rank of \mathbf{mult} is 10. We have the relations

$$\mathbf{mult}(\mathbb{I}_{\{a\}}) + \mathbf{mult}(\mathbb{I}_{\{b\}}) = \mathbf{mult}(\mathbb{I}_{\{a,b\}}),$$

$$\mathbf{mult}(M) - \mathbf{mult}(\mathbb{I}_{\{a,b,c,d\}}) + \mathbf{mult}(\mathbb{I}_{\{b,c,d\}}) - \mathbf{mult}(\mathbb{I}_{\{b,c\}}) - \mathbf{mult}(\mathbb{I}_{\{b,d\}}) = 0,$$

where M is the indecomposable \mathcal{P} module with dimension vector $(1, 2, 1, 1)$.

(2) The hypothesis on \mathcal{P} in Proposition 3.12 implies the weaker property that

$$\bigcup_{f \in \text{Emb}_{\mathcal{X}'_3}^{\mathcal{P}}} \text{Im } f = \mathcal{P}.$$

We note that this weaker property is not sufficient in order to deduce the fact that $\mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}$ is strictly finer than $\mathbf{rk}_{\mathcal{P}}$. Indeed consider the case where \mathcal{P} is given as above. This poset satisfies the weaker property above, but there is no embedding $f : \mathcal{X}'_3 \rightarrow \mathcal{P}$ with $f(1) = a$ and $f(2) = b$. As mentioned above, the family $\{\mathbf{mult}(\mathbb{I}_{\{a\}}), \mathbf{mult}(\mathbb{I}_{\{b\}}), \mathbf{mult}(\mathbb{I}_{\{a,b\}})\}$ is not free, but one easily checks that the family $\{\mathbf{rk}_{\mathcal{P}}(\mathbb{I}_{\{a\}}), \mathbf{rk}_{\mathcal{P}}(\mathbb{I}_{\{b\}}), \mathbf{rk}_{\mathcal{P}}(\mathbb{I}_{\{a,b\}})\}$ is free. Thus the invariants $\mathbf{mult}_{\mathcal{X}'_3, \mathcal{P}}$ and $\mathbf{rk}_{\mathcal{P}}$ cannot be compared in this case. Similar examples occur when \mathcal{X}'_3 is replaced with \mathcal{X}_3'' .

(3) The condition on \mathcal{P} in Proposition 3.12 may seem strong. For instance, it is not satisfied for \mathcal{P} being an $n \times m$ grid. However, one can enlarge the poset \mathcal{P} in order to get the condition, and then restrict to modules with support in \mathcal{P} .

Example 3.14. Consider the poset \mathcal{X}_4 of type D_4

$$\mathcal{X}_4 = \begin{array}{c} & & c \\ & & \swarrow \\ a & \longrightarrow & b \\ & & \searrow \\ & & d \end{array}$$

Then \mathcal{X}_4 has two sincere indecomposable representations, one being the characteristic representation $\mathbb{I}_{\mathcal{X}_4}$, the other one is given by three one-dimensional subspaces in general position, placed at vertices a, c, d , of a two-dimensional vector space at vertex b . Still, their multiplicities can be described in terms of the linear maps of the \mathcal{P} -module M restricted to \mathcal{X}_4 , for example the multiplicity of $\mathbb{I}_{\mathcal{X}_4}$ is given by

$$\mathbf{mult}_{\mathcal{X}_4, \mathcal{P}}(\mathbb{I}_{\mathcal{X}_4}) = \dim(\text{Im } M_{f_a \rightarrow f_b} \cap \text{Im } M_{f_c \rightarrow f_b} \cap \text{Im } M_{f_d \rightarrow f_b}).$$

4. INVARIANTS AND EXACT STRUCTURES

4.1. Exact structures and relative projectives. Exact structures have been introduced in [Qu72] as an axiomatic framework allowing to use methods from homological algebra relative to a fixed class of short exact sequences. Besides this axiomatic approach, there are a number of ways to describe an exact structure \mathcal{E} , such as by certain subfunctors \mathcal{E} of the bifunctor $\text{Ext}_{\mathcal{P}}^1$, or by specifying what are the projective objects relative to \mathcal{E} , or by the Auslander-Reiten sequences belonging to \mathcal{E} .

Let \mathcal{E} be a class of short exact sequences in $\text{mod } \mathcal{P}$. We assume that \mathcal{E} is closed under isomorphisms. If a short exact sequence $\eta = (0 \rightarrow A \xrightarrow{f} E \xrightarrow{g} B \rightarrow 0)$ belongs to \mathcal{E} , we say that η is an (\mathcal{E}) -admissible short exact sequence, and also that f is an (\mathcal{E}) -admissible monomorphism and g is an (\mathcal{E}) -admissible epimorphism. The closure under isomorphisms then implies that the classes of admissible short exact sequences, admissible monomorphisms, and admissible epimorphisms uniquely determine one another. Quillen's definition can then be rephrased as follows:

Definition 4.1. A class \mathcal{E} of short exact sequences in $\text{mod } \mathcal{P}$ is said to be an *exact structure* on $\text{mod } \mathcal{P}$ if all of the following hold:

- (E0) \mathcal{E} contains all split exact sequences.
- (E1) \mathcal{E} is closed under compositions of admissible monomorphisms, that is, if $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are admissible monomorphisms, then $g \circ f : X \rightarrow Z$ is also an admissible monomorphism. Likewise, \mathcal{E} is closed under compositions of admissible epimorphisms.
- (E2) \mathcal{E} is closed under pushouts: If $f : X \rightarrow Y$ is an admissible monomorphism, and $X \xrightarrow{h} W$ is any morphism in \mathcal{A} , then the pushout of f along h yields a short exact sequence in \mathcal{E} . Likewise, \mathcal{E} is closed under pullbacks.

The two extreme examples of exact structures are the *split exact structure* \mathcal{E}^{sp} , which contains only the split short exact sequences and corresponds to the subfunctor 0 of $\text{Ext}_{\mathcal{P}}^1$, and the exact structure \mathcal{E}^{all} , which contains all short exact sequences and corresponds to $\text{Ext}_{\mathcal{P}}^1$ as a subfunctor of itself.

Proposition 4.2. *Let \mathcal{P} and \mathcal{X} be finite posets. Let \mathcal{E} be an exact structure on $\text{mod } \mathcal{X}$ and let $E \subseteq \text{Emb}_{\mathcal{X}}^{\mathcal{P}}$. Let $\Psi_E(\mathcal{E})$ denote the class of short exact sequences*

$$0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$$

in $\text{mod } \mathcal{P}$ which have the property that

$$0 \rightarrow f^*L \rightarrow f^*M \rightarrow f^*N \rightarrow 0$$

is \mathcal{E} -exact for all $f \in E$. Then $\Psi_E(\mathcal{E})$ is an exact structure on $\text{mod } \mathcal{P}$.

Proof. We verify the three points from definition 4.1 for all $f \in E$, using that the restriction functor $f^* : \text{mod } \mathcal{P} \rightarrow \text{mod } \mathcal{X}$ is additive and exact.

- (E0) A split exact sequence is sent by f^* to a split exact sequence, which belongs necessarily to \mathcal{E} , therefore $\Psi(\mathcal{E})$ contains all split exact sequences.
- (E1) This is clear from the definition of $\Psi(\mathcal{E})$ and the fact that (E1) holds for \mathcal{E} .
- (E2) Follows from the fact that (E2) holds for \mathcal{E} and that f^* preserves pushouts since it is an exact functor. \square

Definition 4.3. Let \mathcal{E} be an exact structure on $\text{mod } \mathcal{P}$. We denote by $K_0(\mathcal{E})$ the quotient of $K_0^{\text{sp}}(\mathcal{P})$ by the relations

$$[L] + [N] - [M] \text{ for any short exact sequence } (0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0) \text{ in } \mathcal{E}.$$

The following is then immediate.

Proposition 4.4. *Let \mathcal{X} be a representation-finite poset and \mathcal{P} be a finite poset. Recall that $I = \text{Ind } \mathcal{X}$ and fix $E \subseteq \text{Emb}_{\mathcal{X}}^{\mathcal{P}}$. Then denote by $\mathcal{E}_{\mathcal{P}}^E := \Psi_E(\mathcal{E}_{\mathcal{X}}^{\text{sp}})$, where $\mathcal{E}_{\mathcal{X}}^{\text{sp}}$ is the split exact structure on $\text{mod } \mathcal{X}$. Then we have a factorization*

$$\begin{array}{ccc}
 K_0^{\text{sp}}(\mathcal{P}) & \xrightarrow{\text{mult}_{\mathcal{X}, \mathcal{P}}^E} & \mathbb{Z}^{E \times I} \\
 \parallel & \searrow & \downarrow \sim \\
 & & K_0(\mathcal{E}_{\mathcal{P}}^E) \\
 & \swarrow & \downarrow \\
 K_0^{\text{sp}}(\mathcal{P}) & \xrightarrow{\text{dimh}_{\mathcal{X}, \mathcal{P}}^E} & \mathbb{Z}^{E \times I}
 \end{array}$$

4.2. Relative projectives. The projective objects (relative to \mathcal{E}) are then described as follows:

Definition 4.5. Let \mathcal{E} be an exact structure on $\text{mod } \mathcal{P}$.

- (1) We say an object $P \in \text{mod } \mathcal{P}$ is \mathcal{E} -projective, in symbols $P \in \text{Proj}(\mathcal{E})$, if $g_* = \text{Hom}_{\mathcal{P}}(P, g)$ is an epimorphism for all $Y, Z \in \text{mod } \mathcal{P}$ and for all admissible epimorphisms $g : Y \rightarrow Z$. Equivalently, $P \in \text{Proj}(\mathcal{E})$ if and only if $\text{Hom}_{\mathcal{A}}(P, -)$ sends admissible exact sequences to exact sequences.
- (2) We say an object $I \in \text{mod } \mathcal{P}$ is \mathcal{E} -injective, in symbols $I \in \text{inj}(\mathcal{E})$, if $f_* = \text{Hom}_{\mathcal{P}}(f, I)$ is an epimorphism for all $X, Y \in \text{mod } \mathcal{P}$ and for all admissible monomorphisms $f : X \rightarrow Y$. Equivalently, $I \in \text{inj}(\mathcal{E})$ if and only if $\text{Hom}_{\mathcal{P}}(-, I)$ sends admissible exact sequences to exact sequences.

The following is a classical result of [AS93, DRSS99]. See [BBH24, Section 4] for a summary tailored towards persistence theory.

Proposition 4.6. *Let $\mathcal{U} \subseteq \text{Ind } \mathcal{P}$. Denote by*

$$\mathcal{E}_{\mathcal{U}} := \left\{ (0 \rightarrow M \rightarrow N \xrightarrow{g} L \rightarrow 0) \mid \forall U \in \mathcal{U}, \text{Hom}_{\mathcal{P}}(U, g) \text{ is surjective.} \right\}$$

Then $\mathcal{E}_{\mathcal{U}}$ is an exact structure on $\text{mod } \mathcal{P}$, and we have

$$\text{Proj}(\mathcal{E}_{\mathcal{U}}) = \mathcal{U} \cup \text{Proj}(\mathcal{P}).$$

The following is classical, see [ASS06, Theorem I.6.8].

Proposition 4.7. *Let $f : \mathcal{X} \rightarrow \mathcal{P}$ be an order embedding. Then the restriction functor $f^* : \text{mod } \mathcal{P} \rightarrow \text{mod } \mathcal{X}$ is exact and has a left adjoint $f_! : \text{mod } \mathcal{X} \rightarrow \text{mod } \mathcal{P}$ and a right adjoint $f_* : \text{mod } \mathcal{X} \rightarrow \text{mod } \mathcal{P}$. Moreover we have $f^* f_! = \text{Id}_{\text{mod } \mathcal{X}} = f_* f^*$.*

The functors f^* and $f_!$ are sometimes referred to as the *induction* and *coinduction* functors, respectively. The induction functor preserves projective presentations and the coinduction functor preserves injective copresentations.

Proposition 4.8. *Let \mathcal{X} be a representation-finite poset and \mathcal{P} be a finite poset. Recall that $\mathbf{I} = \text{Ind } \mathcal{X}$ and fix \mathbf{E} a subset of $\text{Emb}_{\mathcal{X}}^{\mathcal{P}}$. Assume that $\mathcal{P} = \bigcup_{f \in \mathbf{E}} \text{Im } f$. Define*

$$\mathcal{H}^{\mathbf{E}} := \{f_! U \mid (f, U) \in \mathbf{E} \times \mathbf{I}\}.$$

Then the set $\mathcal{H}^{\mathbf{E}}$ is a subset of $\text{Ind } \mathcal{P}$ that contains the indecomposable projective modules. And we have

$$\mathcal{E}_{\mathcal{P}}^{\mathbf{E}} = \mathcal{E}_{\mathcal{H}^{\mathbf{E}}} \text{ and } \text{Proj}(\mathcal{E}_{\mathcal{P}}^{\mathbf{E}}) = \mathcal{H}^{\mathbf{E}}.$$

Proof. First note that since $\mathcal{P} = \bigcup_{f \in \mathbf{E}} \text{Im } f$, for any $p \in \mathcal{P}$ there exists $x \in \mathcal{X}$ such that $f(x) = p$. Since the functor $f_!$ sends projectives to projectives, we have that $f_!(P_x) = P_p$, that is the projective P_p is in $\mathcal{H}^{\mathbf{E}}$. Moreover by the adjunction formula we have for any $U \in \text{Ind } \mathcal{X}$

$$\text{End}_{\mathcal{P}}(f_! U) \simeq \text{Hom}_{\mathcal{X}}(U, f^* f_! U) = \text{End}_{\mathcal{X}}(U)$$

which is local since $U \in \text{Ind } \mathcal{X}$. Note that since \mathcal{X} is representation-finite, the module U and hence the module $f_! U$ is even a brick.

Let

$$\epsilon = (0 \longrightarrow L \longrightarrow M \longrightarrow N \longrightarrow 0)$$

be a short exact sequence of $\text{mod } \mathcal{P}$. Then by definition ϵ belongs to $\mathcal{E}_{\mathcal{P}}^{\mathbf{E}}$ if and only if for any $f \in \mathbf{E}$ the short exact sequence in $\text{mod } \mathcal{X}$

$$0 \longrightarrow f^* L \longrightarrow f^* M \longrightarrow f^* N \longrightarrow 0$$

splits. This is equivalent to the fact that for any $U \in \text{Ind } \mathcal{X}$ this exact sequence yields a short exact sequence in $\text{mod } k$

$$0 \longrightarrow \text{Hom}_{\mathcal{X}}(U, f^* L) \longrightarrow \text{Hom}_{\mathcal{X}}(U, f^* M) \longrightarrow \text{Hom}_{\mathcal{X}}(U, f^* N) \longrightarrow 0.$$

By adjunction it is equivalent that for any $(f, U) \in \mathbf{E} \times \mathbf{I}$ the short exact sequence ϵ induces an exact sequence

$$0 \longrightarrow \text{Hom}_{\mathcal{P}}(f_! U, L) \longrightarrow \text{Hom}_{\mathcal{P}}(f_! U, M) \longrightarrow \text{Hom}_{\mathcal{P}}(f_! U, N) \longrightarrow 0,$$

which is equivalent to the fact that $\epsilon \in \mathcal{E}_{\mathcal{H}^{\mathbf{E}}}$. The fact that $\text{Proj}(\mathcal{E}_{\mathcal{H}^{\mathbf{E}}}) = \mathcal{H}^{\mathbf{E}}$ is now a direct consequence of Proposition 4.6. \square

Remark 4.9. Consider the setup of Proposition 4.8. It is an immediate consequence of the adjunction formula that the invariants $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}$ and $\mathbf{dimh}_{\mathcal{P}}^{\mathcal{H}^{\mathbf{E}}}$ are equivalent. In particular, $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}$ (and thus also $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}$ by Proposition 3.6) is a dim-Hom invariant in the sense of Example 2.3(4). In order to prove the stronger property these invariants are *homological* (in the sense of [BBH24, Definition 4.12]), one we need an argument showing that the exact structure $\mathcal{E}_{\mathcal{P}}^{\mathbf{E}} = \mathcal{E}_{\mathcal{H}^{\mathbf{E}}}$ has finite global dimension.

From this proposition we can deduce the following commutative diagram.

$$\begin{array}{ccc}
 K_0^{\text{SP}}(\mathcal{P}) & \xrightarrow{\text{mult}_{\mathcal{X}, \mathcal{P}}^E} & \mathbb{Z}^{E \times I} \\
 \parallel & \searrow & \uparrow \\
 & K_0(\mathcal{E}_{\mathcal{P}}^E) & \xrightarrow{\quad} \mathbb{Z}^{E \times I / \sim_1} \\
 & \nearrow & \downarrow \Phi \\
 K_0^{\text{SP}}(\mathcal{P}) & \xrightarrow{\text{dimh}_{\mathcal{X}, \mathcal{P}}^E} & \mathbb{Z}^{E \times I}
 \end{array}$$

where the equivalence relation \sim_1 on $E \times I$ is defined by

$$(f, U) \sim_1 (g, V) \iff f!U \simeq g!V.$$

Note that, *a priori*, it is not clear that there exists an equivalence relation \sim on $E \times I$ such that Φ induces an isomorphism $\mathbb{Z}^{E \times I / \sim_1} \rightarrow \mathbb{Z}^{E \times I / \sim}$.

Example 4.10. (1) Consider $\mathcal{X} = \mathcal{X}_2$ as in Example 3.7(2). Then the indecomposable \mathcal{X}_2 -modules are $P_1 = \mathbb{I}_{\{1,2\}}$, $P_2 = \mathbb{I}_{\{2\}}$ and $H_{1,2} = \mathbb{I}_{\{1\}}$. The first two modules are projective, and the second is a hook (and is the cokernel of the natural map $P_2 \rightarrow P_1$). Since $f_!$ is right exact and sends projectives to projectives, one obtains

$$\bigcup_{f \in \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}}} \{f!U \mid U \in \text{mod } \mathcal{X}_2\} = \{P_{f_1}, P_{f_2}H_{f_1, f_2} \mid f \in \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}}\}.$$


This recovers the result in [BOO].

(2) Consider $\mathcal{X} = \mathcal{X}'_3$ as in Proposition 3.12. Then for $f \in \text{Emb}_{\mathcal{X}'_3}^{\mathcal{P}}$ we have

$$\{f!U \mid U \in I\} = \{P_{f_1}, P_{f_2}, P_{f_3}, H_{f_1, f_2}, H_{f_1, f_3}, M := \text{Coker}(P_{f_2} \oplus P_{f_3} \rightarrow P_{f_1})\}.$$

Note that each of these is an interval module. The modules H_{f_1, f_2} and H_{f_1, f_3} are hooks, while the module $M = \text{Coker}(P_{f_2} \oplus P_{f_3} \rightarrow P_{f_1})$ satisfies

$$\dim M_x = \begin{cases} k & \text{if } f(1) \leq x \text{ and } (x \not\geq f(2) \text{ or } x \not\geq f(3)) \\ 0 & \text{else.} \end{cases}$$




(3) The situation is surprisingly different for $\mathcal{X} = \mathcal{X}''_3$ as in Proposition 3.12.

Here we have an \mathcal{X}''_3 -module $\mathbb{I}_{\mathcal{X}''_3} = \text{Coker}(P_3 \rightarrow P_1 \oplus P_2)$. The corresponding \mathcal{P} -modules $M = f!\mathbb{I}_{\mathcal{X}''_3} = \text{Coker}(P_{f_3} \rightarrow P_{f_1} \oplus P_{f_2})$ satisfy

$$\dim M_x = \begin{cases} k & \text{if } x \geq f(1) \\ k & \text{if } x \geq f(2) \text{ and } x \not\geq f(3) \\ k & \text{if } x \geq f(3) \text{ and } x \not\geq f(2) \\ k^2 & \text{if } x \geq f(2), x \geq f(3) \text{ and } x \not\geq f(1) \\ 0 & \text{else,} \end{cases}$$

and hence are not interval modules.



4.3. Bases for invariants and relative projectives.

Definition 4.11. Let $\Psi : K_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}^N$ be a finite invariant. A *basis* of Ψ is a set \mathcal{B} of indecomposable \mathcal{P} -modules such that Ψ induces an isomorphism of abelian group

$$\langle [U], U \in \mathcal{B} \rangle \longrightarrow \text{Im } \Psi,$$

where $\langle [U], U \in \mathcal{B} \rangle$ is the free abelian subgroup of $K_0^{\text{sp}}(\mathcal{P})$ generated by the $[U]$ for $U \in \mathcal{B}$.

As a consequence, if \mathcal{B} is a Ψ -basis, then for any module $M \in \text{mod } \mathcal{P}$ there exist unique integers $(\alpha_U)_{U \in \mathcal{B}}$ such that

$$\Psi(M) = \Psi \left(\bigoplus_{U, \alpha_U > 0} U^{\alpha_U} \right) - \Psi \left(\bigoplus_{U, \alpha_U < 0} U^{-\alpha_U} \right).$$

Following the terminology of [BOO], we will refer to the pair $(\bigoplus_{U, \alpha_U > 0} U^{\alpha_U}, \bigoplus_{U, \alpha_U < 0} U^{-\alpha_U})$ as the *minimal Ψ -decomposition of M with respect to \mathcal{B}* .

Example 4.12. (1) The simple \mathcal{P} -modules form a basis of $\mathbf{dim}_{\mathcal{P}}$. Moreover, since $\text{mod } \mathcal{P}$ has finite global dimension, the indecomposable projective \mathcal{P} -modules also form an integer basis of $\mathbf{dim}_{\mathcal{P}}$.
 (2) Let $\mathcal{H} \subseteq \text{Ind } \mathcal{P}$ denote the union of the set of hook modules and the set of indecomposable projectives (see Example 2.1). Then it is shown in [BBH24, BOO] that \mathcal{H} is a **rk**-basis.
 (3) It is shown in [BOO] that the rectangle modules form another **rk**-basis.

For use in Propositions 4.14 and 4.15 below, we recall the following well-known fact. This can also be taken as the definition of the Gabriel quiver containing no oriented cycles.

Lemma 4.13. *Let \mathcal{P} be a finite poset and let \mathcal{U} a finite set of pairwise non-isomorphic indecomposable \mathcal{P} -modules. Then the following are equivalent.*

(1) *If there are nonzero morphisms*

$$X_1 \xrightarrow{f_1} X_2 \xrightarrow{f_2} \dots \xrightarrow{f_{n-1}} X_n \xrightarrow{f_n} X_1$$

with each $X_i \in \mathcal{U}$, then f_j is an isomorphism for all $j = 1, \dots, n$.

(2) *The Gabriel quiver of the algebra $\text{End}_{\mathcal{P}}(\bigoplus_{U \in \mathcal{U}} U)$ contains no oriented cycles.*

Proposition 4.14. *Let \mathcal{X} be a representation-finite poset and \mathcal{P} be a finite poset. Recall that $\mathbf{I} = \text{Ind } \mathcal{X}$ and fix \mathbf{E} a subset of $\text{Emb}_{\mathcal{X}}^{\mathcal{P}}$. Assume that $\bigcup_{f \in \mathbf{E}} \text{Im } f = \mathcal{P}$ and that the Gabriel quiver of the endomorphism algebra $\text{End}_{\mathcal{P}}(\bigoplus_{(f,U) \in \mathbf{E} \times \mathbf{I} / \sim_!} f_!U)$ has no oriented cycle. Then the set $\mathcal{H}_{\mathbf{E}} := \{f_!U \mid (f,U) \in \mathbf{E} \times \mathbf{I} / \sim_!\}$ is a $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}$ -basis (and hence a $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}$ -basis).*

Proof. The proof is a consequence of Theorem 4.22 in [BBH24]. We give an alternative proof here for the convenience of the reader. For (f,U) and (g,V) in $\mathbf{E} \times \mathbf{I} / \sim_!$:

$$\left(\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}(f_!U) \right)_{(g,V)} = \dim \text{Hom}_{\mathcal{P}}(g_!V, f_!U).$$

Hence we have $\left(\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}(f_!U) \right)_{(f,U)} = 1$ and $\left(\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}(f_!U) \right)_{(g,V)} \neq 0$ implies $(g,V) \leq (f,U)$ in the partial order given by the Gabriel quiver of $\text{End}_{\mathcal{P}}(\bigoplus_{(f,U) \in \mathbf{E} \times \mathbf{I} / \sim_!} f_!U)$. We conclude using Lemma 2.8. \square

Proposition 4.15. *Let \mathcal{P} be a connected poset, and \mathcal{X} be one of the following posets*

$$\mathcal{X}'_3 = \begin{array}{c} & & 2 \\ & \nearrow & \\ 1 & & \\ & \searrow & \\ & & 3 \end{array} \quad \mathcal{X}''_3 = \begin{array}{c} 1 & & \\ & \searrow & \\ & & 3 \\ & \nearrow & \\ 2 & & \end{array}$$

Let E be a subset of $\text{Emb}_{\mathcal{X}}^{\mathcal{P}}$ such that $\bigcup_{f \in E} \text{Im } f = \mathcal{P}$. Then the set $\mathcal{H}_E = \{f_!U \mid (f, U) \in E \times I / \sim_!\}$ forms a $\mathbf{dimh}_{\mathcal{X}, \mathcal{P}}^E$ -basis, and hence also a $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^E$ -basis.

Proof. For $\mathcal{X} = \mathcal{X}'_3$ we can use Proposition 5.2 of [BBH24] to see that the objects $f_!U$ are “single-source spreads” since they have an indecomposable projective cover. Then it follows from Lemma 5.9 in [BBH24] that there is no oriented cycle in the Gabriel quiver of $\text{End}_{\mathcal{P}}(\bigoplus_{(f, U) \in E \times I / \sim_!} f_!U)$.

For $\mathcal{X} = \mathcal{X}''_3$, we prove directly that there are no cycle in the quiver of $\text{End}_{\mathcal{P}}(\bigoplus_{(f, U) \in E \times I / \sim_!} f_!U)$. We show that if there are nonzero morphisms

$$X_1 \rightarrow X_2 \rightarrow \cdots \rightarrow X_n \rightarrow X_1$$

with X_i of the form $f_!U$ for $(f, U) \in E \times I$ then $X_j \simeq X_1$ for any $j = 1, \dots, n$. Since the modules $f_!U$ are bricks this ends the proof. There are three types of modules of the form $f_!U$: the projective, the hooks and the modules of the form $\text{Coker}(P_a \rightarrow P_b \oplus P_c)$ with b and c incomparable and $b, c < a$ that we call generalized hooks. First note that there is no nonzero morphism from a hook or a generalized hook to a projective. Since we already know the claim holds if each X_i is projective, we may therefore assume that the X_i are either hooks or generalized hooks. We assume here that all the modules $X_i = \text{Coker}(P_{a_i} \rightarrow P_{b_i} \oplus P_{c_i})$ are generalized hooks, the proof easily generalizes when some of them are hooks.

A nonzero morphism $X_i \rightarrow X_{i+1}$ induces a commutative square

$$\begin{array}{ccc} P_{a_i} & \longrightarrow & P_{b_i} \oplus P_{c_i} \\ \downarrow & & \downarrow \\ P_{a_{i+1}} & \longrightarrow & P_{b_{i+1}} \oplus P_{c_{i+1}} \end{array}$$

The second vertical map is nonzero, hence we may assume $b_i \geq b_{i+1}$. Then we obtain $a \geq b_{i+1}$ and by composing this map with the projection $P_{b_{i+1}} \oplus P_{c_{i+1}} \rightarrow P_{b_{i+1}}$ we obtain that the map $P_{a_i} \rightarrow P_{a_{i+1}}$ is nonzero which implies that $a_i \geq a_{i+1}$. Therefore we have $a_i = a_1$ for all $i = 1, \dots, n$. Now we obtain nonzero morphisms

$$P_{b_1} \oplus P_{c_1} \rightarrow P_{b_2} \oplus P_{c_2} \rightarrow \cdots \rightarrow P_{b_n} \oplus P_{c_n} \rightarrow P_{b_1} \oplus P_{c_1}.$$


This composition is nonzero since the composition of these maps with $P_{a_1} \rightarrow P_{b_1} \oplus P_{c_1}$ is nonzero. Therefore we have a chain $\alpha_1 \geq \alpha_2 \geq \cdots \geq \alpha_n \geq \alpha_1$ with $\alpha_i = b_i$ or $\alpha_i = c_i$. Furthermore in the commutative square above, it is easy to see that if $a_i = a_{i+1}$ and $b_i = b_{i+1}$, then we should have $c_i \geq c_{i+1}$, since b_i and c_i are incomparable by hypothesis. Hence we deduce for each $i = 1, \dots, n$ we have $\{b_i, c_i\} = \{b_1, c_1\}$. Thus all the X_i 's are isomorphic. This ends the proof. \square

5. INTERVAL BASES VIA FAMILIES OF EMBEDDINGS

5.1. Interval modules and the Θ -functor. We highlight that Example 4.12 gives two distinct bases for the invariants \mathbf{dim} and \mathbf{rk} . In each case, one basis comes directly from the interpretation of the relevant invariant as a dim-hom invariant, while the other is somewhat less clear from a homological perspective. In this case, we use poset embeddings to give a unified homological interpretation of the fact that the simple modules form a basis of \mathbf{dim} and the fact that the rectangle modules form a basis of \mathbf{rk} . In doing so, we also uncover new bases of several other invariants.

We begin with an example illustrating that the induction and coinduction functors generally do not send interval modules to interval modules.

Example 5.1. Consider the posets



Let $f : \mathcal{X} \rightarrow \mathcal{P}$ be the poset embedding given by $fa = (2,1)$, $fb = (1,2)$, $fc = (3,2)$, and $fd = (2,3)$. Consider the sincere interval module $\mathbb{I}_{[\mathcal{X}]} \in \text{mod } \mathcal{X}$. Since the induction functor $f_!$ preserves projective presentations and the coinduction functor f_* preserves injective copresentations, we then have

$$\begin{array}{ccccc}
 K & \xrightarrow{1} & K & \longrightarrow & 0 \\
 \uparrow & & \nabla \uparrow & & \uparrow \\
 K & \xrightarrow{\iota_1} & K^2 & \xrightarrow{\nabla} & K \\
 \uparrow & & \iota_2 \uparrow & & \uparrow \\
 0 & \longrightarrow & K & \xrightarrow{1} & K \\
 & & f_! \mathbb{I}_{[\mathcal{X}]} & &
 \end{array}
 \qquad
 \begin{array}{ccccc}
 K & \xrightarrow{1} & K & \longrightarrow & 0 \\
 \uparrow & & \text{pr}_2 \uparrow & & \uparrow \\
 K & \xrightarrow{\Delta} & K^2 & \xrightarrow{\text{pr}_1} & K \\
 \uparrow & & \Delta \uparrow & & \uparrow \\
 0 & \longrightarrow & K & \xrightarrow{1} & K \\
 & & f_* \mathbb{I}_{[\mathcal{X}]} & &
 \end{array}$$

where ι_i denotes the i -th inclusion map, pr_i denotes the i -th projection map, and Δ and ∇ denote the diagonal and co-diagonal maps, respectively. We note that both $f_! \mathbb{I}_{[\mathcal{X}]}$ and $f_* \mathbb{I}_{[\mathcal{X}]}$ are indecomposable but are not interval modules. Moreover, we have that $\text{Hom}_{\mathcal{P}}(f_! \mathbb{I}_{[\mathcal{X}]}, f_* \mathbb{I}_{[\mathcal{X}]}) \cong K$ and that the image of any nonzero map $f_! \mathbb{I}_{[\mathcal{X}]} \rightarrow f_* \mathbb{I}_{[\mathcal{X}]}$ is precisely \mathbb{I}_S for $S = \{(i,j) \in \mathcal{P} \mid 3 \leq i+j \leq 5\}$.

It is shown in [AET, Proposition 4.17] that the final observation in Example 5.1 is not a coincidence. In order to state the general result from [AET], we need the following definitions.

Definition 5.2. Let $S \subseteq \mathcal{P}$. The *convex hull* of \mathcal{P} is the set

$$\overline{S} = \{x \in \mathcal{P} \mid \exists y, z \in S \mid y \leq x \leq z\}.$$

Definition 5.3. Let $f : \mathcal{X} \rightarrow \mathcal{P}$ be a poset embedding and let $U \in \text{mod } \mathcal{X}$. By adjunction and the fact that $f^* f_* = \text{Id}$, there is a natural isomorphism $\text{Hom}_{\mathcal{P}}(f_! U, f_* U) \cong \text{End}_{\mathcal{X}}(U)$. Let $\theta_f U : f_! U \rightarrow f_* U$ denote the image of the identity map $1_U \in \text{End}_{\mathcal{X}}(U)$ under this isomorphism. We denote $\Theta_f U = \text{Im}(\theta_f U)$.

Remark 5.4. (1) It is shown in [AET] that the association Θ_f extends to a functor $\text{mod } \mathcal{X} \rightarrow \text{mod } \mathcal{P}$. We will not use the functoriality of Θ_f in this paper.

(2) Suppose that $U \in \text{mod } \mathcal{X}$ is an interval module. Then $\dim_K \text{End}_{\mathcal{X}}(U) \cong K$ by [DX18, Proposition 14] or [BBH24, Proposition 5.5]. Thus every morphism $f_! U \rightarrow f_* U$ is a scalar multiple of $\theta_f U$ in this case.

We now recall the following, which shows that the functor Θ_f preserves interval modules.

Proposition 5.5. [AET, Proposition 4.17] *Let $f : \mathcal{X} \rightarrow \mathcal{P}$ be a poset embedding and $\mathbb{I}_S \in \text{mod } \mathcal{X}$ an interval module. Then $\Theta_f(\mathbb{I}_S) \cong \mathbb{I}_{\overline{fS}}$.*

The next example uses Proposition 5.5 to explain the relationship between the hook modules and rectangle modules in studying the rank invariant.

Example 5.6. (1) Let $\mathcal{X}_2 = \{1 \rightarrow 2\}$. We recall from Example 3.7 that the three indecomposable \mathcal{X}_2 -modules are $\mathbb{I}_{\{1,2\}} = P_1 = I_2$, $\mathbb{I}_{\{1\}} = H_{1,2} = I_a$, and $\mathbb{I}_{\{2\}} = P_2 = C_{1,2}$. For $f \in \text{Emb}_{\mathcal{X}}^{\mathcal{P}}$, we then compute

$$\begin{aligned}
 f! \mathbb{I}_{\{1\}} &= H_{f1, f2}, & f! \mathbb{I}_{\{2\}} &= P_{f2}, & f! \mathbb{I}_{\{1,2\}} &= P_{f1}, \\
 f* \mathbb{I}_{\{1\}} &= I_{f1}, & f* \mathbb{I}_{\{2\}} &= C_{f1, f2}, & f* \mathbb{I}_{\{1,2\}} &= I_{f2}, \\
 \Theta_f \mathbb{I}_{\{1\}} &= \mathbb{I}_{\{f1\}}, & \Theta_f \mathbb{I}_{\{2\}} &= \mathbb{I}_{\{f2\}}, & \Theta_f \mathbb{I}_{\{1,2\}} &= \mathbb{I}_{\{f1, f2\}}.
 \end{aligned}$$

In particular, we have that $\{\Theta_f U \mid f \in \text{Emb}_{\mathcal{X}}^{\mathcal{P}}, U \in \text{Ind } \mathcal{X}\}$ is the set of rectangle modules. Note that there is some redundancy in this collection: for $x \in \mathcal{P}$ we obtain the (simple) rectangle module $\mathbb{I}_{\{x\}}$ once for every embedding $f : \mathcal{X}_2 \rightarrow \mathcal{P}$ with either $x = f(1)$ or $x = f(2)$.

It follows from [BOO, Corollary 2.7] that $\{\Theta_f U \mid f \in \text{Emb}_{\mathcal{X}}^{\mathcal{P}}, U \in \text{Ind } \mathcal{X}\}$ is a $\mathbf{rk}_{\mathcal{P}}$ -basis, hence also a $\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}$ and $\mathbf{dimh}_{\mathcal{X}_2, \mathcal{P}}$ -basis. It is, however, difficult to reprove it directly without a concrete description of the image of the invariant $\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}$.

Note that we can define an equivalence relation \sim_{Θ} on $\mathbb{E} \times \mathbb{I}$ by setting

$$(f, U) \sim_{\Theta} (g, V) \iff \Theta_f(U) = \Theta_g(V),$$


but the invariant $\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}$ does not factor through $\mathbb{Z}^{\mathbb{E} \times \mathbb{I} / \sim_{\Theta}}$. For example, if $\mathcal{P} = \{a \rightarrow b \rightarrow c\}$, then we have $\Theta_f \mathbb{I}_{\{1\}} = \mathbb{I}_{\{a\}} = \Theta_g \mathbb{I}_{\{1\}}$ for $\text{Im } f = \{a, b\}$ and $\text{Im } g = \{a, c\}$. However, we have for any \mathcal{P} -module M such that $\text{rk } M_{a \rightarrow b} \neq \text{rk } M_{a \rightarrow c}$ that

$$\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}(M)_{(f, I_{\{1\}})} = \dim M_a - \text{rk } M_{a \rightarrow b} \neq \dim M_a - \text{rk } M_{a \rightarrow c} = \mathbf{mult}_{\mathcal{X}_2, \mathcal{P}}(M)_{(g, I_{\{1\}})}.$$

(2) Consider $\mathcal{X} = \mathcal{X}'_3$ or \mathcal{X}''_3 as follows

$$\mathcal{X}'_3 = \begin{array}{c} & & 2 \\ & \nearrow & \\ 1 & & \\ & \searrow & \\ & & 3 \end{array} \qquad \mathcal{X}''_3 = \begin{array}{c} 1 & & \\ & \searrow & \\ & & 3 \\ & \nearrow & \\ 2 & & \end{array}$$

We recall the description of the indecomposable \mathcal{X}'_3 - and \mathcal{X}''_3 -modules from the proof of Proposition 3.12 and Example 4.10. Then the modules $\Theta_f(U)$ are either simples, rectangles, or interval modules with support of the following form:



5.2. Families of embeddings. In the preceding sections, we mainly considered order-embeddings of *one fixed* poset \mathcal{X} into the poset \mathcal{P} , and studied the set of all indecomposables obtained from \mathcal{X} . That turns out to introduce a lot of redundancy and some difficulty to describe the image of the invariant. Hence we study now embeddings of several posets of increasing sizes, but limit attention to only some indecomposables (that have not been obtained from embedding of smaller posets previously). One of the advantage here is that we do not need to assume \mathcal{X} to be of finite representation type.

Definition 5.7. Let \mathcal{P} be a poset. A *family of \mathcal{P} -order embeddings* is a finite set $\mathcal{F} := \{(\mathcal{X}_i, E_i)\}_i$ of with each \mathcal{X}_i a finite poset (not necessarily representation finite) and each E_i a subset of $\text{Emb}_{\mathcal{X}_i}^{\mathcal{P}}$. For $\mathcal{F} = \{(\mathcal{X}_i, E_i)\}_i$ a family of \mathcal{P} -order embeddings, we denote by $\mathbb{E} := \bigsqcup_i E_i$ and we define an invariant $\mathbf{mult}_{\mathcal{F}} : \mathbf{K}_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}^{\mathbb{E}}$ by

$$\text{for } f \in E_i \quad (\mathbf{mult}_{\mathcal{F}}(M))_f := \text{mult}(\mathbb{I}_{\mathcal{X}_i}, f^*(M)).$$

Example 5.8. (1) Let $\mathcal{F} := \{(\mathcal{X}_1, \text{Emb}_{\mathcal{X}_1}^{\mathcal{P}}), (\mathcal{X}_2, \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}})\}$ for \mathcal{X}_1 and \mathcal{X}_2 as in Example 3.7. Then $\mathbb{E} = \text{Emb}_{\mathcal{X}_1}^{\mathcal{P}} \sqcup \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}}$ is in natural bijection with $\{\leq_{\mathcal{P}}\}$ and via this bijection we have $\mathbf{mult}_{\mathcal{F}} = \mathbf{rk}_{\mathcal{P}}$.
 (2) The generalized rank invariant of [KM21] and the various compressed multiplicities of [AENY23b] can both be realized in the form $\mathbf{mult}_{\mathcal{F}}$. More precisely:

- (a) Let \mathfrak{J} be the set of posets which embed into \mathcal{P} as intervals, and for each $\mathcal{X}_i \in \mathfrak{J}$ let $E_i \subseteq \text{Emb}_{\mathcal{X}_i}^{\mathcal{P}}$ be the set of embeddings with interval image. Then, as a result of [CL18, Lemma 3.1], the generalized rank invariant of [KM21] is equal to $\mathbf{mult}_{\mathcal{F}}$ for $\mathcal{F} = \{(\mathcal{X}_i, E_i) \mid \mathcal{X}_i \in \mathfrak{J}\}$.
- (b) Suppose that \mathcal{P} is the product of two (finite) totally ordered sets. Let \mathfrak{J} be the set of connected posets which do not contain any chains of length ≥ 3 and can be embedded into \mathcal{P} . If one computes the “sink-source compression factor” of [AENY23b] with respect to every interval subset of \mathcal{P} , the result is precisely $\mathbf{mult}_{\mathcal{F}}$ for $\mathcal{F} = \{(\mathcal{X}_i, \text{Emb}_{\mathcal{X}_i}^{\mathcal{P}}) \mid \mathcal{X}_i \in \mathfrak{J}\}$. A similar result likewise holds for the “corner-complete compression factors”, while the “total compression factors” coincide with the generalized rank invariant.

See also [BBH24, Section 3.2] and [AENY23b, Example 4.14] for additional discussion.

Proposition 5.9. *Let \mathcal{P} be a finite poset and $\mathcal{F} := \{(\mathcal{X}_i, E_i)\}_i$ be a family of \mathcal{P} -order embeddings. We define an equivalence relation on $E = \bigsqcup_i E_i$ by $f \sim g$ if $\overline{\text{Im } f} = \overline{\text{Im } g}$. Then the invariant $\mathbf{mult}_{\mathcal{F}} : K_0^{\text{sp}}(\mathcal{P}) \rightarrow \mathbb{Z}^E$ induces an isomorphism of abelian groups*

$$\langle [\mathbb{I}_{\overline{\text{Im } f}}], f \in E \rangle \xrightarrow{\sim} \mathbb{Z}^{E/\sim}.$$

Proof. For $f \in E_i$ and $g \in E_j$, we have

$$\left(\mathbf{mult}_{\mathcal{F}}(\mathbb{I}_{\overline{\text{Im } f}}) \right)_g = \text{mult}_{\mathcal{X}_j}(\mathbb{I}_{\mathcal{X}_j}, g^* \mathbb{I}_{\overline{\text{Im } f}}).$$

The \mathcal{X}_j -modules $\mathbb{I}_{\mathcal{X}_j}$ and $g^* \mathbb{I}_{\overline{\text{Im } f}} = \mathbb{I}_{g^{-1}(\overline{\text{Im } f})}$ are both interval modules hence indecomposable. Therefore we obtain $\left(\mathbf{mult}_{\mathcal{F}}(\mathbb{I}_{\overline{\text{Im } f}}) \right)_g$ is 0 or 1 and

$$\begin{aligned} \left(\mathbf{mult}_{\mathcal{F}}(\mathbb{I}_{\overline{\text{Im } f}}) \right)_g = 1 &\Leftrightarrow \mathcal{X}_j = g^{-1}(\overline{\text{Im } f}) \\ &\Leftrightarrow \mathcal{X}_j \subseteq g^{-1}(\overline{\text{Im } f}) \\ &\Leftrightarrow \overline{\text{Im } g} \subseteq \overline{\text{Im } f} \\ &\Leftrightarrow \overline{\text{Im } g} \subseteq \overline{\text{Im } f}. \end{aligned}$$

Then by defining an order relation \leq on E/\sim by

$$g \leq f \Leftrightarrow \overline{\text{Im } g} \subseteq \overline{\text{Im } f},$$

we conclude using Lemma 2.8. \square

It is not clear in general that the invariant $\mathbf{mult}_{\mathcal{F}}$ factors through the injection $\mathbb{Z}^{E/\sim} \rightarrow \mathbb{Z}^E$. However, it can be shown in some cases.

Example 5.10. (1) Consider the setup of Example 5.8(1). Then $\{\mathbb{I}_{\overline{\text{Im } f}} \mid f \in E\}$ is the set of rectangle modules. As previously mentioned, it is shown in [BOO, Corollary 2.7] that this forms a basis of $\mathbf{mult}_{\mathcal{F}}$.

- (2) Consider the setup of Example 5.8(2a) or (2b). In both cases, $\{\mathbb{I}_{\overline{\text{Im } f}} \mid f \in E\}$ is the set of interval modules. Both papers show that these form a basis for the corresponding invariant $\mathbf{mult}_{\mathcal{F}}$ using Möbius inversion arguments. (This is also established for the “corner-complete compression factors” in [AENY23b].)

Generalizing the phenomenon observed for the rank invariant and sink-source compression factors, we have the following.

Corollary 5.11. *Let \mathcal{P} be a finite poset, and $\mathcal{F} = \{(\mathcal{X}_i, E_i)\}$ be a family of \mathcal{P} -order embeddings such that there is no chain of length ≥ 3 in \mathcal{X}_i for each i . Denote $E = \bigsqcup_i E_i$. Then the family*

$$\mathcal{R}_E := \{\mathbb{I}_{\overline{\text{Im } f}} \mid f \in E\} \subseteq K_0^{\text{sp}}(\mathcal{P})$$

forms a basis of $\mathbf{mult}_{\mathcal{F}}$.

Proof. We show that in this case the equivalence relation on E is trivial and conclude by the previous proposition. Indeed let $f \in E_i$ and $g \in E_j$ such that $f \sim g$, that is $\overline{\text{Im } f} = \overline{\text{Im } g}$. Then let $x \in \mathcal{X}_i$. By hypothesis x is either minimal or maximal. Assume first it is minimal. Then $f(x) \in \text{Im } f \subseteq \overline{\text{Im } f} = \overline{\text{Im } g}$, so there exists $y, y' \in \mathcal{X}_j$ with $g(y) \leq f(x) \leq g(y')$. Since $g(y)$ is in $\text{Im } g$, it is in the convex hull of $\text{Im } f$,

but since x is minimal we have $f(x) = g(y)$. By a similar argument, if x is maximal, then $f(x)$ is in $\text{Im } g$. Therefore we obtain $\text{Im } f \subseteq \text{Im } g$, and by symmetry we get $\text{Im } f = \text{Im } g$, that is $f = g$. \square

Remark 5.12. It is an interesting problem to find a representation-theoretic argument generalizing the basis result for the generalized rank invariant established using Möbius inversion in [KM21]. In particular, note that the “zigzags” used in [DKM24] to compute the generalized rank invariant admit injective poset morphisms into \mathcal{P} , but that these morphisms are generally not full as functors between poset categories, and thus not embeddings. It would be interesting to determine the extent to which the theory developed in this paper could be adapted to such a setting.

5.3. Comparing mult-invariants. Let \mathcal{P} be a finite poset, \mathcal{X} be a representation-finite poset and \mathbf{E} be a subset of $\text{Emb}_{\mathcal{X}}^{\mathcal{P}}$. Let $\mathcal{F}_{\mathcal{P}} = \{(\mathcal{X}_i, \mathbf{E}_i)\}_i$ be a family of \mathcal{P} -order embeddings. The aim is here to compare the invariants $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}$ and $\mathbf{mult}_{\mathcal{F}_{\mathcal{P}}}$. We denote by $\mathcal{F}_{\mathcal{X}} = \{(\mathcal{X}_i, \text{Emb}_{\mathcal{X}_i}^{\mathcal{X}})\}_i$ the family of \mathcal{X} -order embeddings of the posets \mathcal{X}_i . The next proposition says that it is enough to compare the invariant $\mathbf{mult}_{\mathcal{X}}$ with the invariant $\mathbf{mult}_{\mathcal{F}_{\mathcal{X}}}$.

Proposition 5.13. (1) *If for any $(\mathcal{X}_i, \mathbf{E}_i) \in \mathcal{F}$ the composition map $\text{Emb}_{\mathcal{X}_i}^{\mathcal{X}} \times \text{Emb}_{\mathcal{X}}^{\mathcal{P}} \rightarrow \text{Emb}_{\mathcal{X}_i}^{\mathcal{P}}$ restricts to a map $\text{Emb}_{\mathcal{X}_i}^{\mathcal{X}} \times \mathbf{E} \rightarrow \mathbf{E}_i$, then*

$$\mathbf{mult}_{\mathcal{F}_{\mathcal{X}}} \geq \mathbf{mult}_{\mathcal{X}} \quad \Rightarrow \quad \mathbf{mult}_{\mathcal{F}_{\mathcal{P}}} \geq \mathbf{mult}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}.$$

(2) *If moreover the restriction $\text{Emb}_{\mathcal{X}_i}^{\mathcal{X}} \times \mathbf{E} \rightarrow \mathbf{E}_i$ is surjective, then*

$$\mathbf{mult}_{\mathcal{F}_{\mathcal{X}}} \simeq \mathbf{mult}_{\mathcal{X}} \quad \Rightarrow \quad \mathbf{mult}_{\mathcal{F}_{\mathcal{P}}} \simeq \mathbf{mult}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}.$$

Proof. (1) By hypothesis there exist integers $\alpha_{(g,U)}$ for any $g \in \bigsqcup_i \text{Emb}_{\mathcal{X}_i}^{\mathcal{X}}$ and $U \in \text{Ind}(\mathcal{X})$ such that for any $M \in \text{mod } \mathcal{X}$

$$\left(\mathbf{mult}_{\mathcal{X}}(M)\right)_U = \sum_{g \in \bigsqcup_i \text{Emb}_{\mathcal{X}_i}^{\mathcal{X}}} \alpha_{(g,U)} \left(\mathbf{mult}_{\mathcal{F}_{\mathcal{X}}}(M)\right)_g,$$

that is

$$\mathbf{mult}_{\mathcal{X}}(U, M) = \sum_{g \in \bigsqcup_i \text{Emb}_{\mathcal{X}_i}^{\mathcal{X}}} \alpha_{(g,U)} \mathbf{mult}_{\mathcal{X}_i}(\mathbb{I}_{\mathcal{X}_i}, g^* M).$$

Now for $(f, U) \in \mathbf{E} \times \text{Ind}(\mathcal{X})$, and $M \in \text{mod } \mathcal{P}$ we have

$$\begin{aligned} \left(\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}(M)\right)_{(f,U)} &= \mathbf{mult}_{\mathcal{X}}(U, f^* M) \\ &= \sum_{g \in \bigsqcup_i \text{Emb}_{\mathcal{X}_i}^{\mathcal{X}}} \alpha_{(g,U)} \mathbf{mult}_{\mathcal{X}_i}(\mathbb{I}_{\mathcal{X}_i}, g^*(f^* M)) \\ &= \sum_{g \in \bigsqcup_i \text{Emb}_{\mathcal{X}_i}^{\mathcal{X}}} \alpha_{(g,U)} \mathbf{mult}_{\mathcal{X}_i}(\mathbb{I}_{\mathcal{X}_i}, (f \circ g)^* M) \\ &= \sum_{h \in \bigsqcup_i \mathbf{E}_i} \beta_{(h,f,U)} \left(\mathbf{mult}_{\mathcal{F}_{\mathcal{P}}}(M)\right)_h \end{aligned}$$

where $\beta_{(h,f,U)} = \alpha_{(g,U)}$ as soon as $h = f \circ g$. Since f is an embedding the value of $\beta_{(h,f,U)}$ is well-defined.

(2) Now assume that there exist integers $\alpha'_{(g,U)}$ for $g \in \bigsqcup_i \text{Emb}_{\mathcal{X}_i}^{\mathcal{X}}$ and $U \in \text{Ind } \mathcal{X}$ such that for any $M \in \text{mod } \mathcal{X}$

$$\left(\mathbf{mult}_{\mathcal{F}_{\mathcal{X}}}(M)\right)_g = \sum_{U \in \text{Ind } \mathcal{X}} \left(\mathbf{mult}_{\mathcal{X}}(M)\right)_U,$$

that is

$$\mathbf{mult}_{\mathcal{X}_i}(\mathbb{I}_{\mathcal{X}_i}, g^* M) = \sum_{U \in \text{Ind}(\mathcal{X})} \alpha'_{(g,U)} \mathbf{mult}_{\mathcal{X}}(U, M).$$

Now for $h \in \mathbf{E}_i$ fix $h_1 \in \text{Emb}_{\mathcal{X}_i}^{\mathcal{X}}$ and $h_2 \in \mathbf{E}$ such that $h = h_2 \circ h_1$. Then we have

$$\begin{aligned} \left(\mathbf{mult}_{\mathcal{F}_{\mathcal{P}}}(M)\right)_h &= \mathbf{mult}_{\mathcal{X}_i}(\mathbb{I}_{\mathcal{X}_i}, h^* M) \\ &= \mathbf{mult}_{\mathcal{X}_i}(\mathbb{I}_{\mathcal{X}_i}, h_1^*(h_2^* M)) \\ &= \sum_{U \in \text{Ind } \mathcal{X}} \alpha'_{(h_1,U)} \mathbf{mult}_{\mathcal{X}}(U, h_2^* M) \\ &= \sum_{f \in \mathbf{E}} \sum_{U \in \text{Ind } \mathcal{X}} \beta'_{(h,f,U)} \left(\mathbf{mult}_{\mathcal{X}, \mathcal{P}}^{\mathbf{E}}(M)\right)_{(f,U)} \end{aligned}$$

where $\beta'_{(h,f,U)} = \alpha'_{(h_1,U)}$ as soon as $f = h_2$. \square

Example 5.14. (1) Let $\mathcal{F} = \{(\mathcal{X}_1, \text{Emb}_{\mathcal{X}_1}^{\mathcal{P}}), (\mathcal{X}_2, \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}})\}$ given as follows.

$$\mathcal{X}_1 = 1 \quad \mathcal{X}_2 = 1 \longrightarrow 2$$

Let $\mathcal{X} = \mathcal{X}_2$. Then we have two embeddings of \mathcal{X}_1 in \mathcal{X} and one of \mathcal{X}_2 to \mathcal{X} . Moreover we have for M in $\text{mod } \mathcal{X}$

$$(\mathbf{mult}_{\mathcal{F}\mathcal{X}}(M))_f = \begin{cases} \dim M_x & \text{if } f \in \text{Emb}_{\mathcal{X}_1}^{\mathcal{X}_2} \text{ with } \text{Im } f = \{x\} \\ \text{rk } M_{1 \rightarrow 2} & \text{if } f \in \text{Emb}_{\mathcal{X}_2}^{\mathcal{X}_2} \end{cases}$$

Then we immediately obtain

$$\mathbf{mult}_{\mathcal{X}} \sim \mathbf{mult}_{\mathcal{F}\mathcal{X}}.$$

It is clear in this case that the composition map $\text{Emb}_{\mathcal{X}_i}^{\mathcal{X}_2} \times \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}}$ is surjective if \mathcal{P} is connected, hence we obtain once again that

$$\mathbf{mult}_{\mathcal{X}_2, \mathcal{P}} \simeq \mathbf{mult}_{\mathcal{F}} = \mathbf{rk}_{\mathcal{P}}.$$

(2) Let \mathcal{F} be the following family of posets :

$$\mathcal{X}_1 = 1 \quad \mathcal{X}_2 = 1 \longrightarrow 2 \quad \mathcal{X}'_3 = 1 \begin{array}{l} \nearrow 2 \\ \searrow 3 \end{array}$$

Set $\mathcal{X} = \mathcal{X}'_3$.

Then we have

$$\text{Emb}_{\mathcal{X}_1}^{\mathcal{X}} = \{f_1, f_2, f_3\}, \quad \text{Emb}_{\mathcal{X}_2}^{\mathcal{X}} = \{f_{12}, f_{13}\}, \quad \text{and } \text{Emb}_{\mathcal{X}'_3}^{\mathcal{X}} = \{\text{id}\}.$$

Then one easily checks that we have

$$\mathbf{mult}_{\mathcal{F}\mathcal{X}} = \Phi \circ \mathbf{mult}_{\mathcal{X}}$$

with

$$\Phi = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$

in the basis $\text{Ind}_{\mathcal{X}} = (\mathbb{I}_{\{1\}}, \mathbb{I}_{\{2\}}, \mathbb{I}_{\{3\}}, \mathbb{I}_{[1,2]}, \mathbb{I}_{[1,3]}, \mathbb{I}_{\mathcal{X}'_3})$.

(3) Let \mathcal{X} be the poset given by

$$\begin{array}{ccc} & b & \\ a & \nearrow & d \\ & c & \end{array}$$

And let \mathcal{F} be the family

$$\mathcal{X}_1 = 1 \quad \mathcal{X}_2 = 1 \longrightarrow 2 \quad \mathcal{X}'_3 = 1 \begin{array}{l} \nearrow 2 \\ \searrow 3 \end{array} \quad \mathcal{X}''_3 = \begin{array}{l} 1 \\ \nearrow 2 \\ \searrow 3 \end{array}$$

Then one can check that

$$|\text{Emb}_{\mathcal{X}_1}^{\mathcal{X}}| = 4, \quad |\text{Emb}_{\mathcal{X}_2}^{\mathcal{X}}| = 5, \quad |\text{Emb}_{\mathcal{X}'_3}^{\mathcal{X}}| = 1, \quad |\text{Emb}_{\mathcal{X}''_3}^{\mathcal{X}}| = 1.$$

The rank of the invariant $\mathbf{mult}_{\mathcal{X}}$ is also 11 since there are 11 indecomposable \mathcal{X} modules. In a similar fashion, one can check that

$$\mathbf{mult}_{\mathcal{X}} \sim \mathbf{mult}_{\mathcal{F}\mathcal{X}}.$$

Using Corollary 5.11 we obtain a basis of the invariant $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$.

(4) Let \mathcal{F} be the family of posets

$$\mathcal{X}_1 = 1 \quad \mathcal{X}_2 = 1 \longrightarrow 2 \quad \mathcal{X}'_3 = 1 \begin{array}{l} \nearrow 2 \\ \searrow 3 \end{array} \quad \mathcal{X}_4 = 1 \longrightarrow 2 \begin{array}{l} \nearrow 3 \\ \searrow 4 \end{array}$$

and let $\mathcal{X} := \mathcal{X}_4$. Then one can check

$$|\text{Emb}_{\mathcal{X}_1}^{\mathcal{X}}| = 4, |\text{Emb}_{\mathcal{X}_2}^{\mathcal{X}}| = 5, |\text{Emb}_{\mathcal{X}'_3}^{\mathcal{X}}| = 2, |\text{Emb}_{\mathcal{X}_4}^{\mathcal{X}}| = 1.$$

The rank of the invariant $\mathbf{mult}_{\mathcal{X}}$ is 12, and we also have an equivalence

$$\mathbf{mult}_{\mathcal{X}} \sim \mathbf{mult}_{\mathcal{F}_{\mathcal{X}}}.$$

(5) Let \mathcal{F} be the following family of posets, and $\mathcal{X} = \mathcal{X}'_4$.

$$\mathcal{X}_1 = 1 \quad \mathcal{X}_2 = 1 \longrightarrow 2 \quad \mathcal{X}'_3 = 1 \begin{array}{l} \nearrow 2 \\ \searrow 3 \end{array} \quad \mathcal{X}'_4 = a \begin{array}{l} \nearrow b \\ \rightarrow c \\ \searrow d \end{array}$$

Then one can check that

$$|\text{Emb}_{\mathcal{X}_1}^{\mathcal{X}}| = 4, |\text{Emb}_{\mathcal{X}_2}^{\mathcal{X}}| = 3, |\text{Emb}_{\mathcal{X}'_3}^{\mathcal{X}}| = 3, |\text{Emb}_{\mathcal{X}'_4}^{\mathcal{X}}| = 1.$$

Here the invariant $\mathbf{mult}_{\mathcal{X}}$ is strictly finer than the invariant $\mathbf{mult}_{\mathcal{F}_{\mathcal{X}}}$.

As a corollary of Example (2) we obtain the following

Corollary 5.15. *Let \mathcal{P} be a finite connected poset, and let \mathcal{X} be one of the following posets*

$$\mathcal{X}'_3 = 1 \begin{array}{l} \nearrow 2 \\ \searrow 3 \end{array} \quad \mathcal{X}''_3 = \begin{array}{l} 1 \\ \searrow 2 \end{array} \begin{array}{l} \nearrow 3 \\ \nearrow 3 \end{array}$$

Let $\mathbf{E} = \text{Emb}_{\mathcal{X}}^{\mathcal{P}}$ and recall that $\mathbf{I} = \text{Ind } \mathcal{X}$. Assume the following

$$(*) \text{ for any } a < b \text{ in } \mathcal{P} \text{ there exists } f \in \mathbf{E} \text{ with } \{a, b\} \in \text{Im } f$$

then the set $\mathcal{R}_{\mathbf{E}} = \{\Theta_f(U) \mid (f, U) \in \mathbf{E} \times \mathbf{I}\}$ forms a basis of $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$.

Proof. We consider the family of embeddings $\mathcal{F} := \{(\mathcal{X}_1, \text{Emb}_{\mathcal{X}_1}^{\mathcal{P}}), (\mathcal{X}_2, \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}}), (\mathcal{X}_3, \text{Emb}_{\mathcal{X}_3}^{\mathcal{P}})\}$ where

$$\mathcal{X}_1 = 1 \quad \mathcal{X}_2 = 1 \longrightarrow 2$$

and $\mathcal{X}_3 = \mathcal{X}$. Because of condition (*), the composition map $\text{Emb}_{\mathcal{X}_2}^{\mathcal{X}} \times \text{Emb}_{\mathcal{X}}^{\mathcal{P}} \rightarrow \text{Emb}_{\mathcal{X}_2}^{\mathcal{P}}$ is surjective. By condition (*) and since \mathcal{P} is connected, the composition map $\text{Emb}_{\mathcal{X}'_3}^{\mathcal{X}} \times \text{Emb}_{\mathcal{X}}^{\mathcal{P}} \rightarrow \text{Emb}_{\mathcal{X}'_3}^{\mathcal{P}}$ is also surjective. Then we can deduce from Example 2.6(2) that $\mathbf{mult}_{\mathcal{X}, \mathcal{P}}$ is equivalent to $\mathbf{mult}_{\mathcal{F}}$. From Corollary 5.11, we deduce that there is a basis of the form

$$\{\mathbb{I}_{\overline{\text{Im } f}} \mid f \in \text{Emb}_{\mathcal{X}'_i}^{\mathcal{P}}, i = 1, 2, 3\}.$$

This basis contains all simples, all rectangles, and all generalized rectangles as described in Example 5.6 (2). One easily check that the set $\mathcal{R}_{\mathbf{E}}$ contains exactly the simples, the rectangles and the generalized rectangles. The fact that it contains all the rectangles comes from condition (*). \square

REFERENCES

- [AENY23a] H. Asashiba, E. G. Escolar, K. Nakashima, and M. Yoshiwaki, *Approximation by interval-decomposables and interval resolutions of persistence modules*, J. Pure Appl. Alg. **227** (2023), no. 10.
- [AENY23b] H. Asashiba, E. G. Escolar, K. Nakashima, and M. Yoshiwaki, *On approximation of 2D persistence modules by interval-decomposables*, Journal of Computational Algebra **6-7** (2023).
- [AET] T. Aoki, E. G. Escolar, and S. Tada, *Summand-injectivity of interval covers and monotonicity of interval resolution global dimensions*. Preprint, 2023. arXiv:2308.14979 [math.RT].
- [ASS06] I. Assem, D. Simson, and A. Skowroński, *Elements of the representation theory of associative algebras, volume 1: techniques of representation theory*, London Math. Soc. Stud. Texts, volume 65, Cambridge University Press, Cambridge, 2006.

- [AS93] M. Auslander and Ø. Solberg, *Relative homology and representation theory 1: Relative homology and homologically finite subcategories*, *Comm. Algebra* **21** (1993), no. 9, 2995-3031.
- [BBH] B. Blanchette, T. Brüstle, and E.J. Hanson, *Exact structures for persistence modules*. Preprint, 2023. arXiv:2308.01790[math.AT].
- [BBH24] B. Blanchette, T. Brüstle, and E.J. Hanson, *Homological approximations in persistence theory*. *Canad. J. Math.* **76** (2024), no. 1, 66-103.
- [BC-B20] M. B. Botnan and W. Crawley-Boevey, *Decomposition of persistence modules*. *Proc. Amer. Math. Soc.* **148** (2020), no. 11, 4581-4596.
- [BLO22] M.B. Botnan, V. Lebovici, and S. Oudot. *On Rectangle-Decomposable 2-Parameter Persistence Modules*. *Discrete Comput Geom* **68** (2022), 1078–1101.
- [BL18] M. Botnan and M. Lesnick, *Algebraic stability of zigzag persistence modules* *Algebr. Geom. Topol.* **18** (2018), no. 6, 3133–3204.
- [BL23] M.B. Botnan and M. Lesnick, *An introduction to multiparameter persistence*. In: (A.B. Buan, H. Krause, and Ø. Solberg, eds.) *Representations of Algebras and Related Structures*, EMS Ser. Congr. Rep., volume 19, Eur. Math. Soc., Zürich, 2023.
- [BOO] M. B. Botnan, S. Oppermann, and S. Oudot, *Signed barcodes for multi-parameter persistence via rank decompositions and rank-exact resolutions*. Preprint, 2022. arXiv:2107.06800v2 [math.AT].
- [BOOS] M. B. Botnan, S. Oppermann, S. Oudot and L. Scoccola, *On the bottleneck stability of rank decompositions of multi-parameter persistence modules*. Preprint, 2022. arXiv:2208.00300 [math.AT].
- [Bü10] T. Bühler, *Exact categories*. *Expo. Math.* **28** (2010), no. 1, 1–69.
- [CZ09] G. Carlsson and A. Zomorodian. *The theory of multidimensional persistence*. *Discrete Comput. Geom.* **42** (2009), no. 1, 71–93.
- [CGRST] W. Chachólski, A. Guidolin, I. Ren, M. Scolamiero, and F. Tombari, *Koszul complexes and relative homological algebra of functors over posets*. Preprint, 2023. arXiv:2209.05923v2 [math.AT]
- [CL18] E. Chambers and D. Letscher, *Persistent homology over directed acyclic graphs*. In (E. Chambers, B. Fasy, and L. Ziegelmeier, editors) *Research in Computational Topology*, Association for Women in Mathematics Series, volume 13, Springer, Cham, 2018.
- [DKM24] T.K. Dey, W. Kim, and F. Mémoli. *Computing generalized rank invariant for 2-parameter persistence modules via zigzag persistence and its applications*. *Discrete Comput. Geom.* **71** (2024), 67-94.
- [DX18] T.K. Dey and C. Xin, *Computing bottleneck distance for 2-D interval decomposable modules*. In *34th International Symposium on Computational Geometry (SoCG 2018)*, LIPIcs. Leibniz Int. Proc. Inform., vol. 99, pp. 32:1-32:15, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2018.
- [DRSS99] P. Dräxler, I. Reiten, S.O. Smalø, and Ø. Solberg, *Exact Categories and Vector Space Categories*, *Trans. Amer. Math. Soc.* **351** (1999).
- [GR92] P. Gabriel and A.V. Roiter, *Algebra VIII: Representations of finite-dimensional algebras*, *Encyclopaedia Math. Sci.*, vol. 73, Springer, Berlin, 1992, with a chapter by B. Keller.
- [HNOX] Y. Hiraoka, K. Nakashima, I. Obayashi, and C. Xu. *Refinement of interval approximations for fully comutative quivers*. Preprint, 2023. arXiv:2310.03649v2 [math.AT].
- [KM21] W. Kim and F. Mémoli, *Generalized persistence diagrams for persistence modules over posets*. *J. Appl. Comput. Topol.* **5** (2021), 533–581.
- [OS] S. Oudot and L. Scoccola, *On the stability of multigraded Betti numbers and Hilbert functions*. Preprint, 2023. arXiv:2112.11901v3 [math.AT].
- [Qu72] D. Quillen, *Higher algebraic K-theory I*. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)* pp. 85–147, *Lecture Notes in Math.*, Vol. 341, 1973.
- [Si10] D. Simson. *Integral bilinear forms, Coxeter transformations and Coxeter polynomials of finite posets*. *Linear Algebra Appl.* **433** (2010), 699-717.

CLAIRE AMIOT, INSTITUT FOURIER, 100 RUE DES MATHS, 38402 SAINT MARTIN D'HÈRES, UNIVERSITÉ GRENOBLE ALPES, INSTITUT UNIVERSITAIRE DE FRANCE

Email address: `claire.amiot@univ-grenoble-alpes.fr`

THOMAS BRÜSTLE, DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE SHERBROOKE, SHERBROOKE, QC, J1K 2R1, CANADA AND BISHOP'S UNIVERSITY, SHERBROOKE, QC, J1M 1Z7, CANADA

Email address: `Thomas.Brustle@USherbrooke.ca`, `tbruestl@bishops.ca`

ERIC J. HANSON, DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27695, USA

Email address: `ejhanso3@ncsu.edu`