



HAL
open science

A NEW CHARACTERIZATION OF SLICES IN WARPED PRODUCTS

Julien Roth

► **To cite this version:**

Julien Roth. A NEW CHARACTERIZATION OF SLICES IN WARPED PRODUCTS. 2024. hal-04720027

HAL Id: hal-04720027

<https://cnrs.hal.science/hal-04720027v1>

Preprint submitted on 3 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A NEW CHARACTERIZATION OF SLICES IN WARPED PRODUCTS

JULIEN ROTH

ABSTRACT. We show that in certain warped products, a starshaped and s -convex hypersurface for which there exists a positive constant c so that the higher order mean curvatures H_s and H_{s-1} satisfy $H_s^{\frac{1}{s}} \leq c \leq H_{s-1}^{\frac{1}{s-1}}$ is a slice.

1. INTRODUCTION

The question of the characterization of round spheres in space forms has a long history. One of the most famous result is the well-known Alexandrov theorem [1] which ensures that a closed embedded hypersurface of the Euclidean space \mathbb{R}^{n+1} with constant mean curvature must be a round sphere. Further, this result has been extended to scalar curvature and then higher order mean curvatures by Ros [9, 10]. Later, Montiel and Ros [8] proved that the Alexandrov theorem for the mean curvature as well as for higher order mean curvatures is also true for hypersurfaces of hyperbolic spaces and half-spheres. There has been a wide litterature one the topic and many other characterizations have been obtained for space forms with different type of conditions. One can refer to [11] and references therein for instance.

More recently, Brendle [3] has obtained an Alexandrov type theorem for hypersurfaces into certain warped products (geodesic sphere are thus replaced by totally umbilical hypersurfaces or slices) that he has extended with Eichmair [4] for higher order mean curvatures. After that, many other characterizations of slices in warped products have been obtained (see [6, 7, 11, 13] for instance).

The aim of this short note is to give a new characterization of slices into warped products. This characterization is not of Alexandrov type, but lies on some inequalities between two consecutive higher order mean curvatures.

The geometric setting is the following. Let $n \geq 2$ be an integer and (M^n, g_M) be a compact Riemannian manifold of dimension n satisfying

$$\text{Ric}_M \geq (n-1)kg,$$

for some constant k . Moreover, let $t_0 > 0$ and $h : [0, t_0) \rightarrow \mathbb{R}$ be a positive function satisfying the following four conditions

$$(H1) \quad h'(0) = 0 \text{ and } h''(0) > 0,$$

$$(H2) \quad h'(t) > 0 \text{ for all } t \in (0, t_0),$$

2010 *Mathematics Subject Classification.* 53C42, 53A07, 49Q10.

Key words and phrases. warped products, geodesic spheres, higher order mean curvatures.

(H3) the function $r \mapsto 2\frac{h''(t)}{h(t)} - (n-1)\frac{k-h'(t)^2}{h(t)^2}$ is non-decreasing on $(0, t_0)$,

(H4) $\frac{h''(t)}{h(t)} + \frac{k-h'(t)^2}{h(t)^2} > 0$ for all $t \in (0, t_0)$.

We consider the warped product P defined by $P = [0, t_0) \times M$ endowed with the metric $g_P = dt^2 \oplus h(t)g_M$. This wide class of Riemannian manifolds includes the three real space forms $\mathbb{M}^{n+1}(\delta)$, that \mathbb{R}^{n+1} if $\delta = 0$, \mathbb{S}_+^{n+1} if $\delta = 1$ or \mathbb{H}^{n+1} if $\delta = -1$ with appropriate choice of the warping function.

Here is the main result of the present note.

Theorem 1.1. *Let $n \geq 2$ and $s \in \{2, \dots, n\}$ are two integers and Σ be a closed, oriented hypersurface embedded into the warped product (P^{n+1}, g_P) satisfying the four conditions (H1)-(H4). If Σ is star-shaped, s -convex and if there exists a constant c so that $H_s^{\frac{1}{s}} \leq c \leq H_{s-1}^{\frac{1}{s-1}}$ then Σ is a slice $\{t_1\} \times M$.*

Remark that, as we will see in the preliminary section, by the s -convexity assumption, we always have $H_s^{\frac{1}{s}} \leq H_{s-1}^{\frac{1}{s-1}}$. The condition here is to assume that some constant c can be inserted between the two quantities $H_s^{\frac{1}{s}}$ and $H_{s-1}^{\frac{1}{s-1}}$, which, by the theorem, forces the hypersurface to be a slice.

Note also that from the s -convexity assumption, if such a constant exists, then it is positive.

Finally, in the case where (P^{n+1}, g_P) is one of the real space forms $\mathbb{M}^{n+1}(\delta)$, we recover the result of Stong [12].

2. PRELIMINARIES

We consider the warped product (P, g_P) of the form given in the introduction and satisfying the four conditions (H1), (H2), (H3) and (H4).

Let Σ be a closed, embedded, connected and oriented hypersurface of (P^{n+1}, g_P) .

We consider the function $f = h'$ and the vector field $X = h\frac{\partial}{\partial t}$. We will also assume that Σ is starshaped, that is, the function $\langle X, \nu \rangle$ has a fixed sign on Σ , where ν is the outward normal unit vector field. In [3], Brendle used the fact that X is a conformal vector field, that is $\mathcal{L}_X g = 2fg$, to obtain the following Hsiung-Minkowski formula for hypersurfaces of these warped products

$$(1) \quad \int_{\Sigma} H \langle X, \nu \rangle dv_g = \int_{\Sigma} f dv_g,$$

where g is the induced metric on Σ and H the mean curvature of Σ .

More generally, the higher order mean curvatures are extrinsic quantities defined from the second fundamental form and generalizing the notion of mean curvature. Up to a normalisation the mean curvature H is the trace of the second fundamental form B :

$$(2) \quad H = \frac{1}{n} \text{tr}(B).$$

In other words, the mean curvature is

$$(3) \quad H = \frac{1}{n} S_1(\kappa_1, \dots, \kappa_n),$$

where S_1 is the first elementary symmetric polynomial and $\kappa_1, \dots, \kappa_n$ are the principal curvatures. Higher order mean curvatures are defined in a similar way for $r \in \{1, \dots, n\}$ by

$$(4) \quad H_r = \frac{1}{\binom{n}{r}} S_r(\kappa_1, \dots, \kappa_n),$$

where S_r is the r -th elementary symmetric polynomial, that is for any n -tuple (x_1, \dots, x_n)

$$(5) \quad S_r(x_1, \dots, x_n) = \sum_{1 \leq i_1 < \dots < i_r \leq n} x_{i_1} \cdots x_{i_r}.$$

Note that of cours, H_1 is nothing else but the mean curvature H .

We recall some classical inequalities between the higher order mean curvatures which are well-known. If $H_s > 0$, then $H_r > 0$ for any $r \in \{1, \dots, s-1\}$ and

$$(6) \quad H_s^{\frac{1}{s}} \leq H_{s-1}^{\frac{1}{s-1}} \leq \dots \leq H_2^{\frac{1}{2}} \leq H.$$

These classical inequalities can be found in [7] for instance where Li, Wei and Xiong extend to warped products the proof of Barbosa and Colares [2] for space forms. Finally, we recall the following general Hsiung-Minkowski type formula in warped product. Namely, if conditions (H1)-(H4) are satisfied and Σ is starshaped, then

$$(7) \quad \int_{\Sigma} f H_{s-1} dv_g \leq \int_{\Sigma} H_s \langle X, \nu \rangle dv_g.$$

This inequality has been first proved by Brendle-Eichmair [4] if Σ is convex but it turned out that the convexity assumption is not necessary and can be replaced by the s -convexity, that is, H_s is positive everywhere, which is a weaker assumption than convexity which means that all the principal curvatures are positive.

Note that (1) and (7) are the generalizations in this class of warped products of the so-called Hsiung-Minkowski formulas [5] for space forms (see [7] for instance). A crucial difference is that in space forms, they are equalities, which is not the case here except for identity (1).

Now, we have all the ingredients to prove Theorem 1.1.

3. PROOF OF THEOREM 1.1

First of all, from assumptions (H1) and (H2), $f = h'$ is a nonnegative function and $\int_{\Sigma} f dv_g > 0$ (in fact f can vanish at most in one point, corresponding to $t = 0$ in the product). Moreover, as mentioned in the preliminary section, since we assume that H_s is a positive function, then H is also positive. Moreover, since Σ is supposed to be starshaped, the support function $\langle X, \nu \rangle$ has fixed sign and so $H \langle X, \nu \rangle$ also has the same fixed sign. Hence, from Brendle identity (1), we have

$$\int_{\Sigma} H \langle X, \nu \rangle dv_g = \int_{\Sigma} f dv_g > 0,$$

and so necessarily, $\langle X, \nu \rangle$ has to be positive.

Second, from the assumptions $H_s^{\frac{1}{s}} \leq c$ and the fact that H_s is positive everywhere,

then the constant c is positive.

Moreover, from the assumption $c \leq H_{s-1}^{\frac{1}{s-1}}$, we have

$$(8) \quad \int_{\Sigma} f H_{s-1} dv_g \geq c^{s-1} \int_{\Sigma} f dv_g,$$

which gives by using Brendle identity (1)

$$(9) \quad \int_{\Sigma} f H_{s-1} dv_g \geq c^{s-1} \int_{\Sigma} H \langle X, \nu \rangle dv_g.$$

On the other hand, from (7), we have

$$(10) \quad \int_{\Sigma} f H_{s-1} dv_g \leq \int_{\Sigma} H_s \langle X, \nu \rangle dv_g.$$

Using the other part of the assumption, $H_s^{\frac{1}{s}} \leq c$, and the fact that $\langle X, \nu \rangle$ is positive everywhere on Σ , we obtain

$$(11) \quad \int_{\Sigma} f H_{s-1} dv_g \leq \int_{\Sigma} c^s \langle X, \nu \rangle dv_g = c^{s-1} \int_{\Sigma} c \langle X, \nu \rangle dv_g.$$

From (6) and the assumption $c \leq H_{s-1}^{\frac{1}{s-1}}$, we obtain immediately that

$$(12) \quad H \geq H_{s-1}^{\frac{1}{s-1}} \geq c.$$

Hence (11), (12) and the positivity of $\langle X, \nu \rangle$ and c give

$$(13) \quad \int_{\Sigma} f H_{s-1} dv_g \leq c^{s-1} \int_{\Sigma} H \langle X, \nu \rangle dv_g.$$

We use the Brendle identity (1) again to obtain from (13)

$$(14) \quad \int_{\Sigma} f H_{s-1} dv_g \leq c^{s-1} \int_{\Sigma} f dv_g.$$

Thus, (8) and (14) together yield to

$$(15) \quad \int_{\Sigma} f H_{s-1} dv_g = c^{s-1} \int_{\Sigma} f dv_g$$

and all inequalities from (8) to (14) are in fact equalities. In particular, we get

$$(16) \quad \int_{\Sigma} f (H_{s-1} - c^{s-1}) dv_g = 0.$$

Since f is a positive function and with the assumption $H_{s-1} \geq c^{s-1}$, we deduce that $H_{s-1} = c^{s-1}$.

On the other hand, from the fact that (11) and (13) are equalities, we also deduce that

$$(17) \quad c^{s-1} \int_{\Sigma} c \langle X, \nu \rangle dv_g = c^{s-1} \int_{\Sigma} H \langle X, \nu \rangle dv_g,$$

or equivalently, since c is a positive constant

$$(18) \quad \int_{\Sigma} (c - H) \langle X, \nu \rangle dv_g = 0.$$

From (18), the fact that $\langle X, \nu \rangle$ is positive and $H \geq c$, we obtain that $H = c$.

Finally, we have $H_{s-1}^{\frac{1}{s-1}} = c = H$ so that equality occurs in (6) and so Σ is totally umbilical.

We finish the proof by using the following argument due to Brendle [3]: if (H1)-(H4) are satisfied, Σ is starshaped and totally umbilical, then Σ is a slice. \square

REFERENCES

- [1] A.D. Alexandrov, *A characteristic property of spheres*, Ann. Mat. Pura Appl., **58** (1962), 303–315.
- [2] J. Barbosa and A. Colares, *Stability of hypersurfaces with constant r -mean curvature*, Ann. Glob. Anal. Geom. **15** (1997), no. 3, 277–297.
- [3] S. Brendle, *Constant mean curvature surfaces in warped product manifolds*, Publ. Math. IHES, **117** (2013) 247–269.
- [4] S. Brendle & M. Eichmair, *Isoperimetric and Weingarten surfaces in the Schwarzschild manifold*, J. Differ. Geom. **94** (2013) 387–407.
- [5] C. C. Hsiung, *Some integral formulas for closed hypersurfaces*, Math. Scand. **2** (1954), 286–294.
- [6] K. K. Kwong, H. Lee & J. Pyo, *Weighted Hsiung-Minkowski formulas and rigidity of umbilical hypersurfaces*, Math. Res. Letters, 25(2), 597–616.
- [7] H. Li, Y. Wei & C. Xiong, *A note on Weingarten hypersurfaces in the warped product manifold*, Int J. Math **25 (14)** (2014), article ID: 1450121.
- [8] S. Montiel & A. Ros, *Compact hypersurfaces : the Alexandrov theorem for higher order mean curvature*, Pitman Monographs Surveys Pure Appl. Math. **52** (1991), 279–296.
- [9] A. Ros, *Compact hypersurfaces with constant scalar curvature and a congruence theorem*, J. Differential Geom. **27(2)** (1988), 215–223.
- [10] A. Ros, *Compact hypersurfaces with constant higher order mean curvatures*, Rev. Mat. Iberoamericana **3(3-4)** (1987), 447–453.
- [11] J. Roth & A. Upadhyay, *On compact embedded Weingarten hypersurfaces in warped products*, J. Math. Anal. Appl. **517 (1)** (2023), article Id: 126593.
- [12] R. E. Stong, *Some characterizations of Riemann n -spheres*, Proc. Amer. Math. Soc. **11** (1960), 945–951. 193–243.
- [13] J. Wu & C. Xia, *On rigidity of hypersurfaces with constant curvature functions in warped product manifolds*, Ann. Glob. Anal. Geom. **46** (2014) 1–22.

(J. ROTH) UNIVERSITÉ GUSTAVE EIFFEL, CNRS, LAMA UMR 8050, F-77447 MARNE-LA-VALLÉE, FRANCE

Email address: julien.roth@univ-eiffel.fr