



**HAL**  
open science

## Serum calprotectin levels do not predict subsequent relapse in rheumatoid arthritis in remission: a post-hoc analysis of STRASS study

Xavier Romand, Margot Clapasson, Minh Vu Chuong, Marie H el ene Paclet, Bruno Fautrel, Athan Baillet

### ► To cite this version:

Xavier Romand, Margot Clapasson, Minh Vu Chuong, Marie H el ene Paclet, Bruno Fautrel, et al.. Serum calprotectin levels do not predict subsequent relapse in rheumatoid arthritis in remission: a post-hoc analysis of STRASS study. *RMD Open*, 2023, 9 (2), pp.e003198. 10.1136/rmdopen-2023-003198. hal-04724625

**HAL Id: hal-04724625**

**<https://cnrs.hal.science/hal-04724625v1>**




Submitted on 6 Jan 2025

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

## LETTER

## Serum calprotectin levels do not predict subsequent relapse in rheumatoid arthritis in remission: a post-hoc analysis of STRASS study

Xavier Romand <sup>1</sup>, Margot Clapasson,<sup>2</sup> Minh Vu Chuong,<sup>1</sup> Marie H el ene Paclet,<sup>1</sup> Bruno Fautrel <sup>3</sup>, Athan Baillet <sup>1</sup>

**To cite:** Romand X, Clapasson M, Chuong MV, *et al*. Serum calprotectin levels do not predict subsequent relapse in rheumatoid arthritis in remission: a post-hoc analysis of STRASS study. *RMD Open* 2023;**9**:e003198. doi:10.1136/rmdopen-2023-003198

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rmdopen-2023-003198>).

Accepted 8 June 2023



  Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>University Grenoble Alpes, CNRS, UMR 5525, T-RAIG, CHU Grenoble Alpes, TIMC, 38000, Grenoble, Auvergne-Rh one-Alpes, France

<sup>2</sup>Department of Rheumatology, Centre Hospitalier Universitaire Grenoble Alpes, Grenoble, France

<sup>3</sup>Assistance Publique H opitaux de Paris, INSERM UMRS 1136, Groupe Hospitalier Universitaire Piti -Salp tri re, Sorbonne Universite, Paris,  le-de-France, France

## Correspondence to

Dr Xavier Romand;  
xromand@chu-grenoble.fr


Dear Editor,

Calprotectin (MRP8/14 or S100A8/A9 protein) is an alarmin intensely released by neutrophils and macrophages in the site of inflammation such as the synovitis. Samantha Louise Smith *et al* showed in a large cohort of patients with active rheumatoid arthritis (RA) that calprotectin is a biomarker of RA disease activity which does not predict, beyond the use of C-reactive protein (CRP), the response to tumour necrosis factor inhibitor (TNFi).<sup>1</sup> Interestingly, a decrease in calprotectin levels at the month 1 and month 3 on TNFi was associated with a moderate-to-good response.<sup>1</sup> The persistence of higher calprotectin level while the patient is in sustained remission may indicate poor control of synovial inflammation, which may lead to subsequent relapse, especially on treatment tapering. Some studies reported that calprotectin in patients with RA in remission could be a useful biomarker to predict change of treatment over 5 years and future relapse.<sup>2 3</sup> The performance of calprotectin to predict clinical response after disease-modifying antirheumatic drugs tapering is still controversial.<sup>4 5</sup>

We conducted a post-hoc analysis of the Spacing of TNF-blocker injections in Rheumatoid Arthritis Study (STRASS) (NCT 00780793).<sup>6</sup> Briefly, patients with RA fulfilling the 1987 American College of Rheumatology criteria, in remission Disease Activity Score 28 (DAS28)<2.6 for more than 6 months with stable joint damage and stable dosage of etanercept or adalimumab for at least 1 year were randomised into two treatment strategies, continue TNFi at a stable dose (maintenance strategy) or injection spacing (spacing strategy). We included all patients with serum samples available and performed

a calprotectin dosage at baseline (IDK Calprotectin ELISA Kit). Relapse during the 18 months was defined as a DAS28>2.6 with DAS28 increase >0.6 compared with the previous visit. The objective was to evaluate if calprotectin serum level during sustained RA remission under TNFi could be associated with the risk of at least one relapse within 18 months of follow-up. T-test, spearman correlation (Rs) and Cox regression analyses were performed. A p value<0.05 was considered statistically significant.

One hundred and twenty-nine patients were included. Sixty-eight were in the maintenance arm and 61 in the spacing arm. One hundred patients (77.5%) were women, mean age was 55.3 ( $\pm$ 11.5) years, 87 were anti-citrullinated protein antibody (78.4%) and 78 (69.6%) rheumatoid factors positive (online supplemental table 1). Levels of baseline calprotectin were modestly correlated with CRP (Rs=0.26, 95% CI 0.06 to 0.40) but not with pain Numeric Rating Scale (NRS), disease activity NRS, erythrocyte sedimentation rate, tender joints count and DAS28-CRP. We did not observe difference in serum calprotectin levels between patients who relapsed during 18 months of follow-up (mean $\pm$ SD, 1564.6 $\pm$ 995.8) and those who did not (1460.2 $\pm$ 1113.7 ng/mL, p=0.59) (figure 1). We also did not find any difference in calprotectin levels in relapsers according to the treatment protocol: maintenance strategy (relapser mean $\pm$ SD 1397.8 $\pm$ 838.3 vs 1423.2 $\pm$ 872.7 ng/mL, p=0.90) or spacing strategy (relapser mean $\pm$ SD 1677.0 $\pm$ 1083.5 vs 1549.1 $\pm$ 1585.7 ng/mL, p=0.77) (figure 1). There was also no difference in calprotectin levels when we study the occurrence of earlier relapse at 3, 6, 9 and 12 months. Calprotectin



**Figure 1** Serum calprotectin levels at baseline in patients with rheumatoid arthritis in sustained remission who relapsed or did not during 18 months of follow-up after continuing the tumour necrosis factor inhibitor at a stable dose (maintenance strategy) or spacing the injections (spacing strategy). Each dot represents one patient. Dark grey dots represent patients with relapse and light grey dots represent patients who do not relapse. Mean $\pm$ SD, t-test.

levels at baseline were not associated with risk of relapse even after adjustment for potential confounders (online supplemental table 2). The main limitation of this post-hoc analysis is the limited number of patients included, which raises a power issue to demonstrate a small mean difference in calprotectin level, which may have contributed to a false-negative result.

Hence, even if we observed that calprotectin was associated with disease activity parameters, we found that calprotectin did not add any information to predict the risk of early or late relapse in patients with RA in sustained remission on TNFi regardless of the treatment strategy used.

**Acknowledgements** We would like to thank Dr Dols for his preliminary assistance in the statistical analysis.

**Contributors** XR, MC and MVC collected the data and performed the calprotectin assay. XR and MC were responsible for data analysis. XR and MC wrote the first draft of the manuscript. MVC, MHP, BF and AB contributed to the conception and study design. BF provided and cared for the study patients. All authors contributed substantially to the interpretation of the data and critically reviewed the work for important intellectual content. All authors approved the final version.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was approved by CPP Paris 6. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Xavier Romand <http://orcid.org/0000-0003-2222-1456>

Bruno Fautrel <http://orcid.org/0000-0001-8845-4274>

Athan Baillet <http://orcid.org/0000-0001-6006-2519>

#### REFERENCES

- Smith SL, Alexander S, Nair N, *et al*. Pre-treatment calprotectin (Mrp8/14) provides no added value to testing CRP alone in terms of predicting response to TNF inhibitors in rheumatoid arthritis in a post hoc analysis. *Ann Rheum Dis* 2023;82:611–20.
- Ramírez J, Cuervo A, Celis R, *et al*. Biomarkers for treatment change and radiographic progression in patients with rheumatoid arthritis in remission: a 5 year follow-up study. *Rheumatology (Oxford)* 2021;60:667–74.
- Inciarte-Mundo J, Ramirez J, Hernández MV, *et al*. Calprotectin strongly and independently predicts relapse in rheumatoid arthritis and polyarticular psoriatic arthritis patients treated with tumor necrosis factor inhibitors: a 1-year prospective cohort study. *Arthritis Res Ther* 2018;20.
- de Moel EC, Rech J, Mahler M, *et al*. Circulating Calprotectin (S100A8/A9) is higher in rheumatoid arthritis patients that relapse within 12 months of tapering anti-rheumatic drugs. *Arthritis Res Ther* 2019;21:268.
- Tweehuysen L, den Broeder N, van Herwaarden N, *et al*. Predictive value of serum Calprotectin (S100A8/A9) for clinical response after starting or tapering anti-TNF treatment in patients with rheumatoid arthritis. *RMD Open* 2018;4.
- Fautrel B, Pham T, Alfaiate T, *et al*. Step-down strategy of spacing TNF-blocker injections for established rheumatoid arthritis in remission: results of the multicentre non-inferiority randomised open-label controlled trial (STRASS: spacing of TNF-blocker injections in rheumatoid arthritis study). *Ann Rheum Dis* 2016;75:59–67.