



HAL
open science

Editorial: The epigenetic control of transposable elements in development and in diseases

Jan Padeken, Michelle S Longworth, Luisa Di Stefano

► To cite this version:

Jan Padeken, Michelle S Longworth, Luisa Di Stefano. Editorial: The epigenetic control of transposable elements in development and in diseases. *Frontiers in Genetics*, 2023, 14, 10.3389/fgene.2023.1282449 . hal-04799620

HAL Id: hal-04799620

<https://cnrs.hal.science/hal-04799620v1>

Submitted on 23 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED AND REVIEWED BY

Michael E. Symonds,
University of Nottingham,
United Kingdom

*CORRESPONDENCE

Luisa Di Stefano,
✉ luisa.di-stefano@univ-tlse3.fr

RECEIVED 24 August 2023

ACCEPTED 25 August 2023

PUBLISHED 04 September 2023

CITATION

Padeken J, Longworth MS and
Di Stefano L (2023), Editorial: The
epigenetic control of transposable
elements in development and in diseases.
Front. Genet. 14:1282449.
doi: 10.3389/fgene.2023.1282449

COPYRIGHT

© 2023 Padeken, Longworth and Di
Stefano. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original author(s)
and the copyright owner(s) are credited
and that the original publication in this
journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Editorial: The epigenetic control of transposable elements in development and in diseases

Jan Padeken¹, Michelle S. Longworth² and Luisa Di Stefano^{3*}

¹Institute of Molecular Biology, Mainz, Germany, ²Cleveland Clinic Lerner Research Institute, Cleveland, OH, United States, ³MCD, Centre de Biologie Intégrative (CBI), Université de Toulouse, CNRS, UPS, Toulouse, France

KEYWORDS

transposable elements, chromatin, development, disease, bioinformatics

Editorial on the Research Topic

The epigenetic control of transposable elements in development and in diseases

A considerable part of many eukaryotic genomes consists of repetitive sequences derived from transposable elements (TEs). TEs are genetic elements that have, or once had, the capability to hop around their host genomes. Through their mobility, they threaten host genomic stability; consistently, uncontrolled TE transposition has been linked to infertility and to diseases such as cancer, hemophilia, and neurodegeneration (McCullers and Steiniger, 2017). An increasing body of evidence shows that TE expression is elevated in cancer, and accumulation of TE transcripts and extrachromosomal DNA derived from TEs has been shown to trigger an innate immune response (Cuellar et al., 2017; Zhao et al., 2018; Heinrich et al., 2019; Simon et al., 2019; Ardeljan et al., 2020).

At the same time, however, TEs can positively impact their host genome. They act as drivers of genomic evolution, influence neighbouring gene expression and have been suggested to shape developmental processes. For example, during Zygotic Genome Activation (ZGA) in early mouse embryonic development, the expression of hundreds of mouse endogenous retroviruses (MERV) is induced. It has been shown that MERVs provide alternative promoter sequences that activate zygotic genes (Macfarlan et al., 2012). More recently LINE1 elements were also implicated in pre-implantation embryonic development by inducing changes in chromatin accessibility and gene transcription (Jachowicz et al., 2017; Percharde et al., 2018). Outstanding questions in the field include whether and when the expression of these TEs is essential for development and the identities of the mechanisms by which TE expression is controlled at various stages of development.

In a study published in this special Research Topic, Ansaloni and colleagues, used multiomic approaches to identify a set of evolutionary young LINE1 whose transcripts are present in the oocyte and which are poised for transcription during ZGA. Through a bioinformatic approach, they attribute their transcription to YY1 activity, paving the way to new research aimed at experimentally validating the role of YY1 in their regulation. Importantly, this work reveals a specific, spatial compartmentalization of MERVs transcription from gene-dense genomic regions and LINE1 from intergenic regions that could coordinately influence the progress of ZGA.

Human pluripotent stem cells (hPSCs) are undifferentiated, self-renewing cells that are present very early after conception, and which can give rise to all differentiated adult cell

types. In this special Research Topic, the review titled “Transposable Elements in Pluripotent Stem Cells and Human Disease,” by [Ma et al.](#), focuses on the various ways in which transposable element (TE) expression and activity affect hPSCs, epigenetic-regulated mechanisms that prevent TE expression in hPSCs, and links between the dysregulation of these mechanisms and human diseases including cancer and neurodegenerative disease. While TEs are largely repressed in most cell types, TE expression and activity is high in hPSCs, and the authors present interesting hypotheses on why this phenomenon occurs. Furthermore, TE sequences influence transcription of both coding and non-coding RNAs in mammalian PSCs, through several different mechanisms, thus impacting PSC homeostasis, pluripotency, and self-renewal. Finally, the authors discuss arguments for the potential contribution of TE sequence fragments in PSCs to tumor development and summarize findings on the epigenetic deregulation of LINE-1 and the endogenous retrovirus, HERV-K in tumor cells and neural cell types present in patients with neurodegenerative disorders.

Another developmental time in which TEs are de-silenced is during germ cell specification when animals epigenetically reprogram their genome to ‘reset’ development. While this is an opportunity for TEs to propagate, it is also a threat to the host progeny’s genome integrity. Taming TE transcription is, therefore, essential for fertility. Eukaryotic genomes have evolved specific molecular machineries to silence TEs transcriptionally and post-transcriptionally. They rely, among others, on the small RNA silencing machinery, DNA transcription factors and chromatin-associated enzymes. One of the chromatin factors involved in TE silencing is the histone demethylase LSD1/KDM1A. Depletion of LSD1 activity induces an increase in the transcription of many TE families in mice ([Macfarlan et al., 2011](#); [Ancelin et al., 2016](#)) humans ([Sheng et al., 2018](#)), and *Drosophila* ([Yang et al., 2019](#); [Lepesant et al., 2020](#)).

In this Research Topic, [Ren and colleagues](#), examine the role of *Caenorhabditis elegans* AMX-1, one of the three worm homologs of the human LSD1 family, in fertility. They report that worms lacking AMX-1 can produce a small number of nonviable eggs. The sterility observed upon AMX-1 depletion is dependent on p53/Cep1 and correlates with an increase in expression of Tc1 and Tc3 transposons but is not dependent on H3K4me2 accumulation. The authors argue that AMX-1 and another LSD1 ortholog, SPR-5, are functionally non-redundant in the germline, and the sterility observed in AMX-1-depleted animals is linked to p53-dependent TE activation. These findings necessitate future studies aimed at validating this hypothesis through genome-wide and genetic approaches.

The repetitive nature and frequent mutations within TE families remain a significant challenge in interpreting genome-wide approaches. Despite a continuous effort in developing new tools that would allow us to reliably map and interpret reads from next-generation sequencing-based approaches, we are still lacking a universal and broadly applicable tool set to interpret transcript levels, chromatin modifications, or protein binding at individual copies of repetitive elements. In this special Research Topic, [Emmanuelle Lerat](#) summarizes these challenges and provides an

overview of the current approaches. She thereby provides a concise list of TE-centric, bioinformatic tools along with brief explanations of their unique features, advantages, and disadvantages.

Many outstanding questions remain to be answered in the field: What are the mechanisms by which TEs are regulated throughout the lifespan of individuals? Which TEs are expressed and when? Do TEs have specific roles in development and in diseases? As research in the field of transposable elements continues to advance, addressing these outstanding questions will provide valuable insights into the roles, regulation, and impacts of TEs on genome evolution, development, and disease.

Author contributions

JP: Writing—original draft, Writing—review and editing. ML: Writing—original draft, Writing—review and editing. LD: Writing—original draft, Writing—review and editing.

Funding

LD research is supported by the French National Research Agency (ANR-20-CE12-0015-01), the CNRS and the INSERM. ML is funded by NIH grant R01GM147138.

Acknowledgments

We would like to thank all the authors for their contributions and the reviewers for their insightful comments. We would also like to thank the Frontiers Editorial Office for giving us the opportunity to host this Research Topic.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

ML declared that she was an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Ancelin, K., Syx, L., Borensztein, M., Ranisavljevic, N., Vassilev, I., Briseño-Roa, L., et al. (2016). Maternal LSD1/KDM1A is an essential regulator of chromatin and transcription landscapes during zygotic genome activation. *Elife* 5, e08851. doi:10.7554/eLife.08851
- Ardeljan, D., Steranka, J. P., Liu, C., Li, Z., Taylor, M. S., Payer, L. M., et al. (2020). Cell fitness screens reveal a conflict between LINE-1 retrotransposition and DNA replication. *Nat. Struct. Mol. Biol.* 27, 168–178. doi:10.1038/s41594-020-0372-1
- Cuellar, T. L., Herzner, A. M., Zhang, X., Goyal, Y., Watanabe, C., Friedman, B. A., et al. (2017). Silencing of retrotransposons by SETDB1 inhibits the interferon response in acute myeloid leukemia. *J. Cell Biol.* 216, 3535–3549. doi:10.1083/jcb.201612160
- Heinrich, M. J., Purcell, C. A., Pruijssers, A. J., Zhao, Y., Spurlock, C. F., Sriram, S., et al. (2019). Endogenous double-stranded Alu RNA elements stimulate IFN-responses in relapsing remitting multiple sclerosis. *J. Autoimmun.* 100, 40–51. doi:10.1016/j.jaut.2019.02.003
- Jachowicz, J. W., Bing, X., Pontabry, J., Bošković, A., Rando, O. J., and Torres-Padilla, M. E. (2017). LINE-1 activation after fertilization regulates global chromatin accessibility in the early mouse embryo. *Nat. Genet.* 49, 1502–1510. doi:10.1038/ng.3945
- Lepesant, J. M. J., Iampietro, C., Galeota, E., Augé, B., Aguirrenbengoa, M., Mercé, C., et al. (2020). A dual role of dLsd1 in oogenesis: regulating developmental genes and repressing transposons. *Nucleic Acids Res.* 48, 1206–1224. doi:10.1093/nar/gkz1142
- Macfarlan, T. S., Gifford, W. D., Agarwal, S., Driscoll, S., Lettieri, K., Wang, J., et al. (2011). Endogenous retroviruses and neighboring genes are coordinately repressed by LSD1/KDM1A. *Genes Dev.* 25, 594–607. doi:10.1101/gad.2008511
- Macfarlan, T. S., Gifford, W. D., Driscoll, S., Lettieri, K., Rowe, H. M., Bonanomi, D., et al. (2012). Embryonic stem cell potency fluctuates with endogenous retrovirus activity. *Nature* 487, 57–63. doi:10.1038/nature11244
- McCullers, T. J., and Steiniger, M. (2017). Transposable elements in Drosophila. *Mob. Genet. Elem.* 7, 1–18. doi:10.1080/2159256X.2017.1318201
- Percharde, M., Lin, C. J., Yin, Y., Guan, J., Peixoto, G. A., Bulut-Karslioglu, A., et al. (2018). A LINE1-nucleolin partnership regulates early development and ESC identity. *Cell* 174, 391–405. doi:10.1016/j.cell.2018.05.043
- Sheng, W., LaFleur, M. W., Nguyen, T. H., Chen, S., Chakravathy, A., Conway, J. R., et al. (2018). LSD1 ablation stimulates anti-tumor immunity and enables checkpoint blockade. *Cell* 174, 549–563. doi:10.1016/j.cell.2018.05.052
- Simon, M., Van Meter, M., Ablueva, J., Ke, Z., Gonzalez, R. S., Taguchi, T., et al. (2019). LINE1 derepression in aged wild-type and SIRT6-deficient mice drives inflammation. *Cell Metab.* 29, 871–885. doi:10.1016/j.cmet.2019.02.014
- Yang, F., Quan, Z., Huang, H., He, M., Liu, X., Cai, T., et al. (2019). Ovaries absent links dLsd1 to HP1a for local H3K4 demethylation required for heterochromatic gene silencing. *Elife* 8, e40806. doi:10.7554/eLife.40806
- Zhao, K., Du, J., Peng, Y., Li, P., Wang, S., Wang, Y., et al. (2018). LINE1 contributes to autoimmunity through both RIG-I- and MDA5-mediated RNA sensing pathways. *J. Autoimmun.* 90, 105–115. doi:10.1016/j.jaut.2018.02.007